Henry Ford Health Henry Ford Health Scholarly Commons

Cardiology Articles

Cardiology/Cardiovascular Research

3-1-2021

Pressure-Volume Analysis Illustrating the Mechanisms of Short-Term Hemodynamic Effects Produced by Premature Ventricular Contractions

Michael I. Brener

Daniel Burkhoff

Mir B. Basir Henry Ford Health, mbasir1@hfhs.org

Mohammad Alqarqaz Henry Ford Health, MALQARQ1@hfhs.org

Follow this and additional works at: https://scholarlycommons.henryford.com/cardiology_articles

Recommended Citation

Brener MI, Burkhoff D, Basir MB, and Alqarqaz M. Pressure-Volume Analysis Illustrating the Mechanisms of Short-Term Hemodynamic Effects Produced by Premature Ventricular Contractions. Circ Heart Fail 2021; 14(3):e007766.

This Article is brought to you for free and open access by the Cardiology/Cardiovascular Research at Henry Ford Health Scholarly Commons. It has been accepted for inclusion in Cardiology Articles by an authorized administrator of Henry Ford Health Scholarly Commons.

UNDER PRESSURE

Pressure-Volume Analysis Illustrating the Mechanisms of Short-Term Hemodynamic Effects Produced by Premature Ventricular Contractions

Michael I. Brener[®], MD; Daniel Burkhoff[®], MD, PhD; Mir Babar Basir, DO; Mohammad Alqarqaz, MD

n 84-year-old man with obstructive coronary artery disease underwent high-risk percutaneous coronary intervention with a high-fidelity conductance catheter seated in the left ventricular apex to assess procedural changes in ventricular contractility via pressure-volume analysis. In the midst of the intervention, the patient developed quadrigeminy (ie, a premature ventricular contraction [PVC] every fourth beat) as shown by the intraventricular electrogram in Figure [A]. Figure [B] shows the corresponding ventricular pressure tracing and illustrates a phenomenon referred to as postextrasystolic (PES) potentiation. Relative to the last sinus beat (labeled beat no. 1), pressure generated on the PVC (beat no. 2) is significantly reduced while it is significantly increased on the PES beat (beat no. 3). The simultaneously recorded pressure-volume loops (Figure [C]) show reduced ventricular filling on the PVC due to the short time interval between the PVC and the prior beat, and the increased ventricular filling on the PES beat due to the longer than normal interval between the PVC and PES beats (referred to as a compensatory pause). More striking than changes in ventricular filling, however, are the marked beat-to-beat changes in contractility evidenced by changes of the slope (referred to as end-systolic elastance) of the respective end-systolic pressure-volume relationships.

Differences in ventricular pressure generation accompanying changes in the beat-to-beat interval are generally attributed to changes in preload caused by variable filling times. Although it is evident that changes in preload contribute to this effect, ventricular contractility is by far the more substantial factor in determining differences in pressure generation. Interval-dependent changes of contractility are the global manifestation of the myocardial force-interval relationship, which relates the force of myocyte contraction to the beat-to-beat time interval.^{1,2} The force-interval relationship, in turn, results from the time interval-dependent availability of calcium for release from the sarcoplasmic reticulum to the myofilaments in a process referred to as calcium restitution.³ Calcium restitution starts at the end of phase 3 of the action potential and follows an exponential time course (with time constant \approx 150-200 ms) to a plateau by \approx 1 second. On a PVC, only a small amount of the total sarcoplasmic reticulum calcium pool is available for release, resulting in reduced contractility. The enhanced contractility on the PES beat results from greater sarcoplasmic reticulum calcium release, reflecting the sum of calcium normally released during a given beat plus calcium leftover from the prior premature beat in combination with the long time for calcium restitution due to the compensatory pause. Here, we describe the hemodynamic effects of a single PVC and PES beat, but these mechanisms also explain the pulse pressure variations observed in other irregular arrhythmias like atrial fibrillation. Although these concepts are well known to students of muscle physiology and cardiac excitation-contraction coupling,⁴ they are not typically part of modern medical school curricula.

Key Words: calcium = coronary artery disease = hemodynamics = percutaneous coronary intervention = sarcoplasmic reticulum

Correspondence to: Michael I. Brener, MD, Columbia University Medical Center, Cardiology, 622 West 168th St, Presbyterian Hospital, Floor 3, Room 347, New York, NY 10032. Email mib2102@cumc.columbia.edu

© 2021 American Heart Association, Inc.

Circulation: Heart Failure is available at www.ahajournals.org/journal/circheartfailure

ARTICLE INFORMATION

Affiliations


Division of Cardiology, Columbia University Medical Center, New York, NY (M.I.B.). Cardiovascular Research Foundation, New York, NY (D.B.). Division of Cardiology, Henry Ford Hospital System, Detroit, MI (M.B.B., M.A.).


Disclosures

None.

REFERENCES

- Burkhoff D, Yue DT, Franz MR, Hunter WC, Sagawa K. Mechanical restitution of isolated perfused canine left ventricles. *Am J Physiol.* 1984;246(1 pt 2):H8–16. doi: 10.1152/ajpheart.1984.246.1.H8
- Yue DT, Burkhoff D, Franz MR, Hunter WC, Sagawa K. Postextrasystolic potentiation of the isolated canine left ventricle. Relationship to mechanical restitution. *Circ Res.* 1985;56:340–350. doi: 10.1161/01.res.56.3.340
- Wier WG, Yue DT. Intracellular calcium transients underlying the shortterm force-interval relationship in ferret ventricular myocardium. *J Physiol.* 1986;376:507–530. doi: 10.1113/jphysiol.1986.sp016167
- 4. Bers DM. Cardiac excitation-contraction coupling. *Nature*. 2002;415:198–205. doi: 10.1038/415198a





A, ECG. **B**, Left ventricular pressure (LVP). **C**, Pressure-volume loops corresponding to beats 1, 2, 3, 4 and 5 as labeled in **B**.