

Baptist Health South Florida

## Scholarly Commons @ Baptist Health South Florida

---

All Publications

---

2020

### HGG-18. CLINICAL EFFICACY OF ONC201 IN THALAMIC H3 K27M-MUTANT GLIOMA

Minesh Mehta

*Baptist Health Medical Group; Miami Cancer Institute, mineshm@baptisthealth.net*

Yazmin Odia

*Baptist Health Medical Group; Miami Cancer Institute; Miami Neuroscience Institute, yazmino@baptisthealth.net*

Matthew Hall

*Baptist Health Medical Group; Miami Cancer Institute, matthewha@baptisthealth.net*

Doured Daghistani

*Baptist Health Medical Group; Miami Cancer Institute, douredd@baptisthealth.net*

Follow this and additional works at: <https://scholarlycommons.baptisthealth.net/se-all-publications>

---

#### Citation

Neuro-Oncology (2020) 22 (3 Suppl):iii347

This Article -- Open Access is brought to you for free and open access by Scholarly Commons @ Baptist Health South Florida. It has been accepted for inclusion in All Publications by an authorized administrator of Scholarly Commons @ Baptist Health South Florida. For more information, please contact [Carrief@baptisthealth.net](mailto:Carrief@baptisthealth.net).

Hala Taha<sup>1,2</sup>, Shahenda El-naggar<sup>2</sup>, and Mohamed El-beltagy<sup>5,2</sup>; <sup>1</sup>National Cancer Institute, Cairo University, Cairo, Egypt, <sup>2</sup>Children Cancer Hospital- Egypt (CCHE-57357), Cairo, Egypt, <sup>3</sup>Faculty of Medicine, Aswan University, Aswan, Egypt, <sup>4</sup>University Hospital of Geneva, Geneva, Switzerland, <sup>5</sup>Kasr El-Ainy School of medicine, Cairo University, Cairo, Egypt

**BACKGROUND:** Data about high-grade glioma (HGG) in very young children ( $\leq 3$  years old at diagnosis) is scarce. **METHODS:** 180 pediatric HGG patients were treated at the Children Cancer Hospital - Egypt (CCHE-57357) between July 2007 and June 2018, with 17 patients aged  $\leq 3$  years at diagnosis. Medical records were retrospectively reviewed for clinical, radiological and histopathological data, treatment received and survival outcome. **RESULTS:** Median age was 29.2 months (range: 2.4 – 35.8 months; males = 9). Most frequent pathological diagnosis was Glioblastoma, WHO grade-IV (n = 11, 64.7%) and one patient had H3-mutant diffuse midline glioma. All patients underwent surgery (gross-total resection, n = 6, 35.3%; subtotal-resection, n = 5, 29.4%; biopsy, n = 6, 35.3%). One patient (age = 7 months) progressed and died before starting adjuvant therapy. All patients  $\leq 1$  year of age (n = 5) received adjuvant chemotherapy (CT) only, older children (n = 11) received adjuvant radiotherapy (RT) (total dose range: 54 – 60 Gy) and CT (CCG-945 protocol). The 1-year overall survival (OS) rate was 47.1%; and event-free survival (EFS) rate was 35.3%. EFS differed between those who received RT and those who did not (1-year EFS 54.5% and 0% respectively, p = 0.001). Compared to older children, anatomical distribution of tumors was significantly different with non-midline locations being the commonest in patients  $\leq 3$  years old (88.2% vs 46.4%, p=0.01). **CONCLUSIONS:** HGG in very young children arise predominantly in non-midline locations and usually lack the H3-mutation. RT seems crucial in the management of pHGG regardless of age subgroup.

#### HGG-18. CLINICAL EFFICACY OF ONC201 IN THALAMIC H3 K27M-MUTANT GLIOMA

Abed Rahman Kawakibi<sup>1</sup>, Rohinton S. Tarapore<sup>2</sup>, Sharon Gardner<sup>3</sup>, Andrew Chi<sup>3</sup>, Sylvia Kurz<sup>3</sup>, Patrick Y. Wen<sup>4</sup>, Isabel Arrillaga-Romany<sup>5</sup>, Tracy T. Batchelor<sup>6</sup>, Nicholas A. Butowski<sup>7</sup>, Ashley Sumrall<sup>8</sup>, Nicole Shonka<sup>9</sup>, Rebecca Harrison<sup>10</sup>, John DeGroot<sup>10</sup>, Minesh Mehta<sup>11</sup>, Yazmin Odia<sup>11</sup>, Matthew D. Hall<sup>11</sup>, Doured Daghistani<sup>11</sup>, Timothy F. Cloughesy<sup>12</sup>, Benjamin M. Ellingson<sup>12</sup>, Yoshie Umemura<sup>1</sup>, Jonathan Schwartz<sup>13</sup>, Vivekanand Yadav<sup>1</sup>, Rodrigo Cartaxo<sup>1</sup>, Ruby Siada<sup>1</sup>, Zachary Miklja<sup>1</sup>, Amy Bruzek<sup>1</sup>, Evan Cantor<sup>1</sup>, Kyle Wierzbicki<sup>1</sup>, Alyssa Paul<sup>1</sup>, Ian Wolfe<sup>1</sup>, Marcia Leaoard<sup>1</sup>, Hugh Garton<sup>1</sup>, Rajen Mody<sup>1</sup>, Patricia L. Robertson<sup>1</sup>, Guangrong Lu<sup>2</sup>, Krystal Merdinger<sup>2</sup>, Sriram Venneti<sup>1</sup>, Wolfgang Oster<sup>2</sup>, Joshua E. Allen<sup>2</sup>, and Carl Koschmann<sup>1</sup>; <sup>1</sup>University of Michigan, Ann Arbor, MI, USA, <sup>2</sup>Oncocotics Inc., Philadelphia, PA, USA, <sup>3</sup>NYU Langone Health, New York, NY, USA, <sup>4</sup>Dana-Farber Cancer Institute, Boston, MA, USA, <sup>5</sup>Massachusetts General Hospital, Boston, MA, USA, <sup>6</sup>Brigham and Women's Hospital, Boston, MA, USA, <sup>7</sup>University of California San Francisco, San Francisco, CA, USA, <sup>8</sup>Levine Cancer Institute, Charlotte, NC, USA, <sup>9</sup>University of Nebraska Medical Center, Omaha, NE, USA, <sup>10</sup>University of Texas MD Anderson Cancer Center, Houston, TX, USA, <sup>11</sup>Miami Cancer Institute, Miami, FL, USA, <sup>12</sup>University of California Los Angeles, Los Angeles, CA, USA, <sup>13</sup>Mayo Clinic, Rochester, MN, USA

ONC201, a bitopic DRD2 antagonist and allosteric ClpP agonist, has shown encouraging efficacy in H3 K27M-mutant glioma. Given that the thalamus has the highest extra-striatal expression of DRD2, we performed an integrated preclinical and clinical analysis of ONC201 in thalamic H3 K27M-mutant glioma. ONC201 was effective in mouse intra-uterine electroporation (IUE)-generated H3 K27M-mutant gliomas, with an *in vitro* IC<sub>50</sub> of 500 nM and 50% prolongation of median survival *in vivo* (p=0.02, n=14). We analyzed thalamic H3 K27M-mutant glioma patients treated with ONC201 on active clinical trials as of 5/22/19 enrollment (n=19 recurrent and 10 post-radiation, non-recurrent; 5–70 years old). As of 12/18/2019, PFS6 and OS12 are 26.3% and 36.8%, respectively, in the recurrent group. For non-recurrent patients, with median follow up of 21.9 months (8.6–26.6) from diagnosis, median PFS or OS have not been reached. This surpasses historical OS of 13.5 months. Best response by RANO includes 1 CR, 3 PR, 4 SD, 8 PD for recurrent patients and 2 PR, 4 SD, 1 PD for non-recurrent patients (4 on-trial patients experienced regressions that are yet unconfirmed responses). Median duration of response for recurrent patients is 14.0 months (2.0–33.1). Furthermore, H3 K27M cell-free tumor DNA in plasma and CSF correlated with MRI response. In summary, single agent ONC201 administered at recurrence, or adjuvantly following radiation, demonstrates promising clinical efficacy in thalamic H3 K27M-mutant glioma patients who currently have no effective treatments following radiation. Investigations are ongoing to assess whether micro-environmental DRD2 expression explains the early exceptional responses in thalamic H3 K27M-mutant glioma.

#### HGG-19. IDENTIFICATION OF NOVEL SUBGROUP-SPECIFIC MIRNA EXOSOMAL BIOMARKERS IN PEDIATRIC HIGH-GRADE GLIOMAS

Lucia Lisa Petrilli<sup>1</sup>, Alessandro Paolini<sup>2</sup>, Angela Galardi<sup>1</sup>, Giulia Pericoli<sup>1</sup>, Marta Colletti<sup>1</sup>, Roberta Ferretti<sup>1</sup>, Virginia Di Paolo<sup>1</sup>, Luisa Pascucci<sup>3</sup>, Hector Peinado<sup>4</sup>, Chris Jones<sup>5</sup>, Antonella Cacchione<sup>1</sup>, Luca De Palma<sup>6</sup>, Marta Alonso<sup>7</sup>, Andrew Moore<sup>8</sup>, Angel Montero Carcaboso<sup>9</sup>, Andrea Carai<sup>10</sup>, Angela Mastronuzzi<sup>11</sup>, Franco Locatelli<sup>1</sup>, Andrea Masotti<sup>2</sup>, Angela Di Giannatale<sup>1</sup>, and Maria Vinci<sup>1</sup>; <sup>1</sup>Department of Onco-hematology, Cell and Gene Therapy, Bambino Gesù Children's Hospital-IRCCS, Rome, Italy, <sup>2</sup>Multifactorial and Complex Phenotypes Research Area, Bambino Gesù Children's Hospital-IRCCS, Rome, Italy, <sup>3</sup>Department of Veterinary Medicine, University of Perugia, Perugia, Italy, <sup>4</sup>Microenvironment & Metastasis Group, Molecular Oncology Program, Spanish National Cancer Research Centre (CNIO), Madrid, Spain, <sup>5</sup>Department of Molecular Pathology, The Institute of Cancer Research, Sutton, United Kingdom, <sup>6</sup>Rare and Complex Epilepsy Unit, Department of Neuroscience, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy, <sup>7</sup>Department of Pediatrics, University Hospital of Navarra, Pamplona, Spain, <sup>8</sup>The University of Queensland Diamantina Institute, Translational Research Institute, Brisbane, Australia, <sup>9</sup>Developmental Tumor Biology Laboratory, Hospital Sant Joan de Déu, Barcelona, Spain, <sup>10</sup>Department of Neuroscience and Neuro-rehabilitation, Neurosurgery Unit, Bambino Gesù Children's Hospital-IRCCS, Rome, Italy, <sup>11</sup>Neuro-oncology Unit, Department of Onco-hematology, Cell and Gene Therapy, Bambino Gesù Children's Hospital-IRCCS, Rome, Italy

Pediatric high-grade gliomas (pHGG) are heterogeneous brain tumors for which new specific diagnostic/prognostic biomarkers are needed. In this study, we aimed to identify new pHGG subgroup specific biomarkers by exploiting exosomes, known vehicles of oncogenic signals. We used plasma from 23 patients (including 6 controls) and conditioned medium from 12 patient-derived cell-lines, representing all locational and molecular subgroups. Upon exosome isolation, total RNA was extracted and miRNAs were assessed using a PCR Panel. Analysis of plasma miRNome showed that tumor exosomal samples were largely clustered together, independently from their locational and/or molecular subgroup. We identified 20 significantly upregulated and 25 downregulated miRNAs compared to controls. Interestingly, 27 miRNAs were expressed only in tumors. Furthermore, the unsupervised clustering showed a clear separation based on locational (hemispheric *vs* pontine) and mutational (WT *vs* H3.3G34R or H3.3G34R *vs* H3K27M) subgroup comparisons, with the identification of distinct miRNomes underlying the key role of location and mutations in defining the pHGG exosomal miRNA profile. This was further confirmed analyzing the miRNome from cell-line derived exosomes. Moreover, we identified a pool of significantly differentially regulated miRNAs in diagnose *vs* relapse and biopsy *vs* autopsy cell-lines. Most importantly, when comparing hemispheric *vs* pontine and H3.3G34R *vs* H3.3K27M, we identified respectively four and three miRNAs equally dysregulated and in common between plasma and cell-lines. Those were strongly associated mainly to transcriptional regulation and targeting *TTC9*, linked to cancer invasion and metastasis. Based on this, we suggest exosomal miRNAs as a powerful new pHGG diagnostic/prognostic tool.

#### HGG-20. DIAGNOSTIC AND BIOLOGICAL ROLE OF METHYLATION PATTERNS IN REPLICATION REPAIR DEFICIENT HIGH GRADE GLIOMAS

Andrew Dodgshun<sup>1,2</sup>, Kohei Fukuoka<sup>3</sup>, Melissa Edwards<sup>4</sup>, Vanessa Bianchi<sup>4</sup>, Alexandra Sexton-Oates<sup>5</sup>, Valerie Larouche<sup>6</sup>, Vanan Magimairajan<sup>7</sup>, Scott Lindhorst<sup>8</sup>, Michal Yalon<sup>9</sup>, Gary Mason<sup>10</sup>, Bruce Crooks<sup>11</sup>, Shlomi Constantini<sup>12</sup>, Maura Massimino<sup>13</sup>, Stefano Chiaravalli<sup>13</sup>, Jagadeesh Ramdas<sup>14</sup>, Warren Mason<sup>15</sup>, Ashraf Shamvil<sup>16</sup>, Roula Farah<sup>17</sup>, An Van Damme<sup>18</sup>, Enrico Opocher<sup>19</sup>, Syed Ahmad Hamid<sup>20</sup>, David Ziegler<sup>21</sup>, David Samuel<sup>22</sup>, Kristina A Cole<sup>23</sup>, Patrick Tomboc<sup>24</sup>, Duncan Stearns<sup>25</sup>, Gregory Thomas<sup>26</sup>, Alexander Lossos<sup>27</sup>, Michael Sullivan<sup>28</sup>, Jordan R Hansford<sup>28</sup>, David Jones<sup>29</sup>, Alan Mackay<sup>30</sup>, Chris Jones<sup>30</sup>, Vijay Ramaswamy<sup>4</sup>, Cynthia Hawkins<sup>4</sup>, Eric Bouffet<sup>4</sup>, and Uri Tabori<sup>4</sup>; <sup>1</sup>Canterbury District Health Board, Christchurch, New Zealand, <sup>2</sup>University of Otago, Christchurch, New Zealand, <sup>3</sup>Jichi Medical University, Shimotsuke, Japan, <sup>4</sup>The Hospital for Sick Children, Toronto, Canada, <sup>5</sup>International Agency for Research on Cancer, Lyon, France, <sup>6</sup>Université Laval, Quebec, Canada, <sup>7</sup>University of Manitoba, Winnipeg, Canada, <sup>8</sup>Medical University of South Carolina, Charleston, SC, USA, <sup>9</sup>Sheba Medical Center, Tel Hashomer, Israel, <sup>10</sup>University of Pittsburgh, Pittsburgh, PA, USA, <sup>11</sup>Dalhousie University, Halifax, Canada, <sup>12</sup>Tel Aviv Sourasky Medical Center, Tel Aviv, Israel, <sup>13</sup>Istituto Nazionale dei Tumori, Milan, Italy, <sup>14</sup>Geisinger Medical Center, Danville, PA, USA, <sup>15</sup>Princess Margaret Cancer Centre, Toronto, Canada, <sup>16</sup>Children's Cancer Hospital, Karachi, Pakistan, <sup>17</sup>Saint George Hospital University Medical Center, Beirut, Lebanon, <sup>18</sup>Université Catholique de Louvain, Brussels, Belgium, <sup>19</sup>University of Padova, Padova,



---

## Neuro-oncology

**PUBLICATION TYPE:** e-Journal

**ISSN:** 1523-5866

**Date:** 1999 - Present

**Publisher:** DUKE UNIVERSITY PRESS

**Language:** English

**Country:** United States of America

**URL:** <http://neuro-oncology.mc.duke.edu>

**Authors:** Society for Neuro-Oncology.

**Licensee:**

**Organization:** Baptist Health South Florida

**Date/Time:** 11 Feb 2021 02:53

---

### Digital Sharing

This permission type is covered.

#### Special terms :

Please note. Any permission obtained through this service only relates to material owned or controlled by Oxford University Press. If the material you wish to use is acknowledged to any other source, you may also need to clear additional permission with the rights holder.

---

Your organization has obtained rights for you to copy and share this title in electronic form.

#### Examples include

- Emailing a copy to my co-workers.
- Storing a copy on an internal shared network.
- Storing a copy on your local hard drive.
  
- Displaying in a presentation to co-workers.
- Distributing in a PowerPoint presentation to co-workers.
- Submitting an electronic copy to regulatory authorities.

The license cannot be used to create a library or collection intended to substantially replace the need to take a subscription or purchase a particular Work.