

University of Mississippi

eGrove

Faculty and Student Publications

Pharmacy, School of

1-1-2019

Phytochemical screening of Pulsatilla species and investigation of their biological activities

Grażyna Łaska

Bialystok University of Technology

Aneta Sienkiewicz

Bialystok University of Technology

Marcin Stocki

Bialystok University of Technology

Jordan K. Zjawiony

University of Mississippi

Vimal Sharma

University of Mississippi

See next page for additional authors

Follow this and additional works at: https://egrove.olemiss.edu/pharmacy_facpubs

Recommended Citation

Łaska G, Sienkiewicz A, Stocki M, Zjawiony JK, Sharma V, Bajguz A, et al. Phytochemical screening of Pulsatilla species and investigation of their biological activities. *Acta Soc Bot Pol.* 2019;88(1):3613. <https://doi.org/10.5586/asbp.3613>

This Article is brought to you for free and open access by the Pharmacy, School of at eGrove. It has been accepted for inclusion in Faculty and Student Publications by an authorized administrator of eGrove. For more information, please contact egrove@olemiss.edu.

Authors

Grażyna Łaska, Aneta Sienkiewicz, Marcin Stocki, Jordan K. Zjawiony, Vimal Sharma, Andrzej Bajguz, Alicja Piotrowska-Nicznyporuk, Melissa Jacob, and Shabana Khan

DOI: 10.5586/asbp.3613

Publication history

Received: 2017-10-10

Accepted: 2018-11-22

Published: 2019-04-01

Handling editor

Wojciech Szypuła, Faculty of Pharmacy, Medical University of Warsaw, Poland

Authors' contributions

GŁ: designed the study and codirected laboratory work; AS: participated in laboratory work; MS: GC-MS analysis; JKZ, VS: conducted chromatographic fractionation; AB, APN: HPLC analysis; MJ, SK: determination of biological activity; GŁ, AS: drafted the manuscript

Funding

This study was supported by a grant No. S/WBiłŚ/5/16 from the Ministry of Science and Higher Education of Poland and a grant No. 1459504 from the National Science Foundation of USA.

Competing interests

No competing interests have been declared.

Copyright notice

© The Author(s) 2019. This is an Open Access article distributed under the terms of the [Creative Commons Attribution License](#), which permits redistribution, commercial and noncommercial, provided that the article is properly cited.

Citation

Łaska G, Sienkiewicz A, Stocki M, Zjawiony JK, Sharma V, Bajguz A, et al. Phytochemical screening of *Pulsatilla* species and investigation of their biological activities. Acta Soc Bot Pol. 2019;88(1):3613. <https://doi.org/10.5586/asbp.3613>

Digital signature

This PDF has been certified using digital signature with a trusted timestamp to assure its origin and integrity. A verification trust dialog appears on the PDF document when it is opened in a compatible PDF reader. Certificate properties provide further details such as certification time and a signing reason in case any alterations made to the final content. If the certificate is missing or invalid it is recommended to verify the article on the journal website.

ORIGINAL RESEARCH PAPER

Phytochemical screening of *Pulsatilla* species and investigation of their biological activities

Grażyna Łaska^{1*}, Aneta Sienkiewicz¹, Marcin Stocki², Jordan K. Zjawiony³, Vimal Sharma³, Andrzej Bajguz⁴, Alicja Piotrowska-Niczypruk⁴, Melissa Jacob⁵, Shabana Khan⁵

¹ Department of Agri-Food Engineering and Environmental Management, Białystok University of Technology, Wiejska 45A, 15-351 Białystok, Poland

² Department of Forestry in Hajnówka, Białystok University of Technology, Wiejska 45A, 15-351 Białystok, Poland

³ Department of BioMolecular Sciences, School of Pharmacy, University of Mississippi, MS 38677, United States

⁴ Department of Plant Biochemistry and Toxicology, University of Białystok, Ciołkowskiego 1J, 15-245 Białystok, Poland

⁵ National Center for Natural Products Research, University of Mississippi, University, MS 38677, United States

* Corresponding author. Email: g.laska@pb.edu.pl

Abstract

The present study aimed to identify biologically active secondary metabolites from the rare plant species, *Pulsatilla patens* subsp. *patens* and the cultivated *P. vulgaris* subsp. *vulgaris*. Chromatographic fractionation of the ethanolic extract of the roots of *P. patens* subsp. *patens* resulted in the isolation of two oleanane-type glycosides identified as hederagenin 3-O-β-D-glucopyranoside (2.7 mg) and hederagenin 3-O-β-D-galactopyranosyl-(1→2)-β-D-glucopyranoside (3.3 mg, patensin). HPLC analysis of the methanolic extract of the crude root of *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* revealed the presence of *Pulsatilla* saponin D (hederagenin 3-O-α-L-rhamnopyranosyl(1→2)-[β-D-glucopyranosyl(1→4)]-α-L-arabinopyranoside). Chromatographic analysis using GC-MS of the silylated methanolic extracts from the leaves and roots of these species identified the presence of carboxylic acids, such as benzoic, caffeic, malic, and succinic acids. The extracts from *Pulsatilla* species were tested for their antifungal, antimicrobial, and antimalarial activities, and cytotoxicity to mammalian cell lines. Both *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* were active against the fungus *Candida glabrata* with the half-maximal inhibitory concentration (IC₅₀) values of 9.37 μg/mL and 11 μg/mL, respectively. The IC₅₀ values for cytotoxicity evaluation were in the range of 32–38 μg/mL for *P. patens* subsp. *patens* and 35–57 μg/mL for *P. vulgaris* subsp. *vulgaris* for each cell line, indicating general cytotoxic activity throughout the panel of evaluated cancer and noncancer cells.

Keywords

pasque flower subspecies; oleanane-type triterpenoid glycosides; carboxylic acids; antifungal and antimicrobial activities; cytotoxicity; medicinal applications

Introduction

Plants are known to produce different biologically active substances, including secondary metabolites. It had been reported that plants from the *Pulsatilla* genus contain ranunculin, anemonin, protoanemonin, triterpenes, and saponins (9%) of mainly the oleanane and lupane-type [1,2]. These metabolites were also subjected to analyses in order to determine the contents of polyphenol compounds, flavonoids, and anthocyanidins

[3,4]. The triterpene saponins were isolated from *P. chinensis* (Bunge) Regel [5], *P. koreana* Nakai [6], *P. cernua* (Thunb.) Bercht. et Opiz. [7], *P. dahurica* (Fisch. ex DC.) Spreng. [8], *P. turczaninowii* Kryl. et Serg. [9], *P. nigricans* Storck [10], *P. pratensis* (L.) Mill. [11], and *P. patens* subsp. *multifida* (G. A. Pritzel) Zämelis [12]. Polyphenolic compounds such as flavonoids and anthocyanidins are produced by *P. montana* subsp. *balcana* (Velen.) Zämelis & Paegle, *P. halleri* subsp. *rhodopaea* (Stoj. et Stef.) K. Krause, and *P. slaviankae* (Zimmer.) Jordanov & Kožuharov [13]. The saponins produced by *Pulsatilla* species show high biological activity across a wide range as an anticancer [14], neuroactive [15], neuroprotective [16], immunomodulating [17], cognitive function enhancing [18], antioxidant [19], antimicrobial [20], and cytotoxic agents [21].

Pulsatilla species are rarely of toxicological importance, as while they are moderately hazardous, they are mutagenic and contain cellular poison [22]. The active ingredient in fresh plant material, a terpenoid glucoside known as ranunculin, is broken down to protoanemonin [23]. Protoanemonin is a reactive compound with an exocyclic methylene group and is mutagenic, because it can bind to the SH-group of proteins and

DNA. Protoanemonin is an irritant lactone that causes allergic dermatitis in human skin [24], and internally, paralysis of the central nervous system and gastrointestinal disturbance [25]. When the plant is dried, the protoanemonin dimerises to form the less toxic anemonin, if it has not already alkylated proteins [23].

Despite the potential toxicity of *Pulsatilla* species, their natural products have been used for centuries in traditional Chinese and Korean medicine for the treatment of many diseases and ailments. They are known as herbal drugs used for the homeopathic treatment of eye ailments, earache, stress, anxiety, tension, skin eruptions, rheumatism, headaches, neuralgia, insomnia, hyperactivity, bacterial skin infections, septicemia, bronchitis, coughs, and asthma [26].

The *Pulsatilla* genus is composed of 38 species in the Northern Hemisphere [23]. The species of *Pulsatilla patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* have not been extensively studied. The existing research focuses primarily on genetic characteristics of the species [27], ecology [28], and various aspects of developmental biology [29]. *Pulsatilla patens* subsp. *patens* is a lowland species of Boreo-meridional-continental distribution [30] and is native to Europe, Russia, Mongolia, China, Canada, and the United States [31] (Fig. 1A).

Pulsatilla patens subsp. *patens* is an endangered plant and is considered critically threatened in the Czech Republic [32], included in the red data books of Germany as an endangered plant [33], vulnerable in Sweden [34], relatively restored in Lithuania, and in decreasing number in the St. Petersburg and Kaliningrad [35] regions of Russia, Latvia [36], and Slovakia [37]. In Finland and Estonia, the populations of *P. patens* subsp. *patens* are considered to be relics [38]. In Poland, *P. patens* subsp. *patens* has been strictly protected since 1958 and requires




Fig. 1 Distribution of *Pulsatilla patens* subsp. *patens* (A) in Europe, Asia, and North America [31], and (B) in Poland [40].




Fig. 2 Distribution of *Pulsatilla vulgaris* subsp. *vulgaris* (A) in Central Europe [31] and (B) in Poland [40].

active protection [39]. On the territory of Poland, *P. patens* subsp. *patens* reaches the western border of its range and the number of its sites clearly decreases towards the west and south [40] (Fig. 1B).

Pulsatilla vulgaris subsp. *vulgaris* is a species occurring across Central Europe, ranging from France in the south to Sweden at its northern limit [31] (Fig. 2A) and is listed as “near threatened” by the International Union for Conservation of Nature [41]. Currently, *P. vulgaris* subsp. *vulgaris* is listed as vulnerable (VU category) in Ukraine, Slovakia [42], Sweden [43], and the United Kingdom [44]. This species is also classified as endangered (EN category) in Germany [45] and Switzerland [46] and as critically endangered (CR category) in Austria [47]. In Poland, it is listed as an extinct (RE category) species [48] and mainly cultivated. *Pulsatilla vulgaris* subsp. *vulgaris* is currently extinct in Poland on natural sites [49]. It was only reported at four sites before 1930 [40] (Fig. 2B) on the habitats of the pine forests of the All. Diczano-Pinion.

The aim of this study was the identification of biologically active secondary metabolites from the rare plant species *Pulsatilla patens* subsp. *patens* and the cultivated *P. vulgaris* subsp. *vulgaris*. In the course of our study, we evaluated the extracts of leaves and roots of these plant species for their antifungal, antimicrobial, and antimalarial activities, and cytotoxicity to mammalian cell lines.

Material and methods

Material collection

The leaves and roots of *P. patens* subsp. *patens* were collected in May 2013–2015 at the Knyszynska Forest near Białystok City, NE

Poland, while *P. vulgaris* subsp. *vulgaris* were obtained from cultivation at the Herbarium “The Herbal Corner” in May 2015, Podlaskie Province, NE Poland. Plant material was identified by Prof. Grażyna Łaska from the Białystok University of Technology, Faculty of Civil and Environmental Engineering, Poland.

Extraction and chromatographical analysis

TLC/CC analysis. For thin layer chromatography (TLC) and column chromatography (CC), ethanolic extracts were prepared from roots of *P. patens* subsp. *patens* (K₂). Dried roots of *P. patens* subsp. *patens* (10 g) were extracted with 1 L of 50% aqueous EtOH at room temperature for 12 h and then filtered. The extraction was repeated three times. The combined filtrates were concentrated in a vacuum evaporator to obtain a syrupy brown residue (0.95 g). Then 5 mg of ethanolic extract from roots of *P. patens* subsp. *patens* (K₂) was transferred to normal and reverse phase (RP) TLC on silica gel with

fluorescence (stationary phase). The following solvent system was used for normal phase TLC: hexanes–ethyl acetate (C_6H_{14} – $C_4H_8O_2$; 60:40), chloroform–methanol ($CHCl_3$ –MeOH; 90:10) (Fig. 3) and for reverse phase TLC: water–methanol (H_2O –MeOH; 30:70), water–methanol (H_2O –MeOH; 70:30). The solvent system H_2O –MeOH (30:70) was found to be best for RP-TLC. A solution of H_2SO_4 –MeOH (5:95) and UV 254 and 366 nm were used as a detecting reagent (Fig. 3). In the next step, an ethanolic extract of the roots of *P. patens* subsp. *patens* was fractionated by CC (0.9 g) using $CHCl_3$ –MeOH gradient (90:10). Compounds were eluted by reverse-phase C_{18} column using the H_2O –MeOH (30:70) solvent system (Fig. 4). The structures of the isolated compounds were determined by 1H and ^{13}C -NMR spectroscopy.




Fig. 3 TLC image of the ethanolic extract of roots (K_2) of *Pulsatilla patens* subsp. *patens* and identification of Compound 1 and 2.




Fig. 4 Column chromatography of the ethanolic extract of *Pulsatilla patens* subsp. *patens* roots and isolation of Compound 1 and 2.

HPLC analysis. The fraction which exhibited the most potent activity was again chromatographed by high performance liquid chromatography (HPLC) using Agilent Technologies 1260 Infinity system. The HPLC profiles were determined at $35^\circ C$ on a COSMOSIL Packed Column 5C-18-MS-II ($4.6 \mu m \times 250 mm$) (Nacalai, USA) eluted with methanol (A) and water (B) as mobile phases in a linear gradient elution with the flow rate of 1 mL/min. The elution program was as follows: 0–8 min, 90% A to 77%

A, 8–25 min, changed to 71% A, 25–40 min, to 60% A, 40–50 min, to 50% A, 50–75 min, to 10% A, 75–80 min, to 0% A [50]. The volume of sample injected was 35 μ L. Fluorescence was monitored by an Agilent 1260 Infinity Fluorescence Light Detector (Agilent Technologies, USA; 210 nm excitation and 460 nm emission wavelength). The analytical data were integrated using the Agilent OpenLAB software. The fraction was purified by HPLC to give *Pulsatilla* saponin D. The peak of *Pulsatilla* saponin D was identified based on retention time with respect to the reference standard (purchased from Toronto Research Chemicals) previously run in an analytical HPLC.

GC-MS analysis. The pressurized solvent extractions were performed using a Büchi E-916 Speed Extractor. The raw materials were extracted with methanol (leaves 32.3 g, and roots, 23.6 g of *P. patens* subsp. *patens*; leaves, 21.8 g and roots, 56.1 g of *P. vulgaris* subsp. *vulgaris*). The extraction time was 30 min, temperature 100°C and pressure 1.2×10^7 Pa. Three extracts were prepared from each raw material. Extracts were filtered through paper filters. The methanol was dried using a rotor evaporator.

Each dry residue (10 mg) was redissolved with 1 mL of pyridine and 200 μ L BSTFA [N,O-bis-(trimethylsilyl)-trifluoroacetamide] added. Next, the mixture was heated at 60°C for 30 min. The solution was analyzed by gas chromatography / mass spectrometry (GC-MS) using an Agilent 7890A gas chromatograph equipped with an Agilent 5975C mass selective detector. The injection of a 1- μ L sample was performed using an Agilent 7693A autosampler. The separation was performed on an HP-5MS (30 m \times 0.25 mm \times 0.25 μ m film thickness) fused silica column at a helium flow rate of 1 mL/min. The injector worked in split mode (1:5), and the injector temperature was 280°C. The initial column temperature was 50°C, rising to 310°C at 3°C/min and the final temperature was held for 10 min. The ion source and quadrupole temperatures were 230°C and 150°C, respectively. The ionization energy (EIMS) was 70 eV. The detection was performed in full scan mode from 41 to 600 amu.

After integration, the percentage of each component in the total ion current (TIC) was calculated. Mass spectral data and calculated retention indices were used to identify the compounds. The mass spectrometric identification was carried out with an automated system of GC-MS data processing supplied by the National Institute of Standards and Technology (NIST). Retention indices of the analytes were determined, taking into account C₁₀–C₄₀n-alkanes retention times and comparing them with the NIST database.

Antimicrobial assay

For the antimicrobial assay, we used the fungi *Candida albicans* ATCC 90028, *C. glabrata* ATCC 90030, *C. krusei* ATCC 6258, *Cryptococcus neoformans* ATCC 90113, and *Aspergillus fumigatus* ATCC 204305 and the bacteria *Staphylococcus aureus* ATCC 29213, methicillin-resistant *S. aureus* ATCC 33591 (MRS), *Escherichia coli* ATCC 35218, *Pseudomonas aeruginosa* ATCC 27853, and *Mycobacterium intracellulare* ATCC 23068 from the American Type Culture Collection (Manassas, VA). The testing was performed using a modified version of the CLSI method (formerly NCCLS) [51–54] except for *M. intracellulare*, where a modified method by Franzblau et al. [55] was used. All samples were diluted in 20% DMSO/saline and then transferred in duplicate to microplates. In order to obtain final target inocula, microbial inocula were prepared by correcting the optical density at 630 nm (OD₆₃₀) of the microbe suspensions in the incubation broth. In each assay, drug controls were used, with ciprofloxacin (ICN Biomedicals, USA) for the bacteria and amphotericin B (ICN Biomedicals, Ohio) for the fungi. The readings were made for all organisms before and after incubation at either 530 nm using the Biotek Powerwave XS plate reader (Bio-Tek Instruments, USA) or 544 excitation / 590 emission, (*M. intracellulare*, *A. fumigatus*) using the Polarstar Galaxy Plate Reader (BMG LabTechnologies, Germany). To determine the minimum fungicidal or bactericidal concentrations required, 5 μ L was removed from each clear well, transferred to agar, and incubated. The MFC/MBC (the lowest test concentration that kills the organism) was also established.

Antimalarial assay

In the antimalarial assay, the chloroquine sensitive (D6) strain of *Plasmodium falciparum* was used and plasmodial LDH activity was measured using the procedure devised by Makler and Hinrichs [56]. To the wells of a 96-well plate containing 10 μ L of diluted sample, a suspension of red blood cells infected with *P. falciparum* of 200 μ L was introduced, together with 2% parasitemia and 2% hematocrit in RPMI 1640 medium supplemented with 10% human serum and 60 μ g/mL Amikacin. Then, the plate was incubated at 37°C for 72 h with 90% N₂, 5% O₂, and 5% CO₂ using a modular incubation chamber. In order to determine the parasitic LDH activity, 20 μ L of the incubation mixture was mixed with 100 μ L of the Malstat reagent (Flow Inc., USA) and incubated at room temperature for 30 min. Then, 20 μ L of a 1:1 mixture of NBT/PES (Sigma, USA) were added and the plate was incubated in the dark for 1 h. To stop the reaction, 100 μ L of a 5% acetic acid solution was added and the absorbance was read at 650 nm. The drug controls were artemisinin and chloroquine. The half-maximal inhibitory concentration (IC₅₀) values were calculated on the basis of the dose response curves of growth inhibition using XLfit 4.2 (IDBS, USA).

Cytotoxicity assay

In the cytotoxicity assay, the root extract was tested against a panel of cancer and non-cancer cell lines. In this assay, we used 96-well tissue culture-treated plates. The cells were seeded at a density of 25,000 cells/well and grown for 24 h. Next, samples of the extract at different concentrations were added and cells were incubated for 48 h. The neutral red method [57] was used to determine the cell viability. The IC₅₀ values were obtained on the basis of dose response curves. Doxorubicin was used as the control drug.

Results

Chromatographic fractionation of the ethanolic extract (0.9 g) of the roots of *P. patens* subsp. *patens* resulted in the isolation of two oleanane-type glycosides identified as hederagenin 3-O- β -D-glucopyranoside (2.7 mg) and hederagenin 3-O- β -D-galactopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside (3.3 mg, patensin) (Fig. 5).





Fig. 5 Structures of two oleanane-type glycosides isolated from *Pulsatilla patens* subsp. *patens*.

HPLC analysis of the silylated methanolic extract of the crude root of *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* revealed the presence of *Pulsatilla* saponin D – hederagenin 3-*O*- α -L-rhamnopyranosyl(1 \rightarrow 2)-[β -D-glucopyranosyl(1 \rightarrow 4)]- α -L-arabinopyranoside (Fig. 6). Analysis of the chromatographic spectra of the extracts from *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* showed similar spectra to the standard solution chromatogram of *Pulsatilla* saponin D. The other chromatographic peaks for the extracts of *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* also showed the presence of triterpenoid saponins, which will need to be investigated further. The *Pulsatilla* saponin D has not been isolated in *Pulsatilla* medicinal plants in Poland until now, with this type of study being conducted here for the first time. *Pulsatilla* saponin D, C₄₇H₇₆O₁₇, has multiple demonstrated biological properties including antifungal, antimicrobial, and cytotoxic activities.



***Pulsatilla* saponin D.** Synonyms: 3 β -[(*O*- β -D-glucopyranosyl-(1 \rightarrow 4))-*O*-[α -L-rhamnopyranosyl-(1 \rightarrow 2)]- α -L-arabinopyranosyl]oxy]-23-hydroxyolean-12-en-28-oic acid; hederacolchiside A; hederagenin 3-*O*-[β -D-glucopyranosyl(1 \rightarrow 4)-[α -L-rhamnopyranosyl(1 \rightarrow 2)]]- α -L-arabinopyranoside; hederagenin 3-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 2)-[β -D-glucopyranosyl-(1 \rightarrow 4)]- α -L-arabinopyranoside; SB 365.

Fig. 6 HPLC chromatogram of standards *Pulsatilla* saponin D – 1 (A); HPLC spectrum of the triterpenoid saponins (*Pulsatilla* saponin D – 1) in *Pulsatilla patens* subsp. *patens* (B); HPLC spectrum of the triterpenoid saponins (*Pulsatilla* saponin D – 1) in *Pulsatilla vulgaris* subsp. *vulgaris* (C) and the structure of *Pulsatilla* saponin D (1) isolated from *Pulsatilla patens* subsp. *patens* and *Pulsatilla vulgaris* subsp. *vulgaris* (D).

Carbohydrates were the main components of the extracts from *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris*. The content of carbohydrates in each extract from *P. patens* subsp. *patens* was over 60% while those from *P. vulgaris* subsp. *vulgaris* were over 90% carbohydrates. The analyzed extracts that were the most abundant were saccharose, glucopyranose and fructofuranose. The extracts from the leaves of these species revealed the presence of carboxylic acids, such as benzoic, caffeic, malic, and succinic acids. Furthermore, glyceric, quinic, and protocatechuic acids were identified in the leaf extracts of *P. patens* subsp. *patens*, while the roots extracts from *P. vulgaris* subsp. *vulgaris* contained the smallest amount of carboxylic acids (Tab. 1).

Microbiological assays of the root extracts showed activity against the fungal pathogen *Candida glabrata*, indicated by the IC₅₀ value of 9.37 µg/mL for *P. patens* subsp. *patens* and 11 µg/mL for *P. vulgaris* subsp. *vulgaris* (Tab. 2, Tab. 3). This result is particularly important as with the increasing use of immunosuppressants, the number of cases of inflammation of mucous membranes and particular systems by *C. glabrata* have significantly increased, especially in people with human immunodeficiency virus. *Candida glabrata* has become the second most important agent responsible for surface (oral cavity, esophagus, vaginal, or urinary duct) or systemic inflammation. The main problem with the inflammation caused by *C. glabrata* is their innate resistance to azole antibiotics that are very effective against other *Candida* species. Treatment of *C. glabrata*-caused inflammation often require the use of amphotericin B that is toxic for humans and can trigger undesired side effects [58]. It has been experimentally shown that the root extracts form *Pulsatilla* species can contribute to new, safer drugs in antifungal therapy against *C. glabrata*.

Tab. 1 Relative chemical composition (% of TIC) of *Pulsatilla patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* extracts.

Group of compounds	t _{ret.} (min)	Fresh leaves of <i>P. patens</i>	Fresh roots of <i>P. patens</i>	t _{ret.} (min)	Fresh leaves of <i>P. vulgaris</i>	Fresh roots of <i>P. vulgaris</i>
Carbohydrates such as:		69.77 ±3.49	88.17 ±4.41		94.14 ±4.71	96.46 ±4.82
α-Fructofuranose	32,494	2.12 ±0.11	3.37 ±0.17	42,998	4.28 ±0.21	1.02 ±0.05
β-Fructofuranose	42,678	2.53 ±0.13	7.68 ±0.38	43,286	4.39 ±0.22	4.19 ±0.21
α-D-Glucopyranose	45,290	13.44 ±0.67	10.19 ±0.51	45,928	16.80 ±0.84	16.89 ±0.84
β-D-Glucopyranose	48,449	17.44 ±0.87	9.42 ±0.47	49,055	19.53 ±0.98	16.81 ±0.84
Saccharose	66,784	14.98 ±0.75	25.74 ±1.29	67,461	23.52 ±1.18	53.65 ±2.28
Carboxylic acids such as:		21.83 ±1.09	3.33 ±0.17		1.13 ±0.06	0.07 ±0.01
Laevulinic acid	-	-	-	15,035	0.05 ±0.01	-
Benzoic acid	19,169	0.03 ±0.01	0.06 ±0.01	19,765	0.04 ±0.01	-
4-Hydroxyvaleric acid	-	-	-	20,763	0.16 ±0.01	-
Succinic acid	22,526	0.34 ±0.02	0.17 ±0.01	23,108	0.07 ±0.01	-
Malic acid	30,260	1.04 ±0.05	1.62 ±0.08	30,861	0.73 ±0.04	0.07 ±0.01
2-Hydroxyglutaric acid	-	-	-	32,448	0.04 ±0.01	-
(E)-Ferulic acid	50,432	0.01 ±0.01	0.02 ±0.01	-	-	-
(E)-Caffeic acid	51,967	1.49 ±0.08	-	52,588	0.06 ±0.01	-
Glyceric acid	23,728	8.46 ±0.42	0.05 ±0.01	-	-	-
3-Phenylactic acid	33,465	-	0.28 ±0.01	-	-	-
Protocatechuic acid	41,965	2.12 ±0.11	-	-	-	-
Shikimic acid	42,394	-	0.22 ±0.01	-	-	-
Quinic acid	44,256	7.58 ±0.38	0.58 ±0.03	-	-	-
Amino acids		2.24 ±0.11	0.58 ±0.03		-	-
Other compounds		5.90 ±0.30	3.86 ±0.19		0.57 ±0.03	1.39 ±0.07
Unidentified compounds		0.17 ±0.01	4.06 ±0.20		4.16 ±0.21	2.08 ±0.10

Tab. 2 The test results of the antimicrobial activity of leaf and root extracts of *Pulsatilla patens* subsp. *patens* and *Pulsatilla vulgaris* subsp. *vulgaris* (primary screen).

Tested strain	Extracts from leaves of <i>P. p.</i> (50 µg/mL)	Extracts from roots of <i>P. p.</i> (50 µg/mL)	Extracts from leaves of <i>P. v.</i> (50 µg/mL)	Extracts from roots of <i>P. v.</i> (50 µg/mL)	Amphotericin B (5 µg/mL)	Ciprofloxacin (1 µg/mL)
<i>C. albicans</i>	2*	27	11	24	100	ND
<i>C. glabrata</i>	0	100	14	98	100	ND
<i>C. krusei</i>	0	17	15	16	100	ND
<i>A. fumigatus</i>	7	3	21	11	99	ND
<i>C. neoformans</i>	0	22	0	40	82	ND
<i>S. aureus</i>	0	8	7	16	ND	89
MRSA	0	10	9	9	ND	94
<i>E. coli</i>	5	11	20	24	ND	96
<i>P. aeruginosa</i>	4	4	7	11	ND	100
<i>M. intracellulare</i>	0	0	0	0	ND	72

* The results in %. ND – not determined; *P. p.* – *Pulsatilla patens*; *P. v.* – *Pulsatilla vulgaris*.

Tab. 3 Dose response (IC₅₀ in µg/mL) results of the roots extracts of *Pulsatilla patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris*.

Test strain	Extracts from roots of <i>P. patens</i>	Extracts from roots of <i>P. vulgaris</i>	Amphotericin B	Ciprofloxacin
<i>C. albicans</i>	NA	NA	0.19	ND
<i>C. glabrata</i>	9.37*	11	0.37	ND
<i>C. krusei</i>	NA	NA	0.67	ND
<i>A. fumigatus</i>	NA	NA	1.17	ND
<i>C. neoformans</i>	42.27	NA	0.18	ND
<i>S. aureus</i>	NA	NA	ND	0.09
MRS	NA	NA	ND	0.08
<i>E. coli</i>	NA	NA	ND	0.01
<i>P. aeruginosa</i>	NA	NA	ND	0.07
<i>M. intracellulare</i>	NA	NA	ND	0.37

* The results in IC₅₀. ND – not determined; NA – not active at 200 µg/mL.

The extracts from the leaves and roots of this plant species showed lower ability to inhibit the growth of the four different fungi (*Candida albicans*, *C. krusei*, *Aspergillus fumigatus*, and *Cryptococcus neoformans*) and other bacteria (*Staphylococcus aureus*, MRSA, *Escherichia coli*, and *Pseudomonas aeruginosa*) pathogenic to humans (Tab. 2, Tab. 3). As far as *C. albicans*, *A. fumigatus*, *E. coli*, and *P. aeruginosa* are concerned, the leaf and root extracts have been found to show biological activity. Although their activity is rather low, the extracts are still highly valuable as *C. albicans* is a frequent fungus pathogen in humans. This species is responsible for the painful inflammation of mucous membranes, e.g., vagina inflammation in women or oral-pharyngeal thrush in AIDS patients. In certain groups of sensitive patients, this species is responsible for serious and life threatening bloodstream inflammation and subsequent inflammation of internal organs [59]. The extracts from *Pulsatilla* species are also attractive in fighting *A. fumigatus*, which is one of the most common saprophytic fungi in the air. With an increasing number of patients treated with immunosuppressants, the incidence of serious and usually lethal invasive aspergillosis has also considerably increased worldwide [60]. Extracts from *Pulsatilla* species are also promising agents for the treatment of *E. coli* caused inflammation. This bacteria strain is the dominant facultative anaerobic

Tab. 4 The test results of the antimalarial activity of leaf and root extracts of *Pulsatilla patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris*.

Tested strain	<i>Plasmodium falciparum</i> (D6 % inhibition)
Chloroquine	97
Chloroquine	94
Chloroquine	98
Extracts from leaves of <i>P. patens</i>	0
Extracts from roots of <i>P. patens</i>	27
Extracts from leaves of <i>P. vulgaris</i>	8
Extracts from roots of <i>P. vulgaris</i>	1

component of the flora of the human large intestine. It can cause inflammation of the urinary tract, sepsis, meningitis, intestinal diseases, and diarrhea, and is difficult to treat as it is resistant to antimicrobial agents [61]. The *Pulsatilla* extracts should also be considered in fighting *P. aeruginosa*, the opportunistic pathogen responsible for hospital infections in patients with immunodeficiency. This pathogen causes inflammation of the lungs related to the use of a respirator, inflammation of the urinary tract, wounds in patients with serious burns, as well as sepsis [62]. Unfortunately, the extracts considered are inactive against *M. intracellulare*.

In our antimalarial assays, the extracts from the leaves and roots showed low activity against the protozoan (27% of inhibition for *P. patens* subsp. *patens* and 1–8% for *P. vulgaris* subsp. *vulgaris*), while the antimalarial drug chloroquine (positive control) showed 94–98% inhibition (Tab. 4).

The root extracts showed cytotoxicity to all the cell lines included in the assay. As shown in Tab. 5, the IC₅₀ for cytotoxicity of *P. patens* subsp. *patens* was in the range of 32–38 µg/mL and for *P. vulgaris* subsp. *vulgaris*, 35–57 µg/mL for each cell line, indicating general cytotoxic activity throughout the panel of cancer and noncancer cells.

Tab. 5 Cytotoxicity of *Pulsatilla patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* roots extracts towards a panel of mammalian cell lines.

Sample name	IC ₅₀ µg/mL					
	SK-MEL	KB	BT-549	SK-OV-3	LLC-PK1	Vero
Root extract of <i>P. patens</i>	34.0	38.0	32.0	32.0	36.0	32.0
Root extract of <i>P. vulgaris</i>	44.0	42.0	57.0	35.0	42.0	39.0
Doxorubicin*	1.7	1.7	2.2	2.3	1.6	>5.0

Cell lines: SK-MEL – skin melanoma; KB – epidermal carcinoma; BT-549 – breast cancer; SK-OV-3 – ovarian cancer; LLC-PK1 – kidney epithelial; Vero – kidney fibroblast. * Positive control drug.

Discussion

The extract of the roots of *P. patens* subsp. *patens* revealed the presence of two oleanane-type glycosides identified as hederagenin 3-O-β-D-glucopyranoside and hederagenin 3-O-β-D-galactopyranosyl-(1→2)-β-D-glucopyranoside (patensin). Structures of both compounds were established via comparing ¹H and ¹³C-NMR spectral data on compounds reported earlier by other authors [63,64]. Patensin, C₄₂H₆₈O₁₄, a triterpenoid glycoside, was also isolated from the ethanolic extract of roots of *P. patens* subsp. *multifida* [63] and five oleanane-type glycosides and two triterpene saponins were also isolated from the roots of *P. patens* subsp. *multifida* [64]. Phytochemical studies of new secondary metabolites from *P. patens* subsp. *multifida* have revealed the presence of a hitherto unknown saponin [12]. This saponin isolated from the methanolic extract of this plant has been found to inhibit the growth of skin cancer. Roots from *P. patens* subsp. *multifida* have been used historically in traditional Chinese medicine [26] as the agent shows anticancer, antimalarial, and antibacterial properties. At present, it is used as a homeopathic drug for the treatment of the ear conditions, exanthema, rheumatism, bronchitis, coughing, asthma, as well as an agent for alleviating stress and anxiety.

The triterpene acid and triterpene glycosides (saponins *Pulsatilla* A and B) were isolated from the plant material of *P. chinensis* [65]. The triterpene acid shows high cytotoxic activity against the malignant cells of lung cancer. The triterpene saponins are used for the treatment of inflammatory conditions [66] and reveal high antiprotozoal


[67], antibacterial [68], antiparasitic [69], antifungal [70], and molluscicidal [71] activities. These saponins have also been used for the treatment of indigestion, premenstrual tension syndrome, and psychosomatic disturbances [72].

Six new saponins, five lupanes, and one oleanane [hederagenin 3-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranosyl-(1 \rightarrow 2)- α -L-arabinopyranoside], along with 11 known saponins, were also isolated from the roots of *P. koreana* [2]. The extracts from *P. koreana* were also determined to contain ranunculin, anemonin, protoanemonins, triterpenes, and saponins (9%). As evidenced in studies on the pharmacological use of components isolated from the extract, its protoanemonins show antifungal properties and antibiotic activity. Moreover, the biologically active compounds isolated from the root extract of *P. koreana* have been found to show anticancer [73], anti-inflammatory [74], antiparasitic [75], and antibacterial [76] activities.

HPLC analysis of the root extract of *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* revealed the presence of *Pulsatilla* saponin D – hederagenin 3-*O*- α -L-rhamnopyranosyl(1 \rightarrow 2)-[β -D-glucopyranosyl(1 \rightarrow 4)]- α -L-arabinopyranoside. Results of numerous phytochemical studies have confirmed high content of triterpenoid saponins of lupane or oleanane-type in the extract of the root of *P. koreana* and *P. chinensis* [2,77] and their chromatographic spectra [50] are similar to those obtained for *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* (Fig. 7A). All structures of triterpenoid saponins (pulsatilloside, *Pulsatilla* saponin, anemoside, hederacolchiside, and cussosaponin) isolated from *P. koreana* roots were elucidated by comparative analysis of spectral data [78].

Analysis of HPLC fingerprints of different species of *Pulsatilla* (*P. koreana*, *P. chinensis*, *P. ambigua*, *P. dahurica*, *P. turczaninowii*) showed significant differences between their HPLC fingerprints, but with 10 common peaks (Fig. 7B). These common peaks not only can be used to classify and authenticate *Pulsatilla* species, but also provide important

A



B




Fig. 7 HPLC-ELSD chromatogram of mixed standards and 15 samples extract of *Pulsatilla koreana* Nakai with spectrum of the triterpenoid saponins (pulsatilloside E, *Pulsatilla* saponin H, anemoside B4, hederacolchiside E, and cussosaponin C-5) (A) [78], and 10 common peaks of different *Pulsatilla* species on HPLC fingerprint study (B) [50].

references for HPLC fingerprints and quality control of *Pulsatilla* medicinal plants [50]. These studies confirmed chromatographic spectra can show differences depending on the *Pulsatilla* species. Significant differences between *P. koreana* and *P. dahurica*, and between *P. turczaninowii* and *P. ambigua* were observed. Slight differences in the chromatographic spectra were also found for *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris*; however, the presence of the triterpenoid saponins and the biological activity of their plant extracts were demonstrated for the first time in this work. This is very important because the triterpenoid saponins, such as *Pulsatilla* saponin D (C₄₇H₇₆O₁₇), have demonstrated multiple biological properties including antifungal, antimicrobial, and cytotoxic activities. These compounds are also produced from the root of several *Pulsatilla* species, including *P. koreana* [1], *P. chinensis* [79], and *P. cernua* [80]. In particular, *Pulsatilla* saponin D shows high cytotoxic activity against cancer cells. *Pulsatilla* saponin D isolated from the root of *P. koreana* showed potent inhibition rate of tumor growth (IR, 82%) at the dose of 6.4 mg/kg on BDF1 mice bearing Lewis lung carcinoma (LLC) cells [1]. This compound is also proving to be promising for the treatment of Alzheimer's disease [15].

It has been reported that the extracts from other species containing triterpenoid saponins, with hederagenin being the active ingredient, showed marked inhibition of serotonin (5-HT), norepinephrine (NE), and dopamine (DA) transporters [81]. Monoamine transporter inhibition studies on pure isolated compounds have not been reported yet, hence the importance of performing CNS studies on isolated compounds from *Pulsatilla* species. In future, we will focus on pure isolated compounds from *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* and submit them to the Psychoactive Drug Screening Program (PDSP), Chapel Hill, NC for monoamine transporter inhibition studies for extracellular monoamines including serotonin (5-HT), norepinephrine (NE), and dopamine (DA) transporters. The two oleanane-type glycosides have been sent to the National Center for the Natural Product Research (NCNPR) at the University of Mississippi to evaluate their antimicrobial activity.

Currently, this preparation, which is called SB31, is undergoing phase II clinical studies in Korea. It was previously reported that SB31 exhibited cytotoxic activity against some human cancer cell lines and potent antitumor activity on a mouse tumor model. An active antitumor constituent from the roots of *P. koreana* was isolated and the main component was found to be deoxypodophyllotoxin (DPT), which exhibited antiangiogenic activity and good antitumor activity against mice bearing LLC cells. However, the DPT content of *Pulsatilla* Radix, which is an active constituent of SB31, is too low for discernible antitumor activity. This suggests that DPT is not the major constituent responsible for the antitumor activity of *P. koreana* roots.

The extracts from *P. cernua* have been found to contain anthocyanidins, as well as triterpene saponins and phytosterols [82]. The natural dyes isolated from *P. cernua* show antiallergic, antifungal, anticancer, and antiulcer activity. Moreover, they improve blood flow in coronary arteries and reduce blood pressure so are attractive substances for the treatment of atherosclerosis. The anthocyanidins isolated from the extracts have been used in medicine as perspiration-promoting agents in the treatment of colds and in the treatment of inflammation of the stomach and intestines.

The flavonoids and tannins isolated from the methanolic extract from *P. nigricans* show considerable antianxiety effect [83]. *Pulsatilla nigricans* is one of the species used in traditional medicine for the treatment of nervous exhaustion and anxiety. Therefore, their high activity permits their application for the treatment of depression. The confirmation of our results shows the extracts from *P. nigricans* can also be used as antifungal agents and to alleviate the weakness of an organism.

Phytochemical studies of the extracts from *P. montana* subsp. *balcana*, *P. halleri* subsp. *rhodopaea*, and *P. slaviankae* have shown polyphenolic compounds, flavonoids, and anthocyanidins [13]. The highest contents of these substances were established in the extracts from *P. montana* subsp. *balcana* and *P. halleri* subsp. *rhodopaea*. The antioxidative properties of *P. slaviankae* were a consequence of the presence of astragalgin, isoquercetin, quercetin, kaempferol, caffeic acid, and isorhamnetin in the aboveground shoots of this species [84]. The natural biologically active substances isolated from this species may play an important role in the protection of humans against breast cancer, colon cancer, prostate cancer, and leukemia. Phytochemical studies of plant material from the species *P. montana* subsp. *balcana* have also revealed the presence of caffeic

acid, chlorogenic acid, gallic acid, methylgallic acid, ferulic acid, and ellagic acid [85]. Our results also identified caffeic acid, and so are in some accordance with the observation of these previous studies. The therapeutic effects related to the presence of these polyphenolic compounds include diastole of blood vessels, and assistance in stabilizing the blood pressure of patients with heart disease and hypertension.

The chemical composition of extracts from the leaves and roots of *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* were significantly different. The leaf extracts, in contrast to the root extracts, contained up to 22% carboxylic acid for *P. patens* subsp. *patens* and 2% for those of *P. vulgaris* subsp. *vulgaris*, whereas the leaf extracts showed lower biological activity. Chemical composition of extracts from dry and raw materials from this species were slightly different (Łaska and Sienkiewicz, unpublished data), while the extracts from fresh material were the most rich with compounds. Upon drying, some compounds were lost through evaporation or destruction.

However, phytochemical analysis of *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris* showed nearly similar results due to the presence of phytochemical constituents. A previous study on the chemotaxonomic characteristics of *Pulsatilla* species revealed that *P. patens* subsp. *patens*, *P. vulgaris* subsp. *vulgaris*, and *P. grandis* Wenderoth contain only delphinidin glycosides, whereas the remaining species [*P. halleri* subsp. *rhodopaea* (Stoj. & Stef.) K. Krause, *P. montana* subsp. *balcana* (Vel.) Zamels, *P. pratensis* (L.) Mill., *P. rubra* (Lamk.) Delarbre, and *P. vulgaris* subsp. *vulgaris* var. *touranginiana* E. G. Camus] contain pelargonidin and delphinidin glycosides [86]. Our studies revealed the presence of the triterpenoid saponins and the biological activity of the extracts from *P. patens* subsp. *patens* and *P. vulgaris* subsp. *vulgaris*. Therefore, isolated saponins have chemotaxonomic features that may be recognized as specific to *Pulsatilla* species.

References

1. Kim Y, Bang SC, Lee JH, Ahn BZ. *Pulsatilla* saponin D: the antitumor principle from *Pulsatilla koreana*. Arch Pharm Res. 2004;27(9):915–918. <https://doi.org/10.1007/BF02975843>
2. Bang SC, Kim Y, Lee JH, Ahn BZ. Triterpenoid saponins from the roots of *Pulsatilla koreana*. J Nat Prod. 2005;68(2):268–272. <https://doi.org/10.1021/np049813h>
3. Seo KH, Jung JW, Thi NN, Lee YH, Baik NI. Flavonoid glycosides from the flowers of *Pulsatilla koreana* Nakai. Natural Product Sciences. 2016;22(1):41–45. <https://doi.org/10.20307/nps.2016.22.1.41>
4. Shumova GS, Savelieva EV, Vladymyrova IN, Tishakova TS. Phenolic compound composition of herb of *Pulsatilla pratensis* (L.) Mill. World Science. 2017;12(28):35–38.
5. Shu Z, Chen Z, Liu YL, Zhu WF, Feng YL, Xu QM, et al. A new oleanane-type triterpenoidal saponin from *Pulsatilla chinensis*. Nat Prod Res. 2013;27(23):2196–2201. <https://doi.org/10.1080/14786419.2013.814052>
6. Yang H, Cho YW, Kim SH, Kim YC, Sung SH. Triterpenoidal saponins of *Pulsatilla koreana* roots. Phytochemistry. 2010;71(16):1892–1899. <https://doi.org/10.1016/j.phytochem.2010.07.013>
7. Fan W, Liu J, Gong Y, Ma J, Zhou N, Xu Y. A new triterpenoid saponin from *Pulsatilla cernua*. Natural Product Sciences. 2013;19(2):150–154.
8. Sun H, Wang Y, Zhang XQ, Zhao SX, Ye WC. Chemical constituents of *Pulsatilla dahurica*. Chem Nat Compd. 2009;45(5):764–765. <https://doi.org/10.1007/s10600-009-9454-3>
9. Xu HJ, Shi XW, Ji X, Du YF, Zhu H, Zhang LT. A rapid method for simultaneous determination of triterpenoid saponins in *Pulsatilla turczaninowii* using microwave-assisted extraction and high performance liquid chromatography-tandem mass spectrometry. Food Chem. 2012;135(1):251–258. <https://doi.org/10.1016/j.foodchem.2012.04.081>
10. Samadder A, Das J, Das S, Khuda-Bukhsh AR. Dihydroxy-isosteviol-methyl-ester, an active biological component of *Pulsatilla nigricans*, reduces arsenic induced cellular dysfunction in testis of male mice. Environ Toxicol Pharmacol. 2012;34(3):743–752.

<https://doi.org/10.1016/j.etap.2012.09.013>

11. Rolski S, Przyborowski L. *Dissertationes Pharmaceuticae*. 1961;13:349–355.
12. Ye WC, Ji NN, Zhao SX, Che CT. A new cytotoxic saponin from *Pulsatilla patens* var. *multifida*. *Pharm Biol*. 2001;39(1):7–10. <https://doi.org/10.1076/phbi.39.1.7.5951>
13. Danova K, Bertoli A, Pistelli L, Dimitrow D, Pistelli L. In vitro culture of Balkan endemic and rare *Pulsatilla* species for conservational purposes and secondary metabolites production. *Bot Serb*. 2009;33(2):157–162.
14. Xu QM, Shu Z, He WJ, Chen LY, Yang SL, Yang G, et al. Antitumor activity of *Pulsatilla chinensis* (Bunge) Regel saponins in human liver tumor 7402 cells in vitro and in vivo. *Phytomedicine*. 2012;19(3–4):293–300. <https://doi.org/10.1016/j.phymed.2011.08.066>
15. Yoo HH, Lee SK, Lim SY, Kim Y, Kang MJ, Kim EJ, et al. LC-MS/MS method for determination of hederacolchiside E, a neuroactive saponin from *Pulsatilla koreana* extract in rat plasma for pharmacokinetic study. *J Pharm Biomed Anal*. 2008;48(5):1425–1429. <https://doi.org/10.1016/j.jpba.2008.09.012>
16. Liu JY, Guan YL, Zou LB, Gong YX, Hua HM, Xu YN, et al. Saponins with neuroprotective effects from the roots of *Pulsatilla cernua*. *Molecules*. 2012;17:5520–5531. <https://doi.org/10.3390/molecules17055520>
17. Dai L, Wang H, Chen Y. The immune-enhancing effect of PcG-A-a glycoprotein isolated from dried root of *Pulsatilla chinensis* (Bunge) Regel. *Chinese Journal of Biochemical Pharmaceutics*. 2000;21:230–231.
18. Seo JS, Yun JH, Baek IS, Leem YH, Kang HW, Cho HK, et al. Oriental medicine Jangwonhwan reduces A β (1–42) level and β -amyloid deposition in the brain of Tg-APPswe/PS1dE9 mouse model of Alzheimer disease. *J Ethnopharmacol*. 2010;128:206–212. <https://doi.org/10.1016/j.jep.2010.01.014>
19. Li HB, Wong CC, Cheng KW, Chen F. Antioxidant properties in vitro and total phenolic contents in methanol extracts from medicinal plants. *LWT – Food Science and Technology*. 2008;41:385–390. <https://doi.org/10.1016/j.lwt.2007.03.011>
20. Lee HS, Beon MS, Kim MK. Selective growth inhibitor toward human intestinal bacteria derived from *Pulsatilla cernua* root. *J Agric Food Chem*. 2001;49:4656–4661. <https://doi.org/10.1021/jf010609z>
21. Xu K, Shu Z, Xu QM, Liu YL, Li XR, Wang YL, et al. Cytotoxic activity of *Pulsatilla chinensis* saponins and their structure-activity relationship. *J Asian Nat Prod Res*. 2013;15(6):680–686. <https://doi.org/10.1080/10286020.2013.790901>
22. Nelson LS, Shih RD, Balick MJ. *Handbook of poisonous and injurious plant*. New York, NY: The New York Botanical Garden; 2007.
23. Wink M, van Wyk BE. *Mind-altering and poisonous plants of the world. A scientifically accurate guide to 1,200 toxic and intoxicating plants*. Portland, OR: Timber Press; 2008.
24. Vance JC. Toxic plants of Minnesota: skin toxicity of the Prairie Crocus (*Anemone patens* L.). *Minn Med*. 1982;65:149–151.
25. Turner NJ. Counter-irritant and other medicinal uses of plants in Ranunculaceae by native peoples in British Columbia and neighbouring areas. *J Ethnopharmacol*. 1984;11:181–200. [https://doi.org/10.1016/0378-8741\(84\)90038-2](https://doi.org/10.1016/0378-8741(84)90038-2)
26. *Encyclopedia of Chinese materia medica*. Shanghai: Shanghai Science and Technology Press; 1977.
27. Szczecińska M, Sramko G, Wołosz K, Sawicki J. Genetic diversity and population structure of the rare and endangered plant species *Pulsatilla patens* (L.) Mill. in east Central Europe. *PLoS One*. 2016;11(3):e0151730. <https://doi.org/10.1371/journal.pone.0151730>
28. Piqueray J, Saad L, Bizoux JP, Mahy G. Why some species cannot colonise restored habitats? The effects of seed and microsite availability. *J Nat Conserv*. 2013;21:189–197. <https://doi.org/10.1016/j.jnc.2012.12.005>
29. Kricsfalusy V. Variations in the life cycle of *Anemone patens* L. (Ranunculaceae) in wild populations of Canada. *Plants (Basel)*. 2016;5(3):E29. <https://doi.org/10.3390/plants5030029>
30. Meusel H, Jäger E, Weinert E. *Vergleichende Chorologie der Zentraleuropäischen Flora*. 1. Karten. Jena: G. Fischer Verl.; 1965.
31. Den virtuella floran – Naturhistoriska riksmuseet [Internet]. 1996 [cited 2017 Aug 1]. Available from: <http://linnaeus.nrm.se/flora/di>

32. Holub J, Procházka F. Red list of vascular plants of the Czech Republic – 2000. *Preslia*. 2000;72:187–230.
33. Röder D, Kiehl K. Population structure and population dynamic of *Pulsatilla patens* (L.) Mill. in relation to vegetation characteristics. *Flora*. 2006;201:499–507. <https://doi.org/10.1016/j.flora.2005.11.001>
34. Gärdenfors U. Population viability analysis in the classification of threatened species: problems and potentials. *Ecol Bull*. 2000;48:181–190.
35. Носков [Noskov] ГА [GA], editor. Красная книга природы Ленинградской области. Растения и грибы [Krasnaja kniga prirody Leningradskoj oblasti. Rastenija i griby]. Vol. 2. Санкт-Петербург [Sankt-Peterbúrg]: Мир [Mir]; 2000.
36. European Nature Information System. Species factsheet for *Pulsatilla patens*. 2005; Available from: <https://eunis.eea.europa.eu/>
37. Průša D, Eliáš MLP, Dítě D, Čačko L, Krása P, Podešva Z, et al. Chránené rastliny Českej a Slovenskej republiky. Brno: Computer Press; 2005.
38. Rassi P, Alanen A, Kanerva T, Mannerkoski I, editors. The 2000 red list of Finnish species. Helsinki: The II Committee for the Monitoring of Threatened Species in Finland, the Ministry of the Environment; 2001.
39. Rozporządzenie Ministra Środowiska z dnia 9 października 2014 r. w sprawie ochrony gatunkowej roślin. *Journal of Laws of the Republic of Poland (Dziennik Ustaw)*, 2014 Oct 9, Item 1409.
40. Zając A, Zając M, editors. Distribution atlas of vascular plants in Poland. Cracow: Laboratory of Computer Chorology, Institute of Botany, Jagiellonian University; 2001.
41. IUCN. The IUCN Red List of Threatened Species. Version 2014.1 [Internet]. 2014 [cited 2014 Jun 12]. Available from: <http://www.iucnredlist.org/>
42. Witkowski ZJ, Król W, Solarz W, editors. Carpathian list of endangered species. Vienna: WWF and Institute of Nature Conservation, Polish Academy of Sciences; 2003.
43. Gärdenfors U. The 2010 red list of Swedish species. Uppsala: ArtDatabanken, SLU; 2010.
44. Cheffings C, Farrell L, editors. The vascular plant red data list for Great Britain. Peterborough: Joint Nature Conservation Committee; 2005.
45. Ludwig G, Schnittler M. Red list of threatened plants in Germany. Bonn: Bundesamt für Naturschutz; 1996.
46. Moser D, Gyğax A, Bäumler B, Wyler N, Palese R. Red list of the threatened ferns and flowering plants of Switzerland. Bern: Bundesamt für Umwelt, Wald und Landschaft, Zentrum des Datenverbundnetzes der Schweizer Flora; 2002.
47. Niklfeld H, Schratt-Ehrendorfer L, editors. Red list of threatened plants of Austria. Vienna: Grüne Reihe, Bundesministerium für Umwelt, Jugend und Familie; 1999.
48. Kaźmierczakowa R, Zarzycki K, Mirek Z. Polish red data book of plants. Cracow: W. Szafer Institute of Botany, Polish Academy of Sciences; 2016.
49. Mirek Z, Piękoś-Mirkowa H, Zając A, Zając M. Flowering plants and pteridophytes of Poland. A checklist. Cracow: W. Szafer Institute of Botany, Polish Academy of Sciences; 2002.
50. Li H, Li H, Hao N, Xu Y, Piao Z. Study on HPLC fingerprint characteristics and chemotaxonomy of *Pulsatilla* medicinal plants. *China Journal of Chinese Materia Medica*. 2011;36(11):1478–1482.
51. National Committee for Clinical Laboratory Standards. Reference method for broth dilution antifungal susceptibility testing of yeasts: approved standard. Wayne, PA: National Committee for Clinical Laboratory Standards; 2002. [NCCLS; vol 22(15)].
52. National Committee for Clinical Laboratory Standards. Reference method for broth dilution antifungal susceptibility testing of yeasts: approved standard. Wayne, PA: National Committee for Clinical Laboratory Standards; 2002. [NCCLS; vol 22(16)].
53. National Committee for Clinical Laboratory Standards. Performance standards for antimicrobial disk susceptibility tests: approved standard. Wayne, PA: National Committee for Clinical Laboratory Standards; 2003. [NCCLS; vol 23(18)].
54. National Committee for Clinical Laboratory Standards. Performance standards for antimicrobial disk susceptibility tests: approved standard. Wayne, PA: National Committee for Clinical Laboratory Standards; 2006. [NCCLS; vol 26(2)].
55. Franzblau SG, Witzig RS, McLaughlin JC, Torres P, Madico G, Hernandez A, et al. Rapid, low-technology MIC determination with clinical *Mycobacterium tuberculosis* isolates by

- using the microplate Alamar Blue assay. *J Clin Microbiol.* 1998;36(2):362–366.
56. Makler MT, Hinrichs DJ. Measurement of the lactate dehydrogenase activity of *Plasmodium falciparum* as an assessment of parasitemia. *Am J Trop Med Hyg.* 1993;48:205. <https://doi.org/10.4269/ajtmh.1993.48.205>
 57. Borenfreund E, Babich H, Martin-Alguacil N. Rapid chemosensitivity assay with human normal and tumor cells in vitro. *In Vitro Cell Dev Biol.* 1990;26:1030. <https://doi.org/10.1007/BF02624436>
 58. Fidel PL, Vazquez JA, Sobel JD. *Candida glabrata*: review of epidemiology, pathogenesis, and clinical disease with comparison to *C. albicans*. *Clin Microbiol Rev.* 1999;12(1):80–96. <https://doi.org/10.1128/CMR.12.1.80>
 59. Kim J, Sudbery P. *Candida albicans*, a major human fungal pathogen. *J Microbiol.* 2011;49(2):171–177. <https://doi.org/10.1007/s12275-011-1064-7>
 60. Latgé JP. *Aspergillus fumigatus* and aspergillosis. *Clin Microbiol Rev.* 1999;12(2):310–350. <https://doi.org/10.1128/9781555815523>
 61. Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev.* 1998;11(1):142–201. <https://doi.org/10.1128/CMR.11.1.142>
 62. Streeter K, Katouli M. *Pseudomonas aeruginosa*: a review of their pathogenesis and prevalence in clinical settings and the environment. *Infect Epidemiol Med.* 2016;2(1):25–32. <https://doi.org/10.18869/modares.iem.2.1.25>
 63. Ye WC, Ou BX, Ji NN, Zhao SX, Ye T, Mc Kervey MA, et al. Patensin, a saponin from *Pulsatilla patens* var. *multifida*. *Phytochemistry.* 1995;39(4):937–939. [https://doi.org/10.1016/0031-9422\(95\)00021-X](https://doi.org/10.1016/0031-9422(95)00021-X)
 64. Ye W, Pan G, Zhang Q, Che CT, Wu H, Zhao S. Five new triterpene saponins from *Pulsatilla patens* var. *multifida*. *J Nat Prod.* 1999;62(2):233–237. <https://doi.org/10.1021/np9802668>
 65. Ye WC, Zhang QW, Hsiao WLW, Zhao SX, Che CT. Novel cytotoxic lupane-type glycosides from *Pulsatilla chinensis*. *Planta Med.* 2002;68:183–186. <https://doi.org/10.1055/s-2002-20254>
 66. Chen Z, Guan Y, Zhou L, Xu Y, Yang M, Liu H. Preparation and characterization of colon-targeted particles of *Pulsatilla chinensis* saponins. *Nat Prod Commun.* 2015;10(2):237–238.
 67. Li LD, Li WC, Liu CW, Shi WJ, Gong PT, Li JH, et al. *Giardia intestinalis*: effects of *Pulsatilla chinensis* extracts on trophozoites. *Parasitol Res.* 2012;111(5):1929–1935. <https://doi.org/10.1007/s00436-012-3035-2>
 68. Li WC, Shi WJ, Gu YF, Chen HL, Chen W, Zhang Y. Antibacterial effect of different extract of *Pulsatilla chinensis* (Bunge) Regel in vitro. *Journal of Traditional Chinese Veterinary Medicine.* 2011;2:38–40.
 69. Ma C. Treatment methods of traditional Chinese medicines against intestinal protozoan infections. In: Mehlhorn H, Zhongdao W, Ye B, editors. *Treatment of human parasitosis in traditional Chinese medicine.* Berlin: Springer; 2014. p. 11–21. (Parasitology Research Monographs; vol 6). https://doi.org/10.1007/978-3-642-39824-7_2
 70. Bi YL, Wang B, Huang BH, Zhang WT, Zhang YH. Antifungal activity of botanical extracts against *Botrytis cinerea* and *Alternaria solani*. *Agricultural Science and Technology.* 2011;12(6):862–864.
 71. Chen YQ, Xu QM, Liu YL, Li XR, Yang SL, Zhuge HX. Laboratory evaluation of the molluscicidal activity of *Pulsatilla chinensis* (Bunge) Regel saponins against the snail *Oncomelania hupensis*. *Biomed Environ Sci.* 2012;25(2):224–229. <https://doi.org/10.3967/0895-3988.2012.02.015>
 72. Chinese Pharmacopoeia Commission. *Pharmacopoeia of the People's Republic of China.* Beijing: People's Medical Publishing House; 2005.
 73. Kim Y, Kim SB, You YJ, Ahn BZ. Deoxypodophyllotoxin; the cytotoxic and antiangiogenic komponent from *Pulsatilla koreana*. *Planta Med.* 2002;68:268–271. <https://doi.org/10.1055/s-2002-23140>
 74. Cheon SA, Choi BK, Jeong CS, Li DW, Lee EB. The anti-inflammatory and analgesic actions of the fractions from *Pulsatilla koreana* root extract. *Korean Journal of Clinical Pharmacy.* 2000;31(2):174–184.
 75. Li W, Sun YN, Yan XY, Yang SY, Lee SJ, Byun HJ, et al. Isolation of nematocidal triterpenoid saponins from *Pulsatilla koreana* root and their activities against *Meloidogyne incognita*. *Molecules.* 2013;18:5306–5316.

<https://doi.org/10.3390/molecules18055306>

76. Chung SW, Chung CH, Lim SB, Kim JK, So EH. The antimicrobial effect of *Pulsatilla koreana* extracts to oral micro-organism. *Journal of the Korean Academy of Periodontology*. 2000;30(3):661. <https://doi.org/10.5051/jkape.2000.30.3.661>
77. Ling Y, Lin Z, Zha W, Lian T, You S. Rapid detection and characterisation of triterpene saponins from the root of *Pulsatilla chinensis* (Bunge) Regel by HPLC-ESI-QTOF-MS/MS. *Phytochem Anal*. 2016;27(3–4):174–183. <https://doi.org/10.1002/pca.2613>
78. Lee KY, Cho YW, Park J, Lee DY, Kim SH, Kim YC, et al. Quality control of *Pulsatilla koreana* based on the simultaneous determination of triterpenoidal saponins by HPLC-ELSD and principal component analysis. *Phytochem Anal*. 2010;21:314–321. <https://doi.org/10.1002/pca.1201>
79. Zhang Y, Bao J, Wang K, Jia X, Zhang C, Huang B, et al. *Pulsatilla* saponin D inhibits autophagic flux and synergistically enhances the anticancer activity of chemotherapeutic agents against HeLa cells. *Am J Chin Med*. 2015;43(8):1657–1670. <https://doi.org/10.1142/S0192415X15500949>
80. Zhang QW, Ye WC, Che CT, Zhao SX. Triterpene saponins from *Pulsatilla cernua*. *Acta Pharmaceutica Sinica*. 2000;35(10):756–759.
81. Jin ZL, Gao N, Zhou D, Chi MG, Yang XM, Xu JP. The extracts of Fructus Akebiae, a preparation containing 90% of the active ingredient hederagenin: serotonin, norepinephrine and dopamine reuptake inhibitor. *Pharmacol Biochem Behav*. 2012;100(3):431–439. <https://doi.org/10.1016/j.pbb.2011.10.001>
82. Xu HJ, Shi XW, Ji X, Du YF, Zhu H, Zhang LT. Qualitative and quantitative determination of nine main active constituents in *Pulsatilla cernua* by high-performance liquid chromatography coupled to electrospray ionization tandem mass spectrometry. *J Sep Sci*. 2011;34:308–316. <https://doi.org/10.1002/jssc.201000660>
83. Goyal S, Kumar S. Anti-anxiety activity studies of various extracts of *Pulsatilla nigricans* Stoerck. *International Journal of Pharmaceutical Sciences and Drug Research*. 2010;2(4):291–293.
84. Nikolova M, Asenov A. Surface flavonoid aglycones in newly studied plant species. *Nat Prod Res*. 2006;20:103–106. <https://doi.org/10.1080/14786410500046463>
85. Arcus M, Lilius G, Schröder V, Taralunga GH, Stoicescu RM, Dumitru IM. Natural compounds with benefits regarding the cardiovascular diseases. *Journal of Environmental Protection and Ecology*. 2015;16(1):333–339.
86. Hegnauer R. *Chemotaxonomie der Pflanzen*. Stuttgart: Birkhäuser Verlag Basel; 1969.