

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

130,000

International authors and editors

155M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Nociceptive TRP Channels and Sex Steroids

*Óscar Enciso-Pablo, Karina Angélica Méndez-Reséndiz,
Tamara Rosenbaum and Sara Luz Morales-Lázaro*

Abstract

Proteins belonging to Transient Receptor Potential (TRP) family are nonselective cation channels that play an essential role in mammalian physiology, functioning as transducers of several environmental signals including those of chemical, thermal and mechanical natures. A subgroup of these receptors is expressed in sensory neurons where they are activated by noxious stimuli and are key players of pain responses in the organism. Some TRP channels are molecular targets for the classical and non-classical effects of sex steroids. This chapter will describe the close relationship between nociceptive TRP channels and sex steroids as well as their impact on nociception and pain-related responses.

Keywords: TRP channels, nociception, steroids, pain, sensory neurons

1. Introduction

The somatosensory system is a complex network of neurons and peripheral receptors that encodes specific information about the state of organisms and their environment, providing them with the ability to detect harmful stimuli [1–3].

According to the International Association for the Study of Pain (IASP), the neuronal process encoding noxious stimuli is known as nociception. A subpopulation of high-threshold sensory neurons known as nociceptors mediates this process by detecting harmful signals from chemical and physical nature **Figure 1** [1, 4]. Nociceptors are neurons with a peripheral axonal branch that innervates cutaneous, articular and visceral afferents, and a central axonal branch that innervates the dorsal horn of the spinal cord. The cell bodies of mammalian nociceptors are located in the dorsal root ganglia (DRG) and in the trigeminal ganglia (TG) and they are classified as medium and small diameter neurons. The axons of these nociceptors are classified as A δ - and C-fibers depending on their diameter, degree of myelination, conduction speed and the type of sensory stimuli that they transmit [1, 4, 5].

The A δ -fibers are characterized by 2–6 μm diameters, are myelinated and present a relatively fast conduction speed between 5 and 30 m/s. The central branch of these fibers reaches the superficial laminae of the dorsal horn. These nociceptor fibers allow the conduction of cold, pressure and heat signals [5].

Remarkably, C-fibers are the ones to mainly establish nociceptive innervation. These are unmyelinated axons with a diameter of 0.4 to 1.2 μm and a conduction speed between 0.5 and 2 m/s. C-fibers innervate laminae I and II of the dorsal horn




Figure 1.

Nociception. Harmful stimuli are transduced along the terminal axon of specialized pseudounipolar neurons (nociceptors), which are classified as unmyelinated C fibers (red) and thinly myelinated A δ fibers (orange); myelinated A β fibers (blue) transduce innocuous stimuli. The cell bodies of these peripheral sensory neurons are located in the dorsal root ganglia and trigeminal ganglia. Nociceptors innervate the epidermis and viscera, transducing noxious stimuli through the activation of several ion channels such as members of the TRP family. Finally, the information is transmitted to second order neurons located in the dorsal horn: C fibers and A δ fibers mainly establish contact with neurons from layers I and II, while tactile A β fibers end mainly in layers III and V.

of the spinal cord and are activated in a polymodal fashion by mechanical, thermal (noxious cold or heat) and chemical stimuli [4, 5].

Diverse injurious stimuli detected by nociceptors are transformed into electrical activity, a process known as transduction where thermal, mechanical, or chemical signals are converted into ion flux through the activation of specialized ion channels [1]. In this way, noxious signals trigger fluctuations in the electrochemical gradient of nociceptors leading to membrane depolarization and subsequent action potentials, as a direct consequence of changes in the activation of ion channels expressed in the nociceptors [6]. Action potentials reach second-order neurons housed in the superficial laminae of the dorsal horn that project towards the spinothalamic and/or spinoparabrachial tracts contributing to the perception of noxious stimuli [2, 3, 7].

The study of the impact of sex steroids on the processing of noxious stimuli has been relevant to our understanding of the underlying differences between females and males. In the last decades, growing evidence related to sexual dimorphism in the perception of harmful signals and the development of pain conditions has been highlighted [8]. Several studies in humans using mechanical, chemical, electrical, or thermal stimuli have suggested that women have more sensitivity to some noxious stimuli than men [8–11]. However, other contrasting reports have concluded that there is no difference in nociception between genders [12, 13]. Even though these differences are still not clear, the most reasonable explanation pinpoints sex steroids as the most relevant effectors. For example, some conditions such as headache, back and temporomandibular pain increase with pubertal development in girls, where the gonadal steroids produce crucial changes in women physiology [14]. Remarkably, differences in the perception of thermal stimuli between different sexes have also been reported, suggesting that women have greater sensitivity to heat than men [15]. In relation to this, an epidemiological study indicated that low testosterone concentrations in women are associated with an increase in the sensation of cold [16], clearly suggesting that steroids influence nociception.

2. Nociceptive TRP channels

The functional properties of nociceptors are critically regulated by the activity of ion channels expressed in their plasma membrane. Some of these channels belong to the Transient Receptor Potential (TRP) family that play vital roles as detectors of a wide range of biological signals. Some TRP channels are crucial receptors within nociceptors since they function as molecular machinery to transduce several noxious sensory stimuli [17].

In mammals, the TRP family consists of 27 members grouped into 6 subfamilies based on their sequence homology as follows: TRPC (canonical), TRPV (vanilloid), TRPM (melastatin), TRPP (polycystin), TRPML (mucolipin) and TRPA (ankyrin) subfamilies. A seventh group is the TRPN (no mechanoreceptor) subfamily which is not found in mammals, although it is found in the fruit fly, in *C. elegans* and zebrafish [18]. These receptors are nonselective cation channels that show an overall tetrameric structure, with each monomer being a six-pass transmembrane (S1-S6) protein with a hydrophilic re-entrant loop between the last two transmembrane segments that give rise to the ion conduction pore in the tetrameric arrangement, **Figure 2**. These channels have cytosolic N- and C- termini which are highly variable between the different members of the TRP subfamilies [19].

Some TRP channels are expressed in nociceptors where they display essential roles in the transduction of several noxious stimuli. Among these nociceptive TRP channels are TRPA1, TRPV1–4, TRPM2–3 and TRPM8 [17] which are activated by noxious thermal, chemical and mechanical signals and hence they are important pain-mediators, **Table 1**.

The **TRPA1 channel**, originally described as a cold sensor [20], is activated by several external compounds such as isothiocyanates contained in natural products such as wasabi, horseradish and mustard oil. Other natural TRPA1 activators are allicin and cinnamaldehyde, compounds found in garlic and cinnamon, respectively, and acrolein is also an exogenously-found compound that activates this ion channel [21–24]. TRPA1's activation by temperature in humans is still controversial, since some reports have described that TRPA1 is an insensitive cold-channel [23, 24],




Figure 2. Overall 3D-structure of a TRP channel. Tetrameric assemble for TRP channels. The colored domains represent conserved domains in these channels: Transmembrane domain (S1-S6); the re-entrant loop forms the ion conduction pore; the pre-S1 helix joins the N-terminal to the S1; TRP box localized in the C-terminus. TM and Cyt: transmembrane and cytosolic regions. (TRPV1 3D- structure, PDB: 3J5P).

Receptor	Temperature	Physical Stimulus	Natural Compounds	Endogenously-produced Compounds
TRPA1	<17°C and noxious heat	Stretch	Allicin, Cinnamaldehyde, Allylisothiocyanate	Androstenedione Lysophosphatidic acid
TRPV1	≥42°C	Osmolality (hyper) Stretch	Capsaicin, Allicin, Resiniferatoxin, Tetrahydrocannabinol	N-arachidonoyl dopamine, Protons Lysophosphatidic acid Arachidonic acid and lipoxygenase-derived products, Anandamide
TRPV2	≥52°C	Osmolality (hypo), Stretch, Shear stress	Cannabidiol	–
TRPV3	30–39°C	–	Camphor, Gingerol, Zingerone, Carvacrol, Eugenol, Thymol	Farnesyl pyrophosphate
TRPV4	30–43°C	Osmolality (hypo), Shear stress	Bisandrographolide A	5,6-epoxyeicosatrienoic acid
TRPM2	>35°C	–	–	Adenosine diphosphate ribose, H ₂ O ₂
TRPM3	≥40°C	Osmolality (hypo)	–	Pregnenolone sulfate
TRPM8	10–28°C	–	Menthol, Eucalyptol	Artemin, Testosterone

Table 1.
Nociceptive TRP channels and their main activators.

while several experiments have strengthened its role as a cold sensor [21, 25–27]. Furthermore, it has also been reported that TRPA1 channel has a redundant role as a noxious heat sensor in the mouse [28]. Indeed, this channel is also indirectly activated by inflammatory mediators like bradykinins and directly by lysophosphatidic acid (LPA, a compound released during injury tissue) and some reactive species produced as the 4-hydroxynonenal [24, 29, 30]. Thus, TRPA1 activation by these endogenous mediators produces inflammatory pain.

Among nociceptive TRP channels some members of the **vanilloid receptors** (i.e., TRPV1–4 channels) are also included. They are activated by warm (TRPV3–V4) to noxious heat (TRPV1–V2) and by several chemical compounds and osmotic changes.

The most studied channel of the vanilloid subfamily is the **TRPV1 channel**, originally described as the capsaicin receptor, since it is specifically activated by this pungent compound found in chili peppers [31]. This channel is also activated by compounds such as allicin, resiniferatoxin (a toxin found in the cactus-like plant *Euphorbia resinifera*) and by animal toxins such as the double-knot (a peptide toxin from the Earth Tiger tarantula), among others [31–33]. Additionally, several reports have demonstrated that TRPV1 activation is achieved by extracellular acid and intracellular basic pHs and heat (≥42°C) [31]. Furthermore, TRPV1 activation is produced by diverse compounds released during inflammation and tissue injury, for example, some arachidonic acid- and lipoxygenase-derived products, LPA and anandamide [34–36]. The high expression of the TRPV1 channel in small and medium-diameter neurons and its polymodal activation through several endogenous noxious stimuli, highlight it as a portal to pain; thus, the understanding of its

regulation is a promising field to develop new strategies in the research field of pain and analgesia.

The vanilloid subfamily contains two less studied than TRPV1 members: **TRPV2 and TRPV3** channels. The former is expressed in medium- and large DRG neurons, activated by temperatures with a threshold of 52°C [37] and by mechanical stretch and osmotic swelling [38]; exogenous compounds like some cannabinoids and endogenous lipids as lysophosphatidyl choline are also TRPV2 activators [39, 40]. Interestingly, the key role of TRPV2 as a noxious sensor is related to mechanical nociception in the somatosensory system, where it is required for the detection of noxious mechanical stimuli [41].

The TRPV3 channel is expressed in keratinocytes and co-expressed with TRPV1 channels in small-diameter sensory neurons. This channel is activated by temperatures between 30 and 39°C [42, 43] and camphor and cannabidiol are exogenous compounds that are also TRPV3 activators [44, 45]. Interestingly, it has been reported that polymorphisms found in the human *TRPV2* and *TRPV3* genes are associated to the susceptibility and severity to fibromyalgia, suggesting the role of these channels in chronic pain disorders [46].

The other vanilloid receptor playing a nociceptive role is the **TRPV4 channel**. This channel is activated by warm temperatures (~30–43°C) and hypotonicity [47–49]. Among the endogenous compounds that are TRPV4 activators are some derivatives of arachidonic acid [50, 51]. Interestingly, this ion channel transduces osmotic stimuli inducing nociception and it is also an important mediator of the pruritogens effects of some compounds like serotonin [52, 53].

Finally, another TRP channels important to nociception are some members of the TRPM subfamily: TRPM2-M3 and TRPM8 channels. Although the **TRPM2 channel** is mainly expressed in immune cells, some expression has also been detected in neurons from DRG and TG [54, 55]. The TRPM2 channel is a warm thermo-sensor, since it is activated at 35°C [56]; other activators for this channel are adenosine 5'-diphosphoribose (ADPR) and H₂O₂ [57, 58]. The activation of this channel through the generation of reactive species (ROS) implicates it as an important mediator of pain perception during stress oxidative conditions, that prevail in some conditions as fibromyalgia and neuropathy diabetic [59, 60].

The **TRPM3 channel**, initially identified in β -pancreatic cells [61] has an important role as a sensor of noxious heat, since it is activated by temperatures $\geq 40^\circ\text{C}$ [62]. The activation of this channel can modulate glucose-induced insulin secretion and also nociception [61, 62]. Its role in the latter is due to high expression in small-diameter sensory neurons from DRG and TG [62] where it also serves as a sensor of noxious chemical stimuli to produce painful responses [62, 63]. Remarkably, TRPM3 together with TRPA1 and TRPV1 channels, form a redundant trio of sensors that mediates noxious heat transduction and provide protection against burn damage [28].

The group of nociceptive TRP channels also comprises the **TRPM8 channel**, a cold-transducer receptor (10–28°C) which is activated by natural compounds as menthol [64, 65]. This ion channel is predominantly expressed in a subpopulation of small-diameter sensory neurons lacking TRPV1 expression [64, 65]. TRPM8 overactivation is related to cold allodynia and hypersensitivity, pain caused by an innocuous cold stimulus and pain produce by enhanced sensitivity to cold stimulus, respectively [66, 67]. Similarly, to other TRP channels, some endogenous compounds such as artemin induces pain through TRPM8 activation [66]. Curiously, a steroid such as testosterone directly binds to this channel and induces its activation [68].

Clearly, the activation of these nociceptive channels is closely related to acute or chronic pain development and interestingly, some of these channels are targets for sex steroids actions, as we will describe here.

2.1 Estrogens and nociceptive TRP channels

Estrogens influence sexual differentiation in women and they have other essential functions in different biological processes as regulators of the cardiovascular system, metabolism, bone resorption and neuronal physiology [69]. These sex steroids comprise estrone (E1) and 17 β estradiol (E2) and estriol (E3) [70]. Similarly, to other steroids, these estrogen's actions are exerted by their interaction with specific nuclear receptors, the estrogen receptors alpha and beta (ER α and ER β) that act as transcription factors to trigger gene regulation through a classical genomic pathway [71]. Furthermore, their actions also are through the activation of a membrane localized receptor, the G Protein-Coupled Estrogen Receptor 1 or GPER1 (previously named GPR30), establishing a non-classical pathway to regulate molecular targets [72, 73]. Several reports have suggested that estrogens influence some painful conditions like fibromyalgia, migraine, irritable bowel syndrome and temporomandibular disorder, all of which are conditions with major prevalence in women than in men [74–78].

Indeed, several experiments made in pain animal models have demonstrated the existence of sexual dimorphism in response to pain. For example, rats from a diabetic neuropathy model exhibited lower mechanical pain thresholds than males. This difference was abolished in gonadectomized animals, highlighting the sex-related differences in diabetic hypersensitivity [79].

These differences could be attributed to estrogen actions on different targets implicated in the transduction of noxious stimuli such as TRP channels, as will be described below.

A close relationship between estrogens and the TRPV1 channel has been demonstrated by several reports. Initially, it was established through behavioral assays that female rats exhibited increased capsaicin pain-related responses as compared to males. Furthermore, ovariectomized rats increased their pain threshold to capsaicin, which was reverted by replacement with E2 in these ovariectomized rats [80]. E2 effects on the TRPV1 nociception have also been evaluated during the rat estrous cycle, where the rats in the proestrus phase showed higher capsaicin-pain responses than the rats in estrus. This observation suggested that high E2 levels exhibited in the proestrus, positively regulate pain associated to TRPV1 activation [80]. These results have also been obtained evaluating the mechanical and thermal pain threshold during the mice estrous cycle, showing that mice in the proestrus displayed lower pain thresholds than the mice in estrus [81]. Remarkably, female TRPV1 knockout mice exhibited comparable mechanical and thermal pain thresholds during the proestrus and estrus [81], strongly suggesting that this channel has a pivotal role in the transduction of these noxious stimuli and that its activation is regulated by estrogen levels. A partial molecular mechanism through which estrogens regulate the capsaicin-evoked pain has been described. Initially, it was suggested that TRPV1 protein levels are downregulated in the DRG from the ER α and ER β knockout mice-model [82], indicating that this ion channel is a molecular target for the effects of estrogen receptors, **Figure 3**. Furthermore, several reports have evidenced that TRPV1 mRNA levels are upregulated by E2. For example, primary cell cultures of rat TG or mice DRG neurons treated with E2 displayed increases in TRPV1 mRNA levels [81, 83]. This TRPV1 mRNA upregulation has also been observed in the hippocampus from ovariectomized rats treated with E2, heightening the allodynia displayed in a rat inflammatory temporomandibular model [84].

The upregulation on TRPV1 gene expression has been also evidenced in synoviocytes, osteoclast precursors and endometrium [85–87]. Furthermore, experiments performed in cultures of human sensory neurons treated with E2 have shown increases in TRPV1 mRNA levels and similar results were obtained when




Figure 3. Schematic representation of the estrogen classical genomic effect in the regulation of TRPV1 gene expression. The inactive estrogen receptors (ER) localized in the cytoplasm form a protein complex with the heat shock protein (HSP). Internalized 17 β estradiol (E2) interacts with the ER, activating and targeting them to the nucleus where they recognize the estrogen response element (ERE), a specific sequence located in the target genes (i.e., TRPV1 gene), resulting in positive regulation of TRPV1 expression.

the sensory neurons were incubated with a specific agonist of the ER β [81, 88]. The E2-dependent effects in the aforementioned cell systems were observed after 24 h of treatment, suggesting that E2 achieves its actions through the classical genomic pathway. This possibility is strengthened by the identification of a putative estrogen response element (ERE) located in the promoter region of the mouse *Trpv1* gene, implying that ER α could interact with this gene regulatory sequence to control TRPV1 gene transcription [89], **Figure 3**. These ER α bindings sites were also identified in the promoter sequences of the *Trpv3–6* genes pointing out that estrogens are regulators of the transcription of these genes [89]. Additionally, it has been determined that different areas of the brains of mice in the proestrus (where the estrogens levels are high) display augmented mRNA levels for the *Trpv2*, *v4* and *v6* genes [89]. In contrast, this report also showed that TRPV1 and TRPV5 mRNA levels are down-regulated in the brain from mice in the proestrus [89].

Additionally, negative E2 effects on TRPV1 activation have been evidenced in ovariectomized rats, which displayed reduced production of gonadal estrogens and augmented neuronal mitochondrial oxidative stress [90]. This was demonstrated using primary cultures of hippocampal and DRG neurons from these rats, determining that the current densities produced through TRPV1 activation are augmented. A similar effect was observed for the TRPA1 and TRPM2 currents [90];

consequently, overactivation of these channels caused rises in cytosolic Ca^{2+} and triggered apoptotic death of these neurons. These effects were prevented in cultures of neurons derived from ovariectomized rats treated with E2 suggesting that this steroid could have neuroprotective actions, avoiding Ca^{2+} overload caused by the activation of some nociceptive TRP channels and preventing neuronal death [90].

Negative E2 actions on TRPV1 regulation have also been shown in primary cultures of rat DRG neurons treated with E2 or with an ER β agonist, displaying decreases in capsaicin-evoked currents while TRPV1 protein levels remained unchanged [91]. This effect was lost using a non-permeable E2 (E2 conjugated to bovine serum albumin) and a slight increase in capsaicin-evoked currents was observed [91]. This result suggests that the role of E2 on TRPV1 activation requires steroid internalization in order to establish an interaction with ER β and to negatively regulate this ion channel [91]. The discrete increase in TRPV1 activation using the non-permeable E2 is in agreement with a previous report which showed that E2 potentiated capsaicin-induced currents in rat DRG primary cultures [92]. Moreover, it was recently described that E2 potentiates TRPV1 activity in an estrogen receptor independent fashion, suggesting that TRPV1 could be an ionotropic receptor for E2 [93]. An interesting outcome from these experimental evidences is that the changes observed in TRPV1 expression are achieved using low E2 concentrations (10–100 nM) [81, 83, 88]; however, the effects on TRPV1 activation need supra-physiological concentrations of the steroid (1–100 μM) [93, 94]. This suggests that low E2 concentrations could modify TRPV1 expression through a classical genomic pathway, whereas a non-classical E2 effect on TRPV1 activation requires high E2 concentrations.

Non-classical E2 actions are produced faster than the genomic effects and are mediated by the interaction of this steroid with the GPER1 [72, 73]. The interaction of E2 with this receptor triggers different signaling pathways, such as those of protein kinase A (PKA) and C (PKC) [72, 95], which could produce TRPV1 phosphorylation. The phosphorylated state of the channel decreases its activation threshold to several stimuli, a process known as sensitization [96]. For instance, a phosphorylated TRPV1 channel can be activated at innocuous heat (i.e., 37°C) [96, 97]. TRPV1 phosphorylation also regulates the channel's interaction with proteins like tubulin [98, 99], a protein association that is important to mediate mechanical gating of TRPV1 [100]. Interruption of the tubulin-TRPV1 complex through the TRPV1 phosphorylation on serine 800 (S800), decreases TRPV1 activation through osmotic stimuli [99, 100]. Interestingly, E2 through its association with GPER1, triggers the activation of PKC ϵ signaling resulting in phosphorylation of TRPV1 in S800 and blocking the interaction of the channel with tubulin [99]; thus, E2 modifies microtubule-dependent TRPV1 mechanical pain sensitization.

The above data exemplify how TRPV1 is a target of classical and non-classical effects of estrogens, highlighting the estrogenic influence on nociception and pain mediated through the TRPV1 channel.

2.1.1 Estrogens upregulate TRPA1 and TRPV1 expression in endometriosis

Endometriosis is a debilitating chronic pelvic pain where the lining layer of the endometrium grows out of the uterus. This condition is highly estrogen-dependent and the lesion displays innervation of nociceptors co-expressing the TRPV1 and TRPA1 channels [101, 102]. Interestingly, mRNA levels for TRPA1 and TRPV1 are upregulated in the biopsies from women with endometriosis [88], thus it is feasible to consider that the overexpression of these channels could mediate the pain produced in this condition. The overexpression of these channels is likely to be through actions of estrogens, since they play a crucial role in the development

of this anomalous pelvic condition. Particularly, it has been demonstrated that female rats treated with diethylstilbestrol, a non-selective estrogen receptor agonist, exhibited upregulation in the TRPV1 and TRPA1 mRNA levels [86] and this expression was found in endometrial cells. Moreover, cultured primary endometrial cells treated with this non-selective agonist also displayed increased TRPA1 and TRPV1 mRNA levels, whereas the treatment with E2 (which is a highly selective agonist of the ER α) only induced the upregulation of the TRPV1 mRNA levels [86]. The data suggests that these nociceptive TRP channels are regulated by the classical effects of estrogen receptors; additionally, these results showed the expression of these channels in non-neuronal cells from the rat endometria. TRPA1 and TRPV1 gene expression also has been demonstrated in human endometria from healthy women; furthermore, endometria samples from women suffering deep infiltrating endometriosis (one of the most severe forms of this condition) exhibit higher TRPV1 and TRPA1 expression, as compared to healthy women [103].

Additionally, it has been reported that hydroxylated estrogens (catechol estrogens) directly activate the TRPA1 channel [104]. This was evidenced using TRPA1-expressing HEK293 cells and whole-cell patch-clamp recordings that showed that 2-hydroxy-estrone (a catechol-estrogen) evoked currents through activation of TRPA1, similarly to another agonist for this channel, cinnamaldehyde. Thus, this hydroxylated estrogen acts as an endogenous agonist for this nociceptive channel [104]. Although, the physiological role of this activation has been demonstrated for insulin secretion in β -pancreatic cells, it would be interesting to explore if the excessive production of catechol-estrogens in endometriosis (as it has been previously reported [105]) affects the function of the TRPA1 channel expressed in endometrial cells and in the free nerve terminals surrounding this area. The expression of these channels in endometrial cells could represent a therapeutic alternative to relieve the main symptoms related to this painful condition; thus, it would be relevant to deepen the studies about the functional roles of TRP channels in these non-neuronal cells.

In addition to the above described evidence, the roles of TRPA1 and TRPV1 in endometriosis have been recently strengthened through the establishment of a mouse endometriosis model [106]. The endometriosis-like lesions induced in these animals displayed the presence of nerve fibers and inflammatory cells around the lesions. Remarkably, these nerve fibers showed expression of TRPA1 and TRPV1 channels [106]. Furthermore, DRG neurons isolated from mice with induced endometriosis exhibited higher Ca²⁺ influx levels than neurons from sham animals, and this effect was through the activation of TRPA1 or TRPV1 channels [106]. The upregulation of these nociceptive channels could be directly associated with spontaneous abdominal pain observed in mice with endometriosis. Interestingly, the treatment of these mice with a synthetic androgen (danazol) or an aromatase inhibitor (letrozole), which are estrogen reducing agents, substantially relieved the spontaneously induced endometriosis pain [106]. These data reinforce the crucial role of estrogens in the development of endometriosis and highlight the importance of the TRPA1 and TRPV1 channels as molecular pain mediators in this debilitating form of abdominal pain.

2.2 Nociceptive TRP channels that are targeted by androgen actions

Androgens are steroids that influence the development of male characteristics and they are produced in the gonads (testicles and ovaries) and adrenal glands [107]. The main endogenously produced androgens are testosterone, androstenedione, dehydroepiandrosterone (DHEA) and dihydrotestosterone (DHT) [108]. Similarly, to other steroids, testosterone and DHT act as ligands for a specific

protein, such as the androgen receptor (AR), while other androgens are weak ligands of the AR. This receptor regulates the expression of several target genes through a classical genomic pathway; furthermore, AR modifies cell physiology through non-classical and non-genomic effects [109].

Indeed, some androgens produce fast- non-genomic effects by interacting with other proteins like some TRP channels and modulating their function. For example, through whole-cell recording electrophysiological experiments, DHEA was shown to decrease capsaicin-dependent currents in dissociated DRG neurons [110]. DHEA effects seem to be direct and specific on TRPV1 channel activation, since with the use of a DHEA stereoisomer the inhibition of channel activation was no longer observed. Additionally, testosterone reduced the activation of TRPV1 to a lesser extent than DHEA [110]. This suggested the presence of a DHEA-binding site in the TRPV1 channel, which should be extracellularly localized, since the steroid was applied to the surface of the neurons during the electrophysiological recordings [110]. However, this remains undetermined and it is possible that DHEA has an antinociceptive role by inhibiting this channel.

The above data exemplified the rapid non-genomic effects of DHEA on TRPV1 activation. Additionally, testosterone seems to negatively regulate the expression of this channel, suggesting that TRPV1 is also a target for the classical actions of androgens. Supporting evidence for this was obtained in dissected TG from castrated male rats with orofacial myositis (induced by the Complete Freund's adjuvant) [111]. The trigeminal TRPV1 mRNA and protein levels from this rat model were upregulated. Interestingly, TRPV1 expression is unaffected in the TG from castrated rats subjected to testosterone replacement [111], suggesting that testosterone prevents the increase in TRPV1 expression, although the molecular mechanism for this regulation is still unknown.

The TRPA1 channel is another molecular target for the non-classical effects of androgens. As a first approach, it was demonstrated that synthetic aromatase inhibitors evoked currents through TRPA1 channels in isolated DRG neurons; however, these compounds were unable to produce currents in DRG neurons from TRPA1 knockout mice, suggesting specific actions of the aromatase inhibitors on the activation of this channel [112]. Indeed, it was demonstrated that the aromatase inhibitors produced pain-like behavior in mice, a conduct not displayed in the TRPA1 knockout mice [112]. Since these compounds increase the levels of androstenedione in plasma, the effects of this androgen on TRPA1 activation have also been studied [113]. The results showed that androstenedione produces currents in mouse DRG neurons that are partially dependent on TRPA1 activation [113]. Furthermore, remanent androstenedione currents were abolished in the presence of the TRPV1 antagonist capsazepine, suggesting that this androgen could activate nociceptive TRPA1 and TRPV1 channels. Although, androstenedione is unable to cause acute pain-like behavior in mice, the animals exhibited mechanical allodynia 30 to 120 min after its intradermal injection into their paws, an effect that was abolished with a TRPA1 antagonist [113]. These data indicate that androstenedione could act as a pain sensitizing molecule leading to the activation of TRPA1.

Finally, the TRPM8 channel is also regulated by androgens. Interestingly, this channel is a target for the classical and nonclassical actions of testosterone. Initial evidence showed that TRPM8 is an androgen-responsive gene since testosterone treatment in some prostate cancer cell lines produced upregulation in the TRPM8 mRNA levels [114, 115]. TRPM8 expression also increases in the urogenital tract from orchietomized rats supplemented with testosterone, suggesting that this steroid positively regulates TRPM8 gene expression [116]. Bioinformatic gene analysis showed that the TRPM8 gene contains putative sites for the androgen

receptor interaction (androgen response elements, ARE) in its promoter sequence. Through chromatin-immunoprecipitation assays, it was confirmed that androgens receptors bind to these AREs located near the start transcription site of the TRPM8 gene [117]. These data indicated that androgens regulate TRPM8 gene expression in a classical genomic fashion.

The TRPM8 channel also acts as an ionotropic testosterone receptor due to direct interaction with this ion channel, resulting in its activation [68, 118]. Experiments have shown that the TRPM8 channel and the androgen receptor compete for testosterone and dihydrotestosterone because activation of TRPM8 by these androgens is lost when both receptors are co-expressed [68]. An extracellularly-located testosterone binding site in the TRPM8 channel has been suggested [68], and by using molecular docking simulation we have identified an additional intracellular testosterone pocket located between the first intracellular loop (pre-S1) and the transmembrane segments 1 and 2 (S1-S2) of the TRPM8 protein [119]. This analysis has also shown that an H-bond is formed between testosterone and an arginine located in the TRP box, similar to the H-bond established between testosterone and an arginine in the aromatase enzyme [119]. The similarity in the testosterone pockets of the aromatase and the TRPM8 proteins could suggest that this is a functional binding site for TRPM8 through which activation by this steroid could be achieved in a non-classical and fast fashion. Interestingly, it has been demonstrated that topical testosterone application on humans produced cold- and pain-sensation; the women resulted more sensitized to the testosterone effects than men [68], suggesting a sexual dimorphism in testosterone-produced nociception through TRPM8 activation.

A third non-classical mechanism for TRPM8 regulation by testosterone has been demonstrated in DRG neurons where the androgen receptor and TRPM8 channels form a protein complex in the plasma membrane [120]. High testosterone concentrations (100 nM) inhibit TRPM8 channel activation and the androgen receptor is translocated to the nucleus [120], **Figure 4**. The evaluation of the TRPM8-mediated cold perception performed in castrated mice or rats revealed a decrease in their threshold for mild-cold perception; interestingly, testosterone




Figure 4. Testosterone regulates the TRPM8-mediated mild-cold sensitivity. The mild-cold stimulus is transduced through the TRPM8 channel located in the cell surface of DRG neurons where it forms a protein complex with the androgen receptor (AR) (upper panel). A high concentration of testosterone (100 nM) decreases activation of the TRPM8 channel and produces translocation of the AR to the nucleus. The physiological consequence of this is the loss of the mild cold sensation (lower panel).

supplementation in these animals inhibited cold perception (temperatures between 16 and 18°C) [120]. This result would suggest that testosterone confers endurance to TRPM8-mediated mild cold perception, which could explain the underlying molecular mechanism for the cold sensation experienced by women presenting lower levels of testosterone [16].

2.3 Progesterone downregulates TRPV1 and TRPV4 channels expression

Progesterone is a steroid indispensable to the maintenance of pregnancy and it also plays important roles in the cardiovascular, immune, nervous systems and in the bones [121]. Progesterone is produced by the granulosa cells and the corpus luteum in the ovaries and, during pregnancy, the placenta is the major source of this steroid. Additionally, progesterone is produced in the nervous system, adrenal glands and testes [122]. The effects of progesterone are produced through nuclear receptors A and B (PRA/B), activating the classical genomic pathway to regulate gene expression. Furthermore, this steroid exerts a non-classical pathway through the activation of several membrane receptors (mPR) and other receptors as the progesterone receptor membrane components 1 and 2 (PGRMC1/2) and by the Sigma-1 Receptor (Sig-1R) [123, 124].

It has been demonstrated that progesterone regulates the expression of some TRP channels; for example, TRPV4 gene expression is negatively regulated by classical genomic progesterone effects [125]. Initial evidence for this was observed in epithelial and vascular cell lines and a recent report showed a correlation between TRPV4 expression and progesterone levels during the menstrual cycle, demonstrating that during the luteal phase when progesterone reaches its highest levels, TRPV4 expression is downregulated in human endometrial biopsies [126]. Furthermore, it has also been reported that TRPV4 expression is downregulated by a synthetic progestin (levonorgestrel) in a human Fallopian tubal epithelial cell line and mouse oviducts which could affect the transport of fertilized oocytes causing ectopic pregnancy [127].

Our group has determined that progesterone impacts on nociception by regulating the expression of TRPV1 channels [128]. Interestingly, this regulation is through a non-classical mechanism that requires an intermediate protein, the Sig-1R, which is a dynamic chaperone mainly localized in the endoplasmic reticulum and that can also be mobilized to the plasma membrane and nuclear envelope [129]. Progesterone decreases the levels of channels localized in the cell surface in HEK293 cells transfected with TRPV1 cDNA [128]. But how does progesterone produce this effect? Our results have shown that this is achieved by disrupting a protein complex formed between TRPV1 channels and the Sig-1R and by affecting the proper folding of TRPV1. Consequently, the channels are degraded by the proteasomal pathway and capsaicin-evoked currents are reduced [128]. The TRPV1 downregulation through progesterone actions was also observed in primary cultures of DRG neurons and we also demonstrated that during mice pregnancy when there are high progesterone levels, the threshold to capsaicin-dependent pain-like behaviors are augmented in comparison to non-pregnant mice, suggesting that progesterone confers protection to pain produced through activation of TRPV1 [128]. This work exemplified the antinociceptive actions that progesterone exerts through the decrease in the levels of TRPV1 channels localized in the plasma membrane of nociceptors and by the reduction of the transduction of harmful signals through this ion channel. Moreover, it has also been reported that progesterone disrupts the protein complex formed between Sig-1R and the TRPA1 and TRPM8 channels [130], suggesting that this steroid could impact the nociception through these nociceptive TRP channels.

3. Conclusion

Nociception is a neuronal process that protects against injurious stimuli that can be endogenously produced during inflammation or tissue injury processes. The transduction of these signals requires the activation of ion channels such as some members of the TRP family. In this chapter we have described how sex steroids influence nociception by regulating some of the TRP channels important for this neuronal process. Here we have described the data showing that estrogens and their nuclear receptors (ER α / β) positively regulate the expression of the TRPV1 and TRPA1 genes through a classical genomic pathway. The reports above discussed also show that progesterone and its nuclear receptors (PRA/B) negatively regulate TRPV4 gene expression.

We have also compiled and discussed the evidence about androstenedione's and testosterone's non-classical effects in the regulation of the activation of TRPA1 and TRPM8 ion channels directly. We have suggested a testosterone binding-pocket in the TRPM8 channel and we have also pinpointed an androstenedione binding-pocket in the C-terminal and pre-S1 regions of TRPV1 and TRPA1 channels [119].

These observations strengthen the notion that androgens act as direct molecular regulators of some nociceptive ion channels function, playing dual roles as nociceptive and antinociceptive molecules. For example, androstenedione has nociceptive actions on TRPA1 and TRPV1 activation and testosterone at low concentrations (picomolar range) can produce cold sensation through the TRPM8 channel, while high testosterone levels could improve resilience to cold sensation inhibiting the currents through TRPM8. Finally, the literature here reviewed shows that the nonclassical sex steroid effects are mediated by membrane receptors or even chaperones such as the Sig-1R, affecting signaling pathways and/or protein stability. This last finding exemplifies the antinociceptive effects of progesterone by which this hormone reduces the transduction of noxious stimuli mediated by TRPV1 channel activation. These data open a research field in which we will deepen our knowledge in the role of molecular sexual dimorphism and the interplay between nociceptive TRP channels and sex steroids.

Acknowledgements

Funding: Este trabajo fue apoyado por el FORDECYT-PRONACES (FORDECYT-PRONACES/64392/2020) del Consejo Nacional de Ciencia y Tecnología. También recibió apoyo de la Dirección General de Asuntos del Personal Académico (DGAPA)-Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica (PAPIIT) IN206819 y de los Estímulos a Investigaciones Médicas Miguel Alemán Valdés.

Conflict of interest

The authors declare no conflict of interest.

Notes/thanks/other declarations

Ó.E.P and K.A.M.R contributed equally to this work. This work comprises the MSc thesis work of Ó.E.P. from the Programa de Maestría y Doctorado en Ciencias Bioquímicas at the Universidad Nacional Autónoma de México and the PhD thesis work of K.A.M.R. student from the Programa de Doctorado en Ciencias Biomédicas at the Universidad Nacional Autónoma de México. We thank Alejandra Llorente for her administrative support.

IntechOpen

IntechOpen


Author details

Óscar Enciso-Pablo, Karina Angélica Méndez-Reséndiz, Tamara Rosenbaum
and Sara Luz Morales-Lázaro*

Department of Cognitive Neuroscience, Institute of Cellular Physiology, UNAM,
México City, México

*Address all correspondence to: saraluzm@ifc.unam.mx

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Tracey, W.D., Jr. *Nociception*. *Curr Biol*. 2017. 27(4): p. R129-R133. 10.1016/j.cub.2017.01.037
- [2] Basbaum, A.I., et al. *Cellular and molecular mechanisms of pain*. *Cell*. 2009. 139(2): p. 267-84. 10.1016/j.cell.2009.09.028
- [3] Woolf, C.J. and Q. Ma. *Nociceptors--noxious stimulus detectors*. *Neuron*. 2007. 55(3): p. 353-64. 10.1016/j.neuron.2007.07.016
- [4] Armstrong, S.A. and M.J. Herr, *Physiology, Nociception*, in *StatPearls*. 2020: Treasure Island (FL).
- [5] Dubin, A.E. and A. Patapoutian. *Nociceptors: the sensors of the pain pathway*. *J Clin Invest*. 2010. 120(11): p. 3760-72. 10.1172/JCI42843
- [6] Waxman, S.G. and G.W. Zamponi. *Regulating excitability of peripheral afferents: emerging ion channel targets*. *Nat Neurosci*. 2014. 17(2): p. 153-63. 10.1038/nn.3602
- [7] Todd, A.J. *Neuronal circuitry for pain processing in the dorsal horn*. *Nat Rev Neurosci*. 2010. 11(12): p. 823-36. 10.1038/nrn2947
- [8] Fillingim, R.B., et al. *Sex, gender, and pain: a review of recent clinical and experimental findings*. *J Pain*. 2009. 10(5): p. 447-85. 10.1016/j.jpain.2008.12.001
- [9] Bartley, E.J. and R.B. Fillingim. *Sex differences in pain: a brief review of clinical and experimental findings*. *Br J Anaesth*. 2013. 111(1): p. 52-8. 10.1093/bja/aet127
- [10] Cairns, B.E. *The influence of gender and sex steroids on craniofacial nociception*. *Headache*. 2007. 47(2): p. 319-24. 10.1111/j.1526-4610.2006.00708.x
- [11] Hashmi, J.A. and K.D. Davis. *Deconstructing sex differences in pain sensitivity*. *Pain*. 2014. 155(1): p. 10-3. 10.1016/j.pain.2013.07.039
- [12] Racine, M., et al. *A systematic literature review of 10 years of research on sex/gender and experimental pain perception - part 1: are there really differences between women and men?* *Pain*. 2012. 153(3): p. 602-18. 10.1016/j.pain.2011.11.025
- [13] Racine, M., et al. *A systematic literature review of 10 years of research on sex/gender and pain perception - part 2: do biopsychosocial factors alter pain sensitivity differently in women and men?* *Pain*. 2012. 153(3): p. 619-35. 10.1016/j.pain.2011.11.026
- [14] LeResche, L., et al. *Relationship of pain and symptoms to pubertal development in adolescents*. *Pain*. 2005. 118(1-2): p. 201-9. 10.1016/j.pain.2005.08.011
- [15] Lue, Y.J., et al. *Thermal pain tolerance and pain rating in normal subjects: Gender and age effects*. *Eur J Pain*. 2018. 22(6): p. 1035-1042. 10.1002/ejp.1188
- [16] Gotmar, A., et al. *Symptoms in peri- and postmenopausal women in relation to testosterone concentrations: data from The Women's Health in the Lund Area (WHILA) study*. *Climacteric*. 2008. 11(4): p. 304-14. 10.1080/13697130802249769
- [17] Mickle, A.D., A.J. Shepherd, and D.P. Mohapatra. *Sensory TRP channels: the key transducers of nociception and pain*. *Prog Mol Biol Transl Sci*. 2015. 131: p. 73-118. 10.1016/bs.pmbts.2015.01.002
- [18] Clapham, D.E., L.W. Runnels, and C. Strubing. *The TRP ion channel family*. *Nat Rev Neurosci*. 2001. 2(6): p. 387-96. 10.1038/35077544

- [19] Rosasco, M.G. and S.E. Gordon, *TRP Channels: What Do They Look Like?*, in *Neurobiology of TRP Channels*, T.L.R. Emir, Editor. 2017: Boca Raton (FL). p. 1-9.
- [20] Story, G.M., et al. *ANKTM1, a TRP-like channel expressed in nociceptive neurons, is activated by cold temperatures*. *Cell*. 2003. 112(6): p. 819-29. 10.1016/S0092-8674(03)00158-2
- [21] Bandell, M., et al. *Noxious cold ion channel TRPA1 is activated by pungent compounds and bradykinin*. *Neuron*. 2004. 41(6): p. 849-57. 10.1016/S0896-6273(04)00150-3
- [22] Bautista, D.M., et al. *Pungent products from garlic activate the sensory ion channel TRPA1*. *Proc Natl Acad Sci U S A*. 2005. 102(34): p. 12248-52. 10.1073/pnas.0505356102
- [23] Jordt, S.E., et al. *Mustard oils and cannabinoids excite sensory nerve fibres through the TRP channel ANKTM1*. *Nature*. 2004. 427(6971): p. 260-5. 10.1038/nature02282
- [24] Bautista, D.M., et al. *TRPA1 mediates the inflammatory actions of environmental irritants and proalgesic agents*. *Cell*. 2006. 124(6): p. 1269-82. 10.1016/j.cell.2006.02.023
- [25] Fajardo, O., et al. *TRPA1 channels mediate cold temperature sensing in mammalian vagal sensory neurons: pharmacological and genetic evidence*. *J Neurosci*. 2008. 28(31): p. 7863-75. 10.1523/JNEUROSCI.1696-08.2008
- [26] Karashima, Y., et al. *TRPA1 acts as a cold sensor in vitro and in vivo*. *Proc Natl Acad Sci U S A*. 2009. 106(4): p. 1273-8. 10.1073/pnas.0808487106
- [27] Moparthy, L., et al. *Human TRPA1 is intrinsically cold- and chemosensitive with and without its N-terminal ankyrin repeat domain*. *Proc Natl Acad Sci U S A*. 2014. 111(47): p. 16901-6. 10.1073/pnas.1412689111
- [28] Vandewauw, I., et al. *A TRP channel trio mediates acute noxious heat sensing*. *Nature*. 2018. 555(7698): p. 662-666. 10.1038/nature26137
- [29] Trevisani, M., et al. *4-Hydroxynonenal, an endogenous aldehyde, causes pain and neurogenic inflammation through activation of the irritant receptor TRPA1*. *Proc Natl Acad Sci U S A*. 2007. 104(33): p. 13519-24. 10.1073/pnas.0705923104
- [30] Kittaka, H., et al. *Lysophosphatidic acid-induced itch is mediated by signalling of LPA5 receptor, phospholipase D and TRPA1/TRPV1*. *J Physiol*. 2017. 595(8): p. 2681-2698. 10.1113/JP273961
- [31] Caterina, M.J., et al. *The capsaicin receptor: a heat-activated ion channel in the pain pathway*. *Nature*. 1997. 389(6653): p. 816-24. 10.1038/39807
- [32] Bohlen, C.J., et al. *A bivalent tarantula toxin activates the capsaicin receptor, TRPV1, by targeting the outer pore domain*. *Cell*. 2010. 141(5): p. 834-45. 10.1016/j.cell.2010.03.052
- [33] Salazar, H., et al. *A single N-terminal cysteine in TRPV1 determines activation by pungent compounds from onion and garlic*. *Nat Neurosci*. 2008. 11(3): p. 255-61. 10.1038/nn2056
- [34] Hwang, S.W., et al. *Direct activation of capsaicin receptors by products of lipoxygenases: endogenous capsaicin-like substances*. *Proc Natl Acad Sci U S A*. 2000. 97(11): p. 6155-60. 10.1073/pnas.97.11.6155
- [35] Nieto-Posadas, A., et al. *Lysophosphatidic acid directly activates TRPV1 through a C-terminal binding site*. *Nat Chem Biol*. 2011. 8(1): p. 78-85. 10.1038/nchembio.712
- [36] Zygmunt, P.M., et al. *Vanilloid receptors on sensory nerves mediate the vasodilator action of anandamide*. *Nature*. 1999. 400(6743): p. 452-7. 10.1038/22761

- [37] Caterina, M.J., et al. *A capsaicin-receptor homologue with a high threshold for noxious heat*. *Nature*. 1999. 398(6726): p. 436-41. 10.1038/18906
- [38] Muraki, K., et al. *TRPV2 is a component of osmotically sensitive cation channels in murine aortic myocytes*. *Circ Res*. 2003. 93(9): p. 829-38. 10.1161/01.RES.0000097263.10220.0C
- [39] De Petrocellis, L., et al. *Effects of cannabinoids and cannabinoid-enriched Cannabis extracts on TRP channels and endocannabinoid metabolic enzymes*. *Br J Pharmacol*. 2011. 163(7): p. 1479-94. 10.1111/j.1476-5381.2010.01166.x
- [40] Monet, M., et al. *Lysophospholipids stimulate prostate cancer cell migration via TRPV2 channel activation*. *Biochim Biophys Acta*. 2009. 1793(3): p. 528-39. 10.1016/j.bbamcr.2009.01.003
- [41] Katanosaka, K., et al. *TRPV2 is required for mechanical nociception and the stretch-evoked response of primary sensory neurons*. *Sci Rep*. 2018. 8(1): p. 16782. 10.1038/s41598-018-35049-4
- [42] Smith, G.D., et al. *TRPV3 is a temperature-sensitive vanilloid receptor-like protein*. *Nature*. 2002. 418(6894): p. 186-90. 10.1038/nature00894
- [43] Xu, H., et al. *TRPV3 is a calcium-permeable temperature-sensitive cation channel*. *Nature*. 2002. 418(6894): p. 181-6. 10.1038/nature00882
- [44] De Petrocellis, L., et al. *Cannabinoid actions at TRPV channels: effects on TRPV3 and TRPV4 and their potential relevance to gastrointestinal inflammation*. *Acta Physiol (Oxf)*. 2012. 204(2): p. 255-66. 10.1111/j.1748-1716.2011.02338.x
- [45] Moqrich, A., et al. *Impaired thermosensation in mice lacking TRPV3, a heat and camphor sensor in the skin*. *Science*. 2005. 307(5714): p. 1468-72. 10.1126/science.1108609
- [46] Park, D.J., et al. *Polymorphisms of the TRPV2 and TRPV3 genes associated with fibromyalgia in a Korean population*. *Rheumatology (Oxford)*. 2016. 55(8): p. 1518-27. 10.1093/rheumatology/kew180
- [47] Guler, A.D., et al. *Heat-evoked activation of the ion channel, TRPV4*. *J Neurosci*. 2002. 22(15): p. 6408-14. 20026679
- [48] Liedtke, W., et al. *Vanilloid receptor-related osmotically activated channel (VR-OAC), a candidate vertebrate osmoreceptor*. *Cell*. 2000. 103(3): p. 525-35. 10.1016/s0092-8674(00)00143-4
- [49] Watanabe, H., et al. *Heat-evoked activation of TRPV4 channels in a HEK293 cell expression system and in native mouse aorta endothelial cells*. *J Biol Chem*. 2002. 277(49): p. 47044-51. 10.1074/jbc.M208277200
- [50] Berna-Erro, A., et al. *Structural determinants of 5',6'-epoxyeicosatrienoic acid binding to and activation of TRPV4 channel*. *Sci Rep*. 2017. 7(1): p. 10522. 10.1038/s41598-017-11274-1
- [51] Watanabe, H., et al. *Anandamide and arachidonic acid use epoxyeicosatrienoic acids to activate TRPV4 channels*. *Nature*. 2003. 424(6947): p. 434-8. 10.1038/nature01807
- [52] Akiyama, T., et al. *Involvement of TRPV4 in Serotonin-Evoked Scratching*. *J Invest Dermatol*. 2016. 136(1): p. 154-160. 10.1038/JID.2015.388
- [53] Alessandri-Haber, N., et al. *Hypotonicity induces TRPV4-mediated nociception in rat*. *Neuron*. 2003. 39(3): p. 497-511. 10.1016/s0896-6273(03)00462-8
- [54] Matsumoto, K., et al. *Role of transient receptor potential melastatin 2 (TRPM2) channels in visceral nociception and hypersensitivity*. *Exp Neurol*.

2016. 285(Pt A): p. 41-50. 10.1016/j.expneurol.2016.09.001

[55] Vandewauw, I., G. Owsianik, and T. Voets. *Systematic and quantitative mRNA expression analysis of TRP channel genes at the single trigeminal and dorsal root ganglion level in mouse*. BMC Neurosci. 2013. 14: p. 21. 10.1186/1471-2202-14-21

[56] Togashi, K., et al. *TRPM2 activation by cyclic ADP-ribose at body temperature is involved in insulin secretion*. EMBO J. 2006. 25(9): p. 1804-15. 10.1038/sj.emboj.7601083

[57] Perraud, A.L., et al. *ADP-ribose gating of the calcium-permeable LTRPC2 channel revealed by Nudix motif homology*. Nature. 2001. 411(6837): p. 595-9. 10.1038/35079100

[58] Wehage, E., et al. *Activation of the cation channel long transient receptor potential channel 2 (LTRPC2) by hydrogen peroxide. A splice variant reveals a mode of activation independent of ADP-ribose*. J Biol Chem. 2002. 277(26): p. 23150-6. 10.1074/jbc.M112096200

[59] Sozbir, E. and M. Naziroglu. *Diabetes enhances oxidative stress-induced TRPM2 channel activity and its control by N-acetylcysteine in rat dorsal root ganglion and brain*. Metab Brain Dis. 2016. 31(2): p. 385-93. 10.1007/s11011-015-9769-7

[60] Yuksel, E., et al. *Involvement of TRPM2 and TRPV1 channels on hyperalgesia, apoptosis and oxidative stress in rat fibromyalgia model: Protective role of selenium*. Sci Rep. 2017. 7(1): p. 17543. 10.1038/s41598-017-17715-1

[61] Wagner, T.F., et al. *Transient receptor potential M3 channels are ionotropic steroid receptors in pancreatic beta cells*. Nat Cell Biol. 2008. 10(12): p. 1421-30. 10.1038/ncb1801

[62] Vriens, J., et al. *TRPM3 is a nociceptor channel involved in the detection of noxious heat*. Neuron.

2011. 70(3): p. 482-94. 10.1016/j.neuron.2011.02.051

[63] Held, K., et al. *Activation of TRPM3 by a potent synthetic ligand reveals a role in peptide release*. Proc Natl Acad Sci U S A. 2015. 112(11): p. E1363-72. 10.1073/pnas.1419845112

[64] Peier, A.M., et al. *A TRP channel that senses cold stimuli and menthol*. Cell. 2002. 108(5): p. 705-15. 10.1016/s0092-8674(02)00652-9

[65] Bautista, D.M., et al. *The menthol receptor TRPM8 is the principal detector of environmental cold*. Nature. 2007. 448(7150): p. 204-8. 10.1038/nature05910

[66] Lippoldt, E.K., et al. *Artemin, a glial cell line-derived neurotrophic factor family member, induces TRPM8-dependent cold pain*. J Neurosci. 2013. 33(30): p. 12543-52. 10.1523/JNEUROSCI.5765-12.2013

[67] Xing, H., et al. *TRPM8 mechanism of cold allodynia after chronic nerve injury*. J Neurosci. 2007. 27(50): p. 13680-90. 10.1523/JNEUROSCI.2203-07.2007

[68] Asuthkar, S., et al. *The TRPM8 protein is a testosterone receptor: II. Functional evidence for an ionotropic effect of testosterone on TRPM8*. J Biol Chem. 2015. 290(5): p. 2670-88. 10.1074/jbc.M114.610873

[69] Barkhem, T., S. Nilsson, and J.A. Gustafsson. *Molecular mechanisms, physiological consequences and pharmacological implications of estrogen receptor action*. Am J Pharmacogenomics. 2004. 4(1): p. 19-28. 10.2165/00129785-200404010-00003

[70] Gruber, C.J., et al. *Production and actions of estrogens*. N Engl J Med. 2002. 346(5): p. 340-52. 10.1056/NEJMra000471

[71] Yasar, P., et al. *Molecular mechanism of estrogen-estrogen receptor signaling*.

Reprod Med Biol. 2017. 16(1): p. 4-20.
10.1002/rmb2.12006

[72] Hammes, S.R. and E.R. Levin. *Extranuclear steroid receptors: nature and actions*. Endocr Rev. 2007. 28(7): p. 726-41. 10.1210/er.2007-0022

[73] Revankar, C.M., et al. *A transmembrane intracellular estrogen receptor mediates rapid cell signaling*. Science. 2005. 307(5715): p. 1625-30. 10.1126/science.1106943

[74] Buse, D.C., et al. *Sex differences in the prevalence, symptoms, and associated features of migraine, probable migraine and other severe headache: results of the American Migraine Prevalence and Prevention (AMPP) Study*. Headache. 2013. 53(8): p. 1278-99. 10.1111/head.12150

[75] van Oosterhout, W.P.J., et al. *Female sex hormones in men with migraine*. Neurology. 2018. 91(4): p. e374-e381. 10.1212/WNL.00000000000005855

[76] Lovell, R.M. and A.C. Ford. *Global prevalence of and risk factors for irritable bowel syndrome: a meta-analysis*. Clin Gastroenterol Hepatol. 2012. 10(7): p. 712-721 e4. 10.1016/j.cgh.2012.02.029

[77] Wan, B., et al. *Gender-Stratified Prevalence of Psychiatric and Pain Diagnoses in a Primary Care Patient Sample with Fibromyalgia*. Pain Med. 2019. 20(11): p. 2129-2133. 10.1093/pm/pnz084

[78] Bagis, B., et al. *Gender difference in prevalence of signs and symptoms of temporomandibular joint disorders: a retrospective study on 243 consecutive patients*. Int J Med Sci. 2012. 9(7): p. 539-44. 10.7150/ijms.4474

[79] Joseph, E.K. and J.D. Levine. *Sexual dimorphism in the contribution of protein kinase C isoforms to nociception in the streptozotocin diabetic rat*. Neuroscience. 2003. 120(4): p. 907-13. 10.1016/s0306-4522(03)00400-7

[80] Lu, Y.C., et al. *17Beta-estradiol mediates the sex difference in capsaicin-induced nociception in rats*. J Pharmacol Exp Ther. 2009. 331(3): p. 1104-10. 10.1124/jpet.109.158402

[81] Payrits, M., et al. *Estradiol Sensitizes the Transient Receptor Potential Vanilloid 1 Receptor in Pain Responses*. Endocrinology. 2017. 158(10): p. 3249-3258. 10.1210/en.2017-00101

[82] Cho, T. and V.V. Chaban. *Expression of P2X3 and TRPV1 receptors in primary sensory neurons from estrogen receptors-alpha and estrogen receptor-beta knockout mice*. Neuroreport. 2012. 23(9): p. 530-4. 10.1097/WNR.0b013e328353fabcb

[83] Yamagata, K., et al. *Estrogens Exacerbate Nociceptive Pain via Up-Regulation of TRPV1 and ANO1 in Trigeminal Primary Neurons of Female Rats*. Endocrinology. 2016. 157(11): p. 4309-4317. 10.1210/en.2016-1218

[84] Wu, Y.W., et al. *17-Beta-estradiol enhanced allodynia of inflammatory temporomandibular joint through upregulation of hippocampal TRPV1 in ovariectomized rats*. J Neurosci. 2010. 30(26): p. 8710-9. 10.1523/JNEUROSCI.6323-09.2010

[85] Chen, F., et al. *Estrogen inhibits RANKL-induced osteoclastic differentiation by increasing the expression of TRPV5 channel*. J Cell Biochem. 2014. 115(4): p. 651-8. 10.1002/jcb.24700

[86] Pohoczky, K., et al. *Estrogen-dependent up-regulation of TRPA1 and TRPV1 receptor proteins in the rat endometrium*. J Mol Endocrinol. 2016. 56(2): p. 135-49. 10.1530/JME-15-0184

[87] Wu, Y.W., et al. *Synovial TRPV1 is upregulated by 17-beta-estradiol and involved in allodynia of inflamed temporomandibular joints in female rats*. Arch Oral Biol. 2015. 60(9): p. 1310-8. 10.1016/j.archoralbio.2015.05.011

- [88] Greaves, E., et al. *Elevated peritoneal expression and estrogen regulation of nociceptive ion channels in endometriosis*. J Clin Endocrinol Metab. 2014. 99(9): p. E1738-43. 10.1210/jc.2014-2282
- [89] Kumar, S., et al. *Transient receptor potential vanilloid 1-6 (Trpv1-6) gene expression in the mouse brain during estrous cycle*. Brain Res. 2018. 1701: p. 161-170. 10.1016/j.brainres.2018.09.005
- [90] Yazgan, Y. and M. Naziroglu. *Ovariectomy-Induced Mitochondrial Oxidative Stress, Apoptosis, and Calcium Ion Influx Through TRPA1, TRPM2, and TRPV1 Are Prevented by 17beta-Estradiol, Tamoxifen, and Raloxifene in the Hippocampus and Dorsal Root Ganglion of Rats*. Mol Neurobiol. 2017. 54(10): p. 7620-7638. 10.1007/s12035-016-0232-5
- [91] Xu, S., et al. *17beta-estradiol activates estrogen receptor beta-signalling and inhibits transient receptor potential vanilloid receptor 1 activation by capsaicin in adult rat nociceptor neurons*. Endocrinology. 2008. 149(11): p. 5540-8. 10.1210/en.2008-0278
- [92] Chen, S.C., T.J. Chang, and F.S. Wu. *Competitive inhibition of the capsaicin receptor-mediated current by dehydroepiandrosterone in rat dorsal root ganglion neurons*. J Pharmacol Exp Ther. 2004. 311(2): p. 529-36. 10.1124/jpet.104.069096
- [93] Ramirez-Barrantes, R., et al. *TRPV1-Estradiol Stereospecific Relationship Underlies Cell Survival in Oxidative Cell Death*. Front Physiol. 2020. 11: p. 444. 10.3389/fphys.2020.00444
- [94] Chen, S.C. and F.S. Wu. *Mechanism underlying inhibition of the capsaicin receptor-mediated current by pregnenolone sulfate in rat dorsal root ganglion neurons*. Brain Res. 2004. 1027(1-2): p. 196-200. 10.1016/j.brainres.2004.08.053
- [95] DeLeon, C., D.Q.-H. Wang, and C.K. Arnatt. *G Protein-Coupled Estrogen Receptor, GPER1, Offers a Novel Target for the Treatment of Digestive Diseases*. 2020. 11(843). 10.3389/fendo.2020.578536
- [96] Lopshire, J.C. and G.D. Nicol. *The cAMP transduction cascade mediates the prostaglandin E2 enhancement of the capsaicin-elicited current in rat sensory neurons: whole-cell and single-channel studies*. J Neurosci. 1998. 18(16): p. 6081-92.
- [97] Joseph, J., et al. *Phosphorylation of TRPV1 S801 Contributes to Modality-Specific Hyperalgesia in Mice*. J Neurosci. 2019. 39(50): p. 9954-9966. 10.1523/JNEUROSCI.1064-19.2019
- [98] Goswami, C., T.B. Hucho, and F. Hucho. *Identification and characterisation of novel tubulin-binding motifs located within the C-terminus of TRPV1*. J Neurochem. 2007. 101(1): p. 250-62. 10.1111/j.1471-4159.2006.04338.x
- [99] Goswami, C., et al. *Estrogen destabilizes microtubules through an ion-conductivity-independent TRPV1 pathway*. J Neurochem. 2011. 117(6): p. 995-1008. 10.1111/j.1471-4159.2011.07270.x
- [100] Prager-Khoutorsky, M., A. Khoutorsky, and C.W. Bourque. *Unique interweaved microtubule scaffold mediates osmosensory transduction via physical interaction with TRPV1*. Neuron. 2014. 83(4): p. 866-78. 10.1016/j.neuron.2014.07.023
- [101] Liu, J., et al. *The expression and functionality of transient receptor potential vanilloid 1 in ovarian endometriomas*. Reprod Sci. 2012. 19(10): p. 1110-24. 10.1177/1933719112443876
- [102] McKinnon, B.D., et al. *Inflammation and nerve fiber interaction in endometriotic pain*. Trends Endocrinol Metab. 2015. 26(1): p. 1-10. 10.1016/j.tem.2014.10.003
- [103] Bohonyi, N., et al. *Local upregulation of transient receptor*

potential ankyrin 1 and transient receptor potential vanilloid 1 ion channels in rectosigmoid deep infiltrating endometriosis. Mol Pain. 2017. 13: p. 1744806917705564. 10.1177/1744806917705564

[104] Ma, W., et al. *Catechol estrogens stimulate insulin secretion in pancreatic beta-cells via activation of the transient receptor potential A1 (TRPA1) channel.* J Biol Chem. 2019. 294(8): p. 2935-2946. 10.1074/jbc.RA118.005504

[105] Hevir, N., M. Ribic-Pucelj, and T. Lanisnik Rizner. *Disturbed balance between phase I and II metabolizing enzymes in ovarian endometriosis: a source of excessive hydroxy-estrogens and ROS?* Mol Cell Endocrinol. 2013. 367(1-2): p. 74-84. 10.1016/j.mce.2012.12.019

[106] Fattori, V., et al. *Nonsurgical mouse model of endometriosis-associated pain that responds to clinically active drugs.* Pain. 2020. 161(6): p. 1321-1331. 10.1097/j.pain.0000000000001832

[107] Mooradian, A.D., J.E. Morley, and S.G. Korenman. *Biological actions of androgens.* Endocr Rev. 1987. 8(1): p. 1-28. 10.1210/edrv-8-1-1

[108] Schiffer, L., W. Arlt, and K.H. Storbeck. *Intracrine androgen biosynthesis, metabolism and action revisited.* Mol Cell Endocrinol. 2018. 465: p. 4-26. 10.1016/j.mce.2017.08.016

[109] Bennett, N.C., et al. *Molecular cell biology of androgen receptor signalling.* Int J Biochem Cell Biol. 2010. 42(6): p. 813-27. 10.1016/j.biocel.2009.11.013

[110] Chen, F., et al. *Partial agonist/antagonist properties of androstenedione and 4-androsten-3beta,17beta-diol.* J Steroid Biochem Mol Biol. 2004. 91(4-5): p. 247-57. 10.1016/j.jsbmb.2004.04.009

[111] Bai, X., X. Zhang, and Q. Zhou. *Effect of Testosterone on TRPV1*

Expression in a Model of Orofacial Myositis Pain in the Rat. J Mol Neurosci. 2018. 64(1): p. 93-101. 10.1007/s12031-017-1009-7

[112] Fusi, C., et al. *Steroid and non-steroidal third-generation aromatase inhibitors induce pain-like symptoms via TRPA1.* Nat Commun. 2014. 5: p. 5736. 10.1038/ncomms6736

[113] De Logu, F., et al. *TRPA1 Mediates Aromatase Inhibitor-Evoked Pain by the Aromatase Substrate Androstenedione.* Cancer Res. 2016. 76(23): p. 7024-7035. 10.1158/0008-5472.CAN-16-1492

[114] Bidaux, G., et al. *Evidence for specific TRPM8 expression in human prostate secretory epithelial cells: functional androgen receptor requirement.* Endocr Relat Cancer. 2005. 12(2): p. 367-82. 10.1677/erc.1.00969

[115] Zhang, L. and G.J. Barritt. *Evidence that TRPM8 is an androgen-dependent Ca²⁺ channel required for the survival of prostate cancer cells.* Cancer Res. 2004. 64(22): p. 8365-73. 10.1158/0008-5472.CAN-04-2146

[116] Yang, Z., et al. *Effect of surgical castration on expression of TRPM8 in urogenital tract of male rats.* Mol Biol Rep. 2012. 39(4): p. 4797-802. 10.1007/s11033-011-1271-7

[117] Asuthkar, S., et al. *TRPM8 channel as a novel molecular target in androgen-regulated prostate cancer cells.* Oncotarget. 2015. 6(19): p. 17221-36. 10.18632/oncotarget.3948

[118] Asuthkar, S., et al. *The TRPM8 protein is a testosterone receptor: I. Biochemical evidence for direct TRPM8-testosterone interactions.* J Biol Chem. 2015. 290(5): p. 2659-69. 10.1074/jbc.M114.610824

[119] Mendez-Resendiz, K.A., et al. *Steroids and TRP Channels: A Close Relationship.* Int J Mol Sci. 2020. 21(11). 10.3390/ijms21113819

- [120] Gkika, D., et al. *Testosterone-androgen receptor: The steroid link inhibiting TRPM8-mediated cold sensitivity*. FASEB J. 2020. 10.1096/fj.201902270R
- [121] Taraborrelli, S. *Physiology, production and action of progesterone*. Acta Obstet Gynecol Scand. 2015. 94 Suppl 161: p. 8-16. 10.1111/aogs.12771
- [122] Goletiani, N.V., D.R. Keith, and S.J. Gorsky. *Progesterone: review of safety for clinical studies*. Exp Clin Psychopharmacol. 2007. 15(5): p. 427-44. 10.1037/1064-1297.15.5.427
- [123] Garg, D., et al. *Progesterone-Mediated Non-Classical Signaling*. Trends Endocrinol Metab. 2017. 28(9): p. 656-668. 10.1016/j.tem.2017.05.006
- [124] Maurice, T., C. Gregoire, and J. Espallergues. *Neuro(active)steroids actions at the neuromodulatory sigma1 (sigma1) receptor: biochemical and physiological evidences, consequences in neuroprotection*. Pharmacol Biochem Behav. 2006. 84(4): p. 581-97. 10.1016/j.pbb.2006.07.009
- [125] Jung, C., et al. *The progesterone receptor regulates the expression of TRPV4 channel*. Pflugers Arch. 2009. 459(1): p. 105-13. 10.1007/s00424-009-0706-7
- [126] De Clercq, K., et al. *Functional expression of transient receptor potential channels in human endometrial stromal cells during the luteal phase of the menstrual cycle*. Hum Reprod. 2015. 30(6): p. 1421-36. 10.1093/humrep/dev068
- [127] Li, C., et al. *TRPV4 is involved in levonorgestrel-induced reduction in oviduct ciliary beating*. J Pathol. 2019. 248(1): p. 77-87. 10.1002/path.5233
- [128] Ortiz-Renteria, M., et al. *TRPV1 channels and the progesterone receptor Sig-1R interact to regulate pain*. Proc Natl Acad Sci U S A. 2018. 115(7): p. E1657-E1666. 10.1073/pnas.1715972115
- [129] Hayashi, T. and T.P. Su. *Sigma-1 receptor chaperones at the ER-mitochondrion interface regulate Ca(2+) signaling and cell survival*. Cell. 2007. 131(3): p. 596-610. 10.1016/j.cell.2007.08.036
- [130] Cortes-Montero, E., et al. *Ligands Exert Biased Activity to Regulate Sigma 1 Receptor Interactions With Cationic TRPA1, TRPV1, and TRPM8 Channels*. Front Pharmacol. 2019. 10: p. 634. 10.3389/fphar.2019.00634