

1
2
3 **Fragmented Management of Long-Term Parenteral Support for Adult**
4
5
6 **Intestinal Failure in Finland**
7

8 Anne Pohju¹, Mikko P. Pakarinen², Taina Sipponen³
9

10
11 *¹Clinical Nutrition Unit, Internal Medicine and Rehabilitation, University of Helsinki and*
12
13 *Helsinki University Hospital, Helsinki, Finland, ²Section of Pediatric Surgery, Pediatric Liver*
14
15 *and Gut Research Group, Children's Hospital, Pediatric Research Center, University of*
16
17 *Helsinki and Helsinki University Hospital, Helsinki, Finland, ³Gastroenterology, Abdominal*
18
19 *Center, University of Helsinki and Helsinki University Hospital, Helsinki, Finland*
20
21
22
23
24

25 **Corresponding author**

26 Anne Pohju

27 Clinical Nutrition Unit

28 Helsinki University Hospital

29 P.O. Box 340

30 FI-00029 HUS

31 FINLAND

32 Email: anne.pohju@helsinki.fi
33
34
35
36
37
38
39
40
41
42
43
44
45

46 **Word count**

47 3678
48
49

50 **Statement of Authorship**

51
52 AP, MP, and TS devised the study protocol, collected the data, analysed the results, drafted the
53
54 manuscript, and approved the final version for submission.
55
56

57 **Funding details**
58
59
60

1
2
3 This research did not receive any specific grant from funding agencies in the public,
4 commercial, or non-profit sectors.
5
6

7 **Fragmented Management of Long-Term Parenteral Support for Adult** 8 **Intestinal Failure in Finland** 9 10 11

12
13
14
15 **Objectives:** Parenteral support (PS) is the first-line therapy for intestinal failure (IF). Optimal patient
16 outcomes require experienced multidisciplinary teams adhering to structured protocols. As practices to
17 provide long-term PS for adult IF patients in Finland are unknown, this cross-sectional nationwide study
18 aimed to evaluate current management of PS for adult IF across the country.
19
20
21
22

23
24 **Materials and Methods:** An internet-based survey was emailed to all Finnish hospitals and hospital-
25 at-home services with the potential to provide PS for adult IF. The survey included 20 items addressing
26 provision of long-term PS for adult IF patients (aged ≥ 18 years). Data were analysed using descriptive
27 statistics.
28
29
30
31

32
33 **Results:** Overall, 52 (47%) of the 111 identified units responded. Of responding units, 38 (73%) had at
34 some point provided long-term (≥ 120 days) PS for adult IF, and 23 (44%) had done so during the
35 preceding year. Only three units currently managed ≥ 3 adult patients. Most (65%) of the respondents
36 worked in a hospital and were either physicians (38%) or dietitians (39%). Only 65% of respondents
37 reported that their unit had an assigned physician responsible for PS provision, and 28% reported that a
38 team was responsible for long-term PS. Only 26% of respondents reported having a written protocol to
39 guide PS management.
40
41
42
43
44
45

46
47 **Conclusions:** Health care providers with very limited experience and a fragmented approach manage
48 most Finnish adult IF patients. Evidence-based protocols and multidisciplinary teams are scarce. The
49 care for adult IF patients on long-term PS needs to be improved in Finland.
50
51
52
53

54
55 **Keywords:** parenteral support; intravenous supplementation; home parenteral nutrition;
56 intestinal failure; short bowel syndrome
57
58
59
60

Introduction

Intestinal failure (IF) is a rare disease defined as a reduction in gut function leading to an inability to maintain health unless parenteral support (PS) is provided [1]. In addition to long-term PS consisting of nutrition and/or fluid and electrolytes, patients require individually tailored medical and nutritional treatment and, in many cases, surgery [2]. To achieve optimal outcomes, specialized teams and structured protocols in managing IF seem to have a key role [3, 4, 5, 6].

Comprehensive reviews of the management of short bowel syndrome, the most common cause of IF, can be found in the literature [7, 8, 9]. More recently, an extensive guideline covering the whole spectrum of IF in adult patients has been published [2]. Guidelines are well adhered to and management strategies for nutritional care in paediatric patients are relatively consistent among specialized centres [10, 11]. The introduction of multidisciplinary intestinal rehabilitation programmes has resulted in improved clinical outcomes of IF, especially in paediatric patient populations [3, 4, 5, 6]. Intestinal rehabilitation programmes require experienced staff before having an impact on outcomes such as patient survival [4, 6, 12].

The need for long-term PS, commonly referred to as home parenteral nutrition (HPN), is often used as a synonym for IF. However, these two terms, IF and HPN, do not necessarily refer to the same patient population. Long-term PS can also be used in some less well-defined conditions, where the gut function is not truly compromised [13], e.g. anorexia in cancer cachexia.

To the best of our knowledge, the clinical practices for long-term PS have not previously been investigated among adult IF patients in Finland. The aim of this cross-sectional study was to evaluate current management of long-term PS for adult IF patients across the country.

Materials and Methods

Study Protocol

An email with a cover letter and a link to an electronic survey was sent to chief physicians of the departments of gastroenterology, gastrointestinal surgery, anaesthesiology, and internal medicine, to clinical dietitians working in all Finnish hospitals, and to chief physicians or head nurses of all public and private hospital-at-home services across Finland. A reminder was sent two weeks later.

Data Collection

The electronic survey included 19 questions regarding the local practices of long-term PS. These were structured multiple-choice questions, but the respondents were able to write free text in comment boxes if they wanted to expand their answers. In addition, the participants were asked whether their unit had provided PS to any patient(s) aged ≥ 18 years for at least 120 consecutive days during the preceding 12 months.

Statistical Analysis

The descriptive data are presented as frequencies and percentages. The IBM SPSS software package (version 24) was used for the analyses.

Ethics Statement

The study was approved by the Medical Research Ethics Committee of Helsinki University Hospital (HUS/751/2017).

Results

Participation Rate and Respondents

An electronic survey was emailed to 306 health care professionals working in 111 separate units identified across Finland. In total, 71 individuals representing 52 different health care providers responded, resulting in an overall response rate of 47% for the units (Table 1). The majority of respondents worked in a hospital and were either physicians or dietitians (Table 2). Responses were obtained from all Finnish hospital districts.

Experience in Patients with Long-Term PS in Finnish Health Care Units

Altogether 28 survey respondents (39%) reported that, to their knowledge, their unit had never treated an adult patient receiving PS for 120 days. These respondents were not asked questions concerning long-term PS. This left 43 participants from 38 units to answer the survey questions pertaining to management of long-term PS in their unit. Twenty-three different units were treating or had treated at least one patient with long-term PS during the preceding year. Only three units (13%) were currently treating three or more adult patients. Therefore, the great majority of the responding units managed currently only one or two patients with long-term PS.

Management of Long-Term PS

The majority of the respondents (65%) reported their patients to have an assigned physician responsible for provision of PS (Table 3). However, it was uncommon to have an assigned nurse or a team responsible for the management of patients on long-term PS (Table 3). Most of the respondents who reported having an assigned management team were working in a hospital-at-home service (8 out of 12 respondents). The remaining four respondents worked in a university hospital.

1
2
3 The most frequently mentioned members of the team were a nurse (10 responses), a
4 physician (usually a specialist in general practice) working in the hospital-at-home service (6
5 responses), a gastroenterologist or specialist in internal medicine (5 responses), and a dietitian
6 (4 responses). No unit had a psychologist or a social worker as an assigned member of the team.
7
8
9

10
11
12 Every third respondent reported that PS is not based on any recommendation or
13 guideline in their unit, or the respondent did not know whether it is (Table 3). When a guideline
14 was used, it was a national or international guideline in 37% and a local guideline in 63% of
15 the units. Most of the respondents reported not having a written protocol on the management
16 of IVS in practice (Table 3).
17
18
19
20
21
22

23
24 Nine respondents reported that they have or have had patients who were compelled to
25 stay admitted in hospital because practical issues prevented the administration of PS at home,
26 even though the health status of the patient would otherwise have allowed the patient to be
27 discharged.
28
29
30
31

32
33 Respondents working in a hospital-at-home service frequently reported carrying out a
34 PS plan created by another unit (Table 4). Otherwise, no uniform pattern was found in the
35 planning of PS. In their comments, the respondents often described that several specialists
36 devise the plan together and/or with the help of a consulting specialist.
37
38
39
40
41

42
43 Less than half of the respondents reported teaching the patient how to administer
44 parenteral infusions independently (Table 3). If the administration of infusions was taught,
45 typically a nurse was responsible for the instruction (18 responses out of 19). Nurses were most
46 frequently (33 responses) responsible for supplementing (e.g. additional vitamins or
47 electrolytes) the PS bags. Only six respondents reported that patients or their relatives did this
48 themselves.
49
50
51
52
53
54

55
56 Great variation was observed in responses to questions about who is responsible for the
57 practical administration of PS and how patients are supplied the PS products or the equipment
58
59
60

1
2
3 for the administration (Tables 5 and 6). Hospital-at-home service was most frequently
4 mentioned to deal with the practical administration of PS as well as to supply the patient with
5
6
7
8
9 parenteral products and equipment.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Only 36% of the respondents reported routine use of infusion pumps for the
administration of PS (Table 7). Approximately half of the respondents stated that infusion pump
and infusion stand were possible to arrange for the patient (Table 6). Only 12 respondents
reported the possibility to provide the patient with a portable infusion pump.

Discussion

Our results suggest that organized multidisciplinary team work and treatment protocols are rare
in the context of adult IF and long-term PS in Finnish health care units. Most of the units
manage only one or two adults with long-term PS. This study is the first step towards enhancing
our understanding of the management of IF in the Finnish adult population.

The prevalence of IF seems to be increasing in the Western countries [14, 15, 16], and
it is likely that a similar development will also occur in Finland. The underlying reasons for the
increase may include e.g. the good overall survival rates [17, 18]. Our findings are helpful in
organizing management of adult IF in Finnish health care units. The urgent need for
standardized management protocols and implementation of evidence-based guidelines as well
as multidisciplinary care was clearly revealed in this survey.

Management of paediatric IF in Finland has been centralized for many years, enabling
continuous development of the programme, and consequently, high-quality outcomes [5, 19].
Furthermore, a Finnish national programme for intestinal transplantation (ITx) has been
ongoing since 2009. Therefore, it is surprising to note that, unlike in other benign chronic

1
2
3 diseases, such as renal failure, management of IF in adults lacks nationwide organization in
4
5 Finland.

6
7
8 In the Nordic countries, Denmark has the longest tradition in managing IF patients [20],
9
10 and the prevalence of IF in Denmark is among the highest in Europe [14]. The first Danish
11
12 patient started HPN already in 1970, and nowadays, the largest Danish referral IF centre has
13
14 over 150 patients in their well-structured HPN programme [21]. Attempts of nationwide
15
16 organization of adult IF care have also taken place in other Nordic countries. Sahlgrenska
17
18 University Hospital in Gothenburg, Sweden, provides surgical treatment of IF, and the unit has
19
20 experience with ITx since 1998 [22]. An initiative to launch a national registry for short bowel
21
22 syndrome in Sweden was described already in 2004 [23]. In Norway, the Oslo University
23
24 Hospital serves as a national referral centre for Norwegian IF patients who are candidates for
25
26 ITx [24].
27
28
29

30
31 Long-term PS can be provided effectively and safely outside the designated IF units [25,
32
33 26]. This will become increasingly necessary as the volume of patients grows. However, as
34
35 recent evidence shows [3, 4, 5, 6], optimal patient outcomes in IF cannot be expected in the
36
37 absence of the required experience and skills of staff and a standardized protocol.
38
39

40
41 To fulfil the above-mentioned requirements, the management of adult IF needs to be
42
43 concentrated in Finland. The concept of centralized care of rare diseases with complex
44
45 treatment is strongly supported by European and Finnish health care authorities. Active co-
46
47 operation between the national referral centre and smaller units across the country is essential.
48
49 Telehealth services might be one means of facilitating this co-operation. Telehealth enables
50
51 regular face-to-face contacts with experts of the specialist IF centre without the burden and cost
52
53 of long-distance travelling [27], which are real concerns for patients living in the sparsely
54
55 populated northern part of Finland.
56
57
58
59
60

1
2
3 Apart from the care given by health care professionals and the technical solutions, peer
4 support and education by patient organizations are also important factors affecting the well-
5 being of patients on long-term PS. A previous case-control study among North American
6 patients receiving HPN suggested that a connection with a national patient organization has a
7 positive effect on clinically important patient outcomes such as quality of life, depression
8 scores, and incidence of catheter-related infections [28]. In Finland, a patient organization for
9 IF is lacking, representing another approach to improve overall outcomes.

10
11
12 Even though this study was limited by the relatively low overall response rate, the
13 geographical coverage of the survey was comprehensive. In addition, over 80% of the
14 university and central hospitals, where the majority of the patients are treated, responded. Less
15 than two-thirds of the respondents had any experience with patients with long-term PS. We
16 speculate that smaller units (i.e. regional hospitals and hospital-at-home services from smaller
17 towns) without any experience with adult IF were less likely to respond. Therefore, our results
18 should reflect clinical reality with reasonable accuracy. Most survey questions covered the
19 practices in the respondent's unit. Individual respondents might have interpreted the concept of
20 a unit in different ways (e.g. as the whole hospital or as a specific unit within the hospital).
21 Therefore, we decided to report the results from single respondents as opposed to units. In
22 addition, the results showed that even professionals working in the same hospital could give
23 different answers. This can be regarded as a consequence of the lack of uniform practices, which
24 indeed was one of the main results of this study.

25
26
27 We are currently in the process of investigating the prevalence of Finnish adult IF, the
28 characteristics of the affected population, and the details of their care (**e.g. incidence of**
29 **catheter-related bloodstream infections**), which were not objectives of this study. On a
30 national level, there is an urgent need for improved care of Finnish adult IF patients requiring
31 long-term PS. The first steps towards this should include devising a national recommendation

1
2
3 on management of adult IF in Finnish language, creating a network of units managing this
4
5 patient group and coordination of care by a national referral centre.
6
7
8
9

10 **Acknowledgements**

11
12 We thank all contributing physicians, dietitians, and nurses for taking the time to answer this
13
14 survey.
15
16

17 **Disclosure of Interest**

18
19 The authors report no conflict of interest.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Pironi L, Arends J, Baxter J, et al. ESPEN endorsed recommendations. Definition and classification of intestinal failure in adults. *Clin Nutr.* 2015 Apr;34(2):171-80. doi: 10.1016/j.clnu.2014.08.017. PubMed PMID: 25311444.
2. Pironi L, Arends J, Bozzetti F, et al. ESPEN guidelines on chronic intestinal failure in adults. *Clin Nutr.* 2016 Apr;35(2):247-307. doi: 10.1016/j.clnu.2016.01.020. PubMed PMID: 26944585.
3. Stanger JD, Oliveira C, Blackmore C, et al. The impact of multi-disciplinary intestinal rehabilitation programs on the outcome of pediatric patients with intestinal failure: a systematic review and meta-analysis. *J Pediatr Surg.* 2013 May;48(5):983-92. doi: 10.1016/j.jpedsurg.2013.02.070. PubMed PMID: 23701771.
4. Oliveira C, de Silva NT, Stanojevic S, et al. Change of Outcomes in Pediatric Intestinal Failure: Use of Time-Series Analysis to Assess the Evolution of an Intestinal Rehabilitation Program. *J Am Coll Surg.* 2016 Jun;222(6):1180-1188 e3. doi: 10.1016/j.jamcollsurg.2016.03.007. PubMed PMID: 27067452.
5. Merras-Salmio L, Pakarinen MP. Refined Multidisciplinary Protocol-Based Approach to Short Bowel Syndrome Improves Outcomes. *J Pediatr Gastroenterol Nutr.* 2015 Jul;61(1):24-9. doi: 10.1097/MPG.0000000000000775. PubMed PMID: 25729889.
6. Avitzur Y, Wang JY, de Silva NT, et al. Impact of Intestinal Rehabilitation Program and Its Innovative Therapies on the Outcome of Intestinal Transplant Candidates. *J Pediatr Gastroenterol Nutr.* 2015 Jul;61(1):18-23. doi: 10.1097/MPG.0000000000000735. PubMed PMID: 25611029.

- 1
2
3 7. DiBaise JK, Young RJ, Vanderhoof JA. Intestinal rehabilitation and the short
4
5 bowel syndrome: part 2. *Am J Gastroenterol*. 2004 Sep;99(9):1823-32. doi: 10.1111/j.1572-
6
7 0241.2004.40836.x. PubMed PMID: 15330926.
- 8
9
10 8. DiBaise JK, Young RJ, Vanderhoof JA. Intestinal rehabilitation and the short
11
12 bowel syndrome: part 1. *Am J Gastroenterol*. 2004 Jul;99(7):1386-95. doi: 10.1111/j.1572-
13
14 0241.2004.30345.x. PubMed PMID: 15233682.
- 15
16
17 9. Nightingale J, Woodward JM, Small B, et al. Guidelines for management of
18
19 patients with a short bowel. *Gut*. 2006 Aug;55 Suppl 4:iv1-12. doi: 10.1136/gut.2006.091108.
20
21 PubMed PMID: 16837533; PubMed Central PMCID: PMCPMC2806687.
- 22
23
24 10. Neelis E, de Koning B, van Winckel M, et al. Wide variation in organisation and
25
26 clinical practice of paediatric intestinal failure teams: an international survey. *Clin Nutr*. 2017
27
28 Nov 22. doi: 10.1016/j.clnu.2017.11.008. PubMed PMID: 29203343.
- 29
30
31 11. Nucci AM, Ellsworth K, Michalski A, et al. Survey of Nutrition Management
32
33 Practices in Centers for Pediatric Intestinal Rehabilitation. *Nutr Clin Pract*. 2018
34
35 Aug;33(4):528-538. doi: 10.1177/0884533617719670. PubMed PMID: 28731841.
- 36
37
38 12. Messing B, Lemann M, Landais P, et al. Prognosis of patients with
39
40 nonmalignant chronic intestinal failure receiving long-term home parenteral nutrition.
41
42 *Gastroenterology*. 1995 Apr;108(4):1005-10. PubMed PMID: 7698566.
- 43
44
45 13. Howard L, Ament M, Fleming CR, et al. Current use and clinical outcome of
46
47 home parenteral and enteral nutrition therapies in the United States. *Gastroenterology*. 1995
48
49 Aug;109(2):355-65. PubMed PMID: 7615183.
- 50
51
52 14. Baxter JP, Gillanders L, Angstmann K, et al. Home parenteral nutrition: An
53
54 international benchmarking exercise. *e-SPEN Journal*. 2012;7(5):e211-e214. doi:
55
56 10.1016/j.clnme.2012.08.001.
- 57
58
59
60

- 1
2
3 15. Bakker H, Bozzetti F, Staun M, et al. Home parenteral nutrition in adults: a
4
5 european multicentre survey in 1997. ESPEN-Home Artificial Nutrition Working Group. Clin
6
7 Nutr. 1999 Jun;18(3):135-40. doi: 10.1054/clnu.1999.0021. PubMed PMID: 10451476; eng.
8
9
10 16. Van Gossum A, Bakker H, De Francesco A, et al. Home parenteral nutrition in
11
12 adults: a multicentre survey in Europe in 1993. Clin Nutr. 1996 Apr;15(2):53-9. PubMed
13
14 PMID: 16843998; eng.
15
16
17 17. Pironi L, Goulet O, Buchman A, et al. Outcome on home parenteral nutrition for
18
19 benign intestinal failure: a review of the literature and benchmarking with the European
20
21 prospective survey of ESPEN. Clin Nutr. 2012 Dec;31(6):831-45. doi:
22
23 10.1016/j.clnu.2012.05.004. PubMed PMID: 22658443.
24
25
26 18. Joly F, Baxter J, Staun M, et al. Five-year survival and causes of death in
27
28 patients on home parenteral nutrition for severe chronic and benign intestinal failure. Clin
29
30 Nutr. 2018 Aug;37(4):1415-1422. doi: 10.1016/j.clnu.2017.06.016. PubMed PMID:
31
32 28701261.
33
34
35 19. Merras-Salmio L, Mutanen A, Ylinen E, et al. Pediatric Intestinal Failure: The
36
37 Key Outcomes for the First 100 Patients Treated in a National Tertiary Referral Center
38
39 During 1984-2017. JPEN J Parenter Enteral Nutr. 2018 Apr 27. doi: 10.1002/jpen.1164.
40
41 PubMed PMID: 29701871.
42
43
44 20. Ugur A, Marashdeh BH, Gottschalck I, et al. Home parenteral nutrition in
45
46 Denmark in the period from 1996 to 2001. Scand J Gastroenterol. 2006 Apr;41(4):401-7. doi:
47
48 10.1080/00365520500441247. PubMed PMID: 16635907.
49
50
51 21. Brandt CF, Hvistendahl M, Naimi RM, et al. Home Parenteral Nutrition in
52
53 Adult Patients With Chronic Intestinal Failure: The Evolution Over 4 Decades in a Tertiary
54
55 Referral Center. JPEN J Parenter Enteral Nutr. 2017 Sep;41(7):1178-1187. doi:
56
57 10.1177/0148607116655449. PubMed PMID: 27323776.
58
59
60

- 1
2
3 22. Gäbel M, Varkey J, Saalman R, et al. Two decades of Experience with Intestinal
4 Transplantation in Gothenburg, Sweden. *Transplantation*. 2017;101(6S2):S55. doi:
5
6 10.1097/01.tp.0000521358.10910.7d. PubMed PMID: 00007890-201706002-00087.
7
8
9
10 23. Kurlberg G, Forssell H, Aly A. National registry of patients with short bowel
11 syndrome. *Transplant Proc*. 2004 Mar;36(2):253-4. doi: 10.1016/j.transproceed.2003.12.021.
12
13 PubMed PMID: 15050125.
14
15
16
17 24. Bental OH, Foss A, Ostensen AB, et al. Intestinal and multivisceral
18 transplantation in patients with chronic intestinal failure. *Tidsskr Nor Laegeforen*. 2012 Mar
19
20 27;132(6):650-4. doi: 10.4045/tidsskr.11.0817. PubMed PMID: 22456144.
21
22
23
24 25. Green CJ, Mountford V, Hamilton H, et al. A 15-year audit of home parenteral
25 nutrition provision at the John Radcliffe Hospital, Oxford. *QJM*. 2008 May;101(5):365-9. doi:
26
27 10.1093/qjmed/hcn011. PubMed PMID: 18292099.
28
29
30
31 26. Freshwater DA, Saadeddin A, Deel-Smith P, et al. Can home parenteral
32 nutrition be provided by non-specialised centres? 2300 weeks of experience at a district
33 general hospital in the United Kingdom. *Clin Nutr*. 2005 Apr;24(2):229-35. doi:
34
35 10.1016/j.clnu.2004.09.015. PubMed PMID: 15784483.
36
37
38
39
40 27. Saqui O, Chang A, McGonigle S, et al. Telehealth videoconferencing:
41 improving home parenteral nutrition patient care to rural areas of Ontario, Canada. *JPEN J*
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure and Table Legends

Table 1. Response rate of the identified units according to the type of health care provider.

	Invited, <i>n</i>	Participated, <i>n</i> (%)
University hospital	11	10 (91)
Central hospital	15	11 (73)
Regional hospital	15	7 (47)
Hospital-at-home service	69	23 (33)
Other	1	1 (100)
Total	111	52 (47)

Table 2. Demographic data on respondents ($n = 71$).

		<i>n</i> (%)
Profession	Physician	27 (38)
	Dietitian	28 (39)
	Nurse	16 (23)
Place of employment	Hospital	46 (65)
	Hospital-at-home service	24 (34)
	Other	1 (1)
Type of hospital for those working in hospital ($n = 46$)	University hospital	18 (39)
	Central hospital	22 (48)
	Regional hospital	6 (13)

Table 3. Management of long-term parenteral support (PS) ($n = 43$).

	Yes	No	I don't know	Missing
The patient has an assigned physician in the unit responsible for PS	28 (65%)	12 (28%)	3 (7%)	-
The patient has an assigned nurse in the unit responsible for PS	7 (16%)	29 (68%)	7 (16%)	-
In the unit, an assigned team is responsible for the management of patients on PS	12 (28%)	30 (70%)	1 (2%)	-
The patient is taught how to independently manage his/her PS by the staff in the unit	19 (44%)	18 (42%)	6 (14%)	-
The PS provided by the unit is based on a recommendation or guideline	24 (63%)	6 (16%)	8 (21%)	5
The unit has a written protocol and/or guideline on the management of PS in practice	10 (26%)	25 (64%)	4 (10%)	4

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 4. Person(s) in charge for the plan of parenteral support (multiple choices possible).

	Number of responses	
	Hospital	Hospital-at-home service
Anaesthesiologist	6	0
Specialist in internal medicine	1	0
Gastroenterologist	1	0
Surgeon	1	0
Dietitian	8	0
The unit carries out a plan devised in another unit	0	12
I don't know	2	0
Someone else	6	6

Table 5. Unit responsible for administration of parenteral support in practice (multiple choices possible).

	Number of responses
Hospital-at-home service	36
Hospital	15
Health centre	6
Patient him-/herself	6
I don't know	1
Some other unit	3

Table 6. Supply of parenteral products and equipment needed for parenteral support (PS; multiple choices possible).

		Number of responses
The patient gets his/her parenteral products from	Hospital-at-home service	32
	Hospital	12
	Health centre	8
	Pharmacy	5
	I don't know	5
	Somewhere else	3
The patient gets PS equipment from	Hospital-at-home service	29
	Hospital	8
	Health centre	7
	Public health care distribution of self-treatment products	8
	I don't know	6
	Somewhere else	0
Our unit provides the following equipment for PS	Infusion pump	21
	Infusion stand	27
	Portable infusion pump	12
	None of these	4

Table 7. Use of an infusion pump for parenteral nutrition ($n = 43$).

	<i>n</i> (%)
Always	14 (36)
Often	5 (13)
Seldom	7 (18)
Never	8 (20)
I don't know	5 (13)
Missing	4