

## Ethnobotanique

### Ethnobotanical survey of medicinal plants used in the traditional treatment of depression and anxiety in Fez-Meknes region, Morocco

A. Amaghnouje<sup>1</sup>, M. Slighoua<sup>1</sup>, I. Es-safi<sup>1</sup>, R. El Jaoudi<sup>2</sup>, A. Elyoubi<sup>1</sup>, D. Bousta<sup>1</sup>, Andriy Grafov<sup>3</sup>.

<sup>1</sup> Laboratory of Neuroendocrinology and Nutritional and climatic Environment, Faculty Of Sciences, University Sidi Mohammed Ben Abdellah, BP 1796-ATLAS, FES, Morocco.

<sup>2</sup> Laboratory of Pharmacology and Toxicology, faculty of medicine and pharmacy, Rabat. Morocco.

<sup>3</sup>Laboratory of Inorganic Chemistry, University of Helsinki, FI

Email : amalsvi17@gmail.com

#### Abstract

Depression and anxiety represent a major mental health problem in the world. The majority of Moroccan people use traditional medicine for their health needs, including various forms of depression and anxiety. The aim of this work is to make an inventory of plant species used in folk medicine for the management of depression and anxiety in Fez-Meknes regions.

Established questionnaires were administered to 243 interviews in 4 communities within Fez-Meknes region. With traditional health practitioners, herbalists and consumer. Plant species belonging to 31 families were reported. The most frequently cited families are Lamiaceae followed by Asteraceae, Apiaceae and Verbenaceae. Leaves were the major plant parts used forming 58% solely and 73 mixed with other parts. This was followed by seeds 17%, flowers 6%, roots 2% and fruits, bark (1% each). The current survey represents a useful documentation, which can use to preserving knowledge on the use of medicinal plants in this region and to explore the phytochemical and pharmacological potential of medicinal plants.

Key words: Ethnobotanical survey, anxiety, depression, medicinal plants, traditional medicine.

## **Enquête ethnobotanique des plantes médicinales utilisées dans le traitement de la dépression et anxiété dans la région Fès-Meknes**

### **Résumé**

La dépression et l'anxiété constituent un problème majeur de santé mentale dans le monde. La majorité des Marocains utilisent la médecine traditionnelle pour répondre à leurs besoins en matière de santé, y compris pour diverses formes de dépression et d'anxiété. Le but de ce travail est de faire un inventaire des espèces de plantes utilisées en médecine traditionnelle pour soigner la dépression et de l'anxiété dans les régions de Fès-Meknès. Des questionnaires établis ont été administrés lors de 243 entretiens dans 4 communautés de cette région, avec les guérisseurs, les herboristes et les consommateurs. 55 espèces de plantes appartenant à 31 familles ont été rapportées. Les familles les plus citées sont les Lamiaceae, suivies des Asteraceae, Apiaceae et Verbenaceae. Les feuilles constituent la partie principale utilisée, seule (58%), et mélangée avec d'autres parties (73 %). Viennent ensuite les graines (17%), les fleurs (6%), les racines (2%) et les fruits et l'écorce (1% chacun). L'enquête actuelle constitue une documentation utile, qui peut servir à préserver les connaissances sur l'utilisation des plantes médicinales dans cette région et à explorer le potentiel phytochimique et pharmacologique des plantes médicinales.

**Mots clés :** Enquête ethnobotanique, anxiété, dépression, plantes médicinales, médecine traditionnelle

## **Introduction**

Anxiety and depression are two disorders mental state characterized by several symptoms. These symptoms are excessive worrying, feeling agitated, restlessness, fatigue, difficulty concentrating, irritability, tense muscles and trouble falling or staying asleep for anxiety and sad, empty, helpless, worthless, irritable for depression.

Indeed, these disorders are among the most common mental disorders in the world [10,22] According to the World Health report (2001), 450 million people suffer from a mental or behavioral disorder. Therefore, 4,5 % of the population, suffer from anxiety and 5,5 % from depression [59].

In Morocco, the prevalence of major depressive disorder for lifetime was 26,5 % conforming to a Moroccan epidemiological study [3]. Additionally, the WHO's report notes that years lived with disability in Morocco amounted to 3,8 % and 7.4% respectively of total number of anxiety and depression [4].

However, the treatment of these disorders in this country is limited on condition of several factors as:

- the cost of treatment;
- the socioeconomic level of the population;
- the use of medicinal plants treatment, currently therapeutic properties are proved scientifically in many diseases such diabetes, cancer... [12].

So Morocco country is known for its biodiversity, their vegetation is estimated to 5200 species including vegetation, 900 species endemic plants [29] and the 600 species are the medicinal floral [66].

The aim of this study was to find out the medicinal plant species commonly used to treat depression and anxiety by the population of Fez-Mekne region, which part of plants are used, how they are obtained, prepared and applied for the treatment of depression and anxiety.

## **Materials and methods**

### **Study area**

The study was conducted in the Fez-Meknes regions, situated in central Morocco (Fig. 1). Fez-Meknes region cover a total land area of 40,075 km<sup>2</sup>, representing 5.7% of the whole land area of the Kingdom of Morocco. Indeed, this region is composed of diversified natural zones divided into several geographical units: the Rif and Pre-Rif in the North, the mountains

of the Middle Atlas, the highlands of Missouri and the plain of Saïis. and it characterized by the diversity of its soils where we can identify three major types of soil. It should be noted that the Saïis plain is well known by rich soils and is full of important agricultural potential. Regarding climate, the region is characterized by a continental climate in the North, cold and humid in mountain areas, and a semi-arid climate in the highlands of Missouri. It should be noted that this variability of the climate leads to a diversity of crops that better adapt to the specificities of this region.

### **Ethnobotanical survey**

An ethnobotanical survey was conducted from December 2016 to June 2017, in 5 stations: Fez, Ain Beida, Ain Taoujdate, Sefrou, Taounate. This survey was conducted in the form of an interview guided by a survey.

A total of 244 people was interviewed for this purpose. Interviews were designed in to record information about the plants used to treat antidepressant/anxiolytic and their local names, methods of preparation, parts of the plant used, administration of medicines and the social identity of the people interviewed (age, sex, ...).

### **Relative frequency of citation**

The relative frequency of citation shows the local importance of each species and it's obtained by dividing the number of informants, who mention the use of the species, also known as the Frequency Citation (FC), by the number of informants participating in the survey[12].  $RFC = \frac{FC}{N}$  ( $0 < RFC < 1$ ).

### **Statistical study**

Statistical analysis of obtained data was performed using IBM spss statistics 22.

## **Results and discussion**

### **Demographic features**

244 informants including traditional healers, herbalists and inhabitants, contributed to the survey.

About 21% of the respondents aged between 18 to 36 years, 36% between 37 to 54 years, and 43% between 55 to 71 years. This age distribution reflects an interesting finding in terms of ethno medicinal knowledge. As similar surveys in the same study region interviews show that elderly people were particularly competent and possess a much greater knowledge of native plants [2, 12]. Similar results have been observed in Niger where ethno-medicinal knowledge in the treatment of mental illness is mainly held by elderly people [3].

74% of respondents are women and 26% are men. Depression and anxiety are more prevalent in women than men [6, 43]. Some evidence suggests that serotonin synthesis in female brain is higher than in the male brain [58].

About one third of the interviewed are illiterate (39%), primary education (7%), secondary education (27%) and university education (25%). This study is affirmative with other ethnobotanical studies carried out in Morocco [4, 12, 27]. In which knowledge on the use of medicinal plants is possessed by illiterate.

Knowledge on traditional medicine has largely been passed orally from generation to generation. The Majority of the people who participate in this survey, acquired the traditional medical knowledge from members of their families mainly grandparents and parents.

### **Medicinal plants and floristic analysis**

In this study we recorded information on a total of 55 medicinal plant species, belonging to 31 families. The plants were summarized in alphabetical order by families and species. Information such as Relative frequency of citation is provided for each species, vernacular name of plants species, mode of preparation and used parts (Table 1, Table 2).

Most recorded species (12 species) were from the Lamiaceae family, Apiaceae (7 species each), Asteraceae (5 species) and Fabaceae (3 species). This finding is in agreement with previous reports where these families were the most represented families in mental illness treatment in Morocco [26]. Dominance of Lamiaceae could be attributed to their abundance in the flora of Morocco (Fennane and Ibn Tattou, 2012).

### **Ethnobotanical indices (Relative frequency citation)**

The most frequently used plants to treat depression and anxiety based on RFC values are *Lippia citriodora* H.B & K. (42,4%), *Salvia officinalis* L (18,5%), *Anthemis nobilis* L. (17,3%), *Origanum Majorana* L. (9,1%), *Lavandula vera* D C. (5,3%). Our results are similar to those described in Morocco in other surveys [26] where *Hypericum perforatum*, *Melissa officinalis*, *Origanum majorana*, *Rosmarinus officinalis*, *Salvia officinalis*, *Hibiscus esculens*, *Citrus aurantium*, *Tilia cordata* and *verbena officinalis* are the plant species most used to treat anxiety[40].

Effectively, the antidepressant-like effect of *Rosmarinus officinalis* L. has been reported in animals [7,50,69], due to its high ursolic acid content [52]. The results of other study showed an involvement of the acid ursolic in the serotonergic and noradrenergic systems [21]. Another work suggested that carnosol and betulinic acid could be responsible for the antidepressant effect of *R. officinalis* extract [51].

Other results showed *Lavandula angustifolia* L. present an antidepressant-like effect in the forced swimming test (FST) after acute treatment and rosmarinic acid was suggested to be one of the active ingredients in this plant [41]. Furthermore, *Anthemis nobilis* L. provides an important antidepressant effect with an anxiolytic property [9].

It is also reported that *Coriandrum sativum* L. seed oil contains linalool (60–70%) as the major essential oil component [53], which presented a significant anxiolytic activity [20, 53].

*Pimpinella anisum* L. has been widely used in traditional medicine to treat a variety of diseases, including the anxiety. This effect is resultant of trans-Anethole [40, 71].

Ginseng (*Panax ginseng*, C.A. Meyer, Araliaceae) has long been used traditionally for the treatment of mental diseases such as anxiety and depression. Ginseng saponins have been to play an important role in its anxiolytic effect [18]. As well *Artemisia absinthium* (Asteraceae) is widely used for depression treatment too. Its effects may be correlated with the presence of antidepressant compounds such as phenolic and flavonoid contents [54]. Additionally *Lactuca sativa* L. belonging to Asteraceae family is one of the most commonly and widely herbal medicines used in the several countries in treatment of insomnia, anxiety, dry coughs. Anxiolytic effect of *Lactuca sativa* is confirmed by some studies [26, 35, 37]. In this context, certain traditional healers have claimed the efficacy of *Glycyrrhiza* species for a variety of pathological conditions as a diuretic, choleric, insecticide, coughs and painful swellings. *Glycyrrhiza glabra* L. plays a very important role in the treatment of depression and inhibiting the serotonin recapture [18, 25]. Likewise *Trigonella foenum graecum* L. (Fabaceae) has a long history of medical use in traditional and modern literature such as antidepressant treatment. This effect is due to 4-hydroxyisoleucine (4-HI), it constitutes about 80% of the total content of free amino acids in *Trigonella foenum graecum* L. seeds [31]. Saffron (*Crocus sativus* L). Contains several volatile compounds such as crocin, picrocrocin, and safranal. It contains also several non-volatile active components, many of them are carotenoids including zeaxanthin, various  $\alpha$ - and  $\beta$ -carotenes and lycopene. Among these molecules, several studies have demonstrated their antidepressant effect [36, 63]. Moreover *Cuminum cyminum* L. is used for treatment of diabetes and prediabetes, muscle and stomach spasms, diarrhea, preventing nausea, cold, infections, and depression. To explore the traditional use of *Cuminum cyminum* L. in the anxiety treatment, a study has demonstrated anxiolytic activity of hydroethanolic extract [38]. Furthermore, the essential oil of *Mentha pulegium* L. is used for different applications such antiseptic, ovarian cancer, sedative and antidepressant like- effect [38]. Additionally, the aerial part of *Ocimum basilicum* L. has been acts mainly on the digestive and nervous systems, stomach cramps, flatulence, colic and

indigestion. It's an antispasmodic, digestive, aromatic, carminative, galactagogue, stomachic and tonic. In the mental disorders, *Ocimum basilicum* L. treated the depression by its essential oil and leaves [1, 64, 75]. Moreover, *Myrtus communis* has been suggested as a hypnotic in traditional medicine. Other previous studies have also indicated its sedative hypnotic-like and anxiolytic properties [34]. This anxiolytic and myorelaxant effects, are explained by involvement of alpha adrenergic pathway [15, 34]. Likewise the active components in *Nigella sativa* L and *Olea europaea* oil are demonstrated to be powerful in treatment of impaired mental. *Olea europaea* oil may produce psychostimulant-like effect. For this reason, this oil exerts an important pharmacological Modulation of dopamine and serotonin level [17]. Furthermore with over 3,000 varieties, tea is the most consumed beverage in the world after water. Tea can be divided into six categories: black, dark, yellow, oolong, white and green. The oral administration of tea extract in rat increased the dopamine and the serotonin [75,73]. However comparative literature analysis suggests 10 species are described in ethnobotanical survey [26, 33] and 20 species are reported for the first time for treating depression and anxiety.

### **Plant parts used**

The obtained results showed that the plant parts such as leaves, stems, roots, bark, fruit, flowers, seeds, and wood were used in treatment. Or, the most commonly used parts are the leaves, solely or mixed with other parts, forming 73% of total users. This was followed by seeds (17%), whole plants (12%), flowers (6%), roots (2%), fruit and stems and bark (1%) (Fig.2).

### **Preparation methods**

As shows the figure 3, the medicinal plants remedies are prepared as decoctions, infusions. Indeed, the infusion (58%) and decoction (22%) preparations in this study are the frequently used methods (Fig. 3). This result is in agreement with other ethnobotanical studies where infusion and decoction are the most frequently modes of preparation [12,11,5,39,68,74].

The great most of the remedies were taken orally (98 %) usually drunk as teas. Similar types of results were obtained in other studies [14,12,70].

### **Conclusion**

This investigation is first meant to focus on ethno-pharmacological knowledge of plants used in the treatment of depression and anxiety disorders in Morocco. The study revealed that traditional medicine continues to play an important role in the healthcare system in Morocco. 58 species belonging to 28 families are used for treating depression and anxiety in Fez-Meknes regions. The current survey represents a useful documentation, which can contribute

to preserving and keeping knowledge on the traditional and popular use of medicinal plants in Morocco.

### **Conflicts of interest**

The authors declare no conflicts of interest.

### **Acknowledgments**

This work was supported by H2020 project, H2020-MSCA-RISE-Marie Skłodowska-Curie Actions (MSCA) Research and Innovation Staff Exchange (RISE), Acronym: VAHVISTUS – Project Number: 734759.



## References

1. Abdoly, M, Farnam, A, Fathiazad, F, (2012) Antidepressant-like activities of *Ocimum basilicum* (sweet Basil) in the forced swimming test of rats exposed to electromagnetic field (EMF) *AJPP* 6: 211–215
2. Abouri, M, EL Mousadik, A, Msanda, F, Boubaker, H, Saadi, B, Cherifi, K (2012) An ethnobotanical survey of medicinal plants used in the Tata Province. Morocco *Int J Med Plant Res* 99–123
3. Ibrahim, J A, Muazzam, I, Jegede, I A, Kunle, O F, Okogun, J I (2007) Ethno-Medicinal Plants and Methods Used by Gwandara Tribe of Sabo Wuse In Niger State, Nigeria, To Treat Mental Illness. *African Journal of Traditional, Complementary and Alternative Medicines* 4: 211–218
4. Zerabruk, S, Yirga, G (2012) Traditional knowledge of medicinal plants in Gindeberet district, Western Ethiopia. *South African Journal of Botany* 78 :165–169
5. Afolayan, A J, Grierson, D S, Mbeng, W O (2014) Ethnobotanical survey of medicinal plants used in the management of skin disorders among the Xhosa communities of the Amathole District, Eastern Cape. *South Africa J Ethnopharmacol* 153: 220–232
6. Albert, P R (2015) Why is depression more prevalent in women? *J Psychiatry Neurosci* 40: 219–221
7. Al-Sereiti, M R, Abu-Amer, K M, Sena, P (1999) Pharmacology of rosemary (*Rosmarinus officinalis* Linn) and its therapeutic potentials *IJEB* 37:02
8. Amsterdam, J D, Li, Y, Soeller, I, Rockwell, K, Mao, J J, Shults, J (2009) A randomized, double-blind, placebo-controlled trial of oral *matricaria recutita* (chamomile) extract therapy of generalized anxiety disorder *J Clin Psychopharmacol* 29: 378–382
9. Amsterdam, J D, Shults, J, Soeller, I, Mao, J J, Rockwell, K, Newberg, A B (2012) Chamomile (*Matricaria recutita*) May Have Antidepressant Activity in Anxious Depressed Humans - An Exploratory Study. *Altern Ther Health Med* 18: 44–49
10. Anseau, M, Dierick, M, Buntinx, F, Cnockaert, P, De Smedt, J, Van Den Haute, M, Vander Mijnsbrugge, D (2004) High prevalence of mental disorders in primary care. *J Affect Disord* 78: 49–55

11. Bahmani, M, Zargaran, A, Rafieian-Kopaei, M, Saki, K (2014) Ethnobotanical study of medicinal plants used in the management of diabetes mellitus in the Urmia, Northwest Iran *Asian Pac J Trop Med* 1:348-54
12. Barkaoui, M, Katiri, A, Boubaker, H, Msanda, F (2017) Ethnobotanical survey of medicinal plants used in the traditional treatment of diabetes in Chtouka Ait Baha and Tiznit (Western Anti-Atlas), Morocco. *J Ethnopharmacol* 198, 338–350
13. Bellakhdar, J (1997) *La Pharmacopée marocaine traditionnelle - Médecine arabe ancienne et savoirs populaires*, Editions Ibis-Press (Paris) et Editions Le Fennec (Casablanca) ed maroc
14. Benarba, B, Belabid, L, Righi, K, Bekkar, A A, Elouissi, M, Khaldi, A, Hamimed, A (2015) Ethnobotanical study of medicinal plants used by traditional healers in Mascara (North West of Algeria). *J Ethnopharmacol* 175: 626–637
15. Birhanie, M W, Walle, B, Rebba, K (2016) Hypnotic effect of the essential oil from the leaves of *Myrtus communis* on mice *Nat Sci Sleep* 8: 267–275
16. Boskabady, M H, Shafei, M N, Saberi, Z, Amini, S (2011) Pharmacological Effects of *Rosa Damascena* Iran *J Basic Med Sci* 14: 295–307
17. Cheema, M A R, Nawaz, S, Gul, S, Salman, T, Naqvi, S, Dar, A, Haleem, D J (2016) Neurochemical and behavioral effects of *Nigella sativa* and *Olea europaea* oil in rats. *Nutritional Neuroscience* 0: 1–10
18. Cho, S, Park, J -H, Pae, A N, Han, D, Kim, D, Cho, N -C, No, K T, Yang, H, Yoon, M, Lee, C, Shimizu, M, Baek, N -I (2012) Hypnotic effects and GABAergic mechanism of licorice (*Glycyrrhiza glabra*) ethanol extract and its major flavonoid constituent glabrol. *Bioorganic & Medicinal Chemistry* 20: 3493–3501
19. Chouksey, D, Upmanyu, N, Pawar, R (2013) Central nervous system activity of *Illicium verum* fruit extracts. *Asian Pacific Journal of Tropical Medicine* 6: 869–875
20. Cioanca, O, Hritcu, L, Mihasan, M, Trifan, A, Hancianu, M (2014) Inhalation of coriander volatile oil increased anxiolytic–antidepressant-like behaviors and decreased oxidative status in beta-amyloid (1–42) rat model of Alzheimer’s disease. *Physiology & Behavior* 131: 68–74

21. Colla, A R S, Oliveira, Á, Pazini, F L, Rosa, J M, Manosso, L M, Cunha, M P, Rodrigues, A L S (2014) Serotonergic and noradrenergic systems are implicated in the antidepressant-like effect of ursolic acid in mice. *Pharmacology Biochemistry and Behavior* 124: 108–116
22. Craven, M A, Bland, R (2013) Depression in primary care: current and future challenges *Can. J Psychiatry* 58: 442–448
23. Dang, H, Chen, Y, Liu, X, Wang, Q, Wang, L, Jia, W, Wang, Y (2009) Antidepressant effects of ginseng total saponins in the forced swimming test and chronic mild stress models of depression. *Progress in Neuro-Psychopharmacology and Biological Psychiatry, Bed nucleus of the stria terminalis: anatomy, physiology, functions* 33: 1417–1424
24. Dhingra, D, Chhillar, R (2012) Antidepressant-like activity of ellagic acid in unstressed and acute immobilization-induced stressed mice. *Pharmacological Reports* 64: 796–807
25. Dhingra, D, Sharma, A (2006) Antidepressant-like activity of *Glycyrrhiza glabra* L in mouse models of immobility tests. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 30 : 449–454
26. Doukkali, Z, Boudida, H, Srfi, A, Taghzouti, K, Cherrah, Y, Alaoui, K (2015) Les plantes anxiolytiques au Maroc. *Études ethnobotanique et ethno-pharmacologique* 13: 306–31
27. Eddouks, M, Maghrani, M, Lemhadri, A, Ouahidi, M -L, Jouad, H (2002) Ethnopharmacological survey of medicinal plants used for the treatment of diabetes mellitus, hypertension and cardiac diseases in the south-east region of Morocco (Tafilalet). *J Ethnopharmacol* 82: 97–103
28. Emamghoreishi, M, Khasaki, M, Aazam, M F (2005) *Coriandrum sativum*: evaluation of its anxiolytic effect in the elevated plus-maze. *Journal of Ethnopharmacology* 96 : 365–370
29. Fennane, M, Ibn Tattou, M (2012) Statistiques et commentaires sur l’inventaire actuel de la flore vasculaire du Maroc *Bulletin de l’Institut Scientifique, Rabat, section Sciences de la Vie*

30. Fern, P F C, es, Azevedo, C N de, Dias, A K L G, Damiatilde, D S, Peregrino, O, Diniz, D R M R, Fern, P M B, es, Melo, D M de F F, Nobrega, D A R (2012) Naturally occurring anxiolytic substances from aromatic plants of genus citrus JMPR 6: 342–347
31. Gaur, V, Bodhankar, S L, Mohan, V, Thakurdesai, P (2012) Antidepressant-like effect of 4-hydroxyisoleucine from *Trigonella foenum graecum* L seeds in mice. *Biomedicine & Aging Pathology* 2: 121–125
32. Ghoshegir, S A, Mazaheri, M, Ghannadi, A, Feizi, A, Babaeian, M, Tanhaee, M, Karimi, M, Adibi, P (2015) *Pimpinella anisum* in the treatment of functional dyspepsia: A double-blind, randomized clinical trial. *J Res Med Sci* 20: 13–21
33. Gutiérrez, G, Laura, S, Reyes Chilpa, R, Bonilla Jaime, H, Gutiérrez, G, Laura, S, Reyes Chilpa, R, Bonilla Jaime, H (2014) Medicinal plants for the treatment of “nervios”, anxiety, and depression in Mexican Traditional Medicine. *Revista Brasileira de Farmacognosia* 245: 91–608
34. Hajiaghaee, R, Faizi, M, Shahmohammadi, Z, Abdollahnejad, F, Naghdibadi, H, Najafi, F, Razmi, A (2016) Hydroalcoholic extract of *Myrtus communis* can alter anxiety and sleep parameters: a behavioural and EEG sleep pattern study in mice and rats. *Pharm Biol* 54: 2141–2148
35. Harsha, S, Anilakumar, K, (2013) Anxiolytic property of *Lactuca sativa*, effect on anxiety behaviour induced by novel food and height Asian Pacific. *Journal of Tropical Medicine* 6: 532–536
36. Hosseinzadeh, H, Noraei, N B (2009) Anxiolytic and hypnotic effect of *Crocus sativus* aqueous extract and its constituents, crocin and safranal, in mice. *Phytother Res* 23: 768–774
37. Ismail, H, Dilshad, E, Waheed, M T, Sajid, M, Kayani, W K, Mirza, B (2016) Transformation of *Lactuca sativa* L. with rol C gene results in increased antioxidant potential and enhanced analgesic, anti-inflammatory and antidepressant activities in vivo 3. *Biotech* 6: 215
38. Jabeen, A, Ramya, B, Soujanya, J, Bhattacharya, B (2017) Evaluation of anxiolytic, muscle relaxant & locomotor activity of *cuminum cyminum*. *Journal of Medicinal Plants Studies* 259–262

39. Kadir, M F, Bin Sayeed, M S, Shams, T, Mia, M M K (2012) Ethnobotanical survey of medicinal plants used by Bangladeshi traditional health practitioners in the management of diabetes mellitus. *Journal of Ethnopharmacology* 144: 605–611
40. Kahloula, K, Slimani, M, Adli, D E H, Rachdi, S, Boumediene, D (2013) Neuro beneficial effects of *Pimpinella anisum* against lead exposure International. *Journal of Green Pharmacy (IJGP)* 7
41. Kageyama, A, Ueno, T, Oshio, M, Masuda, H, Horiuchi, H, Yokogoshi, H (2012) Antidepressant-like Effects of an Aqueous Extract of Lavender (*Lavandula angustifolia* Mill) in Rats. *Food Sci Technol Res* 473 – 479
42. Kalshetti, P B, Alluri, R, Mohan, V, Thakurdesai, P A (2015) Effects of 4-hydroxyisoleucine from Fenugreek Seeds on Depression-like Behavior in Socially Isolated Olfactory Bulbectomized Rats. *Pharmacogn Mag* 11: S388–S396
43. Kessler, R C, McGonagle, K A, Swartz, M, Blazer, D G, Nelson, C B (1993) Sex and depression in the National Comorbidity Survey I: Lifetime prevalence, chronicity and recurrence. *J Affect Disord* 29: 85–96
44. Khan, A W, Khan, A, Ahmed, T (2016) Anticonvulsant, Anxiolytic, and Sedative Activities of *Verbena officinalis*. *Front Pharmacol* 7: 499
45. Khan, R A, Riaz, A (2015) Behavioral effects of citrus limon in rats. *Metab Brain Dis* 30: 589–596
46. Kumar, S, Maheshwari, K K, Singh, V (2008) Central nervous system activity of acute administration of ethanol extract of *Punica granatum* L seeds in mice. *IJEB* 46:811-6
47. Latha, K, Rammohan, B, Sunanda, B P V, Maheswari, M S U, Mohan, S K (2015) Evaluation of anxiolytic activity of aqueous extract of *Coriandrum sativum* Linn in mice: A preliminary experimental study. *Pharmacognosy Res* 7: S47–S51
48. Lopes Campêlo, L M, Gonçalves e Sá, C, de Almeida, A A C, Pereira da Costa, J, Costa Marques, T H, Mendes Feitosa, C, Barros Saldanha, G, Mendes de Freitas, R (2011) Sedative, anxiolytic and antidepressant activities of Citrus limon (Burn) essential oil in mice *Die Pharmazie - An International. Journal of Pharmaceutical Sciences* 66: 623–627

49. López, V, Nielsen, B, Solas, M, Ramírez, M J, Jäger, A K (2017) Exploring Pharmacological Mechanisms of Lavender (*Lavandula angustifolia*) Essential Oil on Central Nervous System Targets. *Front Pharmacol* 19; 8:28
50. Machado, D G, Bettio, L E B, Cunha, M P, Capra, J C, Dalmarco, J B, Pizzolatti, M G, Rodrigues, A L S (2009) Antidepressant-like effect of the extract of *Rosmarinus officinalis* in mice: Involvement of the monoaminergic system. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 33: 642–650
51. Machado, D G, Cunha, M P, Neis, V B, Balen, G O, Colla, A, Bettio, L E B, Oliveira, Á, Pazini, F L, Dalmarco, J B, Simionatto, E L, Pizzolatti, M G, Rodrigues, A L S, (2013) Antidepressant-like effects of fractions, essential oil, carnosol and betulinic acid isolated from *Rosmarinus officinalis* L. *Food Chemistry* 136: 999–1005
52. Machado, D G, Neis, V B, Balen, G O, Colla, A, Cunha, M P, Dalmarco, J B, Pizzolatti, M G, Prediger, R D, Rodrigues, A L S (2012) Antidepressant-like effect of ursolic acid isolated from *Rosmarinus officinalis* L in mice: Evidence for the involvement of the dopaminergic system. *Pharmacology Biochemistry and Behavior* 103: 204–211
53. Mahendra, P, Bisht, S (2011) Anti-anxiety activity of *Coriandrum sativum* assessed using different experimental anxiety models Indian. *J Pharmacol* 43: 574–577
54. Mahmoudi, M, Ebrahimzadeh, M A, Ansaroudi, F, Nabavi, S F, Nabavi, S M (2009) Antidepressant and antioxidant activities of *Artemisia absinthium* L at flowering stage. *African Journal of Biotechnology* 8:24
55. Masoumi-Ardakani, Y, Mahmoudvand, H, Mirzaei, A, Esmailpour, K, Ghazvini, H, Khalifeh, S, Sepehri, G (2017) The effect of *Elettaria cardamomum* extract on anxiety-like behavior in a rat model of post-traumatic stress disorder. *Biomed Pharmacother* 87: 489–495
56. Mesfin, M, Asres, K, Shibeshi, W (2014) Evaluation of anxiolytic activity of the essential oil of the aerial part of *Foeniculum vulgare* Miller in mice. *BMC Complementary and Alternative Medicine* 14: 310
57. Mirza, B, Ikram, H, Bilgrami, S, Haleem, D J, Haleem, M A (2013) Neurochemical and behavioral effects of green tea (*Camellia sinensis*): a model study. *Pak J Pharm Sci* 26: 511–516

58. Nishizawa, S, Benkelfat, C, Young, S N, Leyton, M, Mzengeza, S, de Montigny, C, Blier, P, Diksic, M (1997) Differences between males and females in rates of serotonin synthesis in human brain. *Proc Natl Acad Sci U S A* 94: 5308–5313
59. Oneib, B, Sabir, M, Abda, N, Ouanass, A (2015) Epidemiological study of the prevalence of depressive disorders in primary health care in Morocco. *J Neurosci Rural Pract* 6: 477–480
60. Osanloo, N, Najafi-Abedi, A, Jafari, F, Javid, F, Pirpiran, M, Memar Jafari, M -R, Mousavi Khosravi, S A, Rahimzadeh Behzadi, M, Ranjbaran, M, Sahraei, H (2016) Papaver Rhoeas L Hydroalcoholic Extract Exacerbates Forced Swimming Test-Induced Depression in Mice. *Basic Clin Neurosci* 7: 195–202
61. Park, J -H, Cha, H -Y, Seo, J -J, Hong, J -T, Han, K, Oh, K -W (2005) Anxiolytic-like effects of ginseng in the elevated plus-maze model: Comparison of red ginseng and sun ginseng. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 29: 895–900
62. Perveen, T, Haider, S, Zuberi, N A, Saleem, S, Sadaf, S, Batool, Z (2014) Increased 5-HT Levels Following Repeated Administration of Nigella sativa L (Black Seed) Oil Produce Antidepressant Effects in Rats. *Sci Pharm* 82: 161–170
63. Pitsikas, N, Bouladakis, A, Georgiadou, G, Tarantilis, P A, Sakellaridis, N (2008) Effects of the active constituents of Crocus sativus L, crocins, in an animal model of anxiety. *Phytomedicine* 15: 1135–1139
64. Rabbani, M, Sajjadi, S E, Vaezi, A (2015) Evaluation of anxiolytic and sedative effect of essential oil and hydroalcoholic extract of Ocimum basilicum L and chemical composition of its essential oil. *Res Pharm Sci* 10: 535–543
65. Rabiei, Z, Gholami, M, Rafieian-Kopae, M (2016) Antidepressant effects of Mentha pulegium in mice. *Journal of Pharmacology* 11 : 711–715
66. Rejdali, M (1996) La flore du Maroc : Etat actuel et perspectives de conservation, diversité biologique et valorisation des plantes médicinales. Actes Edition ed pp121-pp122
67. Riaz, A, Khan, R A (2017) Behavioral effects of *Citrus limon* and *Punica granatum* combinations in rats. *Metab Brain Dis* 32: 123–131

68. Salehi Nowbandegani, A, Kiumarcy, S, Rahmani, F, Dokouhaki, M, Khademian, S, Zarshenas, M M, Faridi, P (2015) Ethnopharmacological knowledge of Shiraz and Fasa in Fars region of Iran for diabetes mellitus. *J Ethnopharmacol* 172: 281–287
69. Sasaki, K, El Omri, A, Kondo, S, Han, J, Isoda, H (2013) Rosmarinus officinalis polyphenols produce anti-depressant like effect through monoaminergic and cholinergic functions modulation. *Behavioural Brain Research* 238: 86–94
70. Semanya, S, Potgieter, M, Erasmus, L (2012) Ethnobotanical survey of medicinal plants used by Bapedi healers to treat diabetes mellitus in the Limpopo Province. *South Africa J Ethnopharmacol* 141: 440–445
71. Shahamat, Z, Abbasi-Maleki, S, Motamed, S M (2016) Evaluation of antidepressant-like effects of aqueous and ethanolic extracts of *Pimpinella anisum* fruit in mice. *Avicenna Journal of Phytomedicine* 6: 322
72. Sharma, P K, Singh, V, Ali, M, Kumar, S (2016) Effect of ethanolic extract of *Zingiber officinale* Roscoe on Central Nervous System activity in mice. *IJEB* 54(10):664-669
73. Tabassum, I, Siddiqui, Z N, Rizvi, S J (2010) Effects of *Ocimum sanctum* and *Camellia sinensis* on stress-induced anxiety and depression in male albino *Rattus norvegicus*. *Indian J Pharmacol* 42: 283–288
74. Urso, V, Signorini, M A, Tonini, M, Bruschi, P (2016) Wild medicinal and food plants used by communities living in Mopane woodlands of southern Angola: Results of an ethnobotanical field investigation. *J Ethnopharmacol* 177: 126–139
75. Venâncio, A M, Marchioro, M, Estavam, C S, Melo, M S, Santana, M T, Onofre, A S C, Guimarães, A G, Oliveira, M G B, Alves, P B, Pimentel, H de C, Quintans-Júnior, L J (2011) *Ocimum basilicum* leaf essential oil and (-)-linalool reduce orofacial nociception in rodents: a behavioral and electrophysiological approach. *Revista Brasileira de Farmacognosia* 21: 1043–1051
76. Wang, J, Flaisher-Grinberg, S, Li, S, Liu, H, Sun, L, Zhou, Y, Einat, H (2010) Antidepressant-like effects of the active acidic polysaccharide portion of ginseng in mice. *Journal of Ethnopharmacology* 132: 65–69



77. Wazid, A, Saifuddin, K, Khatoon, M (2015) Anxiolytic activity of aqueous extract of Nerium Oleander flower on experimental animals. *Journal of Pharmacy and Pharmaceutical Sciences* 4: 1712–1730
78. Xu, C, Teng, J, Chen, W, Ge, Q, Yang, Z, Yu, C, Yang, Z, Jia, W (2010) 20(S)-protopanaxadiol, an active ginseng metabolite, exhibits strong antidepressant-like effects in animal tests *Progress in Neuro-Psychopharmacology and Biological Psychiatry. Intracellular Ca-homeostasis in schizophrenia* 34: 1402–1411
79. Yamada, N, Araki, H, Yoshimura, H (2011) Identification of antidepressant-like ingredients in ginseng root (*Panax ginseng* C.A. Meyer) using a menopausal depressive-like state in female mice: participation of 5-HT<sub>2A</sub> receptors. *Psychopharmacology* 216: 589–599



Fig.1 Geographical location of the study area. (Fez - Meknes Regional Directorate)




Figure 2: Plant parts used




Figure 3: Percentage of types of administration

**Table1: Plants use to treat depression/anxiety in Fez-Meknes areas.**

Families and plants pecies	Vernacular name	Ecological distribution	Used parts	Preparation	Mode of administration	RF C (%)	Literature references in relationship of medicinal use of plants
<b>Apiaceae</b>							
<i>Carum carvi</i> L.	Karwiya	Cultivated	seeds	Decoction , Infusion	Oral	3, 7	No
<i>coriandrum sativum</i> L.	qezbor	Cultivated	Seeds, Leaves	Decoction , Infusion	Oral	1,2	[28] [20,28,47,53]
<i>Cuminum cyminum</i> L.	kemmûn	Cultivated	Seeds	Infusion	Oral	4	[38]
<i>Daucus carota</i> L.	ħizzu	Cultivated	Seeds	Raw, Infusion	Oral	1,2	No
<i>Foeniculum vulgare</i> P.Mill	Nâfa'	Cultivated	Seeds	Decoction , Infusion	Oral	4, 9	[56]
<i>petroselinum sativum</i> Mill	ma'dnûs	Cultivated	Leaves	Decoction , jus,	Oral	3,3	No
<i>Pimpinella anisum</i> L.	Habbat halâwa	Cultivated	Seeds	Decoction	Oral	2,1	[32,40,71]
<b>Apocynaceae</b>							
<i>Nerium Oleander</i> L.	ddeflâ	Wild	Stem and leaves and flower	Raw	Nasal	0,4	[77]

<b>Araliaceae</b>							
<i>Panax Ginseng</i> C.A. Mey.	Gensing	Wild	Roots	infusion, Raw	Oral	0,8	[23,61,76, 78,79]
<b>Palmaceae</b>							
<i>Phoenix dactylifera</i> L.	Ttmer	Cultivated	Fruit	<b>Raw</b>	Oral	0,4	No
<b>Astéraceae</b>							
<i>Anthemis nobilis</i> L.	bâbnûj	Wild	Flower And Leaves	Infusion Decoction	Oral	17, 3	[8]
<i>Artemisia absinthium</i> L.	šîba	Cultivated	Leaves	Infusion	Oral	0,8	[54]
<i>Artemisia herba alba</i> Asso Asso.	šîh	Wild	Stem and leaf	Infusion Decoction	Oral	4,1	No
<i>Dittrichia viscosa</i> L.	Magram ane	Wild	Leaves	Infusion Decoction	Oral	1,2	No
<i>Lactuca sativa</i> L.	ḥoss	Cultivated	Raw	Raw	Oral	0, 4	[26,35,37]
<b>Brassicaceae</b>							
<i>Lepidium sativum</i> L.	Habb er- ršâd	Cultivated	Seeds	Infusion	Oral	0, 4	No
<b>Caryophyllaceae</b>							
<i>herniaria hirsute</i> L.	herste lhjer	Wild	whole plant	Infusion	Oral	0,4	No
<b>Maranhtaceae</b>							
<i>Dysphania ambrosioides</i>	Mkhniza	Wild	Leaves	Infusion	Oral	1, 2	No

s L.							
<b>Cupressaceae</b>							
<i>Tetraclinis articulata</i> . Masters.	'ar'ar	Wild	Leaves	Row	Nasal	1,6	No
<b>Fabaceae</b>							
<i>Glycyrrhiza glabra</i> L.	'arq as-sûs	Wild and Cultivated	Stem	Decoction	Oral	0,4	[18,25]
<i>Vicia ervilia</i> L. Willd.	Kersenna	Wild and Cultivated	Seeds	Direct administration	Oral	0,4	No
<i>Trigonella foenum graecum</i> L.	l-halba	Cultivated	Seeds	Decoction, maceration and Mixed with honey	Oral	3,7	[31,42]
<b>Irudaceae</b>							
<i>Crocus sativus</i> L.	Za'afra	Cultivated	Flower	Infusion	Oral	0,8	[36,63]
<b>Juglandaceae</b>							
<i>Juglans regia</i> L.	l-gerga'	Cultivated	Fruit	Direct administration	Oral	0,4	[16,28]
<b>Lauraceae</b>							
<i>Cinnamomum cassia</i> Blume	qarfa	Imported	Bark	Decoction, Infusion	Oral	1, 2	[12,21]
<b>Lythraceae</b>							
<i>Lawsonia inermis</i> L.	l-hennâ	Cultivated	Leaves	Infusion	Oral	1,6	No
<b>Lamiaceae</b>							
<i>Marrubium</i>	merrîwa	Wild	Leav	Decoction	Oral		[33]

<i>vulgare</i> L.			es	, Infusion		0,8	
<i>Mentha pulegium</i> L.	fliyyo	Wild	Leaves	Infusion,	Oral	3,3	[65]
<i>mentha suaveolens</i> Ehr.	Marseta	Cultivated	Leaves	Decoction, Infusion	Oral	2,5	No
<i>Origanum compactum</i> Benth.	za'tar	Wild	Leaves	Decoction, Infusion	Oral	0,8	[26]
<i>Origanum Majorana</i> L.	Merded ūš	Cultivated	Leaves	Decoction, Infusion	Oral	9,1	[26]
<i>Rosmarinus officinalis</i> L.	âzîr	Cultivated	Leaves	Decoction, Infusion	Oral	16,5	[7,50,51,52,69]
<i>Salvia officinalis</i> L.	sâlmiya	Cultivated	Leaves	Decoction, Infusion	Oral	18,5	[26]
<i>Satureja calamintha</i> L. Scheele	Menta	Wild	leaves	Infusion	Oral	0,4	NO
<i>Lavandula vera</i> D C. / <i>Lavandula abrialis</i>	huzâma	Cultivated	Stem and leaf	Decoction, Infusion	Oral	5,3	No
<i>Mentha piperita</i> L.	na'na' abdî	Cultivated	leaves	Infusion	Oral	0,4	[11,49,71]
<i>Mentha spicata</i> L.	na'na'	Cultivated	leaves	Infusion	Oral	1,2	[26]
<i>Ocinum basilicum</i> L.	lahbaq	Cultivated	leaves	Infusion	Oral	0,8	[1,64,75]
<b>Malvaceae</b>							
<i>Hibiscus sabdariffa</i> L.	Karkadiya	Wild	Flower	decoction	Oral	0,8	[32,18]
<b>Myristicaceae</b>							
<i>Myristica fragans</i> houtt.	l-gûza	imported	Fruit	Infusion	Oral	0,4	No

<b>Myrtacées</b>							
<i>Myrtus communis</i> L.	rîhân	Wild	Leaves	Infusion	Oral	0,4	[15,34]
<b>Oleaceae</b>							
<i>Olea europaea</i> L.	zaytûn	Wild and Cultivated	leaves	Infusion	Oral	0,8	[17]
<b>Papaveraceae</b>							
<i>Papaver rhoeas</i> L.	beláman	Wild	Flower	Infusion	Oral	0,4	[60]
<b>Poaceae</b>							
<i>Saccharum officinarum</i> L.	Qasab sakkuri,	Cultivated	Stem	Jus	Oral	0,4	No
<b>Punicaceae</b>							
<i>Punica granatum</i> L.	Qšûr rommân	Cultivated	Peel	decoction	Oral	0,4	[24,46, 66]
<b>Ranunculaceae</b>							
<i>Nigella sativa</i> L.	sânûj	Cultivated	Seeds	Direct administration	Oral	2,9	[62,17]
<b>Rosaceae</b>							
<i>Rosa damascene</i> Mill.	Werd	Cultivated	Flower	Infusion	Oral	0,4	[16]
<b>Rubiaceae</b>							
<i>Coffea arabica</i> L.	qahwa	Imported	Seeds	decoction	Oral	0,8	No
<b>Rutaceae</b>							
<i>Citrus aurantium</i> L.	Lâranj, lernej	Cultivated	Flower	Infusion	Oral	1,2	[30]



<i>Citrus limon</i> L.	Hâmméd	Cultivatéd	Fruit	Jus	Oral	0,8	[45, 48, 67]
<b>Schisandraceae</b>							
<i>Illicium verum</i> Hook.f.	Yansone	Cultivatéd	Seed s	Infusion	Oral	2,5	[19]
<b>Theaceae</b>							
<i>Camellia sinensis</i> L.	Atay	Importéd	leave s	Decoction	Oral	0,8	[57,73]
<b>Verbenaceae</b>							
<i>Lippia citriodora</i> H.B & K.	Iwîza	Cultivatéd	leave s	Decoction , Infusion	Oral	42,4	[26,44]
<b>Zingibéraceae</b>							
<i>Elettaria cardamomum</i> L.	Qa'qolla , hebbel-hal	Importéd	grain	Infusé dans l'eau ou lait, Decoction	Oral	4,1	[55]
<i>Zingiber officinale</i> L.	Skenjbîr	Importéd	Raci ne	Jus	Oral	0,8	[72]
<i>Alpinia officinarum</i> Hance	khoudenjal	Importéd	Raci ne	Decoction	Oral	0,4	No

**Table2:** List of the species according to the different recipes, mode of preparation, used parts and mode of administration.

En-quête	Latin name	Solvents	Used parts	Posologie	Route of administration
1	<i>Lippia citriodora</i> H.B & K.	Water	leaves	1/3 Tsp	Oral

	<i>Salvia officinalis</i> L.		leaves	1/3 Tsp	Oral
	<i>Illicium verum</i> Hook.f.		Seeds	1/3 Tsp	Oral
2	<i>Lippia citriodora</i> H.B & K.	Water	leaves	1/3 Tsp	Oral
	<i>Anthemis nobilis</i> L.		Flower	1/3 Tsp	Oral
	<i>Myrtus communis</i> L.		leaves	1/3 Tsp	Oral
3	<i>Illicium verum</i> Hook.f.	Water	Seeds	1/7 Tsp	Oral
	<i>Lippia citriodora</i> H.B & K.		leaves	1/7 Tsp	Oral
	<i>Origanum Majorana</i> L.		leaves	1/7 Tsp	Oral
	<i>Pimpinella anisum</i> L.		Seeds	1/7 Tsp	Oral
	<i>Carum carvi</i> L.		Seeds	1/7 Tsp	Oral
	<i>Lavandula vera</i> D C. / <i>Lavandula abrialis</i>		leaves+stem	1/7 Tsp	Oral
	<i>Rosmarinus officinalis</i> L.		leaves	1/7 Tsp	Oral
4	<i>Nigella sativa</i> L.	milk	leaves	1/10 Tsp	Oral
	<i>Rosmarinus officinalis</i> L.		leaves	1/10 Tsp	Oral

	<i>Salvia officinalis</i> L.		leaves	1/10 Tsp	Oral
	<i>Origanum Majorana</i> L.		leaves	1/10 Tsp	Oral
	<i>Lippia citriodora</i> H.B & K.		Leaves	1/10 Tsp	Oral
	<i>Carum carvi</i> L.		Seeds	1/10 Tsp	Oral
	<i>Daucus carota</i> L.		Seeds	1/10 Tsp	Oral
	<i>Pimpinella anisum</i> L.		Seeds	1/10 Tsp	Oral
	<i>Anthemis nobilis</i> L.		Flower	1/10 Tsp	Oral
	<i>Apium graveolens</i> L.		Seeds	1/10 Tsp	Oral
5	<i>Lippia citriodora</i> H.B & K.	water	Leaves	1/3 Tsp	Oral
	<i>Salvia officinalis</i> L.		Leaves	1/3 Tsp	Oral
	<i>Mentha pulegium</i> L.		leaves+ Flower	1/3 Tsp	Oral
6	<i>Lippia citriodora</i> H.B & K.	Water	Leaves	1/2 Tsp	Oral
	<i>Salvia officinalis</i> L.		Leaves	1/2 Tsp	Oral
7	<i>Illicium verum</i> Hook.f.	honey	Seeds	1/6 Tsp	Oral
	<i>Lippia citri-</i>		Leaves	1/6 Tsp	Oral

	<i>odora</i> H.B & K.				
	<i>Nigella sativa</i> L.		Seeds	1/6 Tsp	Oral
	<i>Anthemis nobilis</i> L.		Leaves+ Flower	1/6 Tsp	Oral
	<i>Carum carvi</i> L.		Seeds	1/6 Tsp	Oral
	<i>Pimpinella anisum</i> L.		Seeds	1/6 Tsp	Oral