

## ORIGINAL PAPER

# Androgen receptors in areas of the spinal cord and brainstem: A study in adult male cats

Rosa L. Coolen  | Jacqueline C. Cambier | Panagiota I. Spantidea | Els van Asselt | Bertil F. M. Blok

Department of Urology, Erasmus Medical Center, Rotterdam, The Netherlands

**Correspondence**

Rosa Linda Coolen, Department of Urology, Erasmus Medical Center, Rotterdam, The Netherlands.  
Email:

**Abstract**

Sex hormones, including androgens and estrogens, play an important role in autonomic, reproductive and sexual behavior. The areas that are important in these behaviors lie within the spinal cord and brainstem. Relevant dysfunctional behavior in patients with altered androgen availability or androgen receptor sensitivity might be explained by the distribution of androgens and their receptors in the central nervous system. We hypothesize that autonomic dysfunction is correlated with the androgen sensitivity of spinal cord and brainstem areas responsible for autonomic functions. In this study, androgen receptor immunoreactive (AR-IR) nuclei in the spinal cord and brainstem were studied using the androgen receptor antibody PG21 in four uncastrated young adult male cats. A dense distribution of AR-IR nuclei was detected in the superior layers of the dorsal horn, including lamina I. Intensely stained nuclei, but less densely distributed, were found in lamina X and preganglionic sympathetic and parasympathetic cells of the intermediolateral cell column. Areas in the caudal brainstem showing a high density of AR-IR nuclei included the area postrema, the dorsal motor vagus nucleus and the retrotrapezoid nucleus. More cranially, the central linear nucleus in the pons contained a dense distribution of AR-IR nuclei. The mesencephalic periaqueductal gray (PAG) showed a dense distribution of AR-IR nuclei apart from the most central part of the PAG directly adjacent to the ependymal lining. Other areas in the mesencephalon with a dense distribution of AR-IR nuclei were the dorsal raphe nucleus, the retrorubral nucleus, the substantia nigra and the ventral tegmental area of Tsai. It is concluded that AR-IR nuclei are located in specific areas of the central nervous system that are involved in the control of sensory function and autonomic behavior. Furthermore, damage of these AR-IR areas might explain related dysfunction in humans.

**KEYWORDS**

androgen, autonomic function, medulla oblongata, mesencephalon, PG21, spinal cord

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Journal of Anatomy* published by John Wiley & Sons Ltd on behalf of Anatomical Society

## 1 | INTRODUCTION

Androgens play an important role in autonomic, reproductive, and sexual behavior (Cunningham et al., 2012; Santi et al., 2018). The central nervous system areas that are important in these behaviors lie within the spinal cord and brainstem. Relevant dysfunctional behavior in patients can be due to an altered availability of androgens in the central nervous system. This can be caused by androgen deprivation therapy, androgen insensitivity syndrome, hypogonadism, Kennedy's disease, and many other causes. Androgen deprivation therapy is one of the standard treatments of metastasized prostate cancer and may cause cognitive dysfunction, including dementia, impotence, and other side effects (Donovan et al., 2018; McGinty et al., 2014; Nead et al., 2017). Androgen insensitivity syndrome and Kennedy's disease are caused by androgen receptor mutations which alter the structure or function of the receptor (Brinkmann et al., 1995). Hypogonadism is a disorder that results in a decreased production of androgens. It is diagnosed according to the guideline of the European Association of Urology when signs and symptoms of androgen deficiency occur together with consistently low serum testosterone levels (Dohle et al., 2019). One of such symptoms is erectile dysfunction, which is associated with decreased levels of free testosterone (Huang et al., 2019). The consequences of reduced testosterone in the central nervous system are claimed to be sexual dysfunction, such as impotence, and impaired cognition (Bravo et al., 2017; Kawano et al., 2003; Ophoff et al., 2009; Rana et al., 2011; Takov et al., 2018; Traish et al., 2015). On the contrary, increased levels of testosterone are associated with altered behavioral responses such as aggression (Carré & Archer, 2018). To date, the pathophysiology of autonomic dysfunction associated with an altered level of androgens or an altered sensitivity/expression of androgen receptors is not well-known. We hypothesize that autonomic dysfunction in patients with altered levels of androgens is correlated with the androgen sensitivity of spinal cord and brainstem areas that are responsible for autonomic functions. Autonomic reflexes originate in the brainstem. It is, therefore, important to expand the knowledge on the androgen receptor distribution within these areas of the central nervous system.

Androgen receptors are members of a superfamily of ligand-dependent transcription factors. The androgen receptor genes are located on the X chromosome and expressed in most of the organs in the body (Hunter et al., 2018). They are located in an inactive form in the cytoplasm and are activated as a result of strong affinity-binding with androgens (mostly testosterone and dihydrotestosterone) and the subsequent formation of the androgen-androgen receptor complex. The activation of this complex causes translocation to the nucleus, where it binds directly to DNA or through histones or through chromatin remodeling (DNA binding-dependent). The activated complex can also induce activation of secondary messenger pathways including ERK, Akt and MAPK (non-DNA binding dependent) to regulate androgen-regulated genes (Brinkmann et al., 1999; Claessens et al., 2008; Gioeli & Paschal, 2012; Hipkaeo et al., 2004). For an optimal androgen receptor regulation, regulators, directly or indirectly inhibit androgen receptor activation by reducing the concentration of androgens. Those molecules can be cytokines, growth

factors or others. O'Bryant and Jordan showed that there are five different putative cofactors which have the potential to participate in motoneuronal responses to androgens (O'Bryant & Jordan, 2005).

It is known from immunohistochemical receptor studies in various species that androgens may have effects on specific regions of the central nervous system. Studies in rats have shown androgen receptor expression in autonomic behavior related structures of the spinal cord and brainstem such as the area postrema, the intermediolateral cell column, the nucleus of the solitary tract, and the periaqueductal gray (PAG) (Sar & Stumpf, 1975; Simerly et al., 1990). In rhesus and cynomolgus macaques, androgen sensitive structures were found in forebrain areas, but the location of androgen receptor immunoreactive (AR-IR) cells in the spinal cord and brainstem was not investigated in monkeys except for one study that described AR-IR cells in the dorsal raphe (Bethea et al., 2015; Choate et al., 1998; Clancy et al., 1992; Michael et al., 1989). Thus far, the androgen receptor distribution in the spinal cord and brainstem of mammals other than rodents has not been studied. The location of AR-IR cells in the spinal cord and brainstem has not been investigated in cats and humans either. The cat nervous system has a great resemblance to that of humans. Therefore, it is an excellent model for studying the central nervous system.

The present study investigates which areas of the spinal cord and brainstem of adult male cats contain AR-IR nuclei. A comparison is made between the distribution of AR-IR nuclei in the cat and that previously described in the literature on rodents and monkeys. The possible roles of androgen receptors in specific spinal cord and brainstem areas in autonomic function and dysfunction are discussed. Furthermore, the distribution of androgen receptors in the spinal cord and brainstem is compared to the distribution of estrogen receptors in the same areas as described in earlier studies.

## 2 | MATERIALS AND METHODS

### 2.1 | Animals

All surgical procedures, pre- and postoperative care, handling, and housing of the cats were in accordance with the protocols approved by the Committee on Animal Experiments of the Faculty of Medical Sciences of the University of Groningen. Four uncastrated young adult male cats (12–24 months, Harlan) received 6 ml, 6% sodium pentobarbital intraperitoneally, and were perfused transcardially with Saline in 0.01 M Phosphate Buffer (PB, pH 7.4), followed by 4% paraformaldehyde in 0.1 M PB (pH 7.4). The spinal cords and brainstems were removed and post fixed for 2 h at 4°C.

### 2.2 | Immunohistochemistry

Sixty  $\mu\text{m}$  sections of the spinal cord and brainstem were cut on a Vibratome in cold Tris buffered saline (TBS, pH 7.4). One of four sections was pretreated with 0.3%  $\text{H}_2\text{O}_2$  for 30 min, blocked with 5% normal donkey serum for 30 min at room temperature, and

incubated with the primary antibody against androgen receptors, PG21 (donated by Dr Gail S. Prins, University of Chicago, IL, 1.4 µg/ml) diluted in 1% normal donkey serum at 4°C, for three days (Prins et al., 1991). Subsequently, biotinylated donkey anti-rabbit immunoglobulin (Jackson Immuno Research, diluted 1:500 with 1% normal donkey serum) was used as secondary antibody. The sections were incubated for one hour at room temperature, followed by incubation in avidin-biotin-complex-peroxidase (Vectastain, Vector, 1:400 in TBS, 1 h at room temperature). Following antibody incubation, the sections were rinsed with TBS overnight. Finally, the sections were incubated with 0.04% 3,3'-diaminobenzidine tetrahydrochloride (DAB), 0.2% nickel ammonium sulfate, and 0.01% H<sub>2</sub>O<sub>2</sub> in TBS for 5 min, resulting in a dark precipitate. The blocking serum, primary and secondary antibody were diluted in 0.05 M Tris buffer, 0.5 M NaCl, and 0.5% Triton X-100. In order to test the specificity of the AR-IR labeling, appropriate controls were made in a series of sections by omitting the primary antibody to verify the specificity of the AR-IR labeling. The specificity was previously shown using the antibody on multiple positive and negative controls and by performing competition studies (Prins et al., 1991). Another series of sections was counterstained with the Nissl staining method (Avendaño & Verdu, 1992). The sections were mounted on gelatin-covered slides and coverslipped with DePeX (Gurr, BDH laboratory).

### 2.3 | Data analysis

AR-IR nuclei were plotted with a computerized X-Y stage on a Zeiss Axioplan microscope connected to a PC with the NeuroLucida system (MicroBrightField). The drawings of the sections of the spinal cord and brainstem containing plotted nuclei were converted to Encapsulated PostScript (EPS) files and further processed in Adobe Illustrator CC 2018. Example drawings of representative sections at specific levels are presented in the results section (Figures 1 and 3). The results were quantified with a Heidelberg Topaz II + scanner (Kiel, Germany) and Linocolor 5.0 software. The nuclear AR labeling was quantified in a semiquantitative manner. For each area of the spinal cord and the brainstem two aspects of the AR-IR nuclei were determined: the density of the distribution of AR-IR nuclei and the intensity of nuclear staining. The distribution of labeled nuclei and the intensity of

the labeling are presented in Table 1. A Nikon Ts2-FL microscope was used to take bright field photomicrographs. Nissl-stained sections were used to localize the spinal cord nuclei, brainstem nuclei and fiber tracts using the cytoarchitectonic atlas of Berman (Berman, 1968).

## 3 | RESULTS

AR-IR was predominantly found in the nuclei of neurons, but occasionally also in nuclei of glial and ependymal cells. AR-IR cells were present throughout the spinal cord and brainstem. The immunostaining varied from light gray to intense black. AR-IR nuclei in the white matter represented, in all likelihood, glial cells. No staining was found in control sections. Two outcome measures were determined for each spinal cord and brainstem area: the density of the distribution of AR-IR nuclei and the nuclear intensity of the staining. The density of the distribution of AR-IR nuclei and the intensity of nuclear staining in areas of the spinal cord and brainstem are presented in Table 1. These results were consistently found in each of the four cats.


### 3.1 | Spinal cord

The distribution of AR-IR nuclei was moderate to low in density at every level of the spinal cord (Figures 1 and 2). Most densely labeled were the superior layers (layer I, II, and III) of the dorsal horn and the intermediolateral cell column of the thoracic, lumbar and sacral spinal cord, which contain preganglionic sympathetic and parasympathetic motoneurons. The deep layers of the dorsal horn and the ventral horn were sparsely labeled throughout the spinal cord.

Intensely stained nuclei were observed in the central (lamina X) and intermediate zone at all spinal levels, and in the preganglionic sympathetic and parasympathetic cells of the intermediolateral cell column at thoracic, lumbar, and sacral levels.

### 3.2 | Medulla oblongata

The distribution of AR-IR nuclei was dense in the area postrema, the dorsal motor vagus nucleus, and the retrotrapezoid nucleus



**FIGURE 1** AR-IR nuclei in the spinal cord. (a) The laminae of Rexed in a schematic drawing of the 6th cervical segment. (b) Spinal cord sections (C6-S3). Each dot represents one AR-IR nucleus

TABLE 1 Nuclear AR labeling in the spinal cord and brainstem

Region	Distribution of AR-IR nuclei	Intensity of nuclear AR labeling
<b>Spinal cord</b>		
Central zone (lamina X)	+	+++
Dorsal horn, deep layers	+	+
Dorsal horn, superior layers (I,II,III)	++	+++
Intermediate zone	+	+++
Intermediolateral column	+	+++
Ventral horn	+	+
<b>Medulla oblongata</b>		
Ambiguous nucleus	+	+
Area postrema	++++	++++
Cochlear nuclei	++	+
Cuneate nucleus, caudal division	+	++
Dorsal motor vagus nucleus	+++	++
Gracile nucleus, caudal division	-	-
Hypoglossal nucleus	-	-
Inferior central raphe nucleus	+	+
Inferior olive (principal nucleus, medial accessory nucleus, dorsal accessory nucleus)	+	+
Inferior vestibular nucleus	+	++
Lateral reticular nucleus	-	-
Lateral tegmental field	++	+++
Medial tegmental field	+	+
Nucleus raphe magnus	+	+
Nucleus raphe pallidus	-	-
Praeposital hypoglossal nucleus	+	+
Preolivary nucleus	+	++++
Restiform body	-	-
Retroambiguous nucleus	+	+++
Retrotrapezoid nucleus	+++	+++
Solitary tract, medial and lateral nucleus	++	++
Spinal trigeminal nucleus	++	++
Superior olive, lateral and medial nucleus	-	-
Trapezoid body nucleus	-	-
Ventral horn of the caudal brainstem	+	+
Vestibular complex	-	-
<b>Pons and cerebellum</b>		
Central linear nucleus of the raphe	+++	+++
Cerebellum	-	-
Facial nucleus	-	-
Kölliker-Fuse nucleus	++	++
Motor trigeminal nucleus	+	+
Parabrachial nucleus	++	++
Pontine continence center (L-region)	-	-
Pontine micturition center (M-region)	-	-
Superior central raphe nucleus	-	-

(Continues)

TABLE 1 (Continued)

Region	Distribution of AR-IR nuclei	Intensity of nuclear AR labeling
Mesencephalon		
Brachial nucleus of the colliculus inferior	+	+
Central tegmental field	++	+++
Dorsal raphe nucleus	++++	+++
Edinger Westphal nucleus	++	+++
Inferior colliculus	-	-
Medial pretecal area	++	++
Mesencephalic tegmental field	++	+++
Oculomotor nucleus	+	+
Periaqueductal gray (PAG)	++++	++
Red nucleus	-	-
Retrorubral nucleus	+++	+++
Substantia nigra, reticular and compact division	+++	+++
Superior colliculus	++	++
Trochlear nucleus	-	-
Ventral tegmental area of Tsai	+++	++

The table shows the density of the distribution of AR-IR nuclei and the intensity of nuclear AR-IR in the spinal cord and brainstem. The density and intensity were defined as: low (+), medium (++), high (+++), and very high (++++). The description includes the spinal cord, the medulla oblongata, the pons, the cerebellum, and the mesencephalon. Some regions are located in more than one brainstem region (e.g. trigeminal tract). These regions are only referred to in the table in the brainstem region in which the most prominent labeling was found.




FIGURE 2 AR-IR nuclei in the spinal cord. Bright field photomicrographs to illustrate AR-IR labeling in the spinal cord. Enlargement (a):  $\times 10$ ; (b):  $\times 20$ . (a) and (b) The intermediate zone, lamina X and the central canal of spinal segment L7. X, laminae of Rexed

(Figure 3: P7.5, P14.0). A less dense distribution of AR-IR nuclei was found in the cochlear nuclei, the lateral tegmental field, the nucleus of the solitary tract, and the spinal trigeminal nucleus (Figure 3: P7.5-P14.0). AR-IR nuclei in the spinal trigeminal nucleus were mainly located in the outer layer of the pars caudalis. A sparse distribution of AR-IR nuclei was found in the ambiguous nucleus, the cuneate nucleus, the inferior central raphe nucleus, the inferior vestibular nucleus, the medial tegmental field, the nucleus raphe magnus, the praepostial hypoglossal nucleus, the preolivary nucleus, the principal, medial and dorsal accessory inferior olive nucleus, the retroambiguus nucleus and the ventral horn of the caudal brainstem (Figure 3: P3.2-P15.7). The gracile nucleus, the hypoglossal nucleus, the lateral reticular nucleus, the nucleus raphe pallidus, the restiform body, the superior olive, the trapezoid body nucleus,


and the vestibular complex did not contain AR-IR nuclei (Figure 3: P3.2-P15.7).

A high intensity of nuclear AR-IR was observed in the area postrema, the lateral tegmental field, the preolivary nucleus, the retroambiguus nucleus, and the retrotrapezoid nucleus.

### 3.3 | Pons and cerebellum

The central linear nucleus of the raphe had a dense distribution of AR-IR nuclei (Figure 3: P3.2). A less dense distribution was observed in the parabrachial nucleus. The motor trigeminal nucleus showed a sparse distribution of AR-IR nuclei (Figure 3: P3.2). No other areas of the pons, such as the pontine micturition and continence centers,





**FIGURE 3** AR-IR nuclei in the brainstem. Frontal brainstem sections, rostral [A6.1] to caudal [P15.7]. A6.1 corresponds with 6.1 mm anterior (A) and P15.7 corresponds with 15.7 mm posterior (P) to the frontal zero (interaural) plane. Each dot represents one AR-IR neuron. The coordinates refer to corresponding coordinates in Berman's atlas (Berman, 1968). III, oculomotor nucleus; VII, facial nucleus; XII, hypoglossal nucleus; AMB, ambiguus nucleus; AP, area postrema; BC, brachium conjunctivum; BIN, brachial nucleus of the colliculus inferior; BP, brachium pontis; CB, cerebellum; CI, inferior central nucleus; CN, cochlear nuclei; CS, superior central nucleus; CUC, cuneate nucleus, caudal division; CX, external cuneate nucleus; DmX, dorsal motor vagus nucleus; DRN, dorsal raphe nucleus; EW, Edinger Westphal nucleus; FTC, central tegmental field; GRR, gracile nucleus, rostral division; ICX, external nucleus of the inferior colliculus; IN, nucleus interpositus; IO, inferior olive; KF, Kölliker-Fuse nucleus; LC, central linear nucleus of the raphe; LLV, ventral nucleus of the lateral lemniscus; LR, lateral reticular nucleus; LTF, lateral tegmental field; MGP, principal nucleus of the medial geniculate body; MP, medial parabrachial nucleus; MV, motor trigeminal nucleus; P, pyramidal tract; PAG, periaqueductal gray; PC, cerebral peduncle; PH, praeposital hypoglossal nucleus; PR, paramedian reticular nucleus; PUL, pulvinar; RN, red nucleus; RZN, retrotrapezoid nucleus; SC, superior colliculus; SM, medial nucleus of the solitary tract; SN, substantia nigra; SNC, substantia nigra compact division; SpV, spinal trigeminal nucleus; StV, spinal trigeminal tract; TB, trapezoid body; TN, nucleus of the trapezoid body; VIN, inferior vestibular nucleus; VLD, lateral vestibular nucleus; VM, medial vestibular nucleus; VTA, ventral tegmental area of Tsai

the facial nucleus and areas of the cerebellum contained AR-IR nuclei (Figure 3: P3.2, P7.5, Figure 4).

A high intensity of nuclear staining was observed in the central linear nucleus of the raphe. A slightly lower intensity of AR-IR staining was observed in the parabrachial nucleus. The intensity of AR-IR labeling was low in the motor trigeminal nucleus.

### 3.4 | Mesencephalon


The distribution of AR-IR nuclei was dense in the dorsal raphe nucleus, the PAG, the retrorubral nucleus, the substantia nigra (compact and reticular division), and the ventral tegmental area of Tsai (Figure 3: A6.1-A0.6, Figure 4). The PAG showed numerous AR-IR nuclei, throughout all its parts, except for the most central part directly adjacent to the ependymal lining, which did not contain AR-IR nuclei (Figure 3: A6.1-A0.6, Figure 4). Less dense labeling was found in the central tegmental field, the Edinger-Westphal nucleus, the medial pretecal area, the mesencephalic tegmental field, and the superior colliculus (Figure 3: A6.1-A0.6). A sparse distribution of AR-IR nuclei was observed in the brachial nucleus of the inferior colliculus and the oculomotor nucleus

(Figure 3: A6.1-A0.6). AR-IR nuclei were absent in the inferior colliculus, the red nucleus, and the trochlear nucleus (Figure 3: A4.1, A0.6).

Intensely stained nuclei were observed in the central tegmental field, the dorsal raphe nucleus, the Edinger Westphal nucleus, the mesencephalic tegmental field, the retrorubral nucleus, and the substantia nigra. The intensity of nuclear AR-IR staining was less in the brachial nucleus of the colliculus inferior, the medial pretecal area, the oculomotor nucleus, the PAG, the superior colliculus, and the ventral tegmental area of Tsai.

## 4 | DISCUSSION

This study provides the distribution of AR-IR nuclei in the spinal cord and brainstem of the cat. The immunohistochemical results overlap with those of earlier studies in other species such as the rat. Here, we will focus on novel findings and on the overall significance of the distribution of AR-IR nuclei. We firstly describe the similarities and differences with the distribution of androgen receptors as described in other species. Secondly, we discuss the involvement of the areas that contain AR-IR nuclei in autonomic function and dysfunction. And thirdly, the



**FIGURE 4** AR-IR nuclei in the brainstem. Bright field photomicrographs to illustrate AR-IR nuclei in the brainstem. Enlargement (a, c, e, g):  $\times 10$ ; (b, d, f, h):  $\times 20$ . (a) AR labeling in the dorsal raphe nucleus. (b) Enlargement of AR-IR nuclei in the dorsal raphe nucleus. (c) and (d) AR-IR nuclei in the substantia nigra. (e) and (f) AR-IR nuclei in the PAG. (g) and (h) Absence of AR-IR nuclei in the cerebellar cortex. DR, dorsal raphe nucleus; GC, granule cell layer; ML, molecular layer; PAG, periaqueductal gray; SN, substantia nigra; WM, white matter

distribution of androgen receptors in the cat is compared to that of estrogen receptors in the spinal cord and the brainstem.

#### 4.1 | Distribution of androgen receptors in the spinal cord and brainstem of the cat compared to that in other species

The density of AR-IR nuclei in the spinal cord of the cat was moderate. A relatively dense distribution of AR-IR nuclei was observed

in the superior layers of the dorsal horn such as in layer I, in which primary afferent fibers terminate. Nuclei in the intermediate zone, lamina X, and the intermediolateral cell column were also AR-IR. The present study shows that the intermediomedial cell column contains AR-IR nuclei indicating that pre-ganglionic sympathetic and parasympathetic motoneurons might express androgen receptors. These pre-ganglionic motoneurons innervate the pelvic smooth musculature such as those in the bladder and bowel. Some of these AR-IR areas at the lumbosacral level integrate visceral and somatic afferent and efferent information of the pelvic and pudendal nerves (Roppolo

et al., 1985). The results obtained in the present study are consistent with those in studies in the rat (Lumbroso et al., 1996; Ranson et al., 2012). Studies investigating the expression of androgen receptors in the spinal cord of primates could not be found.

In the caudal brainstem, a dense distribution of AR-IR nuclei was found in the area postrema, which is consistent with previous results in the rat (Hamson et al., 2004). The area postrema regulates various autonomic functions such as the control of respiration and blood pressure (Qian & Koon, 1998; Yang et al., 2006). The central linear nucleus of the raphe in the cat contained a dense distribution of AR-IR nuclei, these neurons are serotonin producing neurons that project to the forebrain. The central linear nucleus is hypothesized to play a role in the sleep-wake state of the organism (Trulson et al., 1981). AR-IR nuclei were additionally observed in the dorsal motor vagus nucleus, consistent with previous results in the mouse (Mukudai et al., 2016; Yoon et al., 1996). The retrotrapezoid nucleus, a regulator of respiration automaticity, also contained a dense distribution of AR-IR nuclei (Guyenet et al., 2019). The intensity of AR-IR staining was little to absent in inferior olive, the motor nuclei of some of the cranial nerves (V, VII, IX, and XII), and the vestibular complex. The literature on androgen receptor expression in the motor nuclei of the cranial nerves in rats is conflicting. Hamson et al. did not find AR-IR motoneurons in the dorsal vagus nucleus, the facial nucleus, the hypoglossal nucleus, and the motor trigeminal nucleus (Hamson et al., 2004). In contrast, another previous report in rats did show AR-IR in these motor nuclei (Yu & McGinnis, 2001). Hamson et al. did show AR-IR nuclei in the area postrema, the nucleus ambiguus, the parabrachial nucleus, the nucleus raphe magnus, and the nucleus of the solitary tract in the rat, which is consistent with the present study. In female rhesus monkeys, the uptake of dihydrotestosterone was studied and uptake was identified in the spinal trigeminal nucleus (Sheridan & Weaker, 1982). The present study in cats showed that AR-IR nuclei in the spinal trigeminal nucleus are mainly located in the outer layer of the pars caudalis, an area that is involved in nociception (Patel & Das, 2019). Therefore, we can conclude that higher mammals such as cats and monkeys seem to have a similar distribution of androgen receptors in the motor nuclei of the cranial nerves.

In the mesencephalon, we observed a dense distribution of AR-IR nuclei in the dorsal raphe nucleus in cats. The dorsal raphe nucleus is involved in the sleep-wake state of the organism and it encodes reward signals (Li et al., 2016; McGinty & Harper, 1976). AR-IR neurons in the dorsal raphe nucleus were previously described in male mice, rats and macaques (Bethea et al., 2015; Sheng et al., 2004). In the present study in cats, a high density of AR-IR nuclei was observed in the periphery but not in the center of the PAG (Blok & Holstege, 1994; Faull et al., 2019). In rodents, AR-IR neurons were mainly found in the dorsomedial and medial part of the PAG (Greco et al., 1996; Murphy et al., 1999; Simerly et al., 1990). The PAG regulates autonomic functions, such as cardiovascular control, micturition and respiration, in animals and humans. AR-IR cells in the PAG might modulate autonomic behavioral responses such as sexual and urogenital reflexes that originate in the brainstem in both animals and humans (Blok et al., 1997; Marson & McKenna, 1996;

Marson & Murphy, 2006; Michels et al., 2015). A dense distribution of AR-IR nuclei in the male cat was also observed in the retrorubral nucleus, the substantia nigra, and the ventral tegmental area of Tsai, which plays a role in motivation and reward processing (Morales & Margolis, 2017). Similar results were obtained in the rat using immunohistochemistry, autoradiography, and *in situ* hybridization (Kritzer, 1997; Sar & Stumpf, 1975; Simerly et al., 1990). The substantia nigra in the cat contained AR-IR nuclei in the reticular division as well as the compact division. In the rat, only the compact division showed AR-IR cells (Kritzer, 1997). In conclusion, AR-IR cells in the spinal cord, caudal brainstem, and the mesencephalon are highly conserved throughout species. A dense distribution of AR-IR nuclei was present in regions that are involved in autonomic functions such as cardiovascular function, urogenital function, gastro-intestinal function and respiratory function and in areas involved in processing of motivation and reward.

#### 4.2 | The involvement of androgen receptors in autonomic functions

Several areas with a high density of AR-IR nuclei are involved in respiration. The areas that were identified in the present study are the area postrema, the dorsal motor vagus nucleus, the Kölliker-Fuse nucleus, the nucleus of the solitary tract, the PAG, the parabrachial nucleus, and the retrotrapezoid nucleus. The area postrema has been implicated to alter cardiopulmonary responses in rats (Yang et al., 2006). Nerve fibers originating in the dorsal motor vagus nucleus innervate the muscles of the trachea and lower airways ensuring airway patency (Jordan, 2001). The Kölliker-Fuse nucleus orchestrates the timing of the expiration phase (Barnett et al., 2018; Dutschmann et al., 2021). The PAG plays an important role in the integration of inputs regarding multiple autonomic functions (Faull et al., 2019). Furthermore, the parabrachial nucleus is thought to mediate respiratory rate (Miller et al., 2017). The retrotrapezoid nucleus is a regulator of breathing automaticity (Guyenet et al., 2019). The involvement of androgens is additionally supported by evidence that testosterone replacement in men with sleep apnea alters ventilation responses (Burschtin & Wang, 2016).

The areas involved in cardiovascular function that contained AR-IR nuclei in the present study are: the area postrema, the dorsal motor vagus nucleus, the nucleus of the solitary tract, and the PAG. The area postrema and the nucleus ambiguus have been implicated to control heart rate (Gourine et al., 2016; Xue et al., 2003; Yang et al., 2006). The dorsal motor vagus nucleus controls the excitability of the ventricles of the heart (Gourine et al., 2016). Additionally, in rats it has been shown that the PAG is involved in the modulation of heart rate and blood pressure (Lagatta et al., 2016).

Furthermore, androgens play an important role in sexual function. Here, we described areas in the central nervous system that might be androgen sensitive indicated by a high density of AR-IR nuclei. We showed a high density of AR-IR nuclei in areas involved



in sexual reflexes such as erection and orgasm. These areas include the PAG and the dorsal horn of the sacral spinal cord. The nucleus retroambiguus also contained AR-IR nuclei. It has connections to the PAG and the lumbosacral spinal cord and is thought to be involved in mating behavior. The involvement of androgens in sexual function is supported by evidence that men with a decreased level of testosterone can experience a decrease in sexual desire and arousal, erectile dysfunction, reduced ejaculate, incontinence on orgasm, and reduced orgasm (Basaria et al., 2002; Casey et al., 2012; McGinty et al., 2014; Yeap, 2014). Decreased sexual desire has mostly been attributed to the effect of androgen deprivation in the telencephalon. However, this side effect can also possibly be attributed to an altered sensation of the external genitalia due to decreased androgen receptor activation in the dorsal horn of the sacral spinal cord. We found a relatively dense distribution of AR-IR nuclei in the superficial layers of the dorsal horn, where sensory afferents from areas such as the genitals are located. Similarly, the effects could be mediated by other central nervous system areas with AR-IR nuclei such as the PAG.

#### 4.3 | The distribution of androgen receptors compared to that of estrogen receptors

The distribution of estrogen receptors in the central nervous system is comparable between male and female rodents (Vanderhorst et al., 2005; Zhang et al., 2002). Also the androgen receptor distribution between male and female rodents is similar. However, it is not known whether the distributions of androgen and estrogen receptors overlap. A previous study described the estrogen receptor distribution in the spinal cord and brainstem of ovariectomized female cats. Remarkable overlap with the distribution of androgen receptors in the male cat exists. Estrogen receptors were detected in laminae I, II, V, VII, X, and in sacral pre-ganglionic parasympathetic cells (VanderHorst et al., 2001). This is consistent with the distribution of androgen receptors in the male cat described in the present study. In the caudal brainstem of the female cat, estrogen receptors were present in the area postrema, the parabrachial nucleus, the solitary tract, and the outer layers of the pars caudalis of the spinal trigeminal nucleus (Boers, 2005). In the mesencephalon, estrogen receptors were observed in the brachial nucleus of the colliculus inferior, the dorsal raphe, the PAG, the superior colliculus, and the tegmentum (Boers, 2005). In ovariectomized female rhesus monkeys, a partly overlapping distribution of estrogen receptors was found (Vanderhorst et al., 2009). Estrogen receptors were present in laminae I-V and X of the spinal cord, the pars caudalis of the spinal trigeminal nucleus, the solitary tract and the tegmentum in the caudal brainstem. In the mesencephalon, estrogen receptors were observed in the PAG and the parabrachial nucleus (Vanderhorst et al., 2009). Despite the great overlap between the estrogen receptor distribution in the cat and the monkey, estrogen receptor expression in the monkey was less abundant and less

widespread than that in the cat. Whether this pattern is also observed in the distribution of androgen receptors in male primates is not known.

## 5 | CONCLUSION

In this study, we showed specific areas in the spinal cord and brainstem of the cat in which AR-IR nuclei are present. In the spinal cord of the cat the most dense distribution of AR-IR nuclei was observed in the dorsal horn, which is involved in sensory processing. However, androgen receptor expression was more pronounced in supraspinal areas. AR-IR nuclei in the brainstem of the cat were most prominent in the area postrema, the central linear nucleus, the dorsal motor vagus nucleus, the dorsal raphe nucleus, the PAG, the retrorubral nucleus, the retrotrapezoid nucleus, the substantia nigra, and the ventral tegmental area of Tsai. These areas are important for various autonomic functions such as cardiovascular function, micturition, respiration, and sexual function. These results underline the importance of androgen receptors in the central nervous system and their possible roles in sensory and autonomic functions. For clinical practice this means that patients with an altered availability of androgens or their receptors as is the case in hypogonadism, might experience autonomic dysfunction due to a decreased activation of androgen receptors in the spinal cord and brainstem.

## ACKNOWLEDGMENTS

We thank Dr. Gail S. Prins for her generous donation of the PG21 anti-androgen receptor antibody. We also thank Dr. Jos Th. P. W. van Maarseveen for the NL2EPS Converter program that he wrote. We are grateful to Mr. P. van der Syde for photographic assistance, and Mr. K. van Linschoten and Ms. E. Meijer for their histotechnical work (Department of Human Movement Sciences, Faculty of Medical Sciences Groningen).

## CONFLICT OF INTEREST

The authors declare no conflicts of interest.

## AUTHORS' CONTRIBUTIONS

RC contributed to data analysis/interpretation, drafting of the manuscript, and critical revision of the manuscript. JCC contributed to acquisition of data, data analysis/interpretation, and critical revision of the manuscript. PS and EA contributed to data analysis/interpretation and critical revision of the manuscript. BFMB contributed to the concept and design of the study, acquisition of data, data analysis/interpretation, and critical revision of the manuscript.

## DATA AVAILABILITY STATEMENT

Data available on request from the authors.

## ORCID

Rosa L. Coolen  <https://orcid.org/0000-0002-7229-4867>

## REFERENCES

- Avendaño, C. & Verdu, A. (1992) Area 3a in the cat. I. A reevaluation of its location and architecture on the basis of Nissl, myelin, acetylcholinesterase, and cytochrome oxidase staining. *Journal of Comparative Neurology*, 321, 357–372.
- Barnett, W.H., Jenkin, S.E.M., Milsom, W.K., Paton, J.F.R., Abdala, A.P., Molkov, Y.I., et al. (2018) The Kölliker-Fuse nucleus orchestrates the timing of expiratory abdominal nerve bursting. *Journal of Neurophysiology*, 119, 401–412.
- Basaria, S., Lieb, J., Tang, A.M., DeWeese, T., Carducci, M., Eisenberger, M., et al. (2002) Long-term effects of androgen deprivation therapy in prostate cancer patients. *Clinical Endocrinology*, 56, 779–786.
- Berman, A.L. (1968) The brain stem of the cat. *A cytoarchitectonic atlas with stereotaxic coordinates*, 1–175.
- Bethea, C.L., Phu, K., Belikova, Y. & Bethea, S.C. (2015) Localization and regulation of reproductive steroid receptors in the raphe serotonin system of male macaques. *Journal of Chemical Neuroanatomy*, 66, 19–27.
- Blok, B.F.M. & Holstege, G. (1994) Direct projections from the periaqueductal gray to the pontine micturition center (M-region). An anterograde and retrograde tracing study in the cat. *Neuroscience Letters*, 166, 93–96.
- Blok, B.F., Willemsen, A.T. & Holstege, G. (1997) A PET study on brain control of micturition in humans. *Brain*, 120(Pt 1), 111–121.
- Boers, J. (2005) *The role of the nucleus retroambiguus in the neural control of respiration, vocalization and mating behavior*. University Library Groningen.
- Bravo, G., Massa, H., Rose'meyer, R., Chess-Williams, R., McDermott, C. & Sellers, D.J. (2017) Effect of short-term androgen deficiency on bladder contractility and urothelial mediator release. *Naunyn Schmiedeberg's Archives of Pharmacology*, 390, 547–556.
- Brinkmann, A.O., Blok, L.J., De Ruiter, P.E., Doesburg, P., Steketeer, K., Berrevoets, C.A., et al. (1999) Mechanisms of androgen receptor activation and function. *The Journal of steroid biochemistry and molecular biology*, 69, 307–313.
- Brinkmann, A.O., Jenster, G., Ris-Stalpers, C., van der Korput, J.A., Brüggewirth, H.T., Boehmer, A.L., et al. (1995) Androgen receptor mutations. *The Journal of Steroid Biochemistry and Molecular Biology*, 53, 443–448.
- Burschtin, O. & Wang, J. (2016) Testosterone deficiency and sleep apnea. *Urologic Clinics of North America*, 43, 233–237.
- Carré, J.M. & Archer, J. (2018) Testosterone and human behavior: The role of individual and contextual variables. *Current Opinion in Psychology*, 19, 149–153.
- Casey, R.G., Corcoran, N.M. & Goldenberg, S.L. (2012) Quality of life issues in men undergoing androgen deprivation therapy: A review. *Asian Journal of Andrology*, 14, 226–231.
- Choate, J.V.A., Slayden, O.D. & Resko, J.A. (1998) Immunocytochemical localization of androgen receptors in brains of developing and adult male rhesus monkeys. *Endocrine*, 8, 51–60.
- Claessens, F., Denayer, S., Van Tilborgh, N., Kerkhofs, S., Helsen, C. & Haelens, A. (2008) Diverse roles of androgen receptor (AR) domains in AR-mediated signaling. *Nuclear Receptor Signaling*, 6, e06008.
- Clancy, A.N., Bonsall, R.W. & Michael, R.P. (1992) Immunohistochemical labeling of androgen receptors in the brain of rat and monkey. *Life Sciences*, 50, 409–417.
- Cunningham, R.L., Lumia, A.R. & McGinnis, M.Y. (2012) Androgen receptors, sex behavior, and aggression. *Neuroendocrinology*, 96, 131–140.
- Dohle, G.R., Arver, S., Bettocchi, C., Jones, T.H. & Klisch, S. (2019) *EAU Guidelines. Male hypogonadism*. European Association of Urology.
- Donovan, K.A., Gonzalez, B.D., Nelson, A.M., Fishman, M.N., Zachariah, B. & Jacobsen, P.B. (2018) Effect of androgen deprivation therapy on sexual function and bother in men with prostate cancer: A controlled comparison. *Psycho-oncology*, 27, 316–324.
- Dutschmann, M., Bautista, T.G., Trevizan-Baú, P., Dhingra, R.R. & Furuya, W.I. (2021) The pontine Kölliker-Fuse nucleus gates facial, hypoglossal, and vagal upper airway related motor activity. *Respiratory Physiology and Neurobiology*, 284, 103563.
- Faull, O.K., Subramanian, H.H., Ezra, M. & Pattinson, K.T.S. (2019) The midbrain periaqueductal gray as an integrative and interoceptive neural structure for breathing. *Neuroscience and Biobehavioral Reviews*, 98, 135–144.
- Gioeli, D. & Paschal, B.M. (2012) Post-translational modification of the androgen receptor. *Molecular and Cellular Endocrinology*, 352, 70–78.
- Gourine, A.V., Machhada, A., Trapp, S. & Spyer, K.M. (2016) Cardiac vagal preganglionic neurones: An update. *Autonomic Neuroscience*, 199, 24–28.
- Greco, B., Edwards, D.A., Michael, R.P. & Clancy, A.N. (1996) Androgen receptor immunoreactivity and mating-induced Fos expression in forebrain and midbrain structures in the male rat. *Neuroscience*, 75, 161–171.
- Guyenet, P.G., Stornetta, R.L., Souza, G., Abbott, S.B.G., Shi, Y. & Bayliss, D.A. (2019) The retrotrapezoid nucleus: Central chemoreceptor and regulator of breathing automaticity. *Trends in Neurosciences*, 42, 807–824.
- Hamson, D.K., Jones, B.A. & Watson, N.V. (2004) Distribution of androgen receptor immunoreactivity in the brainstem of male rats. *Neuroscience*, 127, 797–803.
- Hipkæo, W., Wakayama, T., Yamamoto, M. & Iseki, S. (2004) Expression and localization of the transcription factor JunD in the duct system of mouse submandibular gland. *Journal of Histochemistry and Cytochemistry*, 52, 479–490.
- Huang, Y.P., Liu, W., Chen, S.F., Liu, Y.D., Chen, B., Deng, C.H., et al. (2019) Free testosterone correlated with erectile dysfunction severity among young men with normal total testosterone. *International Journal of Impotence Research*, 31, 132–138.
- Hunter, I., Hay, C.W., Esswein, B., Watt, K. & Mcewan, I.J. (2018) Tissue control of androgen action: The ups and downs of androgen receptor expression. *Molecular and Cellular Endocrinology*, 465, 27–35.
- Jordan, D. (2001) Central nervous pathways and control of the airways. *Respiration Physiology*, 125, 67–81.
- Kawano, H., Sato, T., Yamada, T., Matsumoto, T., Sekine, K., Watanabe, T., et al. (2003) Suppressive function of androgen receptor in bone resorption. *Proceedings of the National Academy of Sciences*, 100, 9416–9421.
- Kritzer, M.F. (1997) Selective colocalization of immunoreactivity for intracellular gonadal hormone receptors and tyrosine hydroxylase in the ventral tegmental area, substantia nigra, and retrorubral fields in the rat. *Journal of Comparative Neurology*, 379, 247–260.
- Lagatta, D.C., Ferreira-Junior, N.C., Deolindo, M., Corrêa, F.M. & Resstel, L.B. (2016) Ventrolateral periaqueductal grey matter neurotransmission modulates cardiac baroreflex activity. *European Journal of Neuroscience*, 44, 2877–2884.
- Li, Y., Zhong, W., Wang, D., Feng, Q., Liu, Z., Zhou, J., et al. (2016) Serotonin neurons in the dorsal raphe nucleus encode reward signals. *Nature Communications*, 7, 10503.
- Lumbroso, S., Sandillon, F., Georget, V., Lobaccaro, J.M., Brinkmann, A.O., Privat, A., et al. (1996) Immunohistochemical localization and immunoblotting of androgen receptor in spinal neurons of male and female rats. *European Journal of Endocrinology*, 134, 626–632.
- Marson, L. & McKenna, K.E. (1996) CNS cell groups involved in the control of the ischiocavernosus and bulbospongiosus muscles: A transneuronal tracing study using pseudorabies virus. *Journal of Comparative Neurology*, 374, 161–179.
- Marson, L. & Murphy, A.Z. (2006) Identification of neural circuits involved in female genital responses in the rat: A dual virus and anterograde tracing study. *American Journal of Physiology Regulatory, Integrative and Comparative Physiology*, 291, R419–R428.

- McGinty, D.J. & Harper, R.M. (1976) Dorsal raphe neurons: Depression of firing during sleep in cats. *Brain Research*, 101, 569–575.
- McGinty, H.L., Phillips, K.M., Jim, H.S.L., Cessna, J.M., Asvat, Y., Cases, M.G., et al. (2014) Cognitive functioning in men receiving androgen deprivation therapy for prostate cancer: A systematic review and meta-analysis. *Supportive Care in Cancer*, 22, 2271–2280.
- Michael, R.P., Rees, H.D. & Bonsall, R.W. (1989) Sites in the male primate brain at which testosterone acts as an androgen. *Brain Research*, 502, 11–20.
- Michels, L., Blok, B.F., Gregorini, F., Kurz, M., Schurch, B., Kessler, T.M., et al. (2015) Supraspinal control of urine storage and micturition in men—An fMRI study. *Cerebral Cortex*, 25, 3369–3380.
- Miller, J.R., Zuperku, E.J., Stuth, E.A.E., Banerjee, A., Hopp, F.A. & Stucke, A.G. (2017) A subregion of the parabrachial nucleus partially mediates respiratory rate depression from intravenous remifentanyl in young and adult rabbits. *Anesthesiology*, 127, 502–514.
- Morales, M. & Margolis, E.B. (2017) Ventral tegmental area: cellular heterogeneity, connectivity and behaviour. *Nature Reviews Neuroscience*, 18, 73–85.
- Mukudai, S., Ichi Matsuda, K., Bando, H., Takanami, K., Nishio, T., Sugiyama, Y., et al. (2016) Expression of sex steroid hormone receptors in vagal motor neurons innervating the trachea and esophagus in mouse. *Acta Histochemica et Cytochemica*, 49, 37–46.
- Murphy, A.Z., Shupnik, M.A. & Hoffman, G.E. (1999) Androgen and estrogen ( $\alpha$ ) receptor distribution in the periaqueductal gray of the male rat. *Hormones and Behavior*, 36, 98–108.
- Nead, K.T., Gaskin, G., Chester, C., Swisher-McClure, S., Leeper, N.J. & Shah, N.H. (2017) Association between androgen deprivation therapy and risk of dementia. *JAMA Oncology*, 3, 49–55.
- O'Bryant, E.L. & Jordan, C.L. (2005) Expression of nuclear receptor co-activators in androgen-responsive and-unresponsive motoneurons. *Hormones and Behavior*, 47, 29–38.
- Ophoff, J., Van Proeyen, K., Callewaert, F., De Gendt, K., De Bock, K., Vanden Bosch, A., et al. (2009) Androgen signaling in myocytes contributes to the maintenance of muscle mass and fiber type regulation but not to muscle strength or fatigue. *Endocrinology*, 150, 3558–3566.
- Patel, N.M. & Das, J.M. (2019) *Neuroanatomy, spinal trigeminal nucleus*. StatPearls Publishing.
- Prins, G.S., Birch, L. & Greene, G.L. (1991) Androgen receptor localization in different cell types of the adult rat prostate. *Endocrinology*, 129, 3187–3199.
- Qian, Z.M. & Koon, H.W. (1998) Area postrema is essential for the maintenance of normal blood pressure under cold stress in rats. *Experimental Brain Research*, 121, 186–190.
- Rana, K., Fam, B.C., Clarke, M.V., Pang, T.P., Zajac, J.D. & MacLean, H.E. (2011) Increased adiposity in DNA binding-dependent androgen receptor knockout male mice associated with decreased voluntary activity and not insulin resistance. *American Journal of Physiology-Endocrinology and Metabolism*, 301, E767–E778.
- Ranson, R.N., Connelly, J.H., Santer, R.M. & Watson, A.H.D. (2012) Nuclear expression of PG-21, SRC-1, and pCREB in regions of the lumbosacral spinal cord involved in pelvic innervation in young adult and aged rats. *Anatomy and Cell Biology*, 45, 241–258.
- Roppolo, J.R., Nadelhaft, I. & De Groat, W.C. (1985) The organization of pudendal motoneurons and primary afferent projections in the spinal cord of the rhesus monkey revealed by horseradish peroxidase. *Journal of Comparative Neurology*, 234, 475–488.
- Santi, D., Spaggiari, G., Gilioli, L., Poti, F., Simoni, M. & Casarini, L. (2018) Molecular basis of androgen action on human sexual desire. *Molecular and Cellular Endocrinology*, 467, 31–41.
- Sar, M. & Stumpf, W.E. (1975) Distribution of androgen-concentrating neurons in rat brain. In W. E. Stumpf & L. D. Grant (Eds.), *Anatomical Neuroendocrinology* (pp. 120–133). Karger Publishers.
- Sheng, Z., Kawano, J., Yanai, A., Fujinaga, R., Tanaka, M., Watanabe, Y., et al. (2004) Expression of estrogen receptors ( $\alpha$ ,  $\beta$ ) and androgen receptor in serotonin neurons of the rat and mouse dorsal raphe nuclei; sex and species differences. *Neuroscience Research*, 49, 185–196.
- Sheridan, P.J. & Weaker, F.J. (1982) Androgen receptor systems in the brain stem of the primate. *Brain Research*, 235, 225–232.
- Simerly, R.B., Swanson, L.W., Chang, C. & Muramatsu, M. (1990) Distribution of androgen and estrogen receptor mRNA-containing cells in the rat brain: An in situ hybridization study. *Journal of Comparative Neurology*, 294, 76–95.
- Takov, K., Wu, J., Denvir, M.A., Smith, L.B. & Hadoke, P.W.F. (2018) The role of androgen receptors in atherosclerosis. *Molecular and Cellular Endocrinology*, 465, 82–91.
- Traish, A.M., Melcangi, R.C., Bortolato, M., Garcia-Segura, L.M. & Zitzmann, M. (2015) Adverse effects of 5 $\alpha$ -reductase inhibitors: What do we know, don't know, and need to know? *Reviews in Endocrine and Metabolic Disorders*, 16, 177–198.
- Trulson, M.E., Jacobs, B.L. & Morrison, A.R. (1981) Raphe unit activity during REM sleep in normal cats and in pontine lesioned cats displaying REM sleep without atonia. *Brain Research*, 226, 75–91.
- Vanderhorst, V.G., Gustafsson, J.A. & Ulfhake, B. (2005) Estrogen receptor-alpha and -beta immunoreactive neurons in the brainstem and spinal cord of male and female mice: Relationships to monoaminergic, cholinergic, and spinal projection systems. *Journal of Comparative Neurology*, 488, 152–179.
- Vanderhorst, V.G., Meijer, E. & Holstege, G. (2001) Estrogen receptor-alpha immunoreactivity in parasympathetic preganglionic neurons innervating the bladder in the adult ovariectomized cat. *Neuroscience Letters*, 298, 147–150.
- Vanderhorst, V.G., Terasawa, E. & Ralston, H.J. 3rd (2009) Estrogen receptor-alpha immunoreactive neurons in the brainstem and spinal cord of the female rhesus monkey: Species-specific characteristics. *Neuroscience*, 158, 798–810.
- Xue, B., Gole, H., Pamidimukkala, J. & Hay, M. (2003) Role of the area postrema in angiotensin II modulation of baroreflex control of heart rate in conscious mice. *American Journal of Physiology - Heart and Circulatory Physiology*, 284, H1003–H1007.
- Yang, S.J., Lee, K.Z., Wu, C.H., Lu, K.T. & Hwang, J.C. (2006) Vasopressin produces inhibition on phrenic nerve activity and apnea through V(1A) receptors in the area postrema in rats. *Chinese Journal of Physiology*, 49, 313–325.
- Yeap, B.B. (2014) Hormonal changes and their impact on cognition and mental health of ageing men. *Maturitas*, 79, 227–235.
- Yoon, S.H., Sim, S.S., Hahn, S.J., Rhie, D.J., Jo, Y.H. & Kim, M.S. (1996) Stimulatory role of the dorsal motor nucleus of the vagus in gastrointestinal motility through myoelectromechanical coordination in cats. *Journal of the Autonomic Nervous System*, 57, 22–28.
- Yu, W.H.A. & McGinnis, M.Y. (2001) Androgen receptors in cranial nerve motor nuclei of male and female rats. *Journal of Neurobiology*, 46, 1–10.
- Zhang, J.Q., Cai, W.Q., Zhou, D.S. & Su, B.Y. (2002) Distribution and differences of estrogen receptor beta immunoreactivity in the brain of adult male and female rats. *Brain Research*, 935, 73–80.

**How to cite this article:** Coolen RL, Cambier JC, Spantidea PI, Asselt E, Blok BF. Androgen receptors in areas of the spinal cord and brainstem: A study in adult male cats. *J Anat*. 2021;00:1–11. <https://doi.org/10.1111/joa.13407>