

## Supplementary Material: ECG Scoring for the Evaluation of Therapy-Naïve Cancer Patients to Predict Cardiotoxicity

Julia Pohl, Raluca-Ileana Mincu, Simone M. Mrotzek, Reza Wakili, Amir A. Mahabadi, Sophia K. Potthoff, Jens T. Siveke, Ulrich Keller, Ulf Landmesser, Tienush Rassaf, Markus S. Anker, Matthias Totzeck

**Table S1.** In- and exclusion criteria.

Inclusion Criteria	Exclusion Criteria
- Presentation in cardio-oncology before start of anti-cancer therapy	- Prior cancerous condition with anti-cancer therapy
- Presentation in cardio-oncology for a follow-up routine	

**Table S2.** Applied anti-tumor therapies.

Applied Anti-Tumor Therapy	% of the Patients
Anthracyclines (%)	27%
Doxorubicin (mean dose (mg/m <sup>2</sup> ))	209
Epirubicin (mean dose (mg/m <sup>2</sup> ))	318
Combination therapy with alkylating agents	92%
Antimicrotubule agents (%)	14%
Platin-containing therapies (%)	4%
Antimetabolites (%)	1%
Topoisomerase II inhibitors (%)	13%
Tyrosine kinase inhibitors (%)	3%
Immune checkpoint inhibitors (%)	41%
BRAF/MEK inhibitors (%)	7%
Anti-Her2 compounds (%)	4%
Others (%)	16%