

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

# Resuscitation

journal homepage: [www.elsevier.com/locate/resuscitation](http://www.elsevier.com/locate/resuscitation)

## European Resuscitation Council Guidelines 2021: Ethics of resuscitation and end of life decisions



Spyros D. Mentzelopoulos<sup>a,\*</sup>, Keith Couper<sup>b,c</sup>, Patrick Van de Voorde<sup>d,e</sup>,  
Patrick Druwé<sup>f</sup>, Marieke Blom<sup>g</sup>, Gavin D. Perkins<sup>b</sup>, Ileana Lulic<sup>h</sup>, Jana Djakow<sup>i,j</sup>,  
Violetta Raffay<sup>k,l</sup>, Gisela Lilja<sup>m</sup>, Leo Bossaert<sup>n</sup>

<sup>a</sup> National and Kapodistrian University of Athens Medical School, Athens, Greece

<sup>b</sup> UK Critical Care Unit, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

<sup>c</sup> Warwick Medical School, University of Warwick, Coventry, UK

<sup>d</sup> University Hospital and University Ghent, Belgium

<sup>e</sup> Federal Department Health, Belgium

<sup>f</sup> Ghent University Hospital, Department of Intensive Care Medicine, Ghent, Belgium

<sup>g</sup> Amsterdam University Medical Center, Amsterdam, The Netherlands

<sup>h</sup> Clinical Hospital Merkur, Zagreb, Croatia

<sup>i</sup> Paediatric Intensive Care Unit, NH Hospital, Hořovice, Czech Republic

<sup>j</sup> Department of Paediatric Anaesthesiology and Intensive Care Medicine, University Hospital and Medical Faculty of Masaryk University, Brno, Czech Republic

<sup>k</sup> European University Cyprus, School of Medicine, Nicosia, Cyprus

<sup>l</sup> Serbian Resuscitation Council, Novi Sad, Serbia

<sup>m</sup> Lund University, Skane University Hospital, Department of Clinical Sciences Lund, Neurology, Lund, Sweden

<sup>n</sup> University of Antwerp, Antwerp, Belgium

### Abstract

These European Resuscitation Council Ethics guidelines provide evidence-based recommendations for the ethical, routine practice of resuscitation and end-of-life care of adults and children. The guideline primarily focus on major ethical practice interventions (i.e. advance directives, advance care planning, and shared decision making), decision making regarding resuscitation, education, and research. These areas are tightly related to the application of the principles of bioethics in the practice of resuscitation and end-of-life care.

### Introduction and scope

The purpose of the current European Resuscitation Council Guideline chapter is to provide evidence-based recommendations for the ethical, routine practice of resuscitation and end-of-life care of adults and children. This means maximising the benefit of life-sustaining treatments, while concurrently preventing pertinent harm, and promoting equitable access to best-quality resuscitation care. The chapter should be read in conjunction with other chapters that focus on specific relevant topics; information on e.g. epidemiology, education,

post-resuscitation care and on the ethics of resuscitation of newly born babies (transition at birth), can be found in the dedicated chapters within these guidelines.

We primarily focus on major ethical practice interventions (i.e. advance directives, advance care planning, and shared decision making), decision making regarding resuscitation, education, and research. These areas are tightly related to the application of the principles of bioethics in the practice of resuscitation and end-of-life care. Consensus definitions of core bioethical principles and relevant key terms are included in the online supplement.

\* Corresponding author at: Department of Intensive Care Medicine University of Athens Medical School, Evaggelismos General Hospital, 45-47 Ipsilandou Street, GR-10675 Athens, Greece.

E-mail addresses: [sdmentzelopoulos@yahoo.com](mailto:sdmentzelopoulos@yahoo.com), [sdmentzelopoulos@gmail.com](mailto:sdmentzelopoulos@gmail.com) (S.D. Mentzelopoulos).  
<https://doi.org/10.1016/j.resuscitation.2021.02.017>

We also refer to emerging ethical challenges that resulted from the societal and legal changes associated with the coronavirus disease-19 (COVID-19) pandemic.<sup>1</sup> These changes include new social norms (e.g. social distancing), potential exacerbation of healthcare inequalities, and dissemination of public health misinformation.<sup>1,2</sup>

The chapter content is based on scoping reviews of 22 research questions, and expert opinion. Reviews were originally completed in 2019. Due to the COVID-19 crisis and the postponement of the publication of the 2020 guidelines, we updated each scoping review in mid-2020. Full details of each scoping review, including search strategies, included studies, and study findings are included in the electronic supplement.

Key messages from this chapter are summarised in Fig. 1.

These guidelines were drafted and agreed by the Ethics Writing Group members. The methodology used for guideline development is presented in the Executive summary.<sup>2a</sup> The guidelines were posted for public comment in October 2020. The feedback was reviewed by

the writing group and the guidelines was updated where relevant. The Guidelines were presented to and approved by the ERC General Assembly on 10th December 2020.

## Concise guidelines for clinical practice

### Major interventions aimed at safeguarding autonomy

#### Patient preferences and treatment decisions

Clinicians should:

- Use advance care planning that incorporates shared decision making to improve consistency between patient wishes and treatment.
- Offer advance care planning to all patients at increased risk of cardiac arrest or poor outcome in the event of cardiac arrest.
- Support advance care planning in all cases where it is requested by the patient.



**Fig. 1 – Key messages relating to ethics in Guidelines 2021.**

- Record advance care plans in a consistent manner (e.g. electronic registries, documentation templates etc.).
- Integrate resuscitation decisions with other treatment decisions, such as invasive mechanical ventilation, in overarching advance emergency care treatment plans to increase clarity of treatment goals and prevent inadvertent deprivation of other indicated treatments.
- Clinicians should not offer CPR in cases where resuscitation would be futile.

### Improving communication

- Clinicians should use evidence-based communication interventions to improve end-of-life discussions and support completion of advance directives/advance care plans.
- Clinicians should combine structured end-of-life discussions with video decision aids for shared decision making about end-of-life hospital transfer from nursing homes in systems where this technology is available.
- Clinicians should consider inviting a communication facilitator to join discussions with patients and/or their family when making advance care plans about the appropriateness of life sustaining treatments. This refers to systems where communication facilitators are available.
- Healthcare systems should provide clinicians with communication skills training interventions to improve clinicians' skill and comfort in delivering bad news or supporting patients to define care goals.
- Clinicians should integrate the following patient/family support elements with shared decision making:
  1. Provide information about the patient's status and prognosis in a clear and honest manner. This may be supported by use of a video-support tool.
  2. Seek information about the patient's goals, values, and treatment preferences.
  3. Involve patients/family members in discussions about advance care plans.
  4. Provide empathic statements assuring non-abandonment, symptom control, and decision-making support.
  5. Provide the option of spiritual support.
  6. Where appropriate, explain and apply protocolised patient-centred procedures for treatment withdrawal with concurrent symptom control and patient/family psychological support.
  7. Consider recording meetings with family for the purpose of audit/quality improvement.

### Deciding when to start and when to stop cardiopulmonary resuscitation (CPR)

#### Withholding and Withdrawing CPR

- Systems, clinicians, and the public should consider cardiopulmonary resuscitation (CPR) a conditional therapy.
- Systems should implement criteria for the withholding and termination of CPR for both in-hospital cardiac arrest (IHCA) and out-of-hospital cardiac arrest (OHCA), taking into consideration the specific local legal, organisational, and cultural context.
- Systems should define criteria for the withholding and termination of CPR, and ensure criteria are validated locally. The following criteria may be considered:

- Unequivocal criteria:
  - When the safety of the provider cannot be adequately assured
  - When there is obvious mortal injury or irreversible death
  - When a valid and relevant advance directive becomes available that recommends against the provision of CPR.
- Further criteria to inform decision making:
  - Persistent asystole despite 20 minutes of advanced life support (ALS) in the absence of any reversible cause.
  - Unwitnessed cardiac arrest with an initial non-shockable rhythm where the risk of harm to the patient from ongoing CPR likely outweighs any benefit e.g. absence of return of spontaneous circulation (ROSC), severe chronic co-morbidity, very poor quality of life prior to cardiac arrest.
  - Other strong evidence that further CPR would not be consistent with the patient's values and preferences, or in their best interests.
- Criteria that should not alone inform decision-making e.g.
  - Pupil size
  - CPR duration
  - End-tidal carbon dioxide (CO<sub>2</sub>) value
  - Co-morbid state
  - Initial lactate value
  - Suicide attempt
- Clinicians should clearly document reasons for the withholding or termination of CPR, and systems should audit this documentation.
- Systems should implement criteria for early transport to hospital in cases of OHCA, taking into account the local context, if there are no criteria for withholding/terminating CPR. Transfer should be considered early in the CPR attempt and incorporate patient, event (e.g. distance to hospital, risk of high-priority transport for those involved), and treatment (e.g. risk of suboptimal CPR) factors. Patients who may particularly benefit from early transport include emergency medical services (EMS) witnessed arrest [or by bystander performing high quality basic life support (BLS)] with either ROSC at any moment or ventricular fibrillation/tachycardia (VT/VF) as presenting rhythm and a presumed reversible cause (e.g. cardiac, toxic, hypothermia).
- Systems should implement criteria for inter-hospital transfer of IHCA patients in hospitals where advanced CPR techniques are not offered.
- Clinicians should start CPR in patients who do not meet local criteria for withholding CPR. Treatments may then be tailored as more information becomes available.
- Clinicians should not partake in 'slow codes'.
- During a pandemic, resource demand (e.g. critical care beds, ventilators, staffing, drugs) may significantly exceed resource availability. Healthcare teams should carefully assess each patient's likelihood of survival and/or good long-term outcome and their expected resource use to optimise allocation of resources. Clinicians should not use categorical or blanket criteria (e.g. age thresholds) to determine the eligibility of a patient to receive treatment.
- In systems that offer uncontrolled donation after circulatory death and other systems of organ donation, transparent criteria should be developed for the identification of candidates and process for obtaining consent and organ preservation.

## Bystander CPR

Systems should:

- Recognise the importance of bystander CPR as a core component of the community response to OHCA.
- Recognise bystander CPR as a voluntary act, with no perceived moral or legal obligation to act.
- Support bystanders in minimising the impact on their own health of performing bystander CPR. In the context of transmissible disease (such as COVID-19), bystanders also have a responsibility of preventing further disease transmission to other individuals in the immediate vicinity and the wider community.
- Aim to identify cases where bystander CPR is likely to be beneficial and cases where it is unlikely to be beneficial.
- Never evaluate the value of (bystander) CPR in isolation but as part of the whole system of healthcare within their region. (Bystander) CPR seems feasible in settings where resources and organisation support the integrity of the chain of survival.

## Family presence during resuscitation

Resuscitation teams should offer family members of cardiac arrest patients the opportunity to be present during the resuscitation attempt in cases where this opportunity can be provided safely, and a member of the team can be allocated to provide support to the patient's family. Systems should provide clinicians with training on how best to provide information and support to family members during resuscitation attempts.

## Patient outcomes and ethical considerations

- When making decisions about CPR, clinicians should explore and understand the value that a patient places on specific outcomes.
- Health systems should monitor outcomes following cardiac arrest, and identify opportunities to implement evidence-based interventions to reduce variability in patient outcome.
- Cardiac arrest research should collect core outcomes, as described in the cardiac arrest core outcome set.

## Ethics and emergency research

- Systems should support the delivery of high-quality emergency, interventional and non-interventional research, as an essential component of optimising cardiac arrest outcomes.
- Researchers should involve patients and members of the public throughout the research process, including design, delivery and dissemination of the research.
- For observational research (e.g. in the context of registry data collection and/or DNA biobank data sampling and analyses) we suggest consideration of a deferred and broad consent model, with concurrent implementation of appropriate safeguards aimed at preventing data breaches and patient re-identification.
- Communities or population in which research is undertaken and who bear the risk of research-related adverse events, should be given the opportunity to benefit from its results.
- Researchers must ensure that research has been reviewed and approved by an independent ethical review committee, in line with local law, prior to it being commenced.
- Researchers must respect the dignity and privacy of research subjects and their families.

- Researchers should comply with best practice guidance to ensure transparency of research, including study protocol registration, prompt reporting of results, and data sharing.
- Systems should ensure that funding for cardiac arrest research is proportionate to the societal burden caused by cardiac arrest-associated morbidity and mortality.

## Evidence informing the guidelines

For ethics in relation to the COVID-19 pandemic see "Ethical considerations on resuscitation during the COVID-19 pandemic".<sup>3</sup>

## Major interventions aimed at safeguarding autonomy

The key interventions for safeguarding patient autonomy are advance directives and advance care planning. These interventions should be underpinned by a shared decision-making process.

Variability in terminology, definitions, type and delivery of interventions, and outcome choice makes it challenging to identify and assimilate research evidence in this area.<sup>4,5</sup> In view of this, the writing group developed consensus definitions and statements for advance directives, advance care planning, and shared decision making, which are summarised in Tables 1–3 and the online supplement.

In developing treatment guidelines, we drew on core ethical principles, 29 systematic/scoping reviews, and 49 recent primary research papers. Key systematic reviews and studies are summarised in the supplementary text and Tables S2 and S3. The corresponding rapid reviews 1.1–1.4 are summarised in the respective appendices.

## Advance directives

Effective use of advance directives relies on the accurate and efficient exchange of information about patient values, goals, and preferences, and available treatment options.<sup>4</sup> Consequently, several, structured communication tools (e.g. paper, video, or computer decision aids, and educational interventions) have been developed to facilitate end-of-life decision-making.<sup>4</sup> Evidence from meta-analyses of randomised controlled trials (RCTs), systematic reviews, and recent studies suggests that structured communication tools aid in the completion of advance directives and may increase concordance of end-of-life care with the care desired by the patient.<sup>4,6–12</sup>

Do not attempt CPR (DNACPR) decisions seek to protect patients from receiving invasive treatments they have declined, they have considered futile, or from treatments that are not aligned with the patient's values and preferences.<sup>13</sup> Evidence from 13 RCTs and from 8 nonrandomised studies included in 3 systematic reviews suggests associations of communication interventions with an increased frequency of DNACPR orders.<sup>4,7,8,10</sup>

Four systematic reviews reported mixed findings regarding the impact of advance directives on the documentation of patient's wishes about treatment escalation and resuscitation decision-making.<sup>5,9,10,14</sup> These reviews also highlighted that, in some studies, the making of a DNACPR decision may confer benefit as regards the patient's quality of care through, for example, more adequate pain relief and hydration, and improved response of healthcare providers to clinical deterioration.

Recent evidence from RCTs supports the use of informational video decision support tools in both the nursing home and the in-

**Table 1 – Consensus definition and statements for advance directives.**

## Advance directives \*

- An advance directive is an instrument that relays information concerning an individual's preferences and goals regarding medical procedures and treatments, especially those used for end-of-life care.
- Advance directives intend to extend the patient's autonomy to situations in which he/she is unable to express his/her preferences regarding treatment decisions. They reflect a patient's individual moral, cultural, and religious attitudes. They are represented in three formats: Living Will (or instruction directive), Appointment of a Healthcare Proxy (or proxy directive), and Legal Status of Preferences.
- In principle, advance directives (ADs) must fulfil the following 3 criteria: Existence, Validity (partly realised through periodic review), and Applicability.
- Health care professionals should determine whether their patients have ADs.
- Physicians should respect their patient's ADs and incorporate them into their decision making.
- Physicians should discuss advance directives with their patients.
- Attempts should be made to ascertain patient's wishes (especially patients with terminal diseases) concerning life-sustaining treatments when they are capable of making decisions or, alternatively, from their surrogates when they are not capable of making decisions.
- There are times when advance directives should not be followed. These include situations when the advance directive calls for an action that is prohibited by the country's laws and/or regulations, where there is compelling evidence that the patient may have changed his/her mind since completing the advance directive, when there is compelling evidence suggesting that the patient did not understand the nature of the advance directive he/she completed, or when there is evidence that the patient did not have freedom of choice at the time of drafting.
- If advance directives concern the refusal of a specific treatment, careful interpretation should be made as to whether this should concern similar (but still alternative) treatments or not. For example, a patient may refuse a specific medical or surgical treatment due to certain rare but severe side effects. In such a case, it may not be appropriate to exclude alternative treatments that may exhibit a more favourable safety profile and comparable efficacy relative to the refused treatment.
- Reasons for refusal of standard treatments of a specific disease may not apply after the introduction of new interventions with more favourable safety profiles and increased efficacy. Given the fact of the continuous and rapid progress in clinical practice, old (e.g. >5 years) and non-updated advance directives should be cautiously interpreted in the context of availability of new, safer, and potentially more effective therapies.
- Nonstandard advance directives (e.g. tattoos indicating do-not-attempt cardiopulmonary resuscitation - DNACPR) should not be immediately perceived as legally valid, unless designated so by local law. In countries where the presence of nonstandard advance directives is considered legally valid, CPR administered in conditions where resuscitation is likely to be futile can lead to legal prosecution of the healthcare professional. Concurrently, every effort should be promptly undertaken to clarify whether a valid, pertinent advance directive exists.

\*, Consensus definitions and statements were based on 7 references.<sup>5a–5g</sup>

**Table 2 – Consensus definition and statements for advance care planning.**

## Advance care planning \*

- A process that enables individuals to define goals and preferences for future medical treatment and care, to thoroughly discuss these goals and preferences with family and health-care professionals, and to record and review these preferences if appropriate. The main objective of advance care planning is to help ensure that people receive medical care that is consistent with their values, goals and preferences during serious, chronic and/or acute/life-threatening illness.
- Advance care planning elements may include communication interventions such as information brochures or pamphlets, and video decision support tools.
- Regarding specific population subgroups with morbidity-related decisional incapacity (e.g. patients with dementia, or children with intellectual disability and a life-limiting illness): due to very limited or even completely lacking, relevant evidence, it is still unclear whether advance care planning (based on surrogate decision-making) can positively impact their health-related quality of end-of-life, and also ameliorate the surrogates' psychological burden, symptoms, and distress. In the meantime, advance care planning should still be considered for such patients.
- Advance care plans that are not updated or re-reviewed should be cautiously interpreted in the context of availability of new and improved therapies that might potentially affect patient preferences; patient preferences may also evolve with time independently of available treatment options.
- Patient's cultural background, religious beliefs/religiosity, and associated, possible spiritual needs, should be taken into account/respected in the course of development and reviewing of advance care planning.
- Regarding treatment limitation directives, a recent multicenter observational study suggested that end-of-life treatment limitation practices may be affected both by patients and physician religion.

\*, Consensus definitions and statements were based on 3 references.<sup>5a,5g,64</sup>

hospital setting by reducing the frequency of delivery of interventions that are unlikely to be beneficial.<sup>15–18</sup> Four recent retrospective studies and a point-prevalence review suggest that advance directives and/or DNACPR decisions are associated with reduced use of life-sustaining treatments.<sup>19–24</sup>

Evidence from two systematic reviews suggests that the use of advance directives is associated with reductions in emergency room visits, hospital admissions, health care costs, and more preference for comfort care as opposed to life-sustaining treatments.<sup>4,8</sup> The effect on outcomes such as hospital/ICU length of stay, and patient preference

for end-of-life care is less clear. These mixed findings may be partly attributed to heterogeneity across studies in relation to population, interventions, and the comparator group. Despite study limitations, findings of studies of advance directives generally support the use of structured communication tools.<sup>4</sup>

Symptom control is key to improving the end-of-life experiences of a dying patient.<sup>25</sup> However, 15 RCTs included in a systematic review failed to determine any advance directive benefit on patients' anxiety, depression, pain, psychological well-being and health.<sup>8</sup>

**Table 3 – Consensus definition and statements for shared decision making.****Shared decision making \***

- Shared decision making is a collaborative process that allows patients, or their surrogates, and a possibly/preferably multidisciplinary team of healthcare professionals to reach consensus on which treatment strategies and interventions - including life-support limitation and palliative care- accord with the patient's values, goals, and preferences. Healthcare decisions should take the best available scientific evidence into account. Honest exchange of information should foster the development of trust/partnership between patient/surrogate(s) and clinician(s). Clinicians should be trained in communication skills. Shared decision making practices should be evaluated by research using patient-/surrogate-reported outcomes.
- The shared decision making process should include information exchange, deliberation, and decisions relating to a treatment.
- Shared decision making should preferably be part of the application of current guidelines on family-centred care.
- Shared decision making should take into account any pre-existing, documented patient goals, values, and preferences in the form of either "isolated" advance directives, or advance directives completed in the context of advance care planning.

\*, Consensus definitions and statements were based on 3 references.<sup>4a,5h,64</sup>

In recent guidelines, patient/family satisfaction was considered as a core outcome.<sup>26</sup> Data from eight RCTs included in three systematic reviews indicated that communication interventions aiding the completion of advance directives had no significant effect on patient/family satisfaction with end-of-life care.<sup>4,7,8</sup> However, another 4 RCTs included in one of these reviews, reported an increase in patient/family satisfaction with care associated with a communication intervention.<sup>8</sup>

**Advance care planning**

Advance care planning may be regarded as the state-of-the art procedure for ensuring respect for patient autonomy. It is a dynamic process based on effective and honest communication between the patient and their family, and healthcare professionals (Table 2).

Most studies support the use of advance care planning as a strategy to ensure that end-of-life care is in line with the patient's values and preferences, although there is some inconsistency across the available evidence.<sup>9,27–32</sup> Video-based information and other types of interventions may support the development of advance care plans and thereby increase the concordance between care desired and care received. How effective interventions are in achieving concordance may depend on their nature and on the context in which they are used.<sup>33–36</sup>

Documenting a person's or a patient's updated preferences about end-of-life treatments (including life-sustaining treatments and CPR) is a major objective of advance care planning (Table 2); documented preferences may then be accessed by healthcare professionals to potentially inform treatment decisions. Evidence from six systematic reviews indicates that advance care planning increases documentation of patient preferences.<sup>27,30,32,37–39</sup> Recent studies also reported positive results.<sup>40–42</sup>

We identified mainly positive results on the effect of advance care planning (with or without the aid of communication tools) on the preference for and/or actual use of life-sustaining treatments at the end-of-life. In a meta-analysis of seven RCTs, a video intervention reduced the likelihood of indicating a preference for CPR relative to control.<sup>43</sup> Another systematic review concluded that advance care planning was associated with a reduction in the use of life-sustaining treatments.<sup>9</sup> In contrast, two RCTs and four observational studies included in another systematic review did not report any significant association between communication tools for end-of-life decision making and DNACPR status.<sup>44</sup> Nevertheless, four recent RCTs<sup>45–48</sup> and a cross-sectional survey<sup>49</sup> suggest that advance care planning is associated with less frequent preferences for CPR and/or use of life-sustaining treatments at the end-of-life.

We identified limited supporting evidence for the use of communication tools in the context of advance care planning, to reduce hospitalisations, ICU admissions, and hospital/ICU utilisation among patients who are unlikely to obtain any benefit. Indeed, in meta-analyses that included five RCTs and eight observational studies, communication tools had no effect on ICU length of stay.<sup>44</sup> In addition, although a meta-analysis of three observational studies suggested that communication tools are associated with reduced ICU length of stay of non-survivors, this was not confirmed in one RCT.<sup>44</sup> A meta-analysis of five observational studies suggested that communication tools may be associated with reduced hospital costs. However, one RCT and another two observational studies did not report any effect of communication tools on hospital costs in ICU non-survivors. One RCT reported a reduction in the duration of mechanical ventilation with the use of communication tools, but another two RCTs and two observational studies failed to confirm such benefit.<sup>44</sup>

The effects of advance care planning on hospital/ICU admissions, healthcare resource utilisation, death at preferred location, hospice use, palliative care referrals, healthcare costs, and quality of dying and death are inconsistent between studies.<sup>9,27,30,31,36,38,43,49–58</sup>

Evidence from systematic reviews suggests that advance care planning is associated with improvements in symptom control and quality of life.<sup>9,30,31,37,38,50,59</sup> However, three recent RCTs found no benefit with respect to patients' health-related quality-of-life, physical/functional outcomes, and anxiety or depression.<sup>53,60,61</sup>

A recent cluster RCT in nursing homes reported an advance care planning-associated decrease in family carers' decisional conflict.<sup>62</sup> A cross-sectional survey,<sup>49</sup> and a historically controlled, prospective study<sup>57</sup> reported associations of advance care planning with good quality of end-of-life and decreased suffering in children/adolescents,<sup>49,57</sup> or adults<sup>49</sup> with complex chronic conditions; advance care planning was also associated with reduced parental decisional regret, or lower caregiver burden.

Evidence from five systematic reviews suggests that advance care planning may improve patient/family satisfaction with care.<sup>9,31,37,38,44</sup> However, a recent, multicenter RCT of advanced cancer patients reported that consultation plus early palliative care did not affect family satisfaction with care.<sup>53</sup>

Specific and adequate training of healthcare professionals is key to improving the quality of end-of-life care.<sup>63</sup> The results of 21 studies (RCT,  $n=3$ ) included in a systematic review suggested that communication skills training interventions increase comfort, self-efficacy, and preparedness of healthcare professionals in the delivery of end-of-life care.<sup>37</sup> In an interview-based study included in a systematic review, advance care planning discussions increased

healthcare professionals' confidence in their dealings with the patient.<sup>29</sup>

### Shared decision making

Shared decision making is an individualised, collaborative, multistep process aimed at reaching major and preference-sensitive treatment decisions (Table 3).<sup>64</sup> The concept underpins all patient-focused healthcare decision-making.

Effective communication about end-of-life care relies on a shared decision-making process. Its use has been shown to improve end-of-life care, particularly in relation to concordance between care desired and care received, in a systematic review<sup>65</sup> and most recent studies.<sup>60,66–70</sup> However, studies reporting the effect on quality of care and symptom control have produced conflicting findings.<sup>65</sup>

From a health service perspective, shared decision-making may support the appropriate allocation of resources by ensuring that patient treatment aligns with their values and preferences. Use of interventions based on the concept of effective shared decision-making may be associated with shorter ICU/hospital length-of-stay, selection of palliative care pathways in nursing homes, and reduced health care costs and fewer in-hospital deaths, although evidence from systematic reviews and recent studies is inconsistent.<sup>65,67,69,71–79</sup>

Family members of patients may be impacted by the illness of their loved ones. Up to 50% of family members of critically ill patients experience psychological symptoms, such as acute stress, post-traumatic stress disorder, anxiety, depression, and decisional conflict/regret.<sup>26,80–82</sup> Family support interventions may help to reduce these psychological impacts, as suggested by four systematic reviews,<sup>37,71,72,83</sup> and two recent studies.<sup>84,85</sup> However, some recent studies found that family support interventions did not reduce psychological symptoms in family members.<sup>66,67,75,79</sup>

Patient and family satisfaction is a key objective of patient- and family centred communication and care. Communication in the context of shared decision-making is associated with higher patient/family satisfaction and increased decisional confidence, as suggested by four systematic reviews.<sup>65,71,72,74</sup> Key components of this approach include open, honest, clear, and frequent communication and inclusion of family members in discussions with healthcare professionals.<sup>83</sup> Recordings of clinician-family conferences suggest that communication is often sub-optimal, such that the patient's values and preferences are infrequently elicited.<sup>86</sup> The use of structured communication tools may help to improve communication with families, as suggested by two systematic reviews.<sup>37,44</sup> Furthermore, according to recent studies, communication supported by other strategies such as video decision aids may be associated with improved family satisfaction.<sup>67,75,78</sup>

### Major interventions aimed at safeguarding autonomy and COVID-19

During periods of public health disaster, such as the COVID-19 pandemic, the importance of pre-existing documentation of patient's wishes regarding life-sustaining treatments, including mechanical ventilation and CPR, may increase, especially for overstretched healthcare systems with limited resources.<sup>87–89</sup> In the absence of advance directives, healthcare professionals should actively seek to engage in treatment escalation decisions with patients, especially those at high-risk of death.<sup>90</sup> Ideally, this should apply to all healthcare settings, using digital communications as appropriate and feasible.<sup>87–89</sup> Discussions should focus on eliciting an individual's values and

preferences, rather than asking them to choose a treatment option.<sup>91</sup> Treatment escalation decision-making, such as DNACPR, should be based on an individualised patient assessment that draws upon clinical and scientific evidence,<sup>92</sup> the patient's values and preferences, and the local context, such as resource availability. Decision-making based on single factors such as age, sex, race, religion/ethnicity, intellectual disability, and socioeconomic status is not ethically justifiable.<sup>87,88,92,93</sup> Discrimination according to a patient's COVID-19 status must also be avoided.<sup>94</sup>

Structured advance care planning interventions may include face-to-face conversations between the patient and a healthcare professional over a specified time interval, often with family members present.<sup>46</sup> Clearly, such interventions may be hindered by the need for physical distancing due to healthcare catastrophes like the COVID-19 pandemic. Although digital telecommunication technology may obviate the need for physical presence during a discussion, its availability and the patient's/proxy's capability of adequate use should not always be taken for granted. Concurrently, there may be a perceived need for augmented dissemination and even acceleration of the advance care planning processes to prevent the waste of potentially scarce resources on disproportionate and/or unwanted, aggressive end-of-life treatments.<sup>89</sup> Such upscaling process should be achieved solely through improvements in system organisation and infrastructure, public communication and education, and effective suppression of health misinformation.<sup>1,2</sup> Any form of psychological pressure in the context of categorical discrimination of frail people should be regarded as ethically unacceptable.<sup>92</sup> For emergency department patients at high risk of severe COVID-19 and without advance care plans, a viable alternative may comprise the implementation of an emergency department-based palliative care team committed to high-quality goals of care discussions with the patient and/or proxy. Such interventions may increase the rates of time-sensitive decisions about CPR and other life-sustaining treatments, and comfort care.<sup>95</sup>

Shared decision-making becomes more challenging in situations where face-to-face communication is not feasible. In the context of COVID-19, visiting has been limited in many hospitals and the burden on hospital services may have limited the time available to healthcare professionals to engage in detailed discussions with patients and their families. In these circumstances, use of teleconferences may be an acceptable and feasible approach to maintain patient-centred communication with families and integrate shared decision-making in routine clinical practice.

### Deciding when to start and when to stop CPR

The corresponding rapid reviews 2.1–2.7 are summarised in the respective appendices (pages 167–310 of the online supplement).

### Termination of resuscitation

The 2020 International Liaison Committee on Resuscitation (ILCOR) Consensus on Science and Treatment Recommendation conditionally supported the use of termination of resuscitation (ToR) rules (very-low certainty evidence).<sup>96</sup> In making the recommendation, the ILCOR Education Implementation and Teams task force acknowledged variation in patient values, resources available, and performance of ToR rules across settings. The task force sought to balance the risk that implementation might result in missed survivors, against current variation in practice and improve termination decisions more generally. ToR may also reduce demand on hospital resources and

increase rescuer safety by reducing the number of patients transported to hospital in cardiac arrest.

It is generally agreed that CPR should not be provided to all patients. This viewpoint reflects both medical and ethical perspectives, including the potential harm of CPR (in terms of dignity, victim awareness, deception of relatives, etc.), and the risks of unfavourable outcome in survivors (and likewise burden for caregivers, risks to healthcare providers, medical costs, and preservation of medical resources). Many authors strongly defend the individual's right to die in a society where more and more advanced medical techniques can lengthen life at the potential cost of quality of life and palliative comfort.<sup>97,98</sup> Prolongation of an inevitable dying process should be considered harm (dysthanasia). In practice, it is often challenging to reliably identify which individuals will have a poor outcome in the event of cardiac arrest.<sup>99–103</sup>

Futility has traditionally been described as a likelihood of survival of less than 1%.<sup>104</sup> More recently, this concept has been challenged for not considering either neurological and functional outcome of survivors or broader societal considerations opinions (e.g. utility trade-off).<sup>104–106</sup> Importantly, Van Norman et al. posed relevant questions about fairness of the concept when there is a potential for unconscious bias based on socioeconomic and demographic factors like social status, fear of litigation or the patient gestalt.<sup>107–109</sup> The appreciation of futility is timely and contextual in nature and often also incorporating religious or spiritual beliefs.<sup>110,111</sup> Patients and families may define futility very differently than medical providers. Marked differences are also observed between different providers. Many clinicians lack confidence in making ToR decisions and some report using non-validated or controversial factors as a single reason for terminating CPR.<sup>106,107,112–122</sup> Decision-making becomes even more complex in the context of newer advanced resuscitation technologies.

Defining an unfavourable outcome is challenging. The cut-off of a cerebral performance category (CPC) of 2 may translate to a spectrum of functional outcomes. Moreover, the value of an outcome to an individual will likely be specific to that person.<sup>105</sup> Defining –as a society, healthcare provider or even as a relative– that a certain life no longer is worth living, especially when this becomes balanced against cost or societal interaction, should only be done with the greatest caution as it incorporates a great inherent risk of quickly crossing acceptable ethical boundaries.<sup>123,124</sup> As such, there has been a shift from futility to considering the broader concept of best interests, which rather evaluates burden versus benefit.

Decision-making regarding the withholding or termination of resuscitation exists in a legal framework, which will have primacy over ethical concepts.<sup>125</sup> The ILCOR Education, Implementation, and Teams (EIT) taskforce in their insights highlighted the need to consider local legislation.<sup>96</sup>

There are important differences between the withholding or termination of resuscitation between the in-hospital and out-of-hospital setting. In the out-of-hospital setting, EMS teams often arrive at a scene where CPR is in progress, and then can only decide to withdraw (not withhold) resuscitation efforts. They often have limited information on the patient's previous medical history and their values and preferences, and may be unable to discuss treatment options with family members. As such, where there is uncertainty about the appropriateness of terminating resuscitation, the focus should be on patient treatment with a view to reconsidering the appropriate treatment once the patient's values and preferences, and clinical trajectory are known.<sup>98,102</sup>

The ILCOR COSTR recommends that none of the existing ToR rules should be the sole determinant of when to discontinue resuscitation.<sup>96</sup> ToR rules will inevitably introduce a self-fulfilling prophecy and should be reviewed periodically as new treatments evolve. Intra-arrest factors are not sufficiently reliable to be used in isolation for terminating resuscitation.<sup>126–135</sup> Examples of factors that should not be used alone include serum potassium, end-tidal CO<sub>2</sub>, cardiac standstill on ultrasound, pupillary response/size, temperature, co-morbid status, cause of arrest, no-flow time, low-flow time, and absence of ROSC.

The ILCOR CoSTR summarises several ToR rules.<sup>136,137</sup> Some factors are consistent across tools, such as whether the arrest was witnessed. A key challenge in operationalising these rules stems from uncertainty as to the applicability of rules to other healthcare settings and the challenge in reliably estimating the number of missed potential survivors when applying the rule.<sup>103,104,138–144</sup>

There are specific guidelines for specific subpopulations, such as children.<sup>145</sup> Despite differences in pathophysiology and aetiology, the ethical framework in paediatric cardiac arrest is otherwise similar, although many clinicians may be more cautious in terminating the resuscitation of a child.<sup>99,146,147</sup>

Typical, but not only important for children, is the mandate and role of surrogate decision makers. Time is often limited to come to shared decision-making during cardiac arrest. Moreover, the likelihood of truly informed unbiased consent is low, and it is unclear whether the best interest of the patient might not conflict with the rights and interests of the relatives.<sup>148</sup> Importantly, putting for instance parents in the position to forgo CPR may intensify parental grief and helplessness.<sup>148</sup> Therefore, clinicians should carry the primary professional and moral responsibility for the decision and use a model of informed assent from parents, allowing for respectful disagreement. Nevertheless, local regulation and laws might demand actual guardian's consent.

### Slow code

A 'slow code' is slang for the deceptive practice of purposely delivering sub-optimal CPR with the pretence of attempting to save the patient's life. There is evidence that slow codes continue to be performed both in IHCA and OHCA, even when CPR is considered of no benefit to the patient.<sup>149–151</sup>

Use of the slow code is extremely ethically problematic, although some have advocated for it in certain circumstances.<sup>152,153</sup> Several alternatives have been described that are ethically more acceptable, such as informed non-dissent, tailored code or early advance care planning with open communication. More education on ethics in resuscitation might positively affect this.

### Extracorporeal (E)-CPR

The European Resuscitation Council (ERC) ethics writing group acknowledges the ALS and paediatric life support 2020 ILCOR COSTRs that support the use of E-CPR as a rescue therapy for selected cardiac arrest patients when conventional CPR has failed in settings where E-CPR can be implemented (weak recommendation, very-low certainty of evidence).<sup>154,155</sup> To inform our insights, we further identified 6 systematic reviews,<sup>156–161</sup> four narrative reviews<sup>162–165</sup> and 13 observational studies<sup>68,140,160,166–175</sup> on this topic. Other sources, such as commentaries and ethical dissertations, were considered as indirect information.

The evidence base for the cost-effectiveness and ethical framework of E-CPR is limited. For IHCA, E-CPR may be cost-



effective, provided the programme is limited to specific patient groups. The incremental cost-effectiveness ratio is mainly influenced by the probability of survival, although large variations in in-hospital cost estimates have been reported. Physicians involved should be knowledgeable and provide proper stewardship of available resources. Across 224 North American hospitals that participate in the American Heart Association (AHA) Get-with-the-Guidelines-Resuscitation-registry, fewer than 1% patients received E-CPR between 2000 and 2018, indicating a further need for optimised patient selection and E-CPR implementation strategies.<sup>170,171</sup> One systematic review examined E-CPR in refractory adult OHCA of cardiac origin.<sup>157</sup> They suggested that it is feasible and may increase both neurologically intact survival and organ donation in non-survivors. Implementation in existing EMS systems is challenging and requires detailed protocols for patient selection and transportation.<sup>160,173–175</sup> The Ethics writing group identified an urgent need for more research on patient selection, modifiable outcome variables, risk-benefit, and cost-effectiveness of E-CPR. Such data are crucial for E-CPR programme implementation.

### Organ donation

Patients who sustain a cardiac arrest are an important source of donor organs, mainly because severe neurological injury is a common mode of death.<sup>176–178</sup> There are three pathways by which cardiac arrest patients might donate organs: following confirmation of brainstem death, following withdrawal of life-sustaining treatment leading to circulatory death (controlled donation after circulatory death) or donation where resuscitation attempts to achieve ROSC have been unsuccessful (uncontrolled donation after circulatory death). The Post-Resuscitation Care and ALS sections of the guidelines provide further details on these pathways. This section focuses on the ethics of organ donation.

We included two systematic reviews,<sup>178,179</sup> four narrative reviews,<sup>180–183</sup> five observational studies<sup>184–188</sup> and some additional editorials and ethical dissertations.

Across Europe, there is variability in organ donor rates, availability of organ donation pathways, and law and policy regulating organ donation (World Health Organization Collaborating Centre on Organ Donation and Transplantation 2019). Shortage of available donor organs is an ongoing challenge across Europe and contributes to premature morbidity and mortality in individuals with organ failure. Organ donation provides an opportunity following a tragic event to respect the donor's wish to benefit wider society. For relatives of the donor, consenting to organ donation may provide comfort that their grief has given life to others.<sup>188</sup> Organ donation is generally supported by society, although levels of support vary by cultural group and between individuals.<sup>181,186</sup>

A key issue is the need for both family members and society to maintain trust that donation is considered only when ongoing treatment will not achieve an outcome important to the patient. Examples of safeguards to maintain this trust include respect for the dead-donor rule, a clear division between the clinical and transplantation team, and transparent communication with family members before organ retrieval. A review of attitudes towards organ donation concluded that both general and ethical education may serve to guide policy and to facilitate family member requests and informed consent dialogues.<sup>180</sup> Helping families to understand and accept not only medical and legal criteria for determining death, but also ethical criteria for withdrawing life support, may help them be more comfortable with their decisions.

Uncontrolled donation after circulatory death raises ethical challenges.<sup>185,189</sup> In particular, the time-critical nature of the process usually requires the initiation of organ preservation processes prior to family consultation to maintain organ viability.<sup>186,190,191</sup> Cardiac arrest patients may meet criteria for both uncontrolled donation and E-CPR programmes.<sup>157,159,179</sup> In centres that offer both modalities, uncontrolled donation should be considered only in patients who do not meet clinical criteria for E-CPR, in order to prevent the loss of potentially saveable life.<sup>163</sup> For a more in-depth discussion see the supplement (pages 255–259).

Importantly, several authors suggest that organ-preserving CPR should be considered only for patients who are brain-dead, or in those with evidence of futility, a known wish for organ donation and a specific informed consent from a next of kin.<sup>163,190,192,193</sup>

### Family presence during resuscitation

In our literature search we did not specifically address parental presence during the resuscitation of a child as this is expected to be topic of a specific COSTR from the ILCOR paediatric Taskforce; however, our findings apply equally to this context and we also refer to the 2015 ERC guidelines.<sup>194,195</sup> For family presence during resuscitation, we identified one guideline,<sup>196</sup> two systematic reviews,<sup>197,198</sup> five narrative reviews,<sup>199–203</sup> one RCT,<sup>204</sup> and three observational studies,<sup>205–207</sup> as well as several ethical dissertations and opinion pieces.

The available evidence indicates that family presence during resuscitation does not affect patient outcome but may improve family member psychological outcomes. On this basis, teams should offer family members the option to be present during resuscitation in situations where it is safe, and when the family can be adequately supported.

### CPR after attempted suicide

This guidance is based on one narrative review<sup>208</sup> and on an observational study,<sup>209</sup> with other sources included as indirect evidence.

The 2015 ERC ethics chapter highlighted the challenge of determining whether the patient who has attempted suicide had mental capacity at the time of the suicide attempt.<sup>195</sup> On this basis, the guidance recommended that treatment be started because of the risk of harm if treatment is delayed. Crucial to the decision making is the appreciation of mental capacity. This is defined as sufficient understanding of the nature, purpose and effects of the proffered treatment, and able to comprehend and retain the treatment information; believe the information; and weigh it among other factors to reach a decision.<sup>210</sup> The patient must also be able to communicate and substantiate the decision (see also supplement for our consensus definition of decisional incapacity). Sufficiency of capacity is seen as a spectrum, and the more profound the consequences of the decision, the higher the level of capacity that must be demonstrated.<sup>210</sup>

A specific complex situation is when the patient is not considered competent but has a valid advance directive.<sup>211</sup> A decision to withhold treatment might be viewed as abetting a suicide attempt, but it is reasonable to continue to honour a valid and applicable advance directive. This is because the test of capacity is based on when the advance directive was made, rather than at the time of the suicide attempt.<sup>212</sup> An alternative perspective is that there are competing rights that are sufficient to override a competent decision to refuse treatment. These may include the state's interests in preventing

suicide and the need to protect innocent third parties, such as dependent children and even foetuses.

If the treating healthcare professional is uncertain about the patient's capacity or validity of an advance directive, it is reasonable to provide lifesaving treatment and simultaneously seek urgent ethical or legal advice. Sufficient time should be taken to consider contextual evidence relating to the suicidal behaviour, the nature of the treatment decision and the verification of any documentation.<sup>208</sup>

It is difficult to rapidly judge the context of an attempted suicide and it is suggested that the default should be to initiate life-sustaining treatment.<sup>213</sup> Surrogate decision makers may be unable to represent the views of patients, especially in the setting of attempted suicide. If the patient is stabilised, the quality of ongoing life may not be in line with their values and preferences. The response to the clinical situation should not be dogmatic but proportional to the individual case.<sup>214</sup>

Some authors have suggested that a suicide attempt is not as important as the underlying (disease) process that led to the attempt. In other words, it may be ethical to withhold or withdraw life-sustaining treatment in case of suicide when there is an underlying serious medical condition.<sup>215</sup>

### The ethical framework of bystander CPR

Early bystander CPR improves patient outcomes in OHCA.<sup>216–221</sup> In many countries, systems of trained volunteers and/or first responders, in addition to dispatcher-assisted CPR by lay people, have been implemented. The crucial role of this community response to OHCA is incorporated in the chain of survival and in the ERC guidelines.<sup>222</sup> There are important differences in rates of bystander CPR between countries, regions and even in circumstances or victim characteristics.<sup>217,223–226</sup>

A 2020 ILCOR scoping review explored the individual's willingness to perform bystander CPR.<sup>227</sup> Factors influencing bystander willingness or actual delivery of bystander CPR include emotional factors, patient status (e.g. vomiting), socioeconomic status of the patient, patient sex, physical challenges (e.g. patient positioning, bystander age), and lack of knowledge or confidence.<sup>228,229</sup> Rescuers are more willing to perform compression-only CPR compared to CPR with rescue breaths. Some authors also identified fear of legal consequences as a potential barrier.<sup>230,231</sup> Older bystanders are less likely to start CPR, despite a higher chance of being bystanders to cardiac arrest. Important facilitators include prior knowledge and training, and feeling a moral obligation to act.<sup>232–235</sup>

There are ethical aspects concerning the ILCOR-supported use of smartphone-apps or text-messaging to alert trained lay rescuers to OHCA (strong recommendation, very low-certainty evidence).<sup>236</sup> Regional systems of alerting lay volunteers and/or first responders have many common characteristics but still may vary depending on the local context.<sup>224,225,237–256</sup> Some systems a priori exclude (young) children, traumatic cardiac arrest, intoxication, drowning and/or suicide, unsafe or inaccessible settings and/or nursing homes. Such exclusions are most often not further explained and/or defined. The low sensitivity and specificity of current dispatcher protocols for cardiac arrest recognition results in a high percentage of false positive and false negatives. We identify this as a major issue and consider better case selection a priority. In up to 30% of OHCA, the attending EMS team will not start CPR. A priori identifying these cases is very

difficult but might limit a subsequent ethical conflict between the lay rescuers and the arriving EMS team.

Further key ethical issues in relation to the establishment of these systems include the potential psychological impact on the bystander of attending a cardiac arrest, the potential variability in the skills and competence of volunteers dispatched to OHCA, and the potential impact on the patient's privacy of treatment by a non-professional rescuer. Most authors put higher value on the potential of saving a life than on the possible breach of privacy associated. A survey of North Americans found that most did not object to the implementation of an app-alerted volunteer system in their community nor to receiving crowdsourced help.<sup>249</sup>

An ILCOR review identified only limited evidence of harm to rescuers of performing CPR and/or using an AED. However, in the context of the COVID-19 pandemic, there is a risk of infection transmission to the rescuer. Whilst not performing CPR (or with substantial delay) will reduce the likelihood of a good outcome for many victims, bystanders should try to limit the associated risk of disease transmission by doing CPR.<sup>3,257</sup> The specific risk-benefit will be a function of factors such as the current regional COVID-19 prevalence, the victim's presentation (presumed COVID-19 status), the likelihood that CPR would be effective, the availability of personal protective equipment, and whether the bystander already had previous contact with the victim.

CPR training should better prepare lay rescuers for the various logistical, conceptual, and emotional challenges of resuscitation.<sup>105,258,259</sup> This includes limiting self-doubt, improving knowledge of the exact impact of performing or not performing certain actions and correcting certain misbeliefs.

CPR is promoted as a highly effective treatment both in the popular press and in dedicated media campaigns.<sup>260</sup> Only recently, more discussion about indications and limitations of CPR has started to take place in the public domain.<sup>106</sup> Such discussions, although from a patient and healthcare provider perspective very relevant, are difficult to appreciate for the lay rescuer. The Ethics writing group continues to support the emphasis on bystander CPR as a key link in the chain of survival.

Improved public information about the situations where CPR has a reasonable likelihood of providing clinical benefit and those where not, may be helpful.<sup>105</sup> EMS dispatch centre protocols should seek to better identify patients for whom bystander CPR might be beneficial but also try to identify those for whom it is not. Bystander CPR should never be considered a moral or judicial obligation.

Providing CPR is emotionally challenging for lay rescuers and first responder and, for some, has consequences in terms of family and work life.<sup>253,261,262</sup> The role undertaken by the lay bystander should be acknowledged by both the EMS dispatch centre and the EMS team.<sup>263</sup>

Finally, the ILCOR EIT Taskforce looked at OHCA in resource-limited environments, as many of the statements related to CPR might not be applicable in resource-limited settings.<sup>264</sup> They acknowledged that the feasibility and cost-effectiveness of CPR in OHCA in these settings can be challenged. One could argue that CPR is only ethically acceptable in settings where resources are such that other fundamental parts of the healthcare system are already sufficiently developed. CPR, as with many other healthcare choices, should never be evaluated in isolation but as part of the whole system of healthcare within a country or region. The role and remit of bystander CPR within such a context is obviously far less clear.

## Education, communication and system organisation

### Education of lay persons, persons at risk of cardiac arrest and family

#### *Education about patient's right of autonomy*

Advance care planning discussions led by trained nurse facilitators or social workers are associated with an increase in patient knowledge about advance care planning, significantly more advance care planning discussions with physicians and a higher likelihood to agree to a DNACPR decision.<sup>10,265–268</sup> A patient-centred advance care planning approach increases the congruence in decision-making for future medical treatment between patients and their surrogate, improves satisfaction with the decision-making process and decreases the decisional conflict.<sup>269</sup> A pilot RCT addressing specific cultural factors by a tailored intervention using a bilingual, bicultural patient navigator suggests improved palliative care outcomes for minority groups facing advanced medical illness.<sup>270</sup> A controlled randomised intervention study in the US exploring peer mentoring by patients trained to help other patients with end-of-life planning had a significant influence on the completion of advance directives and the effect was also most prominent among African Americans.<sup>271</sup>

#### *Education about CPR indications, procedures and outcome*

Video decision support tools depicting CPR, resuscitation preference options and different levels of care are associated with higher rates of understanding of the purpose of CPR and resuscitation options.<sup>4,17,272</sup> Educational CPR videos and structured patient-centred interviews can be helpful in decision making with more patients likely to forgo CPR and focus on comfort.<sup>4,45,70,273,274</sup>

### Education of healthcare professionals

#### *DNACPR orders and advance care planning*

More complex interventions involving education of healthcare professionals, education of patients and their caregivers as well as involvement of special teams seem to have greater impact at least on the effectiveness of DNACPR discussions. It is preferable to have these discussions as part of a broader approach such as advance care planning.<sup>275</sup> Some of the background evidence suggests that discrepant interpretations of DNACPR discussion occur with a concerning frequency between physicians and their hospitalised patients.<sup>276</sup> However, there is no direct evidence whether education (and in what form) changes this phenomenon.

#### *Family presence during resuscitation*

A presentation reviewing the literature supporting family presence during resuscitation, open discussion about family presence and a script that could be used to support families during resuscitation are all effective at improving attitudes of nurses and physicians towards family presence.<sup>277–279</sup> The presence of a trained support person may further increase staff comfort with family presence during resuscitation.<sup>196</sup>

### Communication

#### *Advance care planning*

Advance care planning discussions with a trained nurse facilitator, structured nurse-led advance care planning discussions with long-term care residents of a nursing home and their proxies, and a

physician and clinical nurse specialist team processing medical information of the primary physician all improved the consensus about care between patients, their families and the healthcare professionals.<sup>37,265,267,280</sup>

#### *Termination of resuscitation and breaking bad news*

Ambulance personnel feel particularly concerned about the skills required to deliver death notification and communicate with family and bystanders. This unpreparedness is associated with avoidance and distress. Ambulance personnel use distancing and detachment as a coping mechanism and focus on rational or structured behaviours of resuscitation to avoid interaction or empathetic engagement with family and bystanders.<sup>281</sup>

### Patient outcomes and ethical considerations

The outcome of a cardiac arrest can be defined in several ways. Outcomes may be measured at multiple time points from during the cardiac arrest (e.g. end-tidal CO<sub>2</sub>) to hospital discharge (e.g. survival, neurological outcome) and beyond (e.g. survival, neurological outcome, health-related quality of life).<sup>282</sup> A successful resuscitation may be characterised as survival with an acceptable quality of life. This means that long-term outcomes are of particular interest to patients and the resuscitation community.<sup>283,284</sup>

#### *Valuing outcomes*

Traditionally, cardiac arrest outcomes have been clinician-reported, and often dichotomised as good or poor.<sup>283</sup> This dichotomisation often attempts to separate individuals that are functionally independent from those with ongoing dependency or death.

Today, it is understood that cardiac arrest outcome is multifactorial and may include long-term changes in functional, emotional, physical, cognitive and social domains, all associated with health-related quality of life.<sup>283</sup> To make patient-centred decisions about the appropriateness of resuscitation requires clinicians and patients to have a shared understanding of how the patient defines a good outcome. The patient's perspective on outcome may be influenced by factors such as age, religion, societal values, and personal experiences. This should inform decisions about treatments, such as CPR.

Epidemiological data provides information on outcome at the population level.<sup>217,285,286</sup> Outcome for an individual is influenced by patient-level factors such as age, co-morbid status, and aetiology of cardiac arrest. As such, predicting outcome at an individual patient level is challenging. Key challenges for clinicians are effective communication of uncertainty about the likely outcome if an individual has a cardiac arrest, and to ensure that their personal values and preferences do not influence the patient.

Individual autonomy gives individuals the right to decline a treatment but does not obligate a health system to provide a treatment that is either futile or not cost-effective. Publicly funded healthcare systems have limited resources with an expectation that systems use funding in the most effective way. Treatments that do not meet pre-defined cost-effectiveness thresholds may not be made available. To date, few cardiac arrest interventions have been subjected to a health economic evaluation.<sup>287–289</sup>

In recent years organ donation has been highlighted as an important outcome following cardiac arrest.<sup>290</sup> Organ donation provides benefit to the wider health system and society as a clinical and cost-effective treatment for organ failure.

In some cases, eliciting how an individual values a particular outcome may not be possible, such as in the context of young children or individuals with severe cognitive dysfunction. In these circumstances, clinicians should discuss treatment decisions with those close to the individual. Society often places particular value on the life of a child. Clinicians must take care to ensure that any decision is in the individual's best interests. In rare cases where the clinical team and other parties hold discordant views that cannot be reconciled, parties may need to defer decision-making to the legal system.

### Variability in outcome

Variability in outcome following cardiac arrest has been described in both IHCA and OHCA.<sup>217,285,286,291,292</sup> This variability may be between locality, EMS systems, hospitals, regions, and countries. Variation may reflect differences at several levels, including data collection methods, case-mix, and treatment.<sup>293,294</sup> From an ethical perspective, the key concern occurs when variability is caused by differences in treatment or processes of care.

Observational data suggest that females and individuals from socially deprived and ethnic minority groups are less likely to receive bystander CPR and key post-arrest interventions.<sup>295,296</sup> Survey data indicates that both in-hospital systems of care and long-term follow up and rehabilitation differ markedly between hospitals.<sup>291,297–299</sup>

One strategy proposed to improve patient outcomes is centralisation of hospital services across a range of conditions including cardiac arrest.<sup>300,301</sup> This enables the development of clinical expertise and facilitates delivery of specialist interventions, such as primary percutaneous coronary intervention and extracorporeal CPR. There is a concern that centralisation may disadvantage individuals that live in rural areas.

### Research and registry outcomes

Utstein statements describe the outcomes that should be collected by registries. Core outcomes are identified as ROSC, survival at hospital discharge/30-days, and neurological outcome at hospital discharge.<sup>302,303</sup> The inclusion of health-related quality of life and 12-month survival as supplementary outcomes reflects the balance between the importance of these outcomes, and the challenges of collection, such as the associated resource requirement.

In the context of research, differences in the way that outcomes are measured or reported by studies may preclude comparison of results between studies, and limit opportunities for meta-analysis.<sup>304</sup> A systematic review of cardiac arrest literature identified variability in the outcomes reported, differences in outcome definitions, and differences in the timepoint and method used to record outcomes.<sup>282</sup> The patient's perspective on outcome was rarely included.

To address this issue, ILCOR developed a cardiac arrest core outcome set (COSCA) in a process that involved patients, their partners, clinicians and researchers.<sup>305</sup> Core outcome sets describe the key outcomes that should be reported in all clinical trials, thereby ensuring consistency in outcome reporting.<sup>306,307</sup> COSCA identified three outcomes: survival at discharge/30-days; modified Rankin score at discharge/30-days; and health-related quality of life at 180-days/1-year. COSCA supports the collection of detailed measures of specific problems experienced by cardiac arrest survivors, such as fatigue, anxiety and societal participation. These data may improve our knowledge of cardiac arrest survivorship and patient support and rehabilitation in the post-acute phase.

An important challenge for both registries and clinical trials is ensuring a high level of data completeness for outcomes that rely on

patient or proxy engagement, such as health-related quality of life. Response rates vary markedly across trials.<sup>288,308,309</sup> A key concern is that respondents may be systematically different to non-respondents.<sup>310</sup> In cardiac arrest research, survivors with poor outcome are less likely to respond, leading to bias.<sup>311–313</sup> The SPIIRT (Standard Protocol Items: Recommendations for Interventional Trials) PRO (Patient-reported outcome) extension guidelines provide information on including patient reported outcomes in clinical trials.<sup>310</sup>

### Ethics and emergency research

#### Right to self-determination vs. scientific progress

The prognosis after cardiac arrest remains poor.<sup>314–316</sup> Therefore, there is a need for interventional, multicentre, randomised, controlled clinical research aimed at reliably assessing the effects of new and potentially beneficial treatments or validating empirical routine practice treatments of uncertain efficacy.<sup>63,195,317,318</sup> Striking the best balance between respect for autonomy (i.e. the right to self-determination) and beneficence (i.e. improving patient outcomes) or even non-maleficence (i.e. avoiding patient exposure to unproven treatments) has been recognised as one of the greatest challenges of emergency research conduct.<sup>63,195,317,319</sup>

The new European Union Clinical Trials Regulation No. 536/2014, permits the use of deferred consent in drug trials under clearly specified conditions. Tested interventions should be considered of minimal risk/burden for the subject in comparison with the standard treatment for the subject's condition.<sup>317</sup> Thus, the new regulation enables potentially beneficial, low-risk, multicentre, and multinational cardiac arrest research.<sup>195,317,320</sup> Nevertheless, regulatory improvements are still needed as the new regulation does not concern clinical trials evaluating devices.<sup>317</sup> Notably, device-related emergency research may confer considerable benefit in terms of leading to improvements in clinical practice and patient outcomes.<sup>321</sup>

Deferred consent (i.e. obtaining consent from a surrogate and/or patient as soon as possible after enrolment) may be necessary because the therapeutic window is too narrow to obtain a valid pre-enrolment consent.<sup>63,317,322–324</sup> This is considered as an ethically acceptable alternative for low-risk research, ensuring both the possibility of research benefit and respect for patient/family autonomy.<sup>325,326</sup> In contrast, a strict requirement for pre-enrolment consent may delay the initiation of an experimental intervention, thereby hampering its potential benefit to the patient.<sup>327</sup> Another ethically acceptable and legally supported consent model comprises exception to informed consent (EIC) with prior community consultation (and a possibility of prospective opt out for community members).<sup>328–335</sup> The EIC model also mandates obtaining post-enrolment consent.<sup>317</sup>

Both deferred consent and EIC models are limited by the patient's and or next-of-kin's right for consent withdrawal later on, as this may introduce bias in trial results by excluding the data from patients with a more complicated clinical course.<sup>63</sup> This might be partly addressed by regulatory provisions aimed at preventing the exclusion of patient data recorded until the time point of consent revocation.<sup>63</sup>

A recent pragmatic trial of adrenaline (epinephrine) in OHCA used a combination of a deferred consent model with informative press releases before and throughout the study period, a constantly updated trial website during the study period, an electronically supported opt-out option (which requires further evaluation), a pre-specified and realistic approach to inform the patient and request their consent after regaining their decisional capacity, a pre-specified and clear definition of personal and professional legal

representative for patients lacking decisional capacity, a pre-specified method of approach and communication with the legal representative, a clearly specified procedure for consent refusal or revocation, and a pre-specified approach for passive provision of trial information (e.g. through websites or newsletters) to the families of patients who died before their relatives could be contacted.<sup>318</sup> Future research should compare the relative potential benefits (i.e. less emotional stress) and harms (i.e. limited or no knowledge of the patient's trial participation details) of passive versus active provision of information (i.e. more stress but also more knowledge about the patient's trial participation).

During the design phase of the pragmatic adrenaline trial, the main outcomes were specified in collaboration with patient and public representatives.<sup>318</sup> Involvement of all major stakeholders (including patients and representatives of the public) in the iterative development of core outcome sets during study design, as well as conduct and delivery of the research and dissemination of its results is an emerging and promising practice. Indeed, this practice has already been adopted in several fields of research and may comprise patient-centric initiatives such as advocacy group support and involvement, patient advisory panels, and focus groups, interviews with trial participants and staff, questionnaires and Delphi surveys/consensus processes, and consensus meetings.<sup>336–344</sup>

The EIC model is based on the 1996 United States Food and Drug Administration regulation 21 CFR 50.24.<sup>345</sup> Although this regulation seems to provide clearly defined guidance for the conduct of emergency research, several authors have previously attributed to it significant procedural impediments.<sup>346,347</sup> For instance, if a family member is present in an emergency, it may not be feasible for the researcher to explain to them the research protocol, or even the concept of informed consent.<sup>348</sup> Furthermore, a survey with 530 respondents from a community participating in EIC research projects revealed that only 5% of the respondents were aware of ongoing research protocols despite pre-study community consultation. This casts doubt upon the feasibility of adequate dissemination of research information among research-participating communities.<sup>349</sup>

A worrying reduction of cardiac arrest trials of 15% per year between 1992 and 2002 was documented in the United States.<sup>350</sup> Similar worries were articulated for steep reductions of 30–50% in European trials submitted for grants or ethical approval by the end of 2005<sup>351–353</sup>; at that time, European Union Directive 2001/20/EC was in force and its strict interpretation mandated pre-enrolment consent for all types of drug clinical trials.<sup>63,354</sup>

The literature cited above highlights the inherent perplexity of respecting autonomy of patients lacking decisional capacity when enrolling them in emergency clinical research protocols aimed at improving their outcomes. This ongoing ethical dilemma could be partly addressable by advance care planning specifically pertaining to participation in emergency research. However, such care plans should also be immediately accessible by emergency healthcare staff and researchers, even in the setting of OHCA; this may still prove electronic resource-demanding or even impossible in many situations/settings.<sup>355</sup>

Large, national and international registries enable the recording of general population data on the incidence, presumed cause, and outcomes of cardiac arrest. Information about whether a patient collapse was witnessed or not, cardiac arrest location, certain aspects of emergency care organisation (e.g. availability of dispatcher-assisted bystander CPR), patient characteristics (e.g. age, sex, race, and comorbidities), treating hospital characteristics (e.g. bed size and teaching status), downtimes (e.g. time from collapse to first shock),

and treatments administered may also be collected.<sup>356,357</sup> Registry data can be analysed to (1) study regional variation, temporal trends, and predictors of patient outcomes; (2) compare propensity score matched patient subgroups receiving different treatments; and (3) gain insights into the implementation of published evidence and guidelines in routine clinical practice.<sup>316,356,358–361</sup> In addition, DNA biobanks have been established for DNA sequencing in the context of genomic research in sudden cardiac arrest.<sup>362</sup>

Big observational registry/biobank data originate from multiple sources. Such data may have to be linked for the detection of associations between potential predictor variables and patient outcomes.<sup>363</sup> The resulting production of high-quality evidence informing personalised prevention and treatment may contribute to improved outcomes and reduction of healthcare costs.<sup>364</sup> However, these beneficial processes are not free of ethical issues pertaining to privacy (i.e. risk of patient re-identification), genetic discrimination, and moral obligation for disclosure of findings to high-risk patients who decline becoming aware of their genetic test results. There are also challenges around observational data quality and potentially biased results leading to creation of incorrect risk profiles, obtaining consent for data use in an emergency research setting, and use of appropriate safeguards for data protection.<sup>362,365–374</sup>

The current European Union General Data Protection Regulation (GDPR) 2016/680 mandates that specific appropriate safeguards (e.g. safe data storage and encryption, access logging, data enclaves, etc.) be in place for the scientific processing of the data of a natural person. Records of processing activities must be kept by data controllers. A data protection impact assessment may be required to determine and confirm the risks relative to the subject's rights. GDPR compliance of research institutions must be monitored by a designated data protection officer.<sup>362</sup>

The GDPR does not concern anonymous data and data from deceased persons. Nevertheless, there are also stronger conditions concerning consent for the inclusion of personal patient data in research. Notably, a strict requirement for prospective (or pre-collection) informed consent would exclude collection of data from most sudden cardiac arrest patients. This would result in consent bias, skewing of the data, and compromised reliability of research results, with consequent societal harm. Furthermore, excluding collection of data from some incapacitated patients could potentially violate their preference to act in favour of the common good.<sup>362</sup> Therefore, for observational emergency research we suggest that local/regional supervising authorities consider allowing deferred and broad (i.e. for the overall research topic) consent, while concurrently ensuring the implementation of safeguards aimed at preventing data breaches and patient re-identification.<sup>362,375–377</sup> Lastly, and regarding both observational and interventional research, sometimes, it may not be possible to obtain even deferred consent, e.g. the patient dies and no surrogate decision maker can be located, or two surrogates of equal legal standing disagree. In such cases, we suggest consideration of permitting the use of the data collected until the time point of confirmation of the inability to obtain the consent.

### Equal distribution of research benefits and risks

Whenever certain communities or societal groups bear the burden of the risk of research-associated adverse events, they should also have the possibility of enjoying any benefits arising from the research results.<sup>63</sup> Indeed, the use of relevant scientific achievements should not be confined to other privileged populations not participating in the research protocol(s).<sup>63</sup>

### Access to best possible care and respect for patient/family dignity

Enrolment in a research protocol should in no way be linked to the quality or intensity of care. For instance, obtainment of deferred surrogate consent for a given patient's continued participation in a cardiac arrest trial evaluating therapeutic hypothermia should not result in preferential ICU admission of that patient over another patient whose proxy has refused consent.<sup>63</sup>

Researchers should also ensure that the dignity and privacy of the research participant and their family are respected. For example, enrollees in a cardiac arrest trial should be referred to as post cardiac arrest patients rather than cardiac arrests or cardiac arrest victims.<sup>63</sup>

### Study design issues, and transparency of study conduct and reporting of results

Previously identified ethical issues concerning mainly commercial research have prompted the requirement for pre-enrolment registration of trial protocols,<sup>63,378</sup> reporting of any protocol and trial status changes (e.g., temporary suspension) throughout the period of trial conduct, and posting of main results to the trial registry within 12 months of study completion and publication in a peer-reviewed journal after another 12 months.<sup>63,379</sup> At the time of paper submission to a peer-reviewed journal, authors are normally obliged to report on the sponsor's role as well as on their own contributions to the study, and also approve the submission.<sup>63</sup> Furthermore, data sharing policies could be adopted to further enhance research transparency.<sup>63,380</sup>

Another concern pertains to the substantially disproportionate funding favouring commercial research evaluating the efficacy of high-cost, patent protected drugs or devices over the undoubtedly necessary, non-commercial, academic resuscitation research on patent-unprotected, low-cost, widely used drugs of potentially uncertain efficacy, such as adrenaline (epinephrine), or antiarrhythmics.<sup>63,318,381,382</sup> This may partly explain the fact that BLS/ALS guidelines are based on 35–53-fold fewer RCTs/10,000 deaths/year relative to guidelines for acute cardiovascular events and heart failure.<sup>63,381</sup> Governmental, or non-profit organisation, or even mixed public and private/industrial funding of resuscitation research needs therefore to be increased.<sup>63,383</sup> Furthermore, such funding should be fairly and proportionately distributed between studies of in-hospital and pre-hospital interventions, preferably also according to their estimated effect(s) on patient outcomes.<sup>384</sup>

### Emergency research and the COVID-19 mass casualty crisis.

COVID-19 case surges may cause disruption over a wide spectrum of societal and healthcare system activities.<sup>1,385–387</sup> Accordingly, processes and procedures primarily related to interventional research may be hindered or halted. The need for physical distancing may cause cancellation of face-to-face meetings concerning study design (see also above), study protocol approval, and evaluation of the progress of study conduct (by investigators, and data monitoring committees); nevertheless, physical meeting issues can be at least partly addressed by using digital telecommunication technology. Delays in CPR initiation due to donning of personal protective equipment may impact patient outcomes,<sup>385,386,388</sup> and thereby modify the measured effect of concurrent or subsequent, investigational, resuscitative interventions, such as new drug therapies, or temperature/ventilatory management during and/or after resuscitation. In OHCA, increases in the volume of emergency calls in the context of a saturated healthcare system may prolong arrival times of emergency medical services, whereas the potential risk of contracting

the disease while performing chest compressions may reduce bystander CPR rates.<sup>3,389</sup> Again, both latter factors may impact patient outcomes, and ultimately, the results of any ongoing emergency research. Fear of contracting the infection and/or excessive workload may discourage healthcare professionals from participating in research teams or initiating and leading a research project.<sup>89,390–392</sup> Lastly, increases in DNACPR decisions and especially use of blanket CPR exclusion criteria such as age<sup>87,88,385,390,393</sup> may introduce selection bias and hamper the generalisability of the research results, as well as their applicability to normal conditions. Such challenges may be addressable solely through effective governmental policies limiting viral spread and preventing healthcare system overload.

### Future directions

The evidence supporting autonomy-safeguarding interventions exhibits several limitations, such as diversity/variability in definitions of key terms (see also [Tables 1–3](#) and online supplement), evaluated intervention type/design, geographical distribution of studies and characteristics of participating populations (e.g. type of life-limiting illness, religion/religiosity, ethnicity, etc.) specified outcomes and methods of their determination, and reliability of reported results (further details provided in the supplement).

These weaknesses have either precluded the conduct of meta-analyses or increased the heterogeneity of reported meta-analyses results. Accordingly, the certainty of existing evidence has been judged most frequently as low to very low by authors of systematic reviews.<sup>7–10,28,38,39,43,44,72</sup>

As a result, scientific gaps exist regarding the actual effects of advance directives, advance care planning, and shared decision making on patient outcomes. These gaps range from uncertainty about the effect estimates of meta-analyses (in the presence of substantial pertinent literature), to very limited data from nonrandomised studies, and/or even the absence of relevant studies (e.g. the case of healthcare-related quality of life after cardiac arrest; see also online supplement).

Therefore, new, high-quality, and preferably multinational RCTs, based on clear and wide consensus-based definitions of interventions and outcomes are warranted. Observational big data potentially matching the strength of RCT data<sup>394,395</sup> and qualitative research identifying key issues that need to be addressed are also needed.<sup>28,29,59,71,83,396</sup> Further study is also warranted to establish the effectiveness of inter-professional shared decision making, which has been recently recommended by experts for important clinical decisions. Inter-professional shared decision making takes into account the available evidence, the expertise of involved clinicians, and the patient's values goals, and preferences.<sup>397</sup>

Despite the limitations of the currently available, substantial but still heterogenous, body of evidence, the presence of either positive or neutral RCTs on structured communication tools aimed at facilitating the completion of advance care directives and plans suggests a class effect and increased likelihood of benefit compared with usual care.<sup>4,7</sup> Structured, complex, multifaceted interventions in the context of advance care planning and shared decision making may effectively prevent disproportionate/unwanted end-of-life care and accordingly reduce use of healthcare resources.<sup>4,7,44,72,355,398</sup> Future pertinent research should primarily be guided by scientific evidence.

Potentially successful, organisational interventions include: (1) structural educational initiatives of the public (e.g. informational videos, media coverage, and patient-public involvement workshops); (2) systematic training of healthcare professionals in ethics and communication skills<sup>355</sup>; (3) infrastructural initiatives enabling emergency healthcare providers to instantly access and honour the patient's recorded wishes (e.g. establishment of electronic registries/health records and appropriate regulatory provisions<sup>355,398</sup>); (4) public involvement to ensure clarity and acceptability of electronic documents used for the recording of treatment options; (5) immediate availability of adequate palliative care services upon patient/family request – this pertains to paediatric palliative care as well<sup>399</sup>; and (6) continuous monitoring of the quality of care supporting relevant improvement efforts/initiatives.

During a pandemic such as COVID-19, patient/family engagement in advance care planning and shared decision-making should still be feasible as part of remote clinical monitoring and care models (ClinicalTrials.gov NCT04425720).

The Ethics writing group emphasises the importance of thorough societal consultation and debate to provide a context-specific ethical framework for many of the complex resuscitation decisions such as use of extracorporeal CPR or uncontrolled donation after circulatory death.

Systems should continue to evaluate the performance of their decision making with regard to withdrawing or withholding life support, including the potential use of specific ToR rules, the degree of implementation of advance directives, and the number of advanced CPR cases. As technology progresses, it is likely that these concepts will evolve as well.

Systems should try and better define the place and remit of bystanders and first responders, as well as the ethical challenges around bystander CPR particularly in respect of the balance between benefit for the victim and harm to the rescuer.

There is a need to measure and track outcomes that are meaningful to both patients generally and the specific patient being treated.

Future, high-quality research should identify the optimal educational method for healthcare professionals on standardised patient outcome sets, and also evaluate its effect on healthcare professional's understanding of patient's preferences.

Systems should consider educational interventions to introduce the concept of family presence during resuscitation. Future research projects should consider the identification of healthcare personnel best suited to guide families through the resuscitation, and also being able to provide comfort, recognise family distress and participate in debriefing sessions after resuscitation.

More research is needed to determine how best to prepare and support ambulance personnel for the challenges of resuscitation decision-making and patient death, acknowledging the unique contextual demands of the prehospital setting.

We suggest the broadest possible/multinational establishment of harmonised regulations for emergency research aimed at fostering interventional drug and device trials as well as observational studies, while concurrently safeguarding participants' autonomy and protection/integrity of their personal data.

## Conclusions

The Ethics writing group has provided sets of simple and clear recommendations supported by a wealth of systematic reviews, recent RCTs and nonrandomised studies. Despite the generally low certainty about the precision of the effect estimates of several

evaluated meta-analyses, the directions of the effects on patient outcomes clearly favour the use of interventions such as advance care planning, shared decision making, and ToR rules. The writing group also produced three narrative reviews to summarise the existing key evidence/knowledge/issues on education/system organisation, patient outcomes, and ethics of emergency research. Lastly, the writing group has provided a set of consensus definitions of key terms, which could potentially prove useful in both routine clinical practice and the design of future research protocols.

## Conflict of interest

MB declares her role of co-coordinator EU project ESCAPE-NET. GDP reports funding from Elsevier for his role as an editor of the journal Resuscitation. He reports research funding from the National Institute for Health Research (NIHR) in relation to the PARAMEDIC2 trial and the RESPECT project.

## Acknowledgments

The writing group thanks Nele Pauwels, information specialist at Ghent University, Belgium for her support in developing the necessary search strategies. GDP is supported by the NIHR Applied Research Collaboration West Midlands. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.resuscitation.2021.02.017>.

## REFERENCES

1. Bavel JJV, Baicker K, Boggio PS, et al. Using social and behavioural science to support COVID-19 pandemic response. *Nat Hum Behav* 2020;4:460–71.
2. Swire-Thompson B, Lazer D. Public health and online misinformation: challenges and recommendations. *Annu Rev Public Health* 2020;41:433–51.
- 2a. Perkins GD, Graesner JT, Semeraro F, et al. European Resuscitation Council Guidelines 2021 – Executive summary. *Resuscitation* 2021;161.
3. Nolan JP, Monsieurs KG, Bossaert L, et al. European Resuscitation Council COVID-19 guidelines executive summary. *Resuscitation* 2020;153:45–55.
4. Oczkowski SJ, Chung HO, Hanvey L, Mbuagbaw L, You JJ. Communication tools for end-of-life decision-making in ambulatory care settings: a systematic review and meta-analysis. *PLOS ONE* 2016;11:e0150671.
5. Higel T, Alaoui A, Bouton C, Fournier JP. Effect of living wills on end-of-life care: a systematic review. *J Am Geriatr Soc* 2019;67:164–71.
- 5a. Sprung CL, Truog RD, Curtis JR, et al. Seeking worldwide professional consensus on the principles of end-of-life care for the critically ill. The Consensus for Worldwide End-of-Life Practice for Patients in Intensive Care Units (WELPICUS) study. *Am J Respir Crit Care Med* 2014;190(8):855–66. doi:<http://dx.doi.org/10.1164/rccm.201403-0593CC>.

- 5b. Gilbert J, Boag J. Nonstandard Advance Health Care Directives in Emergency Departments: Ethical and Legal Dilemma or Reality: a narrative review. *Adv Emerg Nurs J* 2018;40(4):324–7, doi:http://dx.doi.org/10.1097/TME.0000000000000214.
- 5c. Martin DK, Emanuel LL, Singer PA. Planning for the end of life. *Lancet* 2000;356(9242):1672–6, doi:http://dx.doi.org/10.1016/S0140-6736(00)03168-8.
- 5d. Santonocito C, Ristagno G, Gullo A, Weil MH. Do-not-resuscitate order: a view throughout the world. *J Crit Care* 2013;28(1):14–21, doi:http://dx.doi.org/10.1016/j.jcrrc.2012.07.005.
- 5e. Perez Mdel V, Macchi MJ, Agranatti AF. Advance directives in the context of end-of-life palliative care. *Curr Opin Support Palliat Care* 2013;7(4):406–10, doi:http://dx.doi.org/10.1097/SPC.0000000000000007.
- 5f. Bossaert LL, Perkins GD, Askitopoulou H, et al. European Resuscitation Council Guidelines for Resuscitation 2015: Section 11. The ethics of resuscitation and end-of-life decisions. *Resuscitation* 2015;95:302–11, doi:http://dx.doi.org/10.1016/j.resuscitation.2015.07.033.
- 5g. Andorno R, Biller-Andorno N, Brauer S. Advance health care directives: towards a coordinated European policy? *Eur J Health Law* 2009;16(3):207–27. <http://www.ncbi.nlm.nih.gov/pubmed/19788001>.
- 5h. Davidson JE, Aslakson RA, Long AC, et al. Guidelines for Family-Centered Care in the Neonatal, Pediatric, and Adult ICU. *Crit Care Med* 2017;45(1):103–28, doi:http://dx.doi.org/10.1097/CCM.00000000000002169.
6. Sutter R, Meyer-Zehnder B, Baumann SM, Marsch S, Pargger H. Advance directives in the neurocritically ill: a systematic review. *Crit Care Med* 2020;48:1188–95.
7. Becker C, Lecheler L, Hochstrasser S, et al. Association of communication interventions to discuss code status with patient decisions for do-not-resuscitate orders: a systematic review and meta-analysis. *JAMA Netw Open* 2019;2:e195033.
8. Houben CHM, Spruit MA, Groenen MTJ, Wouters EFM, Janssen DJA. Efficacy of advance care planning: a systematic review and meta-analysis. *J Am Med Assoc* 2014;15:477–89.
9. Brinkman-Stoppelenburg A, Rietjens JA, van der Heide A. The effects of advance care planning on end-of-life care: a systematic review. *Palliat Med* 2014;28:1000–25.
10. Field RA, Fritz Z, Baker A, Grove A, Perkins GD. Systematic review of interventions to improve appropriate use and outcomes associated with do-not-attempt-cardiopulmonary-resuscitation decisions. *Resuscitation* 2014;85:1418–31.
11. Reinhardt JP, Downes D, Cimarolli V, Bomba P. End-of-life conversations and hospice placement: association with less aggressive care desired in the nursing home. *J Soc Work End Life Palliat Care* 2017;13:61–81.
12. Siewiera J, Tomaszewski D, Piechocki J, Kubler A. Withholding and withdrawing life-sustaining treatment: experiences in limiting futile therapy from three Polish intensive care departments. *Adv Clin Exp Med* 2019;28:541–6.
13. Fritz Z, Slowther AM, Perkins GD. Resuscitation policy should focus on the patient, not the decision. *BMJ* 2017;356:j813.
14. Mockford C, Fritz Z, George R, et al. Do not attempt cardiopulmonary resuscitation (DNACPR) orders: a systematic review of the barriers and facilitators of decision-making and implementation. *Resuscitation* 2015;88:99–113.
15. Fokin AA, Wycech J, Katz JK, et al. Palliative care consultations in trauma patients and role of do-not-resuscitate orders: propensity-matched study. *Am J Hosp Palliat Care* 2020;37:1068–75.
16. Mitchell SL, Shaffer ML, Cohen S, Hanson LC, Habtemariam D, Volandes AE. An advance care planning video decision support tool for nursing home residents with advanced dementia: a cluster randomized clinical trial. *JAMA Intern Med* 2018;178:961–9.
17. El-Jawahri A, Mitchell SL, Paasche-Orlow MK, et al. A randomized controlled trial of a CPR and intubation video decision support tool for hospitalized patients. *J Gen Intern Med* 2015;30:1071–80.
18. Merino AM, Greiner R, Hartwig K. A randomized controlled trial of a CPR decision support video for patients admitted to the general medicine service. *J Hosp Med* 2017;12:700–4.
19. Cappell K, Sundaram V, Park A, et al. Advance directive utilization is associated with less aggressive end-of-life care in patients undergoing allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2018;24:1035–40.
20. Fu PK, Tung YC, Wang CY, et al. Early and late do-not-resuscitate (DNR) decisions in patients with terminal COPD: a retrospective study in the last year of life. *Int J Chron Obstruct Pulmon Dis* 2018;13:2447–54.
21. Stream S, Nolan A, Kwon S, Constable C. Factors associated with combined do-not-resuscitate and do-not-intubate orders: a retrospective chart review at an urban tertiary care center. *Resuscitation* 2018;130:1–5.
22. Reuter PG, Agostinucci JM, Bertrand P, et al. Prevalence of advance directives and impact on advanced life support in out-of-hospital cardiac arrest victims. *Resuscitation* 2017;116:105–8.
23. Kuchinad KE, Strowd R, Evans A, Riley WA, Smith TJ. End of life care for glioblastoma patients at a large academic cancer center. *J Neurooncol* 2017;134:75–81.
24. Mills A, Walker A, Levinson M, et al. Resuscitation orders in acute hospitals: a point prevalence study. *Australas J Ageing* 2017;36:32–7.
25. McCarroll CM. Increasing access to palliative care services in the intensive care unit. *Dimens Crit Care Nurs* 2018;37:180–92.
26. Davidson JE, Aslakson RA, Long AC, et al. Guidelines for family-centered care in the neonatal, pediatric, and adult ICU. *Crit Care Med* 2017;45:103–28.
27. MacKenzie MA, Smith-Howell E, Bomba PA, Meghani SH. Respecting choices and related models of advance care planning: a systematic review of published evidence. *Am J Hosp Palliat Care* 2018;35:897–907.
28. O'Halloran P, Noble H, Norwood K, et al. Advance care planning with patients who have end-stage kidney disease: a systematic realist review. *J Pain Symptom Manag* 2018;56:795–807 e18.
29. Voss H, Vogel A, Wagemans AMA, et al. Advance care planning in palliative care for people with intellectual disabilities: a systematic review. *J Pain Symptom Manag* 2017;54:938–60 e1.
30. Weathers E, O'Caoimh R, Cornally N, et al. Advance care planning: a systematic review of randomised controlled trials conducted with older adults. *Maturitas* 2016;91:101–9.
31. Martin RS, Hayes B, Gregorevic K, Lim WK. The effects of advance care planning interventions on nursing home residents: a systematic review. *J Am Med Assoc* 2016;17:284–93.
32. Lim CE, Ng RW, Cheng NC, Cigolini M, Kwok C, Brennan F. Advance care planning for haemodialysis patients. *Cochrane Database Syst Rev* 2016;7:CD010737.
33. Nair R, Kohen SA. Can a patient-directed video improve inpatient advance care planning? A prospective pre-post cohort study. *BMJ Qual Saf* 2019;28:887–93.
34. Wen FH, Chen JS, Chou WC, Chang WC, Hsieh CH, Tang ST. Extent and determinants of terminally ill cancer patients' concordance between preferred and received life-sustaining treatment states: an advance care planning randomized trial in Taiwan. *J Pain Symptom Manag* 2019;58: 1–10e.
35. Jennings LA, Turner M, Keebler C, et al. The effect of a comprehensive dementia care management program on end-of-life care. *J Am Geriatr Soc* 2019;67:443–8.
36. Malhotra C, Sim D, Jaufeerally FR, et al. Impact of a formal advance care planning program on end-of-life care for patients with heart failure: results from a randomized controlled trial. *J Card Fail* 2020;26:594–8.
37. Walczak A, Butow PN, Bu S, Clayton JM. A systematic review of evidence for end-of-life communication interventions: who do they target, how are they structured and do they work? *Patient Educ Couns* 2016;99:3–16.
38. Kavalieratos D, Corbelli J, Zhang D, et al. Association between palliative care and patient and caregiver outcomes: a systematic review and meta-analysis. *JAMA* 2016;316:2104–14.



39. Huber MT, Highland JD, Krishnamoorthi VR, Tang JW. Utilizing the electronic health record to improve advance care planning: a systematic review. *Am J Hosp Palliat Care* 2018;35:532–41.
40. Kang E, Lee J, Choo J, Min J, Yun YH. Randomized controlled trial of advance care planning video decision aid for the general population. *J Pain Symptom Manag* 2020;59:1239–47.
41. Nedjat-Haiem FR, Cadet TJ, Amatya A, Thompson B, Mishra SI. Efficacy of motivational interviewing to enhance advance directive completion in Latinos with chronic illness: a randomized controlled trial. *Am J Hosp Palliat Care* 2019;36:980–92.
42. Denis N, Timsit JF, Giaj Levra M, et al. Impact of systematic advanced care planning in lung cancer patients: a prospective study. *Respir Med Res* 2020;77:11–7.
43. Jain A, Corriveau S, Quinn K, Gardhouse A, Vegas DB, You JJ. Video decision aids to assist with advance care planning: a systematic review and meta-analysis. *BMJ Open* 2015;5:e007491.
44. Oczkowski SJ, Chung HO, Hanvey L, Mbuagbaw L, You JJ. Communication tools for end-of-life decision-making in the intensive care unit: a systematic review and meta-analysis. *Crit Care* 2016;20:97.
45. El-Jawahri A, Paasche-Orlow MK, Matlock D, et al. Randomized controlled trial of an advance care planning video decision support tool for patients with advanced heart failure. *Circulation* 2016;134:52–60.
46. Chan HY, Ng JS, Chan KS, et al. Effects of a nurse-led post-discharge advance care planning programme for community-dwelling patients nearing the end of life and their family members: a randomised controlled trial. *Int J Nurs Stud* 2018;87:26–33.
47. Green MJ, Van Scoy LJ, Foy AJ, Dimmock AEF, Lehman E, Levi BH. Patients with advanced cancer choose less aggressive medical treatment on vignettes after using a computer-based decision aid. *Am J Hosp Palliat Care* 2020;37:537–41.
48. Kizawa Y, Okada H, Kawahara T, Morita T. Effects of brief nurse advance care planning intervention with visual materials on goal-of-care preference of Japanese elderly patients with chronic disease: a pilot randomized-controlled trial. *J Palliat Med* 2020;23:1076–83.
49. DeCoursey DD, Silverman M, Oladunjoye A, Wolfe J. Advance care planning and parent-reported end-of-life outcomes in children adolescents, and young adults with complex chronic conditions. *Crit Care Med* 2019;47:101–8.
50. Kernick LA, Hogg KJ, Millerick Y, Murtagh FEM, Djahit A, Johnson M. Does advance care planning in addition to usual care reduce hospitalisation for patients with advanced heart failure: a systematic review and narrative synthesis. *Palliat Med* 2018;32:1539–51.
51. Klingler C, in der Schmitt J, Marckmann G. Does facilitated advance care planning reduce the costs of care near the end of life? Systematic review and ethical considerations. *Palliat Med* 2016;30:423–33.
52. Overbeek A, Polinder S, Haagsma J, et al. Advance care planning for frail older adults: findings on costs in a cluster randomised controlled trial. *Palliat Med* 2019;33:291–300.
53. Scarpi E, Dall'Agata M, Zagonel V, et al. Systematic vs. on-demand early palliative care in gastric cancer patients: a randomized clinical trial assessing patient and healthcare service outcomes. *Support Care Cancer* 2019;27:2425–34.
54. Pedraza SL, Culp S, Knestrick M, Falkenstine E, Moss AH. Association of physician orders for life-sustaining treatment form use with end-of-life care quality metrics in patients with cancer. *J Oncol Pract* 2017;13:e881–8.
55. Thomas P, Hazif-Thomas C. Discourse on life and Alzheimer's disease. *Soins Gerontol* 2018;23:31–3.
56. Hoell JI, Weber HL, Balzer S, et al. Advance care planning and outcome in pediatric palliative home care. *Oncotarget* 2018;9:17867–75.
57. Chong PH, De Castro Molina JA, Teo K, Tan WS. Paediatric palliative care improves patient outcomes and reduces healthcare costs: evaluation of a home-based program. *BMC Palliat Care* 2018;17:11.
58. Suraarunsumrit P, Nopmaneejumruslers C, Srinonprasert V. Advance care planning (ACP) associated with reduced health care utilization in deceased older patients with advanced stage of chronic diseases. *J Med Asso Thai* 2019;102:801–8.
59. Marsac ML, Kindler C, Weiss D, Ragsdale L. Let's talk about it: supporting family communication during end-of-life care of pediatric patients. *J Palliat Med* 2018;21:862–78.
60. Walczak A, Butow PN, Tattersall MH, et al. Encouraging early discussion of life expectancy and end-of-life care: a randomised controlled trial of a nurse-led communication support program for patients and caregivers. *Int J Nurs Stud* 2017;67:31–40.
61. Overbeek A, Korlage IJ, Jabbarian LJ, et al. Advance care planning in frail older adults: a cluster randomized controlled trial. *J Am Geriatr Soc* 2018;66:1089–95.
62. Brazil K, Carter G, Cardwell C, et al. Effectiveness of advance care planning with family carers in dementia nursing homes: a paired cluster randomized controlled trial. *Palliat Med* 2018;32:603–12.
63. Mentzelopoulos SD, Slowther AM, Fritz Z, et al. Ethical challenges in resuscitation. *Intensive Care Med* 2018;44:703–16.
64. Kon AA, Davidson JE, Morrison W, et al. Shared decision making in ICUs: an american college of critical care medicine and american thoracic society policy statement. *Crit Care Med* 2016;44:188–201.
65. Gonella S, Basso I, Dimonte V, et al. Association between end-of-life conversations in nursing homes and end-of-life care outcomes: a systematic review and meta-analysis. *J Am Med Assoc* 2019;20:249–61.
66. Allen LA, McIlvennan CK, Thompson JS, et al. Effectiveness of an intervention supporting shared decision making for destination therapy left ventricular assist device: the DECIDE-LVAD randomized clinical trial. *JAMA Intern Med* 2018;178:520–9.
67. Hanson LC, Zimmerman S, Song MK, et al. Effect of the goals of care intervention for advanced dementia: a randomized clinical trial. *JAMA Intern Med* 2017;177:24–31.
68. DeMartino ES, Braus NA, Sulmasy DP, et al. Decisions to withdraw extracorporeal membrane oxygenation support: patient characteristics and ethical considerations. *Mayo Clin Proc* 2019;94:620–7.
69. Hsu NC, Huang CC, Chen WC, Yu CJ. Impact of patient-centred and family-centred care meetings on intensive care and resource utilisation in patients with terminal illness: a single-centre retrospective observational study in Taiwan. *BMJ Open* 2019;9:e021561.
70. Wilson ME, Krupa A, Hinds RF, et al. A video to improve patient and surrogate understanding of cardiopulmonary resuscitation choices in the ICU: a randomized controlled trial. *Crit Care Med* 2015;43:621–9.
71. Chen C, Michaels J, Meeker MA. Family outcomes and perceptions of end-of-life care in the intensive care unit: a mixed-methods review. *J Palliat Care* 2020;35:143–53.
72. Lee HW, Park Y, Jang EJ, Lee YJ. Intensive care unit length of stay is reduced by protocolized family support intervention: a systematic review and meta-analysis. *Intensive Care Med* 2019;45:1072–81.
73. Martin RS, Hayes BJ, Hutchinson A, Tacey M, Yates P, Lim WK. Introducing goals of patient care in residential aged care facilities to decrease hospitalization: a cluster randomized controlled trial. *J Am Med Assoc* 2019;20:1318–24 e2.
74. Hinkle LJ, Bosslet GT, Torke AM. Factors associated with family satisfaction with end-of-life care in the ICU: a systematic review. *Chest* 2015;147:82–93.
75. White DB, Angus DC, Shields AM, et al. A randomized trial of a family-support intervention in intensive care units. *N Engl J Med* 2018;378:2365–75.
76. Curtis JR, Trece PD, Nielsen EL, et al. Randomized trial of communication facilitators to reduce family distress and intensity of end-of-life care. *Am J Respir Crit Care Med* 2016;193:154–62.
77. Curtis JR, Nielsen EL, Trece PD, et al. Effect of a quality-improvement intervention on end-of-life care in the intensive care unit: a randomized trial. *Am J Respir Crit Care Med* 2011;183:348–55.
78. Sulmasy DP, Hughes MT, Yenokyan G, et al. The trial of ascertaining individual preferences for loved ones' role in end-of-life decisions (TAILORED) study: a randomized controlled trial to improve

- surrogate decision making. *J Pain Symptom Manag* 2017;54:455–65.
79. Carson SS, Cox CE, Wallenstein S, et al. Effect of palliative care-led meetings for families of patients with chronic critical illness: a randomized clinical trial. *JAMA* 2016;316:51–62.
  80. Lautrette A, Darmon M, Megarbane B, et al. A communication strategy and brochure for relatives of patients dying in the ICU. *N Engl J Med* 2007;356:469–78.
  81. Davidson JE, Jones C, Bienvenu OJ. Family response to critical illness: postintensive care syndrome-family. *Crit Care Med* 2012;40:618–24.
  82. Adelman RD, Tmanova LL, Delgado D, Dion S, Lachs MS. Caregiver burden: a clinical review. *JAMA* 2014;311:1052–60.
  83. DeSanto-Madeya S, Safizadeh P. Family satisfaction with end-of-life care in the intensive care unit: a systematic review of the literature. *Dimens Crit Care Nurs* 2017;36:278–83.
  84. You JJ, Jayaraman D, Swinton M, Jiang X, Heyland DK. Supporting shared decision-making about cardiopulmonary resuscitation using a video-based decision-support intervention in a hospital setting: a multisite before-after pilot study. *CMAJ Open* 2019;7:E630–7.
  85. Sahgal S, Yande A, Thompson BB, et al. Surrogate satisfaction with decision making after intracerebral hemorrhage. *Neurocrit Care* 2020.
  86. Scheunemann LP, Ernecoff NC, Buddadhumaruk P, et al. Clinician-family communication about patients' values and preferences in intensive care units. *JAMA Intern Med* 2019;179:676–84.
  87. Farrell TW, Ferrante LE, Brown T, et al. AGS position statement: resource allocation strategies and age-related considerations in the COVID-19 era and beyond. *J Am Geriatr Soc* 2020;68:1136–42.
  88. Farrell TW, Francis L, Brown T, et al. Rationing limited healthcare resources in the COVID-19 era and beyond: ethical considerations regarding older adults. *J Am Geriatr Soc* 2020;68:1143–9.
  89. Block BL, Smith AK, Sudore RL. During COVID-19 outpatient advance care planning is imperative: we need all hands on deck. *J Am Geriatr Soc* 2020;68:1395–7.
  90. Zheng RJ, Fu Y, Xiang QF, et al. Knowledge, attitudes, and influencing factors of cancer patients toward approving advance directives in China. *Support Care Cancer* 2016;24:4097–103.
  91. Hopkins SA, Lovick R, Polak L, et al. Reassessing advance care planning in the light of covid-19. *BMJ* 2020;369:m1927.
  92. Bledsoe TA, Jokela JA, Deep NN, Snyder Sulmasy L. Universal do-not-resuscitate orders social worth, and life-years: opposing discriminatory approaches to the allocation of resources during the COVID-19 pandemic and other health system catastrophes. *Ann Intern Med* 2020;173:230–2.
  93. White DB, Lo B. A framework for rationing ventilators and critical care beds during the COVID-19 pandemic. *JAMA* 2020;323:1773–4.
  94. Mentzelopoulos SD, Bossaert L, Greif R. Coronavirus disease 2019 and ethical considerations for extracorporeal cardiopulmonary resuscitation. *Resuscitation* 2020;154:127–8.
  95. Lee J, Abruken L, Flores S, et al. Early intervention of palliative care in the emergency department during the COVID-19 pandemic. *JAMA Intern Med* 2020;180:1252–4.
  96. Greif R, Bhanji F, Bigham BL, et al. Education implementation, and teams: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2020;156:A188–239.
  97. Pellegrino ED. Is it ethical to withdraw low-burden interventions in chronically ill patients? *JAMA* 2000;284:1380–2.
  98. Druwe P, Monsieurs KG, Piers R, et al. Perception of inappropriate cardiopulmonary resuscitation by clinicians working in emergency departments and ambulance services: the REAPPROPRIATE international, multi-centre, cross sectional survey. *Resuscitation* 2018;132:112–9.
  99. Munoz MG, Beyda DH. An ethical justification for termination of resuscitation protocols for pediatric patients. *Pediatr Emerg Care* 2017;33:505–15.
  100. Ranola PA, Merchant RM, Perman SM, et al. How long is long enough, and have we done everything we should?—ethics of calling codes. *J Med Ethics* 2015;41:663–6.
  101. Mercurio MR, Murray PD, Gross I. Unilateral pediatric “do not attempt resuscitation” orders: the pros, the cons, and a proposed approach. *Pediatrics* 2014;133(Suppl. 1):S37–43.
  102. Weise KL, Okun AL, Carter BS, et al. Guidance on forgoing life-sustaining medical treatment. *Pediatrics* 2017;140.
  103. Javaudin F, Le Bastard Q, Lascarrou JB, et al. The futility of resuscitating an out-of-hospital cardiac arrest cannot be summarized by three simple criteria. *Resuscitation* 2019;144:199–200.
  104. Grunau B, Scheuermeyer F, Kawano T, et al. North American validation of the Bokutoh criteria for withholding professional resuscitation in non-traumatic out-of-hospital cardiac arrest. *Resuscitation* 2019;135:51–6.
  105. Rosoff PM, Schneiderman LJ. Irrational exuberance: cardiopulmonary resuscitation as fetish. *Am J Bioeth* 2017;17:26–34.
  106. Druwe P, Benoit DD, Monsieurs KG, et al. Cardiopulmonary resuscitation in adults over 80: outcome and the perception of appropriateness by clinicians. *J Am Geriatr Soc* 2020;68:39–45.
  107. Van Norman GA. Decisions regarding forgoing life-sustaining treatments. *Curr Opin Anaesthesiol* 2017;30:211–6.
  108. Ting PS, Chen L, Yang WC, Huang TS, Wu CC, Chen YY. Gender and age disparity in the initiation of life-supporting treatments: a population-based cohort study. *BMC Med Ethics* 2017;18:62.
  109. Chiang WC, Ko PC, Chang AM, et al. Bystander-initiated CPR in an Asian metropolitan: does the socioeconomic status matter? *Resuscitation* 2014;85:53–8.
  110. Ahaddour C, Van den Branden S, Broeckaert B. Between quality of life and hope. Attitudes and beliefs of Muslim women toward withholding and withdrawing life-sustaining treatments. *Med Health Care Philos* 2018;21:347–61.
  111. Chamsi-Pasha H, Albar MA. Do not resuscitate, brain death, and organ transplantation: Islamic perspective. *Avicenna J Med* 2017;7:35–45.
  112. Hansen C, Lauridsen KG, Schmidt AS, Lofgren B. Decision-making in cardiac arrest: physicians' and nurses' knowledge and views on terminating resuscitation. *Open Access Emerg Med* 2019;11:1–8.
  113. Waldrop DP, Waldrop MR, McGinley JM, Crowley CR, Clemency B. Prehospital providers' perspectives about online medical direction in emergency end-of-life decision-making. *Prehosp Emerg Care* 2020;1–15.
  114. Kang M, Kim J, Kim K. Resuscitation duration inequality by patient characteristics in emergency department out-of-hospital cardiac arrest: an observational study. *Clin Exp Emerg Med* 2014;1:87–93.
  115. Wiel E, Di Pompeo C, Segal N, et al. Age discrimination in out-of-hospital cardiac arrest care: a case-control study. *Eur J Cardiovasc Nurs* 2018;17:505–12.
  116. Teefy J, Cram N, Van Zyl T, Van Aarsen K, McLeod S, Dukelow A. Evaluation of the uptake of a prehospital cardiac arrest termination of resuscitation rule. *J Emerg Med* 2020;58:254–9.
  117. Ozer J, Alon G, Leykin D, Varon J, Aharonson-Daniel L, Einav S. Culture and personal influences on cardiopulmonary resuscitation—results of international survey. *BMC Med Ethics* 2019;20:102.
  118. Long AC, Brumback LC, Curtis JR, et al. Agreement with consensus statements on end-of-life care: a description of variability at the level of the provider hospital, and country. *Crit Care Med* 2019;47:1396–401.
  119. Wang CH, Chang WT, Huang CH, et al. Factors associated with the decision to terminate resuscitation early for adult in-hospital cardiac arrest: Influence of family in an East Asian society. *PLOS ONE* 2019;14:e0213168.
  120. Brooks SC, Schmicker RH, Cheskes S, et al. Variability in the initiation of resuscitation attempts by emergency medical services personnel during out-of-hospital cardiac arrest. *Resuscitation* 2017;117:102–8.

121. Phua J, Joynt GM, Nishimura M, et al. Withholding and withdrawal of life-sustaining treatments in low-middle-income versus high-income Asian countries and regions. *Intensive Care Med* 2016;42:1118–27.
122. Campwala RT, Schmidt AR, Chang TP, Nager AL. Factors influencing termination of resuscitation in children: a qualitative analysis. *Int J Emerg Med* 2020;13:12.
123. Obladen M. Despising the weak: long shadows of infant murder in Nazi Germany. *Arch Dis Child Fetal Neonatal Ed* 2016;101:F190–4.
124. Brick C, Kahane G, Wilkinson D, Caviola L, Savulescu J. Worth living or worth dying? The views of the general public about allowing disabled children to die. *J Med Ethics* 2020;46:7–15.
125. Tibballs J. Legal basis for ethical withholding and withdrawing life-sustaining medical treatment from infants and children. *J Paediatr Child Health* 2007;43:230–6.
126. Welbourn C, Efstathiou N. How does the length of cardiopulmonary resuscitation affect brain damage in patients surviving cardiac arrest? A systematic review. *Scand J Trauma Resusc Emerg Med* 2018;26:77.
127. Morrison LJ. Prehospital termination of resuscitation rule. *Curr Opin Crit Care* 2019;25:199–203.
128. Grunau B, Reynolds JC, Scheuermeyer FX, et al. Comparing the prognosis of those with initial shockable and non-shockable rhythms with increasing durations of CPR: informing minimum durations of resuscitation. *Resuscitation* 2016;101:50–6.
129. Matsuyama T, Kitamura T, Kiyohara K, et al. Impact of cardiopulmonary resuscitation duration on neurologically favourable outcome after out-of-hospital cardiac arrest: a population-based study in Japan. *Resuscitation* 2017;113:1–7.
130. Goto Y, Funada A, Goto Y. Relationship between the duration of cardiopulmonary resuscitation and favorable neurological outcomes after out-of-hospital cardiac arrest: a prospective, nationwide, population-based cohort study. *J Am Heart Assoc* 2016;5:e002819.
131. Chien CY, Su YC, Lin CC, Kuo CW, Lin SC, Weng YM. Is 15 minutes an appropriate resuscitation duration before termination of a traumatic cardiac arrest? A case–control study. *Am J Emerg Med* 2016;34:505–9.
132. Murakami N, Kokubu N, Nagano N, et al. Prognostic impact of no-flow time on 30-day neurological outcomes in patients with out-of-hospital cardiac arrest who received extracorporeal cardiopulmonary resuscitation. *Circ J* 2020;84:1097–104.
133. Drennan IR, Case E, Verbeek PR, et al. A comparison of the universal TOR Guideline to the absence of prehospital ROSC and duration of resuscitation in predicting futility from out-of-hospital cardiac arrest. *Resuscitation* 2017;111:96–102.
134. Nagata T, Abe T, Hasegawa M, Hagihara A. Factors associated with the outcome of out-of-hospital cardiopulmonary arrest among people over 80 years old in Japan. *Resuscitation* 2017;113:63–9.
135. Petek BJ, Bennett DN, Ngo C, et al. Reexamination of the UN10 rule to discontinue resuscitation during in-hospital cardiac arrest. *JAMA Netw Open* 2019;2:e194941.
136. Cheung BH, Mercer MP. Prehospital disposition and patient outcomes in cardiac arrest AFTER resuscitation termination protocol change in an urban setting. *Prehosp Disaster Med* 2020;35:285–92.
137. Hreinsson JP, Thorvaldsson AP, Magnusson V, Fridriksson BT, Libungan BG, Karason S. Identifying out-of-hospital cardiac arrest patients with no chance of survival: an independent validation of prediction rules. *Resuscitation* 2020;146:19–25.
138. Winther-Jensen M, Kjaergaard J, Hassager C, et al. Resuscitation and post resuscitation care of the very old after out-of-hospital cardiac arrest is worthwhile. *Int J Cardiol* 2015;201:616–23.
139. Funada A, Goto Y, Maeda T, Teramoto R, Hayashi K, Yamagishi M. Improved survival with favorable neurological outcome in elderly individuals with out-of-hospital cardiac arrest in Japan – a nationwide observational cohort study. *Circ J* 2016;80:1153–62.
140. Fernando SM, Qureshi D, Tanuseputro P, et al. Long-term survival and costs following extracorporeal membrane oxygenation in critically ill children—a population-based cohort study. *Crit Care* 2020;24:131.
141. Smith RJ, Reid DA, Santamaria JD. Frailty is associated with reduced prospect of discharge home after in-hospital cardiac arrest. *Intern Med J* 2019;49:978–85.
142. Nas J, Kleinnibbelink G, Hannink G, et al. Diagnostic performance of the basic and advanced life support termination of resuscitation rules: a systematic review and diagnostic meta-analysis. *Resuscitation* 2020;148:3–13.
143. Ebell MH, Vellinga A, Masterson S, Yun P. Meta-analysis of the accuracy of termination of resuscitation rules for out-of-hospital cardiac arrest. *Emerg Med* 2019;36:479–84.
144. Verhaert DV, Bonnes JL, Nas J, et al. Termination of resuscitation in the prehospital setting: a comparison of decisions in clinical practice vs. recommendations of a termination rule. *Resuscitation* 2016;100:60–5.
145. American College of Surgeons Committee on T. American College of Emergency Physicians Pediatric Emergency Medicine C, National Association of Ems P, American Academy of Pediatrics Committee on Pediatric Emergency M Fallat ME. Withholding or termination of resuscitation in pediatric out-of-hospital traumatic cardiopulmonary arrest. *Pediatrics* 2014;133:e1104–16.
146. Roter VM, Trepels-Kottek S, Heimann K, Brokmann JC, Orlikowsky T, Schoberer M. Adult “termination-of-resuscitation” (TOR)-criteria may not be suitable for children – a retrospective analysis. *Scand J Trauma Resusc Emerg Med* 2016;24:144.
147. Shibahashi K, Sugiyama K, Hamabe Y. Pediatric out-of-hospital traumatic cardiopulmonary arrest after traffic accidents and termination of resuscitation. *Ann Emerg Med* 2020;75:57–65.
148. Clark JD, Dudzinski DM. The culture of dysthanasia: attempting CPR in terminally ill children. *Pediatrics* 2013;131:572–80.
149. Ganz FD, Sharfi R, Kaufman N, Einav S. Perceptions of slow codes by nurses working on internal medicine wards. *Nurs Ethics* 2019;26:1734–43.
150. Einav S, Avidan A, Brezis M, Rubinow A. Attitudes of medical practitioners towards “Do Not Resuscitate” orders. *Med Law* 2006;25:219–28.
151. Vandeplassche S, van de Voorde P. Retrospective population-based study of emergency medical services-attended out-of-hospital cardiac arrests in children in Belgium. *Eur J Emerg Med* 2018;25:400–3.
152. Kelly J. Literature review: decision-making regarding slow resuscitation. *J Clin Nurs* 2007;16:1989–96.
153. Kelly J. Nurses’ and doctors’ perspectives on slow codes. *Nurs Ethics* 2008;15:110–20.
154. Soar J, Berg KM, Andersen LW, et al. Adult advanced life support: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2020;156:A80–A119.
155. Maconochie IK, Aickin R, Hazinski MF, et al. Pediatric life support: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2020;156:A120–55.
156. D’Arrigo S, Cacciola S, Dennis M, et al. Predictors of favourable outcome after in-hospital cardiac arrest treated with extracorporeal cardiopulmonary resuscitation: a systematic review and meta-analysis. *Resuscitation* 2017;121:62–70.
157. Ortega-Deballon I, Hornby L, Shemie SD, Bhanji F, Guadagno E. Extracorporeal resuscitation for refractory out-of-hospital cardiac arrest in adults: a systematic review of international practices and outcomes. *Resuscitation* 2016;101:12–20.
158. Harvey MJ, Gaies MG, Prosser LA. U.S. and international in-hospital costs of extracorporeal membrane oxygenation: a systematic review. *Appl Health Econ Health Policy* 2015;13:341–57.
159. Holmberg MJ, Geri G, Wiberg S, et al. Extracorporeal cardiopulmonary resuscitation for cardiac arrest: a systematic review. *Resuscitation* 2018;131:91–100.
160. Bougouin W, Dumas F, Lamhaut L, et al. Extracorporeal cardiopulmonary resuscitation in out-of-hospital cardiac arrest: a registry study. *Eur Heart J* 2020;41:1961–71.

161. Quality) O.H. Extracorporeal membrane oxygenation for cardiac indications in adults: a health technology assessment. *Ont Health Technol Assess Ser* 2020;20:1–121.
162. Grunau B, Hornby L, Singal RK, et al. Extracorporeal cardiopulmonary resuscitation for refractory out-of-hospital cardiac arrest: the state of the evidence and framework for application. *Can J Cardiol* 2018;34:146–55.
163. Dalle Ave AL, Shaw DM, Gardiner D. Extracorporeal membrane oxygenation (ECMO) assisted cardiopulmonary resuscitation or uncontrolled donation after the circulatory determination of death following out-of-hospital refractory cardiac arrest – an ethical analysis of an unresolved clinical dilemma. *Resuscitation* 2016;108:87–94.
164. Pujara D, Sandoval E, Simpson L, Mallidi HR, Singh SK. The state of the art in extracorporeal membrane oxygenation. *Semin Thorac Cardiovasc Surg* 2015;27:17–23.
165. Singer B, Reynolds JC, Lockey DJ, O'Brien B. Pre-hospital extracorporeal cardiopulmonary resuscitation. *Scand J Trauma Resusc Emerg Med* 2018;26:21.
166. Gravesteijn BY, Schlupe M, Voormolen DC, et al. Cost-effectiveness of extracorporeal cardiopulmonary resuscitation after in-hospital cardiac arrest: a Markov decision model. *Resuscitation* 2019;143:150–7.
167. Meltzer EC, Ivascu NS, Stark M, et al. A survey of physicians' attitudes toward decision-making authority for initiating and withdrawing VA-ECMO: results and ethical implications for shared decision making. *J Clin Ethics* 2016;27:281–9.
168. St-Onge M, Fan E, Megarbane B, Hancock-Howard R, Coyte PC. Venoaerial extracorporeal membrane oxygenation for patients in shock or cardiac arrest secondary to cardiotoxicant poisoning: a cost-effectiveness analysis. *J Crit Care* 2015;30:e7–e14.
169. Dennis M, Zmudzki F, Burns B, et al. Cost effectiveness and quality of life analysis of extracorporeal cardiopulmonary resuscitation (ECPR) for refractory cardiac arrest. *Resuscitation* 2019;139:49–56.
170. Tonna JE, Selzman CH, Girotra S, et al. Patient and institutional characteristics influence the decision to use extracorporeal cardiopulmonary resuscitation for in-hospital cardiac arrest. *J Am Heart Assoc* 2020;9:e015522.
171. Poppe M, Schrieffl C, Steinacher A, et al. Extracorporeal cardiopulmonary resuscitation at the emergency department: a retrospective patient selection evaluation. *Eur J Anaesthesiol* 2020;37:280–5.
172. Yoshida T, Fujitani S, Wakatake H, et al. Exploratory observational study of extracorporeal cardiopulmonary resuscitation for nonshockable out-of-hospital cardiac arrest occurring after an emergency medical services arrival: SOS-KANTO 2012 study report. *J Emerg Med* 2020;58:375–84.
173. Lunz D, Calabro L, Belliati M, et al. Extracorporeal membrane oxygenation for refractory cardiac arrest: a retrospective multicenter study. *Intensive Care Med* 2020;46:973–82.
174. Chonde M, Escajeda J, Elmer J, et al. Challenges in the development and implementation of a healthcare system based extracorporeal cardiopulmonary resuscitation (ECPR) program for the treatment of out of hospital cardiac arrest. *Resuscitation* 2020;148:259–65.
175. McDonald L, Mastoras G, Hickey M, McDonald B, Kwok ESH. Evaluating the potential impact of an emergency department extracorporeal resuscitation (ECPR) program: a health records review. *CJEM* 2020;22:375–8.
176. Cheetham OV, Thomas MJ, Hadfield J, O'Higgins F, Mitchell C, Rooney KD. Rates of organ donation in a UK tertiary cardiac arrest centre following out-of-hospital cardiac arrest. *Resuscitation* 2016;101:41–3.
177. Nolan JP, Ferrando P, Soar J, et al. Increasing survival after admission to UK critical care units following cardiopulmonary resuscitation. *Crit Care* 2016;20:219.
178. Sandroni C, D'Arrigo S, Callaway CW, et al. The rate of brain death and organ donation in patients resuscitated from cardiac arrest: a systematic review and meta-analysis. *Intensive Care Med* 2016;42:1661–71.
179. Ortega-Deballon I, Hornby L, Shemie SD. Protocols for uncontrolled donation after circulatory death: a systematic review of international guidelines, practices and transplant outcomes. *Crit Care* 2015;19:268.
180. DuBois JM, Anderson EE. Attitudes toward death criteria and organ donation among healthcare personnel and the general public. *Prog Transplant* 2006;16:65–73.
181. Shah SK, Kasper K, Miller FG. A narrative review of the empirical evidence on public attitudes on brain death and vital organ transplantation: the need for better data to inform policy. *J Med Ethics* 2015;41:291–6.
182. Escudero D, Otero J, Menendez de Leon B, Perez-Basterrechea M. Organ donation and elective ventilation: a necessary strategy. *Biomed Res Int* 2017;2017:7518375.
183. Dalle Ave AL, Gardiner D, Shaw DM. Cardio-pulmonary resuscitation of brain-dead organ donors: a literature review and suggestions for practice. *Transpl Int* 2016;29:12–9.
184. Weiss MJ, English SW, D'Aragon F, et al. Survey of Canadian intensivists on physician non-referral and family override of deceased organ donation. *Can J Anaesth* 2020;67:313–23.
185. Reed MJ, Lua SB. Uncontrolled organ donation after circulatory death: potential donors in the emergency department. *Emerg Med* 2014;31:741–4.
186. Bruce CM, Reed MJ, MacDougall M. Are the public ready for organ donation after out of hospital cardiac arrest? *Emerg Med* 2013;30:226–31.
187. Dhanani S, Hornby L, Ward R, et al. Vital signs after cardiac arrest following withdrawal of life-sustaining therapy: a multicenter prospective observational study. *Crit Care Med* 2014;42:2358–69.
188. Sque M, Walker W, Long-Sutehall T, Morgan M, Randhawa G, Rodney A. Bereaved donor families' experiences of organ and tissue donation, and perceived influences on their decision making. *J Crit Care* 2018;45:82–9.
189. Dalle Ave AL, Bernat JL. Uncontrolled donation after circulatory determination of death: a systematic ethical analysis. *J Intensive Care Med* 2018;33:624–34.
190. Joffe AR, Carcillo J, Anton N, et al. Donation after cardiocirculatory death: a call for a moratorium pending full public disclosure and fully informed consent. *Philos Ethics Humanit Med* 2011;6:17.
191. Rodriguez-Arias D, Tortosa JC, Burant CJ, Aubert P, Aulisio MP, Youngner SJ. One or two types of death? Attitudes of health professionals towards brain death and donation after circulatory death in three countries. *Med Health Care Philos* 2013;16:457–67.
192. Verheijde JL, Rady MY, McGregor J. Presumed consent for organ preservation in uncontrolled donation after cardiac death in the United States: a public policy with serious consequences. *Philos Ethics Humanit Med* 2009;4:15.
193. Rodriguez-Arias D, Smith MJ, Lazar NM. Donation after circulatory death: burying the dead donor rule. *Am J Bioeth* 2011;11:36–43.
194. Maconochie IK, Bingham R, Eich C, et al. European resuscitation council guidelines for resuscitation 2015: Section 6. Paediatric life support. *Resuscitation* 2015;95:223–48.
195. Bossaert LL, Perkins GD, Askitopoulou H, et al. European resuscitation council guidelines for resuscitation 2015: section 11. The ethics of resuscitation and end-of-life decisions. *Resuscitation* 2015;95:302–11.
196. Oczkowski SJ, Mazzetti I, Cupido C, Fox-Robichaud AE. Canadian critical care S. family presence during resuscitation: a Canadian critical care society position paper. *Can Respir J* 2015;22:201–5.
197. Barreto MDS, Peruzzo HE, Garcia-Vivar C, Marcon SS. Family presence during cardiopulmonary resuscitation and invasive procedures: a meta-synthesis. *Rev Esc Enferm USP* 2019;53:e03435.

198. Oczkowski SJ, Mazzetti I, Cupido C, Fox-Robichaud AE. The offering of family presence during resuscitation: a systematic review and meta-analysis. *J Intensive Care* 2015;3:41.
199. Sak-Dankosky N, Andruszkiewicz P, Sherwood PR, Kvist T. Integrative review: nurses' and physicians' experiences and attitudes towards inpatient-witnessed resuscitation of an adult patient. *J Adv Nurs* 2014;70:957–74.
200. Toronto CE, LaRocco SA. Family perception of and experience with family presence during cardiopulmonary resuscitation: an integrative review. *J Clin Nurs* 2019;28:32–46.
201. Vincent C, Lederman Z. Family presence during resuscitation: extending ethical norms from paediatrics to adults. *J Med Ethics* 2017;43:676–8.
202. Mark K. Family presence during paediatric resuscitation and invasive procedures: the parental experience: an integrative review. *Scand J Caring Sci* 2020.
203. Timmis V. Should family members be present at resuscitation? *Arch Dis Child* 2020;105:506–8.
204. Kenny G, Bray I, Pontin D, Jefferies R, Albarran J. A randomised controlled trial of student nurse performance of cardiopulmonary resuscitation in a simulated family-witnessed resuscitation scenario. *Nurse Educ Pract* 2017;24:21–6.
205. Krochmal RL, Blenko JW, Afshar M, et al. Family presence at first cardiopulmonary resuscitation and subsequent limitations on care in the medical intensive care unit. *Am J Crit Care* 2017;26:221–8.
206. Goldberger ZD, Nallamothu BK, Nichol G, et al. Policies allowing family presence during resuscitation and patterns of care during in-hospital cardiac arrest. *Circ Cardiovasc Qual Outcomes* 2015;8:226–34.
207. Tripon C, Defossez G, Ragot S, et al. Parental presence during cardiopulmonary resuscitation of children: the experience, opinions and moral positions of emergency teams in France. *Arch Dis Child* 2014;99:310–5.
208. Nowland R, Steeg S, Quinlivan LM, et al. Management of patients with an advance decision and suicidal behaviour: a systematic review. *BMJ Open* 2019;9:e023978.
209. Weerasuriya D, Sheikh S, Morgan BW. Do-not-resuscitate orders in fatal toxic exposures: a poison center's review. *West J Emerg Med* 2012;13:294–7.
210. Ryan CJ, Callaghan S. Legal and ethical aspects of refusing medical treatment after a suicide attempt: the Wooltorton case in the Australian context. *Med J Aust* 2010;193:239–42.
211. Sontheimer D. Suicide by advance directive? *J Med Ethics* 2008;34:e4.
212. Philpot SJ. Should an advance care directive refusing life-sustaining treatment be respected after an attempted suicide? Development of an algorithm to aid health care workers. *J Law Med* 2019;26:557–70.
213. Venkat A, Drori J. When to say when: responding to a suicide attempt in the acute care setting. *Narrat Inq Bioeth* 2014;4:263–70.
214. Terman SA. Is the principle of proportionality sufficient to guide physicians' decisions regarding withholding/withdrawing life-sustaining treatment after suicide attempts? *Am J Bioeth* 2013;13:22–4.
215. Brown SM, Elliott CG, Paine R. Withdrawal of nonfutile life support after attempted suicide. *Am J Bioeth* 2013;13:3–12.
216. Song J, Guo W, Lu X, Kang X, Song Y, Gong D. The effect of bystander cardiopulmonary resuscitation on the survival of out-of-hospital cardiac arrests: a systematic review and meta-analysis. *Scand J Trauma Resusc Emerg Med* 2018;26:86.
217. Grasner JT, Wnent J, Herlitz J, et al. Survival after out-of-hospital cardiac arrest in Europe – results of the EuReCa TWO study. *Resuscitation* 2020;148:218–26.
218. Hansen SM, Hansen CM, Folke F, et al. Bystander defibrillation for out-of-hospital cardiac arrest in public vs residential locations. *JAMA Cardiol* 2017;2:507–14.
219. Wissenberg M, Lippert FK, Folke F, et al. Association of national initiatives to improve cardiac arrest management with rates of bystander intervention and patient survival after out-of-hospital cardiac arrest. *JAMA* 2013;310:1377–84.
220. Scquizzato T, Pallanch O, Belletti A, et al. Enhancing citizens response to out-of-hospital cardiac arrest: a systematic review of mobile-phone systems to alert citizens as first responders. *Resuscitation* 2020;152:16–25.
221. Tay PJM, Pek PP, Fan Q, et al. Effectiveness of a community based out-of-hospital cardiac arrest (OHCA) interventional bundle: results of a pilot study. *Resuscitation* 2020;146:220–8.
222. Perkins GD, Olasveengen TM, Maconochie I, et al. European resuscitation council guidelines for resuscitation: 2017 update. *Resuscitation* 2018;123:43–50.
223. Ro YS, Shin SD, Song KJ, et al. Public awareness and self-efficacy of cardiopulmonary resuscitation in communities and outcomes of out-of-hospital cardiac arrest: a multi-level analysis. *Resuscitation* 2016;102:17–24.
224. Ko SY, Ro YS, Shin SD, Song KJ, Hong KJ, Kong SY. Effect of a first responder on survival outcomes after out-of-hospital cardiac arrest occurs during a period of exercise in a public place. *PLOS ONE* 2018;13:e0193361.
225. Smida T, Willson C, Salerno J, Weiss L, Salcido DD. Can you get there from here? An analysis of walkability among PulsePoint CPR alert dispatches. *Resuscitation* 2020;148:135–9.
226. Anto-Ocrah M, Maxwell N, Cushman J, et al. Public knowledge and attitudes towards bystander cardiopulmonary resuscitation (CPR) in Ghana, West Africa. *Int J Emerg Med* 2020;13:29.
227. Matsuyama T, Scapigliati A, Pellis T, Greif R, Iwami T. Willingness to perform bystander cardiopulmonary resuscitation: a scoping review. *Resusc Plus* 2020;4:100043.
228. Brinkrolf P, Bohn A, Lukas RP, et al. Senior citizens as rescuers: is reduced knowledge the reason for omitted lay-resuscitation-attempts? Results from a representative survey with 2004 interviews. *PLOS ONE* 2017;12:e0178938.
229. Dobbie F, MacKintosh AM, Clegg G, Stirzaker R, Bauld L. Attitudes towards bystander cardiopulmonary resuscitation: results from a cross-sectional general population survey. *PLOS ONE* 2018;13:e0193391.
230. Sasson C, Haukoos JS, Bond C, et al. Barriers and facilitators to learning and performing cardiopulmonary resuscitation in neighborhoods with low bystander cardiopulmonary resuscitation prevalence and high rates of cardiac arrest in Columbus, OH. *Circ Cardiovasc Qual Outcomes* 2013;6:550–8.
231. Cheskes L, Morrison LJ, Beaton D, Parsons J, Dainty KN. Are Canadians more willing to provide chest-compression-only cardiopulmonary resuscitation (CPR)? – A nation-wide public survey. *CJEM* 2016;18:253–63.
232. Malta Hansen C, Rosenkranz SM, Folke F, et al. Lay bystanders' perspectives on what facilitates cardiopulmonary resuscitation and use of automated external defibrillators in real cardiac arrests. *J Am Heart Assoc* 2017;6.
233. Tanigawa K, Iwami T, Nishiyama C, Nonogi H, Kawamura T. Are trained individuals more likely to perform bystander CPR? An observational study. *Resuscitation* 2011;82:523–8.
234. Liaw SY, Chew KS, Zulkarnain A, et al. Improving perception and confidence towards bystander cardiopulmonary resuscitation and public access automated external defibrillator program: how does training program help? *Int J Emerg Med* 2020;13:13.
235. Fukushima H, Asai H, Seki T, Takano K, Bolstad F. The effect of 10-min dispatch-assisted cardiopulmonary resuscitation training: a randomized simulation pilot study. *Int J Emerg Med* 2020;13:31.
236. Olasveengen TM, Mancini ME, Perkins GD, et al. Adult basic life support: international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2020;156:A35–79.
237. Ringh M, Rosenqvist M, Hollenberg J, et al. Mobile-phone dispatch of laypersons for CPR in out-of-hospital cardiac arrest. *N Engl J Med* 2015;372:2316–25.
238. Caputo ML, Muschietti S, Burkart R, et al. Lay persons alerted by mobile application system initiate earlier cardio-pulmonary resuscitation: a comparison with SMS-based system notification. *Resuscitation* 2017;114:73–8.

239. Berglund E, Claesson A, Nordberg P, et al. A smartphone application for dispatch of lay responders to out-of-hospital cardiac arrests. *Resuscitation* 2018;126:160–5.
240. Hansen CM, Kragholm K, Granger CB, et al. The role of bystanders, first responders, and emergency medical service providers in timely defibrillation and related outcomes after out-of-hospital cardiac arrest: results from a statewide registry. *Resuscitation* 2015;96:303–9.
241. Sarkisian L, Mickley H, Schakow H, et al. Global positioning system alerted volunteer first responders arrive before emergency medical services in more than four out of five emergency calls. *Resuscitation* 2020;152:170–6.
242. Raun L, Pederson J, Campos L, Ensor K, Persse D. Effectiveness of the dual dispatch to cardiac arrest policy in Houston, Texas. *J Public Health Manag Pract* 2019;25:E13–21.
243. Pijls RWM, Nelemans PJ, Rahel BM, Gorgels APM. Characteristics of a novel citizen rescue system for out-of-hospital cardiac arrest in the Dutch province of Limburg: relation to incidence and survival. *Neth Heart J* 2019;27:100–7.
244. Zijlstra JA, Stieglis R, Riedijk F, Smeekes M, van der Worp WE, Koster RW. Local lay rescuers with AEDs, alerted by text messages, contribute to early defibrillation in a Dutch out-of-hospital cardiac arrest dispatch system. *Resuscitation* 2014;85:1444–9.
245. Stroop R, Kerner T, Strickmann B, Hensel M. Mobile phone-based alerting of CPR-trained volunteers simultaneously with the ambulance can reduce the resuscitation-free interval and improve outcome after out-of-hospital cardiac arrest: a German, population-based cohort study. *Resuscitation* 2020;147:57–64.
246. Boland LL, Formanek MB, Harkins KK, et al. Minnesota heart safe communities: are community-based initiatives increasing pre-ambulance CPR and AED use? *Resuscitation* 2017;119:33–6.
247. Park YM, Shin SD, Lee YJ, Song KJ, Ro YS, Ahn KO. Cardiopulmonary resuscitation by trained responders versus lay persons and outcomes of out-of-hospital cardiac arrest: a community observational study. *Resuscitation* 2017;118:55–62.
248. Nord A, Svensson L, Karlsson T, Claesson A, Herlitz J, Nilsson L. Increased survival from out-of-hospital cardiac arrest when off duty medically educated personnel perform CPR compared with laymen. *Resuscitation* 2017;120:88–94.
249. Dainty KN, Vaid H, Brooks SC. North American public opinion survey on the acceptability of crowdsourcing basic life support for out-of-hospital cardiac arrest with the pulsepoint mobile phone app. *JMIR Mhealth Uhealth* 2017;5:e63.
250. Brooks SC, Simmons G, Worthington H, Bobrow BJ, Morrison LJ. The PulsePoint Respond mobile device application to crowdsource basic life support for patients with out-of-hospital cardiac arrest: challenges for optimal implementation. *Resuscitation* 2016;98:20–6.
251. Smith CM, Wilson MH, Ghorbangholi A, et al. The use of trained volunteers in the response to out-of-hospital cardiac arrest – the GoodSAM experience. *Resuscitation* 2017;121:123–6.
252. Jaffe E, Dadon Z, Alpert EA. Wisdom of the crowd in saving lives: the life guardians app. *Prehosp Disaster Med* 2018;33:550–2.
253. Barry T, Doheny MC, Masterson S, et al. Community first responders for out-of-hospital cardiac arrest in adults and children. *Cochrane Database Syst Rev* 2019;7:CD012764.
254. Krammel M, Lobmeyr E, Sulzgruber P, et al. The impact of a high-quality basic life support police-based first responder system on outcome after out-of-hospital cardiac arrest. *PLOS ONE* 2020;15:e0233966.
255. Svensson A, Elmqvist C, Fridlund B, Rask M, Andersson R, Stening K. Using firefighters as medical first responders to shorten response time in rural areas in Sweden. *Aust J Rural Health* 2020;28:6–14.
256. Harkins B, Smith K, Cameron P, et al. The impact of bystander relation and medical training on out-of-hospital cardiac arrest outcomes. *Resuscitation* 2020;150:72–9.
257. Sayre MR, Barnard LM, Counts CR, et al. Prevalence of COVID-19 in out-of-hospital cardiac arrest: implications for bystander cardiopulmonary resuscitation. *Circulation* 2020;142:507–9.
258. Mausz J, Snobelen P, Tavares W. “Please. Don’t. Die.”: a grounded theory study of bystander cardiopulmonary resuscitation. *Circ Cardiovasc Qual Outcomes* 2018;11:e004035.
259. Rosoff PM, Schneiderman LJ. Response to open peer commentaries on “Irrational Exuberance: Cardiopulmonary Resuscitation as Fetish”. *Am J Bioeth* 2017;17:W1–3.
260. Field RA, Soar J, Nolan JP, Perkins GD. Epidemiology and outcome of cardiac arrests reported in the lay-press: an observational study. *J R Soc Med* 2011;104:525–31.
261. Mathiesen WT, Bjorshol CA, Braut GS, Soreide E. Reactions and coping strategies in lay rescuers who have provided CPR to out-of-hospital cardiac arrest victims: a qualitative study. *BMJ Open* 2016;6:e010671.
262. Zijlstra JA, Beesems SG, De Haan RJ, Koster RW. Psychological impact on dispatched local lay rescuers performing bystander cardiopulmonary resuscitation. *Resuscitation* 2015;92:115–21.
263. Nord-Ljungquist H, Engstrom A, Fridlund B, Elmqvist C. Lone and lonely in a double ambivalence situation as experienced by callers while waiting for the ambulance in a rural environment. *Scand J Caring Sci* 2020;34:566–74.
264. Schnaubelt S, Monsieurs KG, Semeraro F, et al. Clinical outcomes from out-of-hospital cardiac arrest in low-resource settings – a scoping review. *Resuscitation* 2020;156:137–45.
265. Schwartz CE, Wheeler HB, Hammes B, et al. Early intervention in planning end-of-life care with ambulatory geriatric patients: results of a pilot trial. *Arch Intern Med* 2002;162:1611–8.
266. Doorenbos AZ, Levy WC, Curtis JR, Dougherty CM. An intervention to enhance goals-of-care communication between heart failure patients and heart failure providers. *J Pain Symptom Manag* 2016;52:353–60.
267. Houben CHM, Spruit MA, Luyten H, et al. Cluster-randomised trial of a nurse-led advance care planning session in patients with COPD and their loved ones. *Thorax* 2019;74:328–36.
268. Engelhardt JB, Rizzo VM, Della Penna RD, et al. Effectiveness of care coordination and health counseling in advancing illness. *Am J Manag Care* 2009;15:817–25.
269. Briggs LA, Kirchhoff KT, Hammes BJ, Song MK, Colvin ER. Patient-centered advance care planning in special patient populations: a pilot study. *J Prof Nurs* 2004;20:47–58.
270. Fischer SM, Cervantes L, Fink RM, Kutner JS. Apoyo con Carino: a pilot randomized controlled trial of a patient navigator intervention to improve palliative care outcomes for Latinos with serious illness. *J Pain Symptom Manag* 2015;49:657–65.
271. Perry E, Swartz J, Brown S, Smith D, Kelly G, Swartz R. Peer mentoring: a culturally sensitive approach to end-of-life planning for long-term dialysis patients. *Am J Kidney Dis* 2005;46:111–9.
272. McCannon JB, O’Donnell WJ, Thompson BT, et al. Augmenting communication and decision making in the intensive care unit with a cardiopulmonary resuscitation video decision support tool: a temporal intervention study. *J Palliat Med* 2012;15:1382–7.
273. Volandes AE, Ferguson LA, Davis AD, et al. Assessing end-of-life preferences for advanced dementia in rural patients using an educational video: a randomized controlled trial. *J Palliat Med* 2011;14:169–77.
274. Epstein AS, Volandes AE, Chen LY, et al. A randomized controlled trial of a cardiopulmonary resuscitation video in advance care planning for progressive pancreatic and hepatobiliary cancer patients. *J Palliat Med* 2013;16:623–31.
275. Sharma RK, Szmuiłowicz E, Ogunseitan A, et al. Evaluation of a mastery learning intervention on hospitalists’ code status discussion skills. *J Pain Symptom Manag* 2017;53:1066–70.
276. Deep KS, Griffith CH, Wilson JF. Communication and decision making about life-sustaining treatment: examining the experiences of resident physicians and seriously-ill hospitalized patients. *J Gen Intern Med* 2008;23:1877–82.
277. Mian P, Warchal S, Whitney S, et al. Impact of a multifaceted intervention on nurses’ and physicians’ attitudes and behaviors toward family presence during resuscitation. [see comment]. *Crit Care Nurse* 2007;27:52–61.

278. Pye S, Kane J, Jones A. Parental presence during pediatric resuscitation: the use of simulation training for cardiac intensive care nurses. *J Spec Pediatr Nurs* 2010;15:172–5.
279. Feagan LM, Fisher NJ. The impact of education on provider attitudes toward family-witnessed resuscitation. *J Emerg Nurs* 2011;37:231–9.
280. Morrison RS, Chichin E, Carter J, Burack O, Lantz M, Meier DE. The effect of a social work intervention to enhance advance care planning documentation in the nursing home. *J Am Geriatr Soc* 2005;53:290–4.
281. Anderson N, Slark J, Gott M. How are ambulance personnel prepared and supported to withhold or terminate resuscitation and manage patient death in the field? A scoping review. *Australas J Paramed* 2019;16.
282. Whitehead L, Perkins GD, Clarey A, Haywood KL. A systematic review of the outcomes reported in cardiac arrest clinical trials: the need for a core outcome set. *Resuscitation* 2015;88:150–7.
283. Sawyer KN, Camp-Rogers TR, Kotini-Shah P, et al. Sudden cardiac arrest survivorship: a scientific statement from the American Heart Association. *Circulation* 2020;141:e654–85.
284. Sawyer KN, Camp-Rogers TR, Kotini-Shah P, et al. Sudden Cardiac Arrest Survivorship: A Scientific Statement From the American Heart Association. *Circulation*;0:CIR.0000000000000747.
285. Nolan JP, Soar J, Smith GB, et al. Incidence and outcome of in-hospital cardiac arrest in the United Kingdom National Cardiac Arrest Audit. *Resuscitation* 2014;85:987–92.
286. Hawkes C, Booth S, Ji C, et al. Epidemiology and outcomes from out-of-hospital cardiac arrests in England. *Resuscitation* 2017;110:133–40.
287. Moolaert VR, Goossens M, Heijnders IL, Verbunt JA, Heugten CM. Early neurologically focused follow-up after cardiac arrest is cost-effective: a trial-based economic evaluation. *Resuscitation* 2016;106:30–6.
288. Gates S, Lall R, Quinn T, et al. Prehospital randomised assessment of a mechanical compression device in out-of-hospital cardiac arrest (PARAMEDIC): a pragmatic, cluster randomised trial and economic evaluation. *Health Technol Assess* 2017;21:1–176.
289. Achana F, Petrou S, Madan J, et al. Cost-effectiveness of adrenaline for out-of-hospital cardiac arrest. *Crit Care* 2020;24:579.
290. Orioles A, Morrison WE, Rossano JW, et al. An under-recognized benefit of cardiopulmonary resuscitation: organ transplantation. *Crit Care Med* 2013;41:2794–9.
291. Chan PS, Krein SL, Tang F, et al. Resuscitation practices associated with survival after in-hospital cardiac arrest: a nationwide survey. *JAMA Cardiol* 2016;1:189–97.
292. Couper K, Kimani PK, Gale CP, et al. Patient, health service factors and variation in mortality following resuscitated out-of-hospital cardiac arrest in acute coronary syndrome: analysis of the Myocardial Ischaemia National Audit Project. *Resuscitation* 2018;124:49–57.
293. Lilford R, Mohammed MA, Spiegelhalter D, Thomson R. Use and misuse of process and outcome data in managing performance of acute medical care: avoiding institutional stigma. *Lancet* 2004;363:1147–54.
294. Perkins GD, Brace-McDonnell SJ, Group OP, The UK. Out of Hospital Cardiac Arrest Outcome (OHCAO) project. *BMJ Open* 2015;5:e008736.
295. Blewer AL, McGovern SK, Schmicker RH, et al. Gender disparities among adult recipients of bystander cardiopulmonary resuscitation in the public. *Circ Cardiovasc Qual Outcomes* 2018;11:e004710.
296. Brown TP, Booth S, Hawkes CA, et al. Characteristics of neighbourhoods with high incidence of out-of-hospital cardiac arrest and low bystander cardiopulmonary resuscitation rates in England. *Eur Heart J Qual Care Clin Outcomes* 2019;5:51–62.
297. Edelson DP, Yuen TC, Mancini ME, et al. Hospital cardiac arrest resuscitation practice in the United States: a nationally representative survey. *J Hosp Med* 2014;9:353–7.
298. Tirkkonen J, Nurmi J, Olkkola KT, Tenhunen J, Hoppu S. Cardiac arrest teams and medical emergency teams in Finland: a nationwide cross-sectional postal survey. *Acta Anaesthesiol Scand* 2014;58:420–7.
299. Carberry J, Couper K, Yeung J. The implementation of cardiac arrest treatment recommendations in English acute NHS trusts: a national survey. *Postgrad Med J* 2017;93:653–9.
300. Soar J, Maconochie I, Wyckoff MH, et al. 2019 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2019;145:95–150.
301. Yeung J, Matsuyama T, Bray J, Reynolds J, Skrifvars MB. Does care at a cardiac arrest centre improve outcome after out-of-hospital cardiac arrest? – A systematic review. *Resuscitation* 2019;137:102–15.
302. Perkins GD, Jacobs IG, Nadkarni VM, et al. Cardiac Arrest and Cardiopulmonary Resuscitation Outcome Reports: Update of the Utstein Resuscitation Registry Templates for Out-of-Hospital Cardiac Arrest: A Statement for Healthcare Professionals From a Task Force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian and New Zealand Council on Resuscitation, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Southern Africa, Resuscitation Council of Asia); and the American Heart Association Emergency Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. *Resuscitation* 2015;96:328–40.
303. Nolan JP, Berg RA, Andersen LW, et al. Cardiac Arrest and Cardiopulmonary Resuscitation Outcome Reports: Update of the Utstein Resuscitation Registry Template for In-Hospital Cardiac Arrest: A Consensus Report From a Task Force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian and New Zealand Council on Resuscitation, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Southern Africa, Resuscitation Council of Asia). *Resuscitation* 2019;144:166–77.
304. Chalmers I, Glasziou P. Avoidable waste in the production and reporting of research evidence. *Lancet* 2009;374:86–9.
305. Haywood K, Whitehead L, Nadkarni VM, et al. COSCA (Core Outcome Set for Cardiac Arrest) in adults: an advisory statement from the international liaison committee on resuscitation. *Resuscitation* 2018;127:147–63.
306. Clarke M. Standardising outcomes for clinical trials and systematic reviews. *Trials* 2007;8:39.
307. Williamson PR, Altman DG, Blazeby JM, et al. Developing core outcome sets for clinical trials: issues to consider. *Trials* 2012;13:132.
308. Tiainen M, Vaahersalo J, Skrifvars MB, Hastbacka J, Gronlund J, Pettila V. Surviving out-of-hospital cardiac arrest: the neurological and functional outcome and health-related quality of life one year later. *Resuscitation* 2018;129:19–23.
309. Bohm M, Lilja G, Finnbogadottir H, et al. Detailed analysis of health-related quality of life after out-of-hospital cardiac arrest. *Resuscitation* 2019;135:197–204.
310. Calvert M, Kyte D, Mercieca-Bebber R, et al. Guidelines for inclusion of patient-reported outcomes in clinical trial protocols: the SPIRIT-PRO extension. *JAMA* 2018;319:483–94.
311. Andrew E, Mercier E, Nehme Z, Bernard S, Smith K. Long-term functional recovery and health-related quality of life of elderly out-of-hospital cardiac arrest survivors. *Resuscitation* 2018;126:118–24.
312. Caro-Codon J, Rey JR, Lopez-de-Sa E, et al. Long-term neurological outcomes in out-of-hospital cardiac arrest patients treated with targeted-temperature management. *Resuscitation* 2018;133:33–9.
313. Nehme Z, Andrew E, Bernard S, Smith K. Sex differences in the quality-of-life and functional outcome of cardiac arrest survivors. *Resuscitation* 2019;137:21–8.
314. Grasner JT, Lefering R, Koster RW, et al. EuReCa ONE-27 Nations, ONE Europe ONE Registry: a prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in Europe. *Resuscitation* 2016;105:188–95.

315. Okubo M, Schmicker RH, Wallace DJ, et al. Variation in survival after out-of-hospital cardiac arrest between emergency medical services agencies. *JAMA Cardiol* 2018;3:989–99.
316. Kiguchi T, Okubo M, Nishiyama C, et al. Out-of-hospital cardiac arrest across the World: first report from the International Liaison Committee on Resuscitation (ILCOR). *Resuscitation* 2020;152:39–49.
317. Mentzelopoulos SD, Mantzanas M, van Belle G, Nichol G. Evolution of European Union legislation on emergency research. *Resuscitation* 2015;91:84–91.
318. Perkins GD, Ji C, Deakin CD, et al. A randomized trial of epinephrine in out-of-hospital cardiac arrest. *N Engl J Med* 2018;379:711–21.
319. Mentzelopoulos SD, Bossaert L, Raffay V, et al. A survey of key opinion leaders on ethical resuscitation practices in 31 European Countries. *Resuscitation* 2016;100:11–7.
320. Perkins GD, Bossaert L, Nolan J, et al. Proposed revisions to the EU clinical trials directive—comments from the European Resuscitation Council. *Resuscitation* 2013;84:263–4.
321. Lascarrou JB, Merdji H, Le Gouge A, et al. Targeted temperature management for cardiac arrest with nonshockable rhythm. *N Engl J Med* 2019;381:2327–37.
322. Group CTM. Research in emergency situations: with or without relatives consent. *Emerg Med* 2004;21:703.
323. Booth MG. Informed consent in emergency research: a contradiction in terms. *Sci Eng Ethics* 2007;13:351–9.
324. Kompanje EJ, Maas AI, Menon DK, Kesecioglu J. Medical research in emergency research in the European Union member states: tensions between theory and practice. *Intensive Care Med* 2014;40:496–503.
325. Lecouturier J, Rodgers H, Ford GA, et al. Clinical research without consent in adults in the emergency setting: a review of patient and public views. *BMC Med Ethics* 2008;9:9.
326. Kamarainen A, Silfvast T, Saarinen S, Virta J, Virkkunen I. Conduct of emergency research in patients unable to give consent—experiences and perceptions of patients, their consent providing next of kin, and treating physicians following a prehospital resuscitation trial. *Resuscitation* 2012;83:81–5.
327. Roberts I, Prieto-Merino D, Shakur H, Chalmers I, Nicholl J. Effect of consent rituals on mortality in emergency care research. *Lancet* 2011;377:1071–2.
328. Biros MH, Sargent C, Miller K. Community attitudes towards emergency research and exception from informed consent. *Resuscitation* 2009;80:1382–7.
329. Goldstein JN, Espinola JA, Fisher J, Pallin DJ, Camargo CA. Public opinion of a stroke clinical trial using exception from informed consent. *Int J Emerg Med* 2010;3:385–9.
330. Burns KE, Magyarody NM, Duffett M, Nisenbaum R, Cook DJ. Attitudes of the general public toward alternative consent models. *Am J Crit Care* 2011;20:75–83.
331. Silbergleit R, Biros MH, Harney D, Dickert N, Baren J, Investigators N. Implementation of the exception from informed consent regulations in a large multicenter emergency clinical trials network: the RAMPART experience. *Acad Emerg Med* 2012;19:448–54.
332. Nelson MJ, Deiorio NM, Schmidt TA, Zive DM, Griffiths D, Newgard CD. Why persons choose to opt out of an exception from informed consent cardiac arrest trial. *Resuscitation* 2013;84:825–30.
333. Dickert NW, Mah VA, Biros MH, et al. Consulting communities when patients cannot consent: a multicenter study of community consultation for research in emergency settings. *Crit Care Med* 2014;42:272–80.
334. Callaway CW. Studying community consultation in exception from informed consent trials. *Crit Care Med* 2014;42:451–3.
335. Fehr AE, Pentz RD, Dickert NW. Learning from experience: a systematic review of community consultation acceptance data. *Ann Emerg Med* 2015;65:162–71000.
336. Stergiopoulos S, Michaels DL, Kunz BL, Getz KA. Measuring the impact of patient engagement and patient centrality in clinical research and development. *Ther Innov Regul Sci* 2020;54:103–16.
337. D'Souza R, Hall C, Sermer M, Siu S, Silversides C. Development of a core outcome set for studies on cardiac disease in pregnancy (COSCarP): a study protocol. *Trials* 2020;21:300.
338. Ingoe HMA, Eardley W, Rangan A, Hewitt C, McDaid C. An international multi-stakeholder delphi consensus exercise to develop a core outcomes set (COS) for surgical fixation of rib fractures. *Injury* 2020;51:224–9.
339. Webbe JWH, Duffy JMN, Afonso E, et al. Core outcomes in neonatology: development of a core outcome set for neonatal research. *Arch Dis Child Fetal Neonatal Ed* 2020;105:425–31.
340. Vat LE, Finlay T, Jan Schuitmaker-Warnaar T, et al. Evaluating the “return on patient engagement initiatives” in medicines research and development: a literature review. *Health Expect* 2020;23:5–18.
341. Koffman J, Yorganci E, Murtagh F, et al. The AMBER care bundle for hospital inpatients with uncertain recovery nearing the end of life: the ImproveCare feasibility cluster RCT. *Health Technol Assess* 2019;23:1–150.
342. Ngaage D, Mitchell N, Dean A, et al. Feasibility study of early outpatient review and early cardiac rehabilitation after cardiac surgery: mixed-methods research design—a study protocol. *BMJ Open* 2019;9:e035787.
343. Rapport F, Auton E, Warren C, Braithwaite J. Addressing clinical equipoise for hearing devices: the qualitative COACH (q-COACH) study protocol for Australian stakeholder involvement in the design of a randomised controlled trial. *BMJ Open* 2019;9:e030100.
344. Lewis RK. Mental health strategies for prevention and intervention: community perspectives. *J Prev Interv Community* 2019;1–4.
345. Sanders AB, Hiller K, Duldner J. Researchers' understanding of the federal guidelines for waiver of and exception from informed consent. *Acad Emerg Med* 2005;12:1045–9.
346. Kowey P, Ornato J. Resuscitation research and emergency waiver of informed consent. *Resuscitation* 2000;47:307–10.
347. Sayre MR, White LJ, Brown LH, McHenry SD. National EMSRAWT. The National EMS Research Agenda executive summary. Emergency Medical Services. *Ann Emerg Med* 2002;40:636–43.
348. Hsieh M, Dailey MW, Callaway CW. Surrogate consent by family members for out-of-hospital cardiac arrest research. *Acad Emerg Med* 2001;8:851–3.
349. McClure KB, Delorio NM, Gunnels MD, Ochsner MJ, Biros MH, Schmidt TA. Attitudes of emergency department patients and visitors regarding emergency exception from informed consent in resuscitation research, community consultation, and public notification. *Acad Emerg Med* 2003;10:352–9.
350. Nichol G, Steen P, Herlitz J, et al. International Resuscitation Network Registry: design, rationale and preliminary results. *Resuscitation* 2005;65:265–77.
351. Hartmann M, Hartmann-Vareilles F. The clinical trials directive: how is it affecting Europe's noncommercial research? *PLoS Clin Trials* 2006;1:e13.
352. McMahon AD, Conway DI, Macdonald TM, McInnes GT. The unintended consequences of clinical trials regulations. *PLoS Med* 2009;3:e1000131.
353. Bosch X. Europe's restrictive rules strangling clinical research. *Nat Med* 2005;11:1260.
354. Liddell K, Kompanje EJ, Lemaire F, et al. Recommendations in relation to the EU clinical trials directive and medical research involving incapacitated adults. *Wien Klin Wochenschr* 2006;118:183–91.
355. Tolle SW, Teno JM. Lessons from oregon in embracing complexity in end-of-life care. *N Engl J Med* 2017;376:1078–82.
356. Grasner JT, Masterson S. EuReCa and international resuscitation registries. *Curr Opin Crit Care* 2015;21:215–9.
357. Girotra S, Nallamothu BK, Spertus JA, et al. Trends in survival after in-hospital cardiac arrest. *N Engl J Med* 2012;367:1912–20.
358. Merchant RM, Berg RA, Yang L, et al. Hospital variation in survival after in-hospital cardiac arrest. *J Am Heart Assoc* 2014;3:e000400.
359. Hagihara A, Hasegawa M, Abe T, Nagata T, Wakata Y, Miyazaki S. Prehospital epinephrine use and survival among patients with out-of-hospital cardiac arrest. *JAMA* 2012;307:1161–8.



360. Nakahara S, Tomio J, Takahashi H, et al. Evaluation of pre-hospital administration of adrenaline (epinephrine) by emergency medical services for patients with out of hospital cardiac arrest in Japan: controlled propensity matched retrospective cohort study. *BMJ* 2013;347:f6829.
361. Donnino MW, Saliccioli JD, Howell MD, et al. Time to administration of epinephrine and outcome after in-hospital cardiac arrest with non-shockable rhythms: retrospective analysis of large in-hospital data registry. *BMJ* 2014;348:g3028.
362. Bak MAR, Blom MT, Tan HL, Willems DL. Ethical aspects of sudden cardiac arrest research using observational data: a narrative review. *Crit Care* 2018;22:212.
363. Mumma BE, Diercks DB, Danielsen B, Holmes JF. Probabilistic linkage of prehospital and outcomes data in out-of-hospital cardiac arrest. *Prehosp Emerg Care* 2015;19:358–64.
364. Priori SG. Genetic testing to predict sudden cardiac death: current perspectives and future goals. *Indian Heart J* 2014;66(Suppl. 1): S58–60.
365. Docherty AB, Lone NI. Exploiting big data for critical care research. *Curr Opin Crit Care* 2015;21:467–72.
366. Pullman D, Hodgkinson K. Genetic knowledge and moral responsibility: ambiguity at the interface of genetic research and clinical practice. *Clin Genet* 2006;69:199–203.
367. Badcock D, Kelly AM, Kerr D, Reade T. The quality of medical record review studies in the international emergency medicine literature. *Ann Emerg Med* 2005;45:444–7.
368. Kaji AH, Schriger D, Green S. Looking through the retrospectroscope: reducing bias in emergency medicine chart review studies. *Ann Emerg Med* 2014;64:292–8.
369. Nishiyama C, Brown SP, May S, et al. Apples to apples or apples to oranges? International variation in reporting of process and outcome of care for out-of-hospital cardiac arrest. *Resuscitation* 2014;85: 1599–609.
370. Manrai AK, Funke BH, Rehm HL, et al. Genetic misdiagnoses and the potential for health disparities. *N Engl J Med* 2016;375: 655–65.
371. Tu JV, Willison DJ, Silver FL, et al. Impracticability of informed consent in the Registry of the Canadian Stroke Network. *N Engl J Med* 2004;350:1414–21.
372. Freeman BD, Bolcic-Jankovic D, Kennedy CR, et al. Perspectives of decisional surrogates and patients regarding critical illness genetic research. *AJOB Empir Bioeth* 2016;7:39–47.
373. Offerman SR, Nishijima DK, Ballard DW, Chetipally UK, Vinson DR, Holmes JF. The use of delayed telephone informed consent for observational emergency medicine research is ethical and effective. *Acad Emerg Med* 2013;20:403–7.
374. Fox EE, Bulger EM, Dickerson AS, et al. Waiver of consent in noninterventonal, observational emergency research: the PROMMTT experience. *J Trauma Acute Care Surg* 2013;75: S3–8.
375. Fang J, Kapral MK, Richards J, Robertson A, Stamplecoski M, Silver FL. The Registry of Canadian Stroke Network: an evolving methodology. *Acta Neurol Taiwan* 2011;20:77–84.
376. Thorogood A, Zawati MH. International guidelines for privacy in genomic biobanking (or the unexpected virtue of pluralism). *J Law Med Ethics* 2015;43:690–702.
377. Auffray C, Balling R, Barroso I, et al. Making sense of big data in health research: towards an EU action plan. *Genome Med* 2016; 8:71.
378. Bodenheimer T. Uneasy alliance—clinical investigators and the pharmaceutical industry. *N Engl J Med* 2000;342:1539–44.
379. Moorthy VS, Karam G, Vannice KS, Kieny MP. Rationale for WHO's new position calling for prompt reporting and public disclosure of interventional clinical trial results. *PLoS Med* 2015;12:e1001819.
380. Kiley R, Peatfield T, Hansen J, Reddington F. Data sharing from clinical trials – a research funder's perspective. *N Engl J Med* 2017;377:1990–2.
381. Ornato JP, Becker LB, Weisfeldt ML, Wright BA. Cardiac arrest and resuscitation: an opportunity to align research prioritization and public health need. *Circulation* 2010;122:1876–9.
382. Kudenchuk PJ, Brown SP, Daya M, et al. Amiodarone, lidocaine, or placebo in out-of-hospital cardiac arrest. *N Engl J Med* 2016;374:1711–22.
383. Tierney WM, Meslin EM, Kroenke K. Industry support of medical research: important opportunity or treacherous pitfall? *J Gen Intern Med* 2016;31:228–33.
384. Ong MEH, Perkins GD, Cariou A. Out-of-hospital cardiac arrest: prehospital management. *Lancet* 2018;391:980–8.
385. Chan PS, Berg RA, Nadkarni VM. Code blue during the COVID-19 pandemic. *Circ Cardiovasc Qual Outcomes* 2020;13:e006779.
386. Kramer DB, Lo B, Dickert NW. CPR in the Covid-19 Era – an ethical framework. *N Engl J Med* 2020;383:e6.
387. Mahase E, Kmiotowicz Z. Covid-19: doctors are told not to perform CPR on patients in cardiac arrest. *BMJ* 2020;368:m1282.
388. Shao F, Xu S, Ma X, et al. In-hospital cardiac arrest outcomes among patients with COVID-19 pneumonia in Wuhan, China. *Resuscitation* 2020;151:18–23.
389. Baldi E, Sechi GM, Mare C, et al. Out-of-hospital cardiac arrest during the covid-19 outbreak in Italy. *N Engl J Med* 2020;383:496–8.
390. Van de Voorde P, Bossaert L, Mentzelopoulos S, et al. Ethics of resuscitation and end-of-life decisions. *Notf Rett Med* 2020;23: 263–7.
391. Docherty AB, Harrison EM, Green CA, et al. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study. *BMJ* 2020;369:m1985.
392. Grasselli G, Pesenti A, Cecconi M. Critical care utilization for the COVID-19 outbreak in Lombardy Italy: early experience and forecast during an emergency response. *JAMA* 2020;323:1545–6.
393. Kandori K, Narumiya H, Iizuka R. Extracorporeal cardiopulmonary resuscitation should not be performed on confirmed or suspected COVID-19 patients. *Resuscitation* 2020;153:6–7.
394. Concato J, Shah N, Horwitz RJ. Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med* 2000;342:1887–92.
395. Barnish MS, Turner S. The value of pragmatic and observational studies in health care and public health. *Pragmat Obs Res* 2017;8:49–55.
396. Howes C. Caring until the end: a systematic literature review exploring Paediatric Intensive Care Unit end-of-life care. *Nurs Crit Care* 2015;20:41–51.
397. Michalsen A, Long AC, DeKeyser Ganz F, et al. Interprofessional shared decision-making in the ICU: a systematic review and recommendations from an expert panel. *Crit Care Med* 2019;47: 1258–66.
398. Hawkes CA, Fritz Z, Deas G, et al. Development of the recommended summary plan for emergency care and treatment (ReSPECT). *Resuscitation* 2020;148:98–107.
399. Taylor J, Booth A, Beresford B, Phillips B, Wright K, Fraser L. Specialist paediatric palliative care for children and young people with cancer: a mixed-methods systematic review. *Palliat Med* 2020;34:731–75.