

Editorial



Interplay between Saturated Free Fatty Acids and mmLDL Induces Inflammation in LPS-stimulated Macrophages

Jeong Su Park , PhD¹, and Soo Han Bae , PhD^{1,2}¹Severance Biomedical Science Institute, Yonsei University College of Medicine, Seoul, Korea²Brain Korea 21 PLUS Project for Medical Science, Yonsei University, Seoul, Korea

► See the article "*Nlrp3*, *Csf3*, and *Edn1* in Macrophage Response to Saturated Fatty Acids and Modified Low-Density Lipoprotein" in volume 51 on page 68.

Received: Sep 2, 2020

Accepted: Sep 9, 2020

Correspondence to


Soo Han Bae, PhD

Severance Biomedical Science Institute, Yonsei University College of Medicine, 50-1, Yonsei-ro, Seodaemun-gu, Seoul 03722, Korea.
E-mail: soohanbae@yuhs.ac

Copyright © 2021. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Jeong Su Park <https://orcid.org/0000-0003-4551-4294>Soo Han Bae <https://orcid.org/0000-0002-8007-2906>

Conflict of Interest

The authors have no financial conflicts of interest.

Author Contributions

Supervision: Bae SH; Writing - original draft: Park JS; Writing - review & editing: Park JS. The contents of the report are the author's own views and do not necessarily reflect the views of the *Korean Circulation Journal*.We have followed an article titled "Nlrp3, Csf3, and Edn1 in Macrophage Response to Saturated Fatty Acids and Modified Low-Density Lipoprotein" by Lee *et al.*¹ with profound interest. The study showed that saturated fatty acids, palmitic acid (PA) in particular, and minimally modified low-density lipoprotein (mmLDL) induce the secretion of pro-inflammatory cytokines through activation of the extracellular signal-regulated kinase/mitogen-activated protein kinase pathway in lipopolysaccharide (LPS)-stimulated macrophages. Further, the authors identified several genes targeted by PA or mmLDL via RNA sequencing of LPS-stimulated macrophages. Although this study provides insights into the detailed molecular mechanism underlying the interplay between metabolic stress and inflammation, the insights from this study must be analyzed in the context of several concerns that have been discussed below.NLR family pyrin domain containing 3 (NLRP3) is an essential regulator of adaptive immune response and its expression is induced by endogenous cytokines or microbial molecules through binding to Toll-like receptor (TLR), interleukin-1 receptor (IL-1R), and tumor necrosis factor receptor (TNFR).² Activation of NLRP3 via K⁺ efflux, Ca²⁺ signaling, and mitochondrial reactive oxygen species (mtROS) triggers caspase-1 activation and subsequent secretion of IL-1 β and IL-18.^{2,3} As NLRP3 expression is known to be regulated by post-translational modifications (PTM),^{2,4} the increase in NLRP3 messenger RNA (mRNA) level is not sufficient to promote its activation. However, this aspect has been overlooked in this study, with the authors only reporting an increase in NLRP3 at the mRNA level in LPS-stimulated macrophages treated with PA or mmLDL. Therefore, it is crucial to estimate the NLRP3 protein levels to gain comprehensive insights into the action of PA or mmLDL in LPS-stimulated macrophages.Further, this study identified several genes (*Nlrp3*, *Csf3*, and *Edn1*) associated with inflammation in LPS-treated macrophages that are targeted by PA or mmLDL.¹ In this context, it must be noted that the activation of inflammatory responses is closely related with diverse sets of metabolic disease conditions, and the intricate interplay between inflammation and metabolic stress is mechanistically complex. In particular, PA, a free fatty

acid (FFA), has been widely reported to be a key regulator of insulin resistance and NLRP3 inflammasome formation in obesity models.^{5,6)} Recent research has highlighted the role of PA as a TLR agonist that leads to the direct activation of the NLRP3 inflammasome and onset of insulin resistance in obesity models.^{7,8)} Further, PA has been shown to induce insulin resistance through regulation of the nuclear factor (NF)- κ B/insulin receptor substrate 1 axis in hepatocytes (Huh7 cells) and liver of high-fat diet-fed mouse.⁵⁾ Therefore, the interplay between FFA-mediated activation of inflammation and the pathophysiology of metabolic disease must be considered while interpreting results from the study under consideration.

mmLDL—a product of progressive low-density lipoprotein (LDL) oxidation—and oxidized LDL have been recognized as biomarkers of inflammation in human macrophages.⁹⁾ The authors have previously reported that mmLDL induces the secretion of chemokines (including chemokine [C-X-C motif] ligand 2) and TNF α in LPS-treated macrophages in the presence of PA.¹⁰⁾ Here, they report the regulation of genes by mmLDL in LPS-stimulated macrophages and elucidate the detailed mechanism underlying mmLDL-mediated inflammatory response to LPS in macrophages. Taken together, these insights are expected to contribute to the development of approaches for the treatment of inflammatory diseases.¹⁾

REFERENCES

1. Youk H, Kim M, Lee CJ, et al. *Nlrp3*, *Csf3*, and *Edn1* in macrophage response to saturated fatty acids and modified low-density lipoprotein. *Korean Circ J* 2021;51:68-80.
[CROSSREF](#)
2. He Y, Hara H, Núñez G. Mechanism and regulation of NLRP3 inflammasome activation. *Trends Biochem Sci* 2016;41:1012-21.
[PUBMED](#) | [CROSSREF](#)
3. Kelley N, Jeltema D, Duan Y, He Y. The NLRP3 inflammasome: an overview of mechanisms of activation and regulation. *Int J Mol Sci* 2019;20:3328.
[PUBMED](#) | [CROSSREF](#)
4. Juliana C, Fernandes-Alnemri T, Kang S, Farias A, Qin F, Alnemri ES. Non-transcriptional priming and deubiquitination regulate NLRP3 inflammasome activation. *J Biol Chem* 2012;287:36617-22.
[PUBMED](#) | [CROSSREF](#)
5. Wang Y, Cao F, Wang Y, Yu G, Jia BL. Silencing of SAA1 inhibits palmitate- or high-fat diet induced insulin resistance through suppression of the NF- κ B pathway. *Mol Med* 2019;25:17.
[PUBMED](#) | [CROSSREF](#)
6. Ding S, Xu S, Ma Y, Liu G, Jang H, Fang J. Modulatory mechanisms of the NLRP3 inflammasomes in diabetes. *Biomolecules* 2019;9:850.
[PUBMED](#) | [CROSSREF](#)
7. Xu L, Wang W, Zhang X, et al. Palmitic acid causes insulin resistance in granulosa cells via activation of JNK. *J Mol Endocrinol* 2019;62:197-206.
[PUBMED](#) | [CROSSREF](#)
8. Fusco R, Siracusa R, Genovese T, Cuzzocrea S, Di Paola R. Focus on the role of NLRP3 inflammasome in diseases. *Int J Mol Sci* 2020;21:4223.
[PUBMED](#) | [CROSSREF](#)
9. Lara-Guzmán OJ, Gil-Izquierdo Á, Medina S, et al. Oxidized LDL triggers changes in oxidative stress and inflammatory biomarkers in human macrophages. *Redox Biol* 2018;15:1-11.
[PUBMED](#) | [CROSSREF](#)
10. Ann SJ, Kim KK, Cheon EJ, et al. Palmitate and minimally-modified low-density lipoprotein cooperatively promote inflammatory responses in macrophages. *PLoS One* 2018;13:e0193649.
[PUBMED](#) | [CROSSREF](#)