


RESEARCH ARTICLE

Open Access



The influence of social media on recruitment to surgical trials

Carly Nichola Bisset^{1*} , Ben Carter², Jennifer Law³, Jonathan Hewitt⁴, Kat Parmar⁵, Susan Joan Moug¹ and on behalf of the ELF Study Group Collaborative Authorship

Abstract

Background: Social media has changed the way surgeons communicate worldwide, particularly in dissemination of trial results. However, it is unclear if social media could be used in recruitment to surgical trials. This study aimed to investigate the influence of Twitter in promoting surgical recruitment in The Emergency Laparotomy and Frailty (ELF) Study.

Methods: The ELF Study was a UK-based, prospective, observational cohort that aimed to assess the influence of frailty on 90-day mortality in older adults undergoing emergency surgery. A power calculation required 500 patients to be recruited to detect a 10% change in mortality associated with frailty. A 12-week recruitment period was selected, calculated from information submitted by participating hospitals and the numbers of emergency surgeries performed in adults aged > 65 years. A Twitter handle was designed (@ELFStudy) with eye-catching logos to encourage enrolment and inform the public and clinicians involved in the study. Twitter Analytics and Twitonomy (Digonomy Pty Ltd) were used to analyse user engagement in relation to patient recruitment.

Results: After 90 days of data collection, 49 sites from Scotland, England and Wales recruited 952 consecutive patients undergoing emergency laparotomy, with data logged into a database created on REDCap. Target recruitment ($n = 500$) was achieved by week 11.

A total of 591 tweets were published by @ELFStudy since its conception, making 218,136 impressions at time of writing. The number of impressions (number of times users see a particular tweet) prior to March 20th 2017 (study commencement date) was 23,335 (343.2 per tweet), compared to the recruitment period with 114,314 impressions (256.3 per tweet), ending June 20th 2017. Each additional tweet was associated with an increase in recruitment of 1.66 (95%CI 1.36 to 1.97; $p < 0.001$).

Conclusion: The ELF Study over-recruited by nearly 100%, reaching over 200,000 people across the U.K. Branding enhanced tweet aesthetics and helped increase tweet engagement to stimulate discussion and healthy competition amongst clinicians to aid trial recruitment. Other studies may draw from the social media experiences of the ELF Study to optimise collaboration amongst researchers.

Trial registration: This study is registered online at www.clinicaltrials.gov (registration number [NCT02952430](https://clinicaltrials.gov/ct2/show/study/NCT02952430)) and has been approved by the National Health Service Research Ethics Committee.

Keywords: Social media, Recruitment, Surgical trials

* Correspondence: cbisset@nhs.net

¹Department of General Surgery, Royal Alexandra Hospital, Paisley PA2 9PN, UK

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Social media is the term used to describe the interactive, web-based free applications, that have been instrumental in increasing accessibility of information by allowing instantaneous worldwide communication. The reach of social media in modern society is extensive, with an estimated 2.46 billion users of any social media application reported in 2017 [1].

One of these applications is Twitter (Twitter Inc., San Francisco, CA, USA): an online micro-blogging site which limits messages (or 'tweets') to 280 characters or less, and may be attached to an image or website link and can connect to other notable Twitter users/institutions by 'tagging' and thereby including other users. Twitter is an ideal social media modality to convey concise messages on a public platform, is free at point of service and is often used professionally to represent the views of that user or institution. Social media usage is often associated with increasing engagement with professional bodies and surgical societies such as the Association of Coloproctologists' of Great Britain and Ireland (ACPGBI) [2]. There is also some evidence to suggest that the age of the user will affect platform preference in social media usage, for example amongst emergency medicine physicians, trainees tended to use Facebook and YouTube, whilst older, more experienced physicians used Twitter and LinkedIn [3]. Whether or not this is true of general surgery is yet to be established.

In surgery, the role of social media is evolving, with reported beneficial effects including: facilitating patient education, sharing information on new guidelines or published research and increasing collaboration amongst stakeholders such as: patients, clinicians, industry, trainees and educational institutions [4, 5].

These benefits to research raise the question of the influence of social media on performing research, in particular: recruitment to trials. It is anticipated that recruitment to research trials using social media may substantially increase recruitment numbers in a cost-effective manner [6]. Little is known in this area, however social media outlets such as Twitter could facilitate recruitment specifically to surgical trials and engagement in research, such as that seen in the National Audit of Small Bowel Obstruction study (NASBO) [7]. This study demonstrated that tweets which involved images and tagged other users, led to higher levels of engagement with tweets and maintained collaborator engagement. Unfortunately within surgical research, use of social media is variable amongst different specialties in the UK including: colorectal surgeons (3.1%) [2], vascular surgeons (4.8%) [7] and plastic surgeons (22%) [8], which overall seems low in comparison to the estimated 1 in 5 doctors in the UK using Twitter on a daily basis [9].

The Emergency Laparotomy and Frailty (ELF) Study was a multi-disciplinary, multicentre prospective cohort

study undertaken in the UK in 2017 [10]. Using Twitter (@ElfStudy) to advertise for site registration and patient recruitment, the ELF Study achieved its target recruitment early, reporting significant over-recruitment at completion. This study considers the influence of social media on ELF Study's recruitment success, offering insight on how clinical trial groups may use social media-based branding strategies to promote recruitment to surgical trials.

Methods

The ELF Study investigated the relationship to pre-operative frailty and mortality at 90-days in older adults undergoing emergency laparotomy in the U.K. (trial registration number NCT02952430). Consecutive patient recruitment was performed for 3 months (20th March 2017 - 20th June 2017) with each patient subsequently followed up for 90-days. Target recruitment was calculated with 500 patients anticipated to detect a 10% change in mortality associated with frailty. Ethical approval was given by the National Health Service Research Ethics Committee (Black Country Research Committee, November 2016; 16/WM/0500), with central registration at the Health Research Authority (HRA) for English sites, NHS Research Scotland Permissions Coordinating Centre (NRSPCC) for Scottish sites and Health and Care Research Permissions Service for Welsh sites [11]. Patient consent was not necessary for this study.

The ELF Study was a trainee-led initiative, led by the North West Research Collaborative (surgical trainees) and the Older Persons Surgical Outcomes Collaboration (OPSOC: surgeons and geriatricians). The principal investigators at each site were either general surgery registrars or consultant surgeons in a hospital performing emergency abdominal surgery and peri-operative care. Principal investigators led local teams of trainees, but each team required a consultant general surgeon to ensure collaborators acted in accordance with local clinical governance and guidelines.

The eligibility criteria for the ELF Study is listed below:

- Patients aged > 65 years
- Patients who undergo an expedited, urgent or emergency abdominal procedure on the gastrointestinal tract, including:
 - Open, laparoscopic or laparoscopic-assisted procedures
 - Procedures involving the stomach, small or large bowel, or rectum for conditions such as: perforation, ischaemia, abdominal abscess, bleeding or obstruction
 - Washout/evacuation of intraperitoneal haematoma or abscess (except when due to appendicitis or cholecystitis)

- Bowel resection/repair due to incarcerated umbilical, inguinal and femoral hernias, or acute incisional hernias
- Laparotomy or laparoscopy with inoperable pathology (i.e. peritoneal or hepatic metastases)
- Laparoscopic/open adhesiolysis
- Return to theatre for repair of substantial dehiscence of major abdominal wound (i.e. 'burst abdomen'), or any major post-operative complication

The Twitter handle @ElfStudy was opened (8th January 2017) with eye-catching and consistent 'branding' used to advertise for site registration prior to the recruitment period (10 weeks prior to the date of study commencement). This branding took the form of a catchy and consistent colour scheme (red and green), and used a study-specific Twitter account (@ELFStudy). This Twitter account provided regular public updates on: study recruitment requirements, news on data collection sparking regional and hospital-based light-hearted competition, sharing links to related research articles of interests and presentations by the steering group on provisional ELF Study results at general surgery, geriatrics and anaesthetic research meetings.

Definitions

Twitter Analytics and an online analytics tool (Twitonomy; Digonomy Pty Ltd) were used to examine potential patterns or changes in user engagement in relation to weekly progression of the study, with definitions described in Table 1.

Outcome and exposure

Exposure of social media was defined in three ways: the total number of tweets by @ELFStudy; the total number of the impressions and engagements of each tweet (a marker of the reach of each tweet and interactions); and the maximum number of impressions and engagements with a single tweet in the week. Cumulative recruitment (i.e. weekly recruitment of patients) and number of tweets were our primary measures of outcome and exposure. The final week was excluded (week 13 - study end date), as some sites had collated their recruitment numbers and reported them in Week 13 to a database set up using REDCap, rather than contemporaneously over the course of the trial. The overall denominator for the number of recruiting surgeons and trainees cannot be calculated using a social media approach, as the number of impressions may include irrelevant individuals, groups or organisations.

The pre-trial phase and recruitment phase were selected for comparison as the content of tweets changed between the two periods. In the pre-trial phase, most tweet content was based upon individual sites signing up to the trial for patient recruitment, with general

Table 1 Definitions used in Twitter Analytics

| Analytical Tool | Definition |
|-----------------|---|
| Impressions | Number of times users see a particular tweet Total number of times a tweet was seen by individual users on Twitter |
| Engagements | Total number of times a tweet was interacted with (including users opening a link, re-tweeting, replies to tweets, 'likes') |
| Re-tweet | The forwarding of an original tweet to another user's profile, sharing information with other users |
| Likes | Users indicate they agree with the sentiment of tweet content |
| Followers | Number of users who receive updates on a particular account's posts (i.e. the greater number of followers, the greater the likelihood of higher tweet engagement) |
| Tags / Tagging | Including another user in Tweet content and/or directing a message at another user, often increasing the engagement of the Tweet e.g. @ELFStudy |
| Hashtags | 'Tagging' groups a collection of tweets with similar content so that users may find a common thread of tweets on a similar topic e.g. #colorectalsurgery, #surgicaltraining |
| User | An individual or an organisation with a Twitter account |
| Handle | The username beginning with "@" which is unique to that Twitter account e.g. @ELFStudy |

information on how to become involved in the ELF Study. In the recruitment phase, tweet content became more focussed on maintaining recruitment momentum, with site-specific and whole-study updates which encouraged competition. Tweets from the recruitment phase included: site league tables, photographs from local recruiters (for example, containing 'elf' toys facilitating the branding strategy) and general trial progress updates.

Statistical analysis

Tweets published by @ELFStudy were divided into two time phases to allow comparison to weekly recruitment figures: 1) pre-trial commencement (pre-trial phase), 2) trial commencement and data collection (recruitment phase – date selected for commencement March 20th 2017 and date of first patient recruited). Tweets after these time phases were excluded.

For the 12 weeks of the study, we fitted a linear regression to model cumulative recruitment. To assess the additional impact of the social media campaign we fitted: sum of engagement; and total cumulative number of tweets as two independent analyses. The different models were compared using the adjusted coefficient to determination (adjusted r^2) summary. The slope parameter was presented alongside the 95% Confidence interval and p -value.

Results

Study recruitment

A total of 52 sites registered with 49 submitting data to a database set up in REDCap. There was a total of 161 collaborators (54 consultants, 107 trainees). The target recruitment ($n = 500$) was achieved by week 11 with recruitment increasing exponentially from week 11 to almost double the intended target by week 12 ($n = 952$) (Table 2). A total of 926 tweets were published by @ELFStudy during the pre-trial (phase 1) and recruitment phases (phase 2). Site recruitment of patients meeting the inclusion criteria was performed by junior doctors and consultants. Interested users on Twitter could choose to 'follow' the @ELFStudy account, which is open to the public, to receive updates on the study.

Impressions and engagements

The number of phase 1 tweets was 69, resulting in 23,335 impressions (343.2 per tweet) versus 437 tweets published in phase 2 generating 114,314 impressions (256.3 per tweet) (Fig. 1). The number of engagements during phase 1 was 1124, compared to 4523 engagements in phase 2, with the recruitment phase seeing four times the number of tweet engagements of that seen in the pre-trial phase. This suggests that more users were reached and were more engaging with tweets published by @ELFStudy during the recruitment phase, possibly reflecting increased overall recruitment of patients (Fig. 2).

Week 6 marked the time period where recruitment exceeded the target prediction, however this did not bear correlation to the number of tweet engagements and impressions (Fig. 3).

Interestingly from week 11, the number of impressions and engagements markedly increased perhaps reflecting the 'final push' effect of individual sites logging data.

This is reflected by the overall number of tweets published by @ELFStudy, which predictably had a large amount of activity at the beginning and latter stages of the recruitment phase, with a moderate number of tweets published in the mid-way point (Fig. 3).

Data analysis

We present the aggregate results of the social media data and weekly recruitment data (Table 2). This table demonstrates a weekly temporal effect, where there is an increasing number of patients recruited on a weekly basis and increasing tweet engagement. After fitting the two linear regressions, it was clear that the cumulative number of tweets was associated with the cumulative recruitment (Fig. 4). It was found that the adjusted- $r^2 = 0.94$. and each additional tweet was associated with an increase in recruitment of 1.66 (95% CI 1.36, to 1.97; $p < 0.001$). There was no association between the cumulative engagement and cumulative recruitment.


Discussion

Social media has revolutionised academia and the distribution of information. Not only do social media outlets such as Twitter inform users of trial updates, it also has a potential role in changing clinical practice with trial conclusion. The vast online network connects users of the platform across the globe instantaneously to discuss a common theme and share experiences, particularly with the use of hashtags which group together messages that may be followed more easily by users who search for that hashtag item (e.g. #colorectalresearch).

Part of the successful social media recruitment drive to this study was undoubtedly due to the non-professional marketing strategy of the ELF Study, which played upon a Christmas theme in keeping with its


Table 2 Weekly recruitment comparison with tweet engagement


| Week | Weekly recruitment | Weekly number of tweets | Weekly Total Impressions | Weekly Total Engagements | Cumulative Patient Recruitment | Cumulative number of tweets |
|------|--------------------|-------------------------|--------------------------|--------------------------|--------------------------------|-----------------------------|
| 1 | 30 | 30 | 7938 | 553 | 30 | 30 |
| 2 | 28 | 55 | 8142 | 384 | 58 | 85 |
| 3 | 41 | 61 | 11,261 | 470 | 99 | 146 |
| 4 | 23 | 8 | 1027 | 18 | 122 | 154 |
| 5 | 49 | 56 | 8033 | 448 | 171 | 210 |
| 6 | 56 | 20 | 3960 | 164 | 227 | 230 |
| 7 | 67 | 33 | 2528 | 50 | 294 | 263 |
| 8 | 69 | 23 | 4254 | 103 | 363 | 286 |
| 9 | 45 | 30 | 12,020 | 426 | 408 | 316 |
| 10 | 62 | 20 | 7459 | 182 | 470 | 336 |
| 11 | 37 | 15 | 6780 | 160 | 507 | 351 |
| 12 | 100 | 35 | 12,979 | 552 | 607 | 386 |



namesake. Consistent use of a Christmas colour scheme and themed logo made the study more visible amongst its peers and was easily identifiable when presented at meetings as oral and poster presentations (Additional Figures 1 and 2). This reflects standardised marketing strategies by other major industries which recognise that consumer participation increases engagement quality and experiences (<https://sysomos.com/inside-twitter/twitter-statistics/> (accessed 28/2/19)). A professional marketing strategy was not considered due to high anticipated costs, within a small funding budget from the Bowel Disease Research Fund (charitable

status). Twitter provides a free at point-of-access online platform which has a large colorectal surgeon and trainee membership, and users can possess multiple accounts (e.g. a personal account, and an organisation account such as @ELFStudy) [2]. With increasing popularity of Twitter profiles for research trials, it is becoming more difficult to ‘stand out’ and have a visible online presence [12]. Creative strategies in advertising which engage consumers recognise: ‘play on words’ appeal (e.g. ELF), animation (e.g. use of GIFs and other illustrations) and social appeal (e.g. as part of the increasing spread of multi-centre, multi-author






collaborative studies) as being highly influential factors which increase consumerism [13, 14].

The successes of the ELF Study in recruiting patients ahead of time, and in excess, may be reflected by the findings of Nowotarski et al., who found that of 179 adults recruited for a medical trial, only 30% were aware of information providing clinical trial websites (e.g. clinicaltrials.gov). Once these adults were informed of the online network available for trial recruitment, 81% showed interest in engaging with relevant clinical trials [15]. Whilst our study did not directly investigate patient information and consent to surgical trials, there is some evidence that social media may facilitate the education of patients and clinicians on relevant clinical trials. The open manner with which information is received through social media may increase engagement, as users have a direct contact (or a ‘face’) they can seek further information

from. Site recruitment will also have been facilitated by ‘word-of-mouth’ marketing, via presentations and networking at surgical and geriatric medicine conferences. The open nature of social media is further enhanced by collaboration and sharing of information (including sharing of presentations at conferences), and allows multi-centre studies to communicate with individual sites and collectively with all sites to address problems with trial protocol, trial recruitment and to instil a sense of ‘healthy competition’, where sites could compete for optimal trial recruitment and participation.

Twitter was the only social media modality used for the ELF Study and this paper, given the high level of professional usage within the outlet, in comparison to other platforms which have not been examined as part of this study, which many users have for personal and not professional reasons [2]. As is true with all social



media modalities which are instantaneous, in comparison to traditional industrial media (e.g. printed news, television), communications such as tweets may be altered at any time and ‘deleted’, meaning that social media data may be viewed as less ‘permanent’ than industrial level media [16]. In order to ethically manage this and moderate pages, several boards have drawn consensus on moderating the use of social media platforms for professional purposes such as the General Medical Council (UK). There is increasing governance over content of social media posts in medicine, particularly relating to patient confidentiality, or other violations of professionalism such as: inappropriate contact with patients, declaring conflicts of interest and maintaining professionalism amongst colleagues and other users [17, 18].

As recruitment data was collected per week, daily analysis of the effect of tweets on patient recruitment was not possible. It is likely that the large number of impressions @ELFStudy generated (a surrogate marker for study ‘reach’) included individuals and organisations who may not have an interest in peri-operative patient care. The number of impressions may also be higher than the actual number of users seeing an individual tweet, particularly where users have multiple accounts (for example, a personal and an organisation account). It is not currently possible to identify which users met the intended ‘target audience’ for recruitment, or which users contribute to tweet impressions. However, it is also possible that Twitter users in our group share an ‘echo chamber’; where Twitter updates are seen and shared by like-minded users, contributing to confirmation bias.

Conclusion

The ELF Study is a multi-disciplinary, multi-centre trial whose success was undoubtedly assisted by social media and non-professional marketing. Eye-catching logos and consistent colour schemes enhanced tweet aesthetics and helped increase tweet engagement to stimulate discussion and healthy competition amongst clinicians to aid trial recruitment.

Social media usage in surgical research is becoming standard, however its role in other medical specialties is still to be explored and utilised; including geriatric medicine where there is no current evidence to demonstrate clinician engagement and patterns of usage. The ELF Study achieved target recruitment within half of the predicted time period, a feat not to be underestimated. Other studies may draw from the social media experiences of the ELF Study, including: regular updates to inform participants of study progress, successful ‘branding’ playing on the study acronym and ultimately engaging a multi-disciplinary platform of: physicians,

surgeons and anaesthetists to collaborate and produce high quality peri-operative outcomes data.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12874-020-01072-1>.

Additional file 1: Figure S1. The Elf Study logo as an example of the branding strategy. **Figure S2.** Example of tweet media content using the branding strategy

Abbreviations

ELF: Emergency Laparotomy and Frailty Study; ACPGBI: Association of Coloproctology of Great Britain and Ireland; NASBO: National Audit of Small Bowel Obstruction study; HRA: Health Research Authority; NRSPPC: NHS Research Scotland Permissions Coordinating Centre

Acknowledgements

This work was presented in poster format at the Association of Coloproctologists’ of Great Britain and Ireland Annual Meeting in July 2019, with the abstract published in *Colorectal Disease*. <https://onlineibrary.wiley.com/doi/full/10.1111/codi.14770>

The ELF Study Group* collaborative authorship (contributing authors by site in alphabetical order).

| Site | Contributors |
|---------------------------------------|-----------------------|
| Addenbrooke’s Hospital | Bryony Ross |
| | Julia Oleksiewicz |
| | Nicola Fearnhead |
| Blackpool Victoria Hospital | Christopher Jump |
| | Jemma Boyle |
| | Alex Shaw |
| | Jonathan Barker |
| Bristol Royal Infirmary | Jane Hughes |
| | Jonathan Randall |
| | Isileli Tonga |
| | James Kynaston |
| | Matthew Boal |
| Countess of Chester Hospital | Nicola Eardley |
| | Elizabeth Kane |
| | Harriet Reader |
| | Sunanda Roy Mahapatra |
| Croydon University Hospital | Michael Garner-Jones |
| | Jessica Juliana Tan |
| | Said Mohamed |
| Doncaster Royal Infirmary | Rina George |
| | Ed Whiteman |
| East Cheshire NHS Trust | Kamran Malik |
| | Christopher J Smart |
| | Monica Bogdan |
| East Lancashire Hospitals (Blackburn) | Madhu Parna Chaudhury |
| | Videha Sharma |
| | Daren Subar |

(Continued)

| | |
|--|--|
| Furness General Hospital (Barrow) | Panna Patel Sok-Moi Chok Evelyn Lim |
| Glan Clywd Hospital | Vedamurthy Adhiyaman Glesni Davies |
| Glasgow Royal Infirmary | Ellen Ross Rudra Maitra Colin W Steele Campbell Roxburgh |
| Gloucestershire Royal Hospital | Shelly Griffiths Natalie S Blencowe Emily N Kirkham |
| Lorn and Islands Hospital (Oban) | John S Abraham Kirsty Griffiths |
| Maidstone and Tunbridge Wells NHS Trust | Yasser Abdulaal Muhammad Rafaih Iqbal Munir Tarazi |
| Manchester Royal Infirmary | James Hill Azam Khan Ian Farrell |
| Mid Essex / Broomfield | Gemma Conn Jugal Patel Hyder Reddy |
| Mid Yorkshire NHS Hospitals | Janahan Sarveswaran Lakshmanan Arunachalam Afaq Malik |
| Milton Keynes University Hospital NHS Foundation Trust | Luca Ponchietti Krystian Pawelec Yan Mei Goh Parveen Vitish-Sharma Ahmed Saad |
| Musgrove Park (Taunton) | Edward Smyth Amy Crees Louise Merker Nahida Bashir |
| Newport | Gethin Williams Jennifer Hayes Kelly Walters Rhiannon Harries Rahulpreet Singh |
| Ninewells Hospital (Dundee) | Nikola A Henderson Francesco M Polignano |
| Queen Alexandra Hospital (Portsmouth) | Ben Knight Louise Alder Alexandra Kenchington |

(Continued)

| | |
|---|---|
| Queen Elizabeth Hospital (Birmingham) | Yan Li Goh Ilaria Dicurzio Ewen Griffiths |
| Queen Elizabeth University Hospital (Glasgow) | Ahmed Alani Katrina Knight Patrick MacGoey Guat Shi Ng |
| Royal Albert Edward Infirmary (Wigan) | Naomi Mackenzie Ishaan Maitra |
| Royal Alexandra Hospital (Paisley) | Susan Moug Kelly Ong |
| Royal Berkshire Hospital (Reading) | Daniel McGrath Emanuele Gammeri Guillame Lafaurie |
| Royal Bolton Hospital | Gemma Faulkner Gabriele Di Benedetto Julia McGovern Bharathi Subramanian |
| Royal Devon and Exeter Hospital NHS | Sunil Kumar Narang Jennifer Nowers Neil J Smart Ian R Daniels |
| Royal Free Hospital (Hampstead site) | Massimo Varcada Tanzeela Gala |
| Royal Glamorgan Hospital | Julie Cornish Zoe Barber |
| Royal Infirmary of Edinburgh | Stephen O'Neill Richard McGregor Andrew G Robertson Simon Paterson-Brown |
| Royal Lancaster Infirmary | Thomas Raymond |
| Royal London Hospital | Mohamed A Thaha William J English Cillian T Forde Heidi Paine Alpa Morawala |
| Royal Preston Hospital | Ravindra Date Patrick Casey Thomas Bolton Xuan Gleaves Joshua Fasuyi |
| Royal Surrey County Hospital, Guildford | Sanja Durakovic Matt Dunstan Sophie Allen Angela Riga |

(Continued)

| | |
|--|--------------------------|
| Salford Royal NHS Foundation Trust | Jonathan Epstein |
| | Lyndsay Pearce |
| | Emily Gaines |
| | Anthony Howe |
| | Halima Choonara |
| Southmead (North Bristol) | Ffion Dewi |
| | Joanne Bennett |
| | Emile King |
| | Kathryn McCarthy |
| Swansea | Greg Taylor |
| | Dean Harris |
| | Hari Nageswaran |
| | Amy Stimpson |
| Tameside Hospital | Kamran Siddiqui |
| | Lay In Lim |
| University Hospital Crosshouse, Kilmarnock | Christopher Ray |
| | Laura Smith |
| | Gillian McColl |
| University Hospital of South Manchester | Mohammed Rahman |
| | Aaron Kler |
| | Abhi Sharma |
| | Kat Parmar |
| University Hospital of Wales (Cardiff) | Neil Patel |
| | Perry Crofts |
| | Claudio Baldari |
| | Rhys Thomas |
| | Michael Stechman |
| Western General Hospital (Edinburgh) | Roland Aldridge |
| | James O'Kelly |
| | Graeme Wilson |
| Weston general Hospital | Nicholas Gallegos |
| Whiston | Ramya Kalaiselvan |
| | Rajasundaram Rajaganesan |
| William Harvey Hospital (East Kent) | Aliya Mackenzie |
| | Prashant Naik |
| | Kaushiki Singh |
| | Harinath Gandraspulli |
| Wirral University Teaching Hospital | Jeremy Wilson |
| | Kate Hancorn |
| | Amir Khawaja |
| | Felix Nicholas |
| | Thomas Marks |
| Wrexham | Cameron Abbott |
| | Susan Chandler |

Authors' contributions

CB, BC, KP, JH and SM analysed and interpreted social media data. CB, BC, JL and SM were major contributors in writing the manuscript. BC was responsible for statistical analysis. All authors read and approved the final manuscript.

Funding

The ELF Study was funded by the Bowel Disease Research Foundation (BDRF). The funding body had no influence on the design of the study, data collection and analysis or manuscript submission.

Availability of data and materials

The datasets used and/or analysed during the current study are available via Twitter Analytics and Twitonomy, or from the corresponding author on reasonable request.

Ethics approval and consent to participate

Ethical approval was obtained from National Health Service Research Ethics Committee (Black Country Research Committee: November 2016; 16/WM/0500). The study was registered centrally with the Health Research Authority (England), the NHS Research Scotland Permissions Co-ordinating Centre (Scotland), and the Health and Care Research Permissions Service (Wales). The ELF study was registered online at www.clinicaltrials.gov (NCT02952430). The ELF Study did not require consent for participation as it was a prospective audit of practice, with no direct clinical impact upon participants.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of General Surgery, Royal Alexandra Hospital, Paisley PA2 9PN, UK. ²Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK. ³Department of General Surgery, Blackpool Victoria Hospital, Blackpool, UK. ⁴Department of Population Medicine, Cardiff University, Cardiff, UK. ⁵Department of General Surgery, Manchester Royal Infirmary, Manchester, UK.

Received: 27 February 2020 Accepted: 1 July 2020

Published online: 28 July 2020

References

1. Statista. Number of social network users worldwide from 2010 to 2021. <https://www.statista.com/statistics/278414/number-of-worldwide-social-network-users/> (accessed 21/1/20).
2. Macdonald JJ, Bisset C, Coleman MG, Speake D, Brady RRW. Contemporary use of social media by consultant colorectal surgeons. *Colorectal Dis.* 2015; 17(2):165–71.
3. Pearson D, Bond MC, Kegg J, et al. Evaluation of social media use by emergency medicine residents and faculty. *West J Emerg Med.* 2015;16: 715–20.
4. Steele SR, Arshad S, Bush R, Dasani S, Cologne K, Bleier JIS, Raphaeli T, Kelz RR. Social media is a necessary component of surgical practice. *Surgery.* 2015;158(3):857–62.
5. Wexner SD, Petrucci AM, Brady RR, Ennis-O'Connor M, Fitzgerald JE, Mayol J. Social media in colorectal surgery. *Colorectal Dis.* 2017;19(2):105–14.
6. Kapp JM, Peters C, Oliver DP. Research recruitment using Facebook advertising: big potential, big challenges. *J Cancer Educ.* 2013;28:134–7. <https://doi.org/10.1007/s13187-012-0443-z>.
7. Sayers AE, Lee MJ, Smart N, Fearnhead NS, on behalf of the NASBO Steering Group. Optimising collaborator recruitment and maintaining engagement via social media during large multicentre studies: lessons learned from the National Audit of Small Bowel Obstruction (NASBO). *Colorectal Dis.* 2018; 20(12):1142–50.

8. Cochrane AR, Macdonald JJ, Brady RRW. Social media use among United Kingdom vascular surgeons: a cross-sectional study. *Ann Vasc Surg.* 2016;33:252–7.
9. Mabvuure NT, Rodrigues J, Klimach S, et al. A cross sectional study of the presence of United Kingdom (UK) plastic surgeons on social media. *J Plast Reconstruct Aesthet Surg.* 2014;67:362–7.
10. Medical Protection Society (MPS) press release (14 June 2012). Accessed (25/09/2014) via: www.medicalprotection.org/Default.aspx?DN=43fb99ca-937e-4c30-86ae-393054e3e99f.
11. Parmar KL, Pearce L, Farrell I, et al. Influence of frailty in older patients undergoing emergency laparotomy: a UK-based observational study. *BMJ Open.* 2017;7:e017928. <https://doi.org/10.1136/bmjopen-2017-017928>.
12. <https://sysomos.com/inside-twitter/twitter-statistics/> (accessed 28/2/19).
13. Ashley C, Tuten T. Creative strategies in social media marketing: an exploratory study of branded social content and consumer engagement. *Psychol Mark.* 2014;32(1):15–27.
14. Sheehan K, Morrison D. The creativity challenge: media confluence and its effects on the evolving advertising industry. *J Interact Advert.* 2009;9:40–3.
15. Doorn JV, Lemon K, Mittal V, Nass S, Pick D, Pimer P, et al. Customer engagement behavior: theoretical foundations and research directions. *J Serv Res.* 2010;13:253–66.
16. Schmitt B. The consumer psychology of brands. *J Consum Psychol.* 2012;22:7–17.
17. Nowotarski M. Do not steal my avatar! Challenges of social network patents [internet]. IP Watchdog; January 23, 2011. <http://www.ipwatchdog.com/2011/01/23/dont-steal-my-avatar-challenges-of-social-networking-patents/id=14531/> (accessed 28/2/19).
18. Indes JE, Gates L, Mitchell EL, Muhs BE. Social media in vascular surgery. *J Vasc Surg.* 2013;57(4):1159–62.
19. <https://www.gmc-uk.org/ethical-guidance/ethical-guidance-for-doctors/doctors-use-of-social-media/doctors-use-of-social-media> (accessed 28/2/19).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

