Northumbria Research Link

Citation: Sangiorgi, Francesca, Bijl, Peter, Passchier, Sandra, Salzmann, Ulrich, Schouten, Stefan, Pross, Jorg, Escutia, Carlotta and Brinkhuis, Henk (2015) Middle Miocene environmental and climatic evolution at the Wilkes Land margin, East Antarctica. In: EGU General Assembly 2015, 12-17 April 2015, Vienna.

URL:

This version was downloaded from Northumbria Research Link: http://nrl.northumbria.ac.uk/21498/

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: http://nrl.northumbria.ac.uk/policies.html

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

www.northumbria.ac.uk/nrl



Geophysical Research Abstracts Vol. 17, EGU2015-15194, 2015 EGU General Assembly 2015 © Author(s) 2015. CC Attribution 3.0 License.



Middle Miocene environmental and climatic evolution at the Wilkes Land margin, East Antarctica

Francesca Sangiorgi (1), Peter Bijl (1), Sandra Passchier (2), Ulrich Salzmann (3), Stefan Schouten (4), Jörg Pross (5), Carlota Escutia (6), Henk Brinkhuis (1,4)

(1) Marine Palynology, Department of Earth Sciences, Faculty of Geology, Laboratory of Palaeobotany and Palynology, Utrecht University, Budapestlaan 4, 3584CD Utrecht, The Netherlands, (2) Department of Earth and Environmental Studies, Montclair State University, 1 Normal Ave, 252 Mallory Hall, Montclair, NJ 07043, (3) Geography and Environment, School of Built and Natural Environment, Northumbria University, Ellison Building Newcastle upon Tyne, NE1 8ST, UK, (4) NIOZ Royal Netherlands Institute for Sea Research, Landsdiep 4, 1797 SZ 't Horntje (Texel), The Netherlands, (5) Institute of Earth Sciences, University of Heidelberg, Im Neuenheimer Feld 234, 69120 Heidelberg, Germany, (6) Instituto Andaluz de Ciencias de la Tierra, CSIC-Universidad de Granada, Granada, Spain

Integrated Ocean Drilling Program (IODP) Expedition 318 successfully drilled a Middle Miocene ($\sim 17-12.5$ Ma) record from the Wilkes Land Margin at Site U1356A (63°18.6138'S, 135°59.9376'E), located at the transition between the continental rise and the abyssal plain at 4003 mbsl.

We present a multiproxy palynological (dinoflagellate cyst, pollen and spores), sedimentological and organic geochemical (TEX86, MBT/CBT) study, which unravels the environmental and climate variability across the Miocene Climatic Optimum (MCO, ~17-15 Ma) and the Mid Miocene Climate Transition (MMCT).

Several independent lines of evidence suggest a relatively warm climate during the MCO. Dinocyst and pollen assemblage diversity at the MCO is unprecedented for a Neogene Antarctic record and indicates a temperate, sea ice-free marine environment, with woody sub-antarctic vegetation with elements of forest/shrub tundra and peat lands along the coast. These results are further confirmed by relatively warm TEX86-derived Sea Surface Temperatures and mild MBT-derived continental temperatures, and by the absence of glacially derived deposits and very few ice-rafted clasts. A generally colder but highly dynamic environment is suggested for the interval 15-12.5 Ma.