

*Note di Matematica* **22**, n. 2, 2003, 83–92.

# Topological diagonalizations and Hausdorff dimension

**Tomasz Weiss**

*Institute of Mathematics, Akademia Podlaska  
08-119 Siedlce, Poland  
tomaszweiss@go2.pl*

**Boaz Tsaban**

*Einstein Institute of Mathematics, Hebrew University of Jerusalem,  
Givat Ram, Jerusalem 91904, Israel  
tsaban@math.huji.ac.il, <http://www.cs.biu.ac.il/~tsaban>*

Received: 13/01/2003; accepted: 03/09/2003.

**Abstract.** The Hausdorff dimension of a product  $X \times Y$  can be strictly greater than that of  $Y$ , even when the Hausdorff dimension of  $X$  is zero. But when  $X$  is countable, the Hausdorff dimensions of  $Y$  and  $X \times Y$  are the same. Diagonalizations of covers define a natural hierarchy of properties which are weaker than “being countable” and stronger than “having Hausdorff dimension zero”. Fremlin asked whether it is enough for  $X$  to have the strongest property in this hierarchy (namely, being a  $\gamma$ -set) in order to assure that the Hausdorff dimensions of  $Y$  and  $X \times Y$  are the same.

We give a negative answer: Assuming the Continuum Hypothesis, there exists a  $\gamma$ -set  $X \subseteq \mathbb{R}$  and a set  $Y \subseteq \mathbb{R}$  with Hausdorff dimension zero, such that the Hausdorff dimension of  $X + Y$  (a Lipschitz image of  $X \times Y$ ) is maximal, that is, 1. However, we show that for the notion of a *strong*  $\gamma$ -set the answer is positive. Some related problems remain open.

**Keywords:** Hausdorff dimension, Gerlits-Nagy  $\gamma$  property, Galvin-Miller strong  $\gamma$  property.

**MSC 2000 classification:** primary: 03E75; secondary: 37F20, 26A03.

## Introduction

The Hausdorff dimension of a subset of  $\mathbb{R}^k$  is a derivative of the notion of Hausdorff *measures* [4]. However, for our purposes it will be more convenient to use the following equivalent definition. Denote the diameter of a subset  $A$  of  $\mathbb{R}^k$  by  $\text{diam}(A)$ . The *Hausdorff dimension* of a set  $X \subseteq \mathbb{R}^k$ ,  $\dim(X)$ , is the infimum of all positive  $\delta$  such that for each positive  $\epsilon$  there exists a cover  $\{I_n\}_{n \in \mathbb{N}}$  of  $X$  with

$$\sum_{n \in \mathbb{N}} \text{diam}(I_n)^\delta < \epsilon.$$

From the many properties of Hausdorff dimension, we will need the following easy ones.

**1 Lemma.**

- (1) If  $X \subseteq Y \subseteq \mathbb{R}^k$ , then  $\dim(X) \leq \dim(Y)$ .
- (2) Assume that  $X_1, X_2, \dots$  are subsets of  $\mathbb{R}^k$  such that  $\dim(X_n) = \delta$  for each  $n$ . Then  $\dim(\bigcup_n X_n) = \delta$ .
- (3) Assume that  $X \subseteq \mathbb{R}^k$  and  $Y \subseteq \mathbb{R}^m$  is such that there exists a Lipschitz surjection  $\phi : X \rightarrow Y$ . Then  $\dim(X) \geq \dim(Y)$ .
- (4) For each  $X \subseteq \mathbb{R}^k$  and  $Y \subseteq \mathbb{R}^m$ ,  $\dim(X \times Y) \geq \dim(X) + \dim(Y)$ .

Equality need not hold in item (4) of the last lemma. In particular, one can construct a set  $X$  with Hausdorff dimension zero and a set  $Y$  such that  $\dim(X \times Y) > \dim(Y)$ . On the other hand, when  $X$  is countable,  $X \times Y$  is a union of countably many copies of  $Y$ , and therefore

$$\dim(X \times Y) = \dim(Y). \quad (1)$$

Having Hausdorff dimension zero can be thought of as a notion of smallness. Being countable is another notion of smallness, and we know that the first notion is not enough restrictive in order to have Equation 1 hold, but the second is.

Notions of smallness for sets of real numbers have a long history and many applications – see, e.g., [11]. We will consider some notions which are weaker than being countable and stronger than having Hausdorff dimension zero.

According to Borel [3], a set  $X \subseteq \mathbb{R}^k$  has *strong measure zero* if for each sequence of positive reals  $\{\epsilon_n\}_{n \in \mathbb{N}}$ , there exists a cover  $\{I_n\}_{n \in \mathbb{N}}$  of  $X$  such that  $\text{diam}(I_n) < \epsilon_n$  for all  $n$ . Clearly strong measure zero implies Hausdorff dimension zero. It does not require any special assumptions in order to see that the converse is false. A perfect set can be mapped onto the unit interval by a uniformly continuous function and therefore cannot have strong measure zero.

**2 Proposition (folklore).** *There exists a perfect set of reals  $X$  with Hausdorff dimension zero.*

PROOF. For  $0 < \lambda < 1$ , denote by  $C(\lambda)$  the Cantor set obtained by starting with the unit interval, and at each step removing from the middle of each interval a subinterval of size  $\lambda$  times the size of the interval (So that  $C(1/3)$  is the canonical middle-third Cantor set, which has Hausdorff dimension  $\log 2 / \log 3$ .) It is easy to see that if  $\lambda_n \nearrow 1$ , then  $\dim(C(\lambda_n)) \searrow 0$ .

Thus, define a special Cantor set  $C(\{\lambda_n\}_{n \in \mathbb{N}})$  by starting with the unit interval, and at step  $n$  removing from the middle of each interval a subinterval of size  $\lambda_n$  times the size of the interval. For each  $n$ ,  $C(\{\lambda_n\}_{n \in \mathbb{N}})$  is contained in a union of  $2^n$  (shrunk) copies of  $C(\lambda_n)$ , and therefore  $\dim(C(\{\lambda_n\}_{n \in \mathbb{N}})) \leq \dim(C(\lambda_n))$ .  $\square$

As every countable set has strong measure zero, the latter notion can be thought of an “approximation” of countability. In fact, Borel conjectured in [3] that every strong measure zero set is countable, and it turns out that the usual axioms of mathematics (ZFC) are not strong enough to prove or disprove this conjecture: Assuming the Continuum Hypothesis there exists an uncountable strong measure zero set (namely, a Luzin set), but Laver [10] proved that one cannot prove the existence of such an object from the usual axioms of mathematics.

The property of strong measure zero (which depends on the metric) has a natural topological counterpart. A topological space  $X$  has *Rothberger’s property  $C''$*  [13] if for each sequence  $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$  of covers of  $X$  there is a sequence  $\{U_n\}_{n \in \mathbb{N}}$  such that for each  $n$   $U_n \in \mathcal{U}_n$ , and  $\{U_n\}_{n \in \mathbb{N}}$  is a cover of  $X$ . Using Scheepers’ notation [15], this property is a particular instance of the following selection hypothesis (where  $\mathfrak{U}$  and  $\mathfrak{V}$  are any collections of covers of  $X$ ):

$S_1(\mathfrak{U}, \mathfrak{V})$ : For each sequence  $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$  of members of  $\mathfrak{U}$ , there is a sequence  $\{U_n\}_{n \in \mathbb{N}}$  such that  $U_n \in \mathcal{U}_n$  for each  $n$ , and  $\{U_n\}_{n \in \mathbb{N}} \in \mathfrak{V}$ .

Let  $\mathcal{O}$  denote the collection of all open covers of  $X$ . Then the property considered by Rothberger is  $S_1(\mathcal{O}, \mathcal{O})$ . Fremlin and Miller [5] proved that a set  $X \subseteq \mathbb{R}^k$  satisfies  $S_1(\mathcal{O}, \mathcal{O})$  if, and only if,  $X$  has strong measure zero with respect to each metric which generates the standard topology on  $\mathbb{R}^k$ .

But even Rothberger’s property for  $X$  is not strong enough to have Equation 1 hold: It is well-known that every Luzin set satisfies Rothberger’s property (and, in particular, has Hausdorff dimension zero).

**3 Lemma.** *The mapping  $(x, y) \mapsto x + y$  from  $\mathbb{R}^2$  to  $\mathbb{R}$  is Lipschitz.*

PROOF. Observe that for nonnegative reals  $a$  and  $b$ ,  $(a - b)^2 \geq 0$  and therefore  $a^2 + b^2 \geq 2ab$ . Consequently,

$$a + b = \sqrt{a^2 + 2ab + b^2} \leq \sqrt{2(a^2 + b^2)} = \sqrt{2}\sqrt{a^2 + b^2}.$$

Thus,

$$|(x_1 + y_1) - (x_2 + y_2)| \leq \sqrt{2}\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} \text{ for all } (x_1, y_1), (x_2, y_2) \in \mathbb{R}^2.$$

$\square$

Assuming the Continuum Hypothesis, there exists a Luzin set  $L \subseteq \mathbb{R}$  such that  $L + L$ , a Lipschitz image of  $L \times L$ , is equal to  $\mathbb{R}$  [9].

We therefore consider some stronger properties. An open cover  $\mathcal{U}$  of  $X$  is an  $\omega$ -cover of  $X$  if each finite subset of  $X$  is contained in some member of the cover, but  $X$  is not contained in any member of  $\mathcal{U}$ .

$\mathcal{U}$  is a  $\gamma$ -cover of  $X$  if it is infinite, and each element of  $X$  belongs to all but finitely many members of  $\mathcal{U}$ . Let  $\Omega$  and  $\Gamma$  denote the collections of open  $\omega$ -covers and  $\gamma$ -covers of  $X$ , respectively. Then  $\Gamma \subseteq \Omega \subseteq \mathcal{O}$ , and these three classes of covers introduce 9 properties of the form  $S_1(\mathfrak{U}, \mathfrak{V})$ . If we remove the trivial ones and check for equivalences [9, 20], then it turns out that only six of these properties are really distinct, and only three of them imply Hausdorff dimension zero:

$$S_1(\Omega, \Gamma) \rightarrow S_1(\Omega, \Omega) \rightarrow S_1(\mathcal{O}, \mathcal{O}).$$

The properties  $S_1(\Omega, \Gamma)$  and  $S_1(\Omega, \Omega)$  were also studied before.  $S_1(\Omega, \Omega)$  was studied by Sakai [14], and  $S_1(\Omega, \Gamma)$  was studied by Gerlits and Nagy in [8]: A topological space  $X$  is a  $\gamma$ -set if each  $\omega$ -cover of  $X$  contains a  $\gamma$ -cover of  $X$ . Gerlits and Nagy proved that  $X$  is a  $\gamma$ -set if, and only if,  $X$  satisfies  $S_1(\Omega, \Gamma)$ . It is not difficult to see that every countable space is a  $\gamma$ -set. But this property is not trivial: Assuming the Continuum Hypothesis, there exist uncountable  $\gamma$ -sets [7].

$S_1(\Omega, \Omega)$  is closed under taking finite powers [9], thus the Luzin set we used to see that Equation 1 need not hold when  $X$  satisfies  $S_1(\mathcal{O}, \mathcal{O})$  does not rule out that possibility that this Equation holds when  $X$  satisfies  $S_1(\Omega, \Omega)$ . However, in [2] it is shown that assuming the Continuum Hypothesis, there exist Luzin sets  $L_0$  and  $L_1$  satisfying  $S_1(\Omega, \Omega)$ , such that  $L_0 + L_1 = \mathbb{R}$ . Thus, the only remaining candidate for a nontrivial property of  $X$  where Equation 1 holds is  $S_1(\Omega, \Gamma)$  ( $\gamma$ -sets). Fremlin (personal communication) asked whether Equation 1 is indeed provable in this case. We give a negative answer, but show that for a yet stricter (but nontrivial) property which was considered in the literature, the answer is positive.

The notion of a strong  $\gamma$ -set was introduced in [7]. However, we will adopt the following simple characterization from [20] as our formal definition. Assume that  $\{\mathfrak{U}_n\}_{n \in \mathbb{N}}$  is a sequence of collections of covers of a space  $X$ , and that  $\mathfrak{V}$  is a collection of covers of  $X$ . Define the following selection hypothesis.

$S_1(\{\mathfrak{U}_n\}_{n \in \mathbb{N}}, \mathfrak{V})$ : For each sequence  $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$  where  $\mathcal{U}_n \in \mathfrak{U}_n$  for each  $n$ , there is a sequence  $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$  such that  $\mathcal{U}_n \in \mathfrak{U}_n$  for each  $n$ , and  $\{\mathcal{U}_n\}_{n \in \mathbb{N}} \in \mathfrak{V}$ .

A cover  $\mathcal{U}$  of a space  $X$  is an  $n$ -cover if each  $n$ -element subset of  $X$  is contained in some member of  $\mathcal{U}$ . For each  $n$  denote by  $\mathcal{O}_n$  the collection of all open  $n$ -covers of a space  $X$ . Then  $X$  is a *strong  $\gamma$ -set* if  $X$  satisfies  $S_1(\{\mathcal{O}_n\}_{n \in \mathbb{N}}, \Gamma)$ .

In most cases  $S_1(\{\mathcal{O}_n\}_{n \in \mathbb{N}}, \mathfrak{V})$  is equivalent to  $S_1(\Omega, \mathfrak{V})$  [20], but not in the case  $\mathfrak{V} = \Gamma$ : It is known that for a strong  $\gamma$ -set  $G \subseteq \{0, 1\}^{\mathbb{N}}$  and each  $A \subseteq \{0, 1\}^{\mathbb{N}}$  of measure zero,  $G \oplus A$  has measure zero too [7]; this can be contrasted with Theorem 5 below. In Section 2 we show that Equation 1 is provable in the case that  $X$  is a strong  $\gamma$ -set, establishing another difference between the notions

of  $\gamma$ -sets and strong  $\gamma$ -sets, and giving a positive answer to Fremlin's question under a stronger assumption on  $X$ .

## 1 The product of a $\gamma$ -set and a set of Hausdorff dimension zero

**4 Theorem.** *Assuming the Continuum Hypothesis (or just  $\mathfrak{p} = \mathfrak{c}$ ), there exist a  $\gamma$ -set  $X \subseteq \mathbb{R}$  and a set  $Y \subseteq \mathbb{R}$  with Hausdorff dimension zero such that the Hausdorff dimension of the algebraic sum*

$$X + Y = \{x + y : x \in X, y \in Y\}$$

(a Lipschitz image of  $X \times Y$  in  $\mathbb{R}$ ) is 1. In particular,  $\dim(X \times Y) \geq 1$ .

Our theorem will follow from the following related theorem. This theorem involves the Cantor space  $\{0, 1\}^{\mathbb{N}}$  of infinite binary sequences. The Cantor space is equipped with the product topology and with the product measure.

**5 Theorem (Bartoszyński and Reclaw [1]).** *Assume the Continuum Hypothesis (or just  $\mathfrak{p} = \mathfrak{c}$ ). Fix an increasing sequence  $\{k_n\}_{n \in \mathbb{N}}$  of natural numbers, and for each  $n$  define*

$$A_n = \{f \in \{0, 1\}^{\mathbb{N}} : f \upharpoonright [k_n, k_{n+1}) \equiv 0\}.$$

If the set

$$A = \bigcap_{m \in \mathbb{N}} \bigcup_{n \geq m} A_n$$

has measure zero, then there exists a  $\gamma$ -set  $G \subseteq \{0, 1\}^{\mathbb{N}}$  such that the algebraic sum  $G \oplus A$  is equal to  $\{0, 1\}^{\mathbb{N}}$  (where  $\oplus$  denotes the modulo 2 coordinate-wise addition).

Observe that the assumption in Theorem 5 holds whenever  $\sum_n 2^{-(k_{n+1}-k_n)}$  converges.

**6 Lemma.** *There exists an increasing sequence of natural numbers  $\{k_n\}_{n \in \mathbb{N}}$  such that  $\sum_n 2^{-(k_{n+1}-k_n)}$  converges, and such that for the sequence  $\{B_n\}_{n \in \mathbb{N}}$  defined by*

$$B_n = \left\{ \sum_{i \in \mathbb{N}} \frac{f(i)}{2^{i+1}} : f \in \{-1, 0, 1\}^{\mathbb{N}} \text{ and } f \upharpoonright [k_n, k_{n+1}) \equiv 0 \right\}$$

for each  $n$ , the set

$$Y = \bigcap_{m \in \omega} \bigcup_{n \geq m} B_n$$

has Hausdorff dimension zero.

PROOF. Fix a sequence  $p_n$  of positive reals which converges to 0. Let  $k_0 = 0$ . Given  $k_n$  find  $k_{n+1}$  satisfying

$$3^{k_n} \cdot \frac{1}{2^{p_n(k_{n+1}-2)}} \leq \frac{1}{2^n}.$$

Clearly, every  $B_n$  is contained in a union of  $3^{k_n}$  intervals such that each of the intervals has diameter  $1/2^{k_{n+1}-2}$ . For each positive  $\delta$  and  $\epsilon$ , choose  $m$  such that  $\sum_{n \geq m} 1/2^n < \epsilon$  and such that  $p_n < \delta$  for all  $n \geq m$ . Now,  $Y$  is a subset of  $\bigcup_{n \geq m} B_n$ , and

$$\sum_{n \geq m} 3^{k_n} \left( \frac{1}{2^{k_{n+1}-2}} \right)^\delta < \sum_{n \geq m} 3^{k_n} \left( \frac{1}{2^{k_{n+1}-2}} \right)^{p_n} < \sum_{n \geq m} \frac{1}{2^n} < \epsilon.$$

Thus, the Hausdorff dimension of  $Y$  is zero.  $\square$

The following lemma concludes the proof of Theorem 4.

**7 Lemma.** *There exists a  $\gamma$ -set  $X \subseteq \mathbb{R}$  and a set  $Y \subseteq \mathbb{R}$  with Hausdorff dimension zero such that  $X + Y = \mathbb{R}$ . In particular,  $\dim(X + Y) = 1$ .*

PROOF. Choose a sequence  $\{k_n\}_{n \in \mathbb{N}}$  and a set  $Y$  as in Lemma 6. Then  $\sum_n 2^{-(k_{n+1}-k_n)}$  converges, and the corresponding set  $A$  defined in Theorem 5 has measure zero. Thus, there exists a  $\gamma$ -set  $G$  such that  $G \oplus A = \{0, 1\}^{\mathbb{N}}$ . Define  $\Phi : \{0, 1\}^{\mathbb{N}} \rightarrow \mathbb{R}$  by

$$\Phi(f) = \sum_{i \in \mathbb{N}} \frac{f(i)}{2^{i+1}}.$$

As  $\Phi$  is continuous,  $X = \Phi[G]$  is a  $\gamma$ -set of reals. Assume that  $z$  is a member of the interval  $[0, 1]$ , let  $f \in \{0, 1\}^{\mathbb{N}}$  be such that  $z = \sum_i f(i)/2^{i+1}$ . Then  $f = g \oplus a$  for appropriate  $g \in G$  and  $a \in A$ . Define  $h \in \{-1, 0, 1\}^{\mathbb{N}}$  by  $h(i) = f(i) - g(i)$ . For infinitely many  $n$ ,  $a \upharpoonright [k_n, k_{n+1}) \equiv 0$  and therefore  $f \upharpoonright [k_n, k_{n+1}) \equiv g \upharpoonright [k_n, k_{n+1})$ , that is,  $h \upharpoonright [k_n, k_{n+1}) \equiv 0$  for infinitely many  $n$ . Thus,  $y = \sum_i h(i)/2^{i+1} \in Y$ , and for  $x = \Phi(g)$ ,

$$x + y = \sum_{i \in \mathbb{N}} \frac{g(i)}{2^{i+1}} + \sum_{i \in \mathbb{N}} \frac{h(i)}{2^{i+1}} = \sum_{i \in \mathbb{N}} \frac{g(i) + h(i)}{2^{i+1}} = \sum_{i \in \mathbb{N}} \frac{f(i)}{2^{i+1}} = z.$$

This shows that  $[0, 1] \subseteq X + Y$ . Consequently,  $X + (Y + \mathbb{Q}) = (X + Y) + \mathbb{Q} = \mathbb{R}$ . Now, observe that  $Y + \mathbb{Q}$  has Hausdorff dimension zero since  $Y$  has.  $\square$

## 2 The product of a strong $\gamma$ -set and a set of Hausdorff dimension zero

**8 Theorem.** *Assume that  $X \subseteq \mathbb{R}^k$  is a strong  $\gamma$ -set. Then for each  $Y \subseteq \mathbb{R}^l$ ,  $\dim(X \times Y) = \dim(Y)$ .*

PROOF. The proof for this is similar to that of Theorem 7 in [7]. It is enough to show that  $\dim(X \times Y) \leq \dim(Y)$ .

**9 Lemma.** *Assume that  $Y \subseteq \mathbb{R}^l$  is such that  $\dim(Y) < \delta$ . Then for each positive  $\epsilon$  there exists a large cover  $\{I_n\}_{n \in \mathbb{N}}$  of  $Y$  (i.e., such that each  $y \in Y$  is a member of infinitely many sets  $I_n$ ) such that  $\sum_n \text{diam}(I_n)^\delta < \epsilon$ .*

PROOF. For each  $m$  choose a cover  $\{I_n^m\}_{n \in \mathbb{N}}$  of  $Y$  such that  $\sum_n \text{diam}(I_n^m)^\delta < \epsilon/2^m$ . Then  $\{I_n^m : m, n \in \mathbb{N}\}$  is a large cover of  $Y$ , and  $\sum_{m,n} \text{diam}(I_n^m)^\delta < \sum_n \epsilon/2^m = \epsilon$ .  $\square$

**10 Lemma.** *Assume that  $Y \subseteq \mathbb{R}^l$  is such that  $\dim(Y) < \delta$ . Then for each sequence  $\{\epsilon_n\}_{n \in \mathbb{N}}$  of positive reals there exists a large cover  $\{A_n\}_{n \in \mathbb{N}}$  of  $Y$  such that for each  $n$   $A_n$  is a union of finitely many sets,  $I_1^n, \dots, I_{m_n}^n$ , such that  $\sum_j \text{diam}(I_j^n)^\delta < \epsilon_n$ .*

PROOF. Assume that  $\{\epsilon_n\}_{n \in \mathbb{N}}$  is a sequence of positive reals. By Lemma 9, there exists a large cover  $\{I_n\}_{n \in \mathbb{N}}$  of  $Y$  such that  $\sum_n \text{diam}(I_n)^\delta < \epsilon_1$ . For each  $n$  let  $k_n = \min\{m : \sum_{j \geq m} \text{diam}(I_j)^\delta < \epsilon_n\}$ . Take

$$A_n = \bigcup_{j=k_n}^{k_{n+1}-1} I_j.$$

 $\square$ 

Fix  $\delta > \dim(Y)$  and  $\epsilon > 0$ . Choose a sequence  $\{\epsilon_n\}_{n \in \mathbb{N}}$  of positive reals such that  $\sum_n 2n\epsilon_n < \epsilon$ , and use Lemma 10 to get the corresponding large cover  $\{A_n\}_{n \in \mathbb{N}}$ .

For each  $n$  we define an  $n$ -cover  $\mathcal{U}_n$  of  $X$  as follows. Let  $F$  be an  $n$ -element subset of  $X$ . For each  $x \in F$ , find an open interval  $I_x$  such that  $x \in I_x$  and

$$\sum_{j=1}^{m_n} \text{diam}(I_x \times I_j^n)^\delta < 2\epsilon_n.$$

Let  $U_F = \bigcup_{x \in F} I_x$ . Set

$$\mathcal{U}_n = \{U_F : F \text{ is an } n\text{-element subset of } X\}.$$

As  $X$  is a strong  $\gamma$ -set, there exist elements  $U_{F_n} \in \mathcal{U}_n$ ,  $n \in \mathbb{N}$ , such that  $\{U_{F_n}\}_{n \in \mathbb{N}}$  is a  $\gamma$ -cover of  $X$ . Consequently,

$$X \times Y \subseteq \bigcup_{n \in \mathbb{N}} (U_{F_n} \times A_n) \subseteq \bigcup_{n \in \mathbb{N}} \bigcup_{x \in F_n} \bigcup_{j=1}^{m_n} I_x \times I_j^n$$

and

$$\sum_{n \in \mathbb{N}} \sum_{x \in F_n} \sum_{j=1}^{m_n} \text{diam}(I_x \times I_j^n)^\delta < \sum_n n \cdot 2\epsilon_n < \epsilon.$$

$\square$  *QED*

### 3 Open problems

There are ways to strengthen the notion of  $\gamma$ -sets other than moving to strong  $\gamma$ -sets. Let  $\mathcal{B}_\Omega$  and  $\mathcal{B}_\Gamma$  denote the collections of *countable Borel*  $\omega$ -covers and  $\gamma$ -covers of  $X$ , respectively. As every open  $\omega$ -cover of a set of reals contains a countable  $\omega$ -subcover [9], we have that  $\Omega \subseteq \mathcal{B}_\Omega$  and therefore  $\mathsf{S}_1(\mathcal{B}_\Omega, \mathcal{B}_\Gamma)$  implies  $\mathsf{S}_1(\Omega, \Gamma)$ . The converse is not true [17].

**11 Problem.** Assume that  $X \subseteq \mathbb{R}$  satisfies  $\mathsf{S}_1(\mathcal{B}_\Omega, \mathcal{B}_\Gamma)$ . Is it true that for each  $Y \subseteq \mathbb{R}$ ,  $\dim(X \times Y) = \dim(Y)$ ?

We conjecture that assuming the Continuum Hypothesis, the answer to this problem is negative. We therefore introduce the following problem. For infinite sets of natural numbers  $A, B$ , we write  $A \subseteq^* B$  if  $A \setminus B$  is finite. Assume that  $\mathcal{F}$  is a family of infinite sets of natural numbers. A set  $P$  is a *pseudointersection* of  $\mathcal{F}$  if it is infinite, and for each  $B \in \mathcal{F}$ ,  $A \subseteq^* B$ .  $\mathcal{F}$  is *centered* if each finite subcollection of  $\mathcal{F}$  has a pseudointersection. Let  $\mathfrak{p}$  denote the minimal cardinality of a centered family which does not have a pseudointersection. In [17] it is proved that  $\mathfrak{p}$  is also the minimal cardinality of a set of reals which does not satisfy  $\mathsf{S}_1(\mathcal{B}_\Omega, \mathcal{B}_\Gamma)$ .

**12 Problem.** Assume that the cardinality of  $X$  is smaller than  $\mathfrak{p}$ . Is it true that for each  $Y \subseteq \mathbb{R}$ ,  $\dim(X \times Y) = \dim(Y)$ ?

Another interesting open problem involves the following notion [18, 19]. A cover  $\mathcal{U}$  of  $X$  is a  $\tau$ -cover of  $X$  if it is a large cover, and for each  $x, y \in X$ , one of the sets  $\{U \in \mathcal{U} : x \in U \text{ and } y \notin U\}$  or  $\{U \in \mathcal{U} : y \in U \text{ and } x \notin U\}$  is finite. Let  $\mathsf{T}$  denote the collection of open  $\tau$ -covers of  $X$ . Then  $\Gamma \subseteq \mathsf{T} \subseteq \Omega$ , therefore  $\mathsf{S}_1(\{\mathcal{O}_n\}_{n \in \mathbb{N}}, \Gamma)$  implies  $\mathsf{S}_1(\{\mathcal{O}_n\}_{n \in \mathbb{N}}, \mathsf{T})$ .

**13 Problem.** Assume that  $X \subseteq \mathbb{R}$  satisfies  $\mathsf{S}_1(\{\mathcal{O}_n\}_{n \in \mathbb{N}}, \mathsf{T})$ . Is it true that for each  $Y \subseteq \mathbb{R}$ ,  $\dim(X \times Y) = \dim(Y)$ ?

It is conjectured that  $\mathsf{S}_1(\{\mathcal{O}_n\}_{n \in \mathbb{N}}, \mathsf{T})$  is strictly stronger than  $\mathsf{S}_1(\Omega, \mathsf{T})$  [20]. If this conjecture is false, then the results in this paper imply a negative answer to Problem 13.

Another type of problems is the following: We have seen that the assumption that  $X$  is a  $\gamma$ -set and  $Y$  has Hausdorff dimension zero is not enough in order to prove that  $X \times Y$  has Hausdorff dimension zero. We also saw that if  $X$  satisfies a



stronger property (strong  $\gamma$ -set), then  $\dim(X \times Y) = \dim(Y)$  for all  $Y$ . Another approach to get a positive answer would be to strengthen the assumption on  $Y$  rather than  $X$ .

If we assume that  $Y$  has strong measure zero, then a positive answer follows from a result of Scheepers [16] (see also [21]), asserting that if  $X$  is a strong measure zero metric space which also has the Hurewicz property, then for each strong measure zero metric space  $Y$ ,  $X \times Y$  has strong measure zero. Indeed, if  $X$  is a  $\gamma$ -set then it has the required properties.

Finally, the following question of Krawczyk remains open.

**14 Problem.** Is it consistent (relative to ZFC) that there are uncountable  $\gamma$ -sets but for each  $\gamma$ -set  $X$  and each set  $Y$ ,  $\dim(X \times Y) = \dim(Y)$ ?

## References

- [1] T. BARTOSZYŃSKI, I. RECLAW: *Not every  $\gamma$ -set is strongly meager*, Contemporary Mathematics **192** (1996), 25–29.
- [2] T. BARTOSZYŃSKI, S. SHELAH, B. TSABAN: *Additivity properties of topological diagonalizations*, Journal of Symbolic Logic, **68**, (2003), 1254–1260. (Full version: <http://arxiv.org/abs/math.LO/0112262>)
- [3] É. BOREL: *Sur la classification des ensembles de mesure nulle*, Bulletin de la Société Mathématique de France **47** (1919), 97–125.
- [4] K. FALCONER: *The geometry of fractal sets*, Cambridge University Press, 1990.
- [5] D.H. FREMLIN, A.W. MILLER: *On some properties of Hurewicz, Menger and Rothberger*, Fundamenta Mathematica **129** (1988), 17–33.
- [6] F. GALVIN, J. MYCIELSKI, R. SOLOVAY: *Strong measure zero sets*, Notices of the American Mathematical Society (1973), A–280.
- [7] F. GALVIN, A.W. MILLER:  *$\gamma$ -sets and other singular sets of real numbers*, Topology and its Applications **17** (1984), 145–155.
- [8] J. GERLITS, Zs. NAGY: *Some properties of  $C(X)$ , I*, Topology and its applications **14** (1982), 151–161.
- [9] W. JUST, A. W. MILLER, M. SCHEEPERS, P. SZEPTYCKI: *Combinatorics of open covers II*, Topology and Its Applications, **73** (1996), 241–266.
- [10] R. LAVER: *On the consistency of Borel’s conjecture*, Acta Mathematica **137** (1976), 151–169.
- [11] A.W. MILLER: *Special subsets of the real line*, in: Handbook of Set Theoretic Topology (eds. K. Kunen and J.E. Vaughan), 201–233, North Holland, Amsterdam: 1984.
- [12] A. NOWIK, M. SCHEEPERS, T. WEISS: *The algebraic sum of sets of real numbers with strong measure zero sets*, The Journal of Symbolic Logic **63** (1998), 301–324.
- [13] F. ROTHBERGER: *Sur des familles indénombrables de suites de nombres naturels, et les problèmes concernant la propriété  $C$* , Proceedings of the Cambridge Philosophical Society **37** (1941), 109–126.

- [14] M. SAKAI: *Property  $C''$  and function spaces*, Proceedings of the American Mathematical Society **104** (1988), 917–919.
- [15] M. SCHEEPERS: *Combinatorics of open covers I: Ramsey Theory*, Topology and its Applications **69** (1996), 31–62.
- [16] M. SCHEEPERS: *Finite powers of strong measure zero sets*, Journal of Symbolic Logic **64**, (1999), 1295–1306.
- [17] M. SCHEEPERS, B. TSABAN: *The combinatorics of Borel covers*, Topology and its Applications **121** (2002), 357–382.
- [18] B. TSABAN: *A topological interpretation of  $\mathfrak{t}$* , Real Analysis Exchange **25** (1999/2000), 391–404.
- [19] B. TSABAN: *Selection principles and the minimal tower problem*, Note di Matematica, this volume.
- [20] B. TSABAN: *Strong  $\gamma$ -sets and other singular spaces*, submitted.
- [21] B. TSABAN, T. WEISS: *Products of special sets of real numbers*, Real Analysis Exchange, to appear.