THE EPIDEMIOLOGY OF OSTEOPOROSIS

IN THE FRAIL INSTITUTIONALIZED ELDERLY

BY

JANE ZOCHLING

BMedSci, MBBS, MMed (Clin Epi), FRACP

A THESIS IN SATISFACTION OF THE REQUIREMENT FOR THE DEGREE OF DOCTOR OF PHILOSOPHY (MEDICINE)

INSTITUTE OF BONE AND JOINT RESEARCH DEPARTMENT OF RHEUMATOLOGY FACULTY OF MEDICINE, THE UNIVERSITY OF SYDNEY DECEMBER 2003

STATEMENT OF ORIGINALITY

I declare that this submission is my own work. To the best of my knowledge and belief it contains no material previously published or written by another person. No part of this thesis has been used to obtain any other degree.

Jane Zochling

ACKNOWLEDGEMENTS

This thesis would not have been possible without the support, guidance and encouragement of my two supervisors, Professor Philip Sambrook and Associate Professor Lyn March. Prof. Sambrook's assistance with content, concepts and timeliness has been invaluable, and A/Prof. March's optimism and lateral thinking have kept the project moving forward. Thank you to you both.

Thanks also go to Dr Tuan Nguyen for his unequalled expertise in statistical methods, and Dr. Markus Seibel for his time and resources for vitamin D analysis. Mrs Jennifer Schwarz and Dr Charles Chen have worked tirelessly in data collection and collation, and made my task so much easier in so many ways. My appreciation also extends to the staff of the Rheumatology Department at the Royal North Shore Hospital in Sydney, who have supported me through my doctorate studies with their tolerance and friendship.

I am grateful for the financial support of the National Health and Medical Research Council and the Royal North Shore Hospital (Lincoln Scholarship). Finally but by no means less importantly, I would like to extend my thanks and love to my family and friends who kept me going with 'Is it finished yet?', always believing that one day the answer would be 'Yes'.

PUBLICATIONS RESULTING FROM THIS THESIS

Peer reviewed papers

Zochling J, Sitoh YY, Lau TC, Cameron ID, Cumming RG, Lord SR, Schwarz J, Trube A, March LM, Sambrook PN. Quantitative ultrasound of the calcaneus and falls risk in the institutionalized elderly: sex differences and relationship to vitamin D status. Osteoporos Int 2002;13(11):882-887.

Zochling J, Nguyen TV, March LM, Sambrook PN. Quantitative ultrasound measurements of bone: measurement error discordance, and their effects on longitudinal studies. Osteoporos Int 2004:Feb13 (online first)

Presentations and Abstracts

Zochling J, Nguyen T, Sambrook PN. Quantitative ultrasound variance in elderly institutionalized women. Calcif Tiss Int 2003;72(4):408-9

Zochling J, Lau TC, Schwarz JM, March L, Cummings R, Cameron I, Lord S, Sambrook PN. Fracture incidence in the frail elderly. Int Med J 2002;32(Suppl.):A26.

Zochling J, Lau TC, Schwarz JM, March LM, Cumming RG, Cameron ID, Sambrook PN. Fracture incidence in the frail elderly. Arthritis Rheum 2001;44(Suppl.):S356

Zochling JM, March LM, Cumming RG, Cameron ID, Sambrook PN. A novel approach to estimating the prevalence of osteoporosis in the elderly. ANZBMS 2001 Lau TC, Brnabic AB, Zochling J, Sitoh YY, Lord SJ, Schwarz JM, March LM, Cameron ID, Cumming RG, Sambrook PN. A prospective study of risk factors for falls in elderly institutionalized individuals. Int Med J 2002;32(Suppl.):A25, A47.

Sitoh YY, Zochling J, Lau TC, Schwartz JM, Brnabic AJ, March L, Cameron ID, Cumming RG, Lord S, Sambrook P. Hypovitaminosis D is common, and more severe in winter, in institutionalized older people in Sydney. Gerontology 2001;47(Suppl.1):561

Awards and Scholarships

Lincoln Scholarship 2000 NH&MRC Scholarship 2001-2003

ABSTRACT

As our population ages, the proportion of frail elderly people requiring assisted accommodation in aged care facilities is increasing. This population is at high risk of falls and fractures, which bring significant morbidity and mortality. The prevalence of osteoporosis also increases with age, but there have been few studies of bone density in residents of hostels and nursing homes. This thesis looked at the prevalence of osteoporosis and falls in elderly people in residential care, to define the size of the problem and identify risk factors for low bone density and falling, with particular reference to vitamin D levels.

Two thousand and five men and women aged between 65 and 104 years were enrolled in the Falls and Fracture Risk in the Elderly Epidemiology (FREE) study between 1999 and 2003. The key findings from analysis of this population were firstly, that quantitative ultrasound (QUS) measures were higher in men than women independent of age, and that in men there was no significant decline in either BUA or VOS, but in women BUA declined by over 3% per decade and VOS by 1% per decade. Both ultrasound machines used in the study were shown to be reliable, with precision unaffected by advanced age. QUS was found to be sensitive to longitudinal change even in this frail elderly cohort.

Vitamin D deficiency was found in the majority of elderly aged care facility residents but supplementation conferred higher serum 25-OH-vitamin D levels. Vitamin D levels were not shown to be related to BUA, VOS or the risk of falling in this

vi

population. Serum parathyroid hormone might be important in determining future falls risk.

In summary, the results of this thesis give an important insight into the prevalence of osteoporosis and falls in the frail elderly, and how these might be predicted. Future study of prospective fracture rates in this group will then be able to assess relative risk factors for osteoporotic fracture, and identify those individuals who might benefit from directed fracture prevention strategies.

TABLE OF CONTENTS

CHAP	FER 1: INTRODUCTION AND BACKGROUND	1
1.1 I	introduction	2
1.2 A	Assessment of Bone	3
1.2.1	Quantitative Assessment	3
1.2.	.1.1 Dual-Energy X-ray Absorptiometry (DEXA)	3
1.2.	.1.2 Quantitative Ultrasound (QUS)	5
1.2.	.1.3 Quantitative computed tomography (QCT)1	0
1.2.2	Biochemical markers1	0
1.3 H	Bone changes with Age and Gender1	4
1.3.1	Bone in childhood1	4
1.3.2	Age-related changes in bone1	5
1.3.3	Anatomy and bone strength 1	7
1.3.4	Bone turnover)
1.3.5	Associations with BMD)
1.3.6	Fractures	l
1.4 H	Falls in the Elderly25	,
1.5 H	Fractures in the Elderly	3

CHAPTER 2: STUDY DESIGN AND DESCRIPTION OF

2.1	Aims	37
2.2	Study Design	38
2.3	Measurements	40
2.4	Follow Up	46

2.5	Population Characteristics48
2.6	Discussion55

CHAPTER 3: VARIANCE OF QUANTITATIVE

ULTRASOUND MEASUREMENT IN THE FRAIL

INSTITUTIONALIZED ELDERLY......58

3.1	In	troduction	
3.2	Μ	lethods	60
3.2	.1	Measurements	60
3.2	.2	Statistical methods	61
3.3	R	esults	64
3.3	.1	Demographics	64
3.3	.2	Reproducibility	65
3.3	.3	Concordance between CUBA and QUS-2	69
3.3	.4	Longitudinal measurements	71
3.4	D	iscussion	74

CHAPTER 4: PREVALENCE OF OSTEOPOROSIS IN THE

	INSTITUTIONALIZED ELDERLY	80
4.1	Introduction	81
4.2	Methods	83
4.3	Results	85
4.4	Discussion	93

CHAPTER 5: BIOCHEMICAL PROFILE OF THE FRAIL

INSTITUTIONALIZED ELDERLY......97

5.1	In	troduction	98
5.2	Μ	ethods1	00
5.2	.1	Subjects10	00
5.2	.2	Biochemical measurements1	00
5.2	.3	Statistical analysis	01
5.3	Re	esults1	02
5.3	.1	Baseline demographics1	02
5.3	.2	Distribution of 25-OH-vitamin D levels)4
5.3	.3	Determinants of 25-OH-vitamin D levels10	05
5.3	.4	Serum 25-OH-vitamin D and associated biochemical parameters10)9
5.3	.5	Independent predictors of 25-OH-vitamin D levels1	12
5.3	.6	Vitamin D supplementation1	15
5.4	Di	scussion1	18

CHAPTER 6: VITAMIN D DEFICIENCY AND

	OSTEOPOROSIS	123
6.1 I	ntroduction	124
6.2 N	1ethods	
6.2.1	Subjects	126
6.2.2	Measurements	126
6.2.3	Statistical analysis	126
6.3 R	esults	127

6.4	D	iscussion	141
6.3	.7	Independent predictors of BUA and VOS	139
6.3	.6	Hypovitaminosis D and QUS measures	.137
6.3	.5	Univariate associations with BUA and VOS	.132
6.3	.4	Quantitative ultrasound and 25-OH-vitamin D	.129
6.3	.3	Vitamin D supplementation	.129
6.3	.2	Demographic determinants of BUA and VOS	.127
6.3	.1	Baseline demographics	.127

CHAPTER 7: VITAMIN D AND PARATHYROID HORMONE AS

A PREDICTOR OF FALLING IN THE

INSTITUTIONALIZED ELDERLY......145

7.1	In	ntroduction	146
7.2	Μ	Iethods	148
7.2	.1	Subjects	148
7.2	.2	Measurements	148
7.2	.3	Statistical analysis	148
7.3	R	esults	149
7.3	.1	Distribution of falls	149
7.3	.2	Serum 25-OH-vitamin D and balance	150
7.3	.3	Survival analysis	154
7.4	Di	iscussion	158

CHAPTER 8:	SUMMARY	AND FUTURE	DIRECTIONS	163
8.1 Introducti	on		•••••	164

8.2	Summary and implications of each study165
8.3	Future directions170
REF	ERENCES
APP	ENDIX199

LIST OF TABLES

Chapter 1

Table 1.1:	Markers of bone turnover	.11
Table 1.2:	Risk factors for falling in community-dwelling elderly persons	.26

Chapter 2

Table 2.1:	Participation of residents from hostels and nursing homes	48
Table 2.2:	Resident classification scores (RCS) for men and women	.50
Table 2.3:	Baseline characteristics of the FREE cohort	51
Table 2.4:	Odds ratios (OR) for demographic measures for females compared to	
	males	52
Table 2.5:	Odds ratios (OR) for demographic parameters for nursing home residents	
	compared to hostel residents	53

Table 3.1	Descriptive statistics of the study population and duplicate ultrasound
]	measures, given as mean and standard deviation64
Table 3.2	Analysis of reproducibility of CUBA BUA, CUBA VOS and QUS-2 BUA
	in an elderly institutionalized population
Table 3.3:	Mean, between-subjects standard deviation (bSD), within-subjects
	standard deviation (wSD) and within-subject coefficient of variation
	(wCV) stratified by age group67
Table 3.4:	Comparison of BUA (dB/MHz) and T-scores measured by CUBA and
	QUS-2 instruments
Table 3.5:	T-score classifications for CUBA and QUS-2 instruments70

Table 3.6: Cha	anges in BUA and	VOS (N=56), meas	sured by CUBA	72

Chapter 4

Table 4.1: Characteristics of hostel and nursing home populations, divided by gender86
Table 4.2: Independent predictors of BUA in men and women

Chapter 5

Table 5.1: 1	Demographics of institutionalized elderly cohort with biochemical
1	measures102
Table 5.2:	Biochemical measures grouped by gender and institution, given as mean
-	+/- standard error of the mean (range), median103
Table 5.3:	Classification of residents with low 25-OH-vitamin D levels 105
Table 5.4:	Number of residents recruited each season for men and women108
Table 5.5:	Mean 25-OH-vitamin D level (standard error of the mean) for each season108
Table 5.6:	Independent determinants (mean) of serum25-OH-vitamin D113
Table 5.7:	Independent determinants of hypovitaminosis D (25-OH-vitamin D <=30
1	nmol/L)114
Table 5.8:	Frequency of bone-active medication use among men and women living in
6	aged care facilities115
Table 5.9:	Comparison of residents on vitamin D supplementation at baseline with
t	those not receiving supplements116

Table 6.1:	Mean 25-OH-vitamin D levels and QUS values for men and women in	
	hostels and nursing homes	.127

Table 6.2:	Number (%) of men and women classified by mobility128
Table 6.3:	Comparison of residents on vitamin D supplementation at baseline with
	those not receiving supplements
Table 6.4:	Mean QUS and 25-OH-vitamin D levels for each category of mobility in
	men and women
Table 6.5:	Unadjusted relationship between hypovitaminosis D and QUS variables
	(where the mean difference is the magnitude of QUS in 25-OH-vitamin D
	replete residents less QUS in residents with 25-OH-vitamin $D < 30$
	nmol/L)138
Table 6.6:	Unadjusted relationship between hypovitaminosis D and QUS variables
	(where the mean difference is the magnitude of QUS in 25-OH-vitamin D
	replete4 residents less QUS in residents with 25-OH-vitamin $D < 22$
	nmol/L)
Table 6.7:	Multivariate analysis of BUA and VOS 139
Table 6.8:	Multivariate analysis of BUA (dB/MHz) in men and women140
Table 6.9:	Multivariate analysis of VOS (m/s) in men and women140

Table 7.1:	Incidence of falls at 6 and 12 months in men and women	49
Table 7.2:	Relationship between mean 25-OH-vitamin D levels and balance1	50
Table 7.3:	Relationship between mean 25-OH-vitamin D levels and function, as	
	graded by the implicit review score1	51
Table 7.4:	Baseline characteristics and status with respect to falls at 6 months	52
Table 7.5:	Baseline characteristics and status with respect to falls at 12 months1	53
Table 7.6:	Univariate associations with time to first fall1	55

Table 7.7:	Survival analysis for time to first fall	156
Table 7.8:	Survival analysis for time to first fall in subgroup of residents aged <90	
	years with RCS>=5 and able to walk	157

LIST OF FIGURES

Chapter 3

Figure 3.1:	Difference between duplicate measures for BUA in women measured	
	by (a) CUBA, and (b) QUS-2	.68
Figure 3.2:	Percentage of individuals with significant change in BUA and VOS over	
	2.2 years using CUBA (N=56)	.73

Chapter 4

Figure 4.1:	BUA (dB/MHz) vs age (years) in (a) women and (b) men
Figure 4.2:	BUA changes with age, in a 'normal' healthy population and the
	institutionalized elderly
Figure 4.3:	BUA in institutionalized elderly women, compared with a cohort of
	independent older women taken from the DOES study, related to age,
	and with (b) separate regression lines

Figure 5.1:	Distribution of serum 25-OH-vitamin D and PTH for males and females107
Figure 5.2:	Changes in 25-OH-vitamin D levels with season in males and females109
Figure 5.3:	Parathyroid hormone levels related to 25-OH-vitamin D110
Figure 5.4:	Mean PTH values relating to 25-OH-vitamin D quintiles111
Figure 5.5:	Median PTH values relating to 25-OH-vitamin D quintiles112

Chapter 6

Figure 6.1:	Mean BUA and VOS vs 25-OH-vitamin D in men
Figure 6.2:	Mean BUA and VOS vs 25-OH-vitamin D in women131
Figure 6.3:	Mean BUA and VOS vs 25-OH-vitamin D in the whole cohort132
Figure 6.4:	Box and whisker plots of BUA and VOS related to level of mobility in
	the whole cohort135
Figure 6.5:	Box and whisker plots of 25-OH-vitamin D levels related to level of
	mobility in the whole cohort136
Figure 6.6:	Mean BUA and VOS values relating to 25-OH-vitamin D quintiles137

Figure 7.1:	Independent determinants of balance	150
Figure 7.2:	Independent determinants of implicit review score	151
Figure 7.3:	Survival curve for time to first fall	155

LIST OF ABBREVIATIONS

1,25-OH-Vit D	1-25 dihydroxy Vitamin D
25-OH-Vit D	25 hydroxy Vitamin D
ANOVA	Analysis of Variance
bALP	Bone-specific alkaline phosphatase isoenzyme
bSD	Between-subjects standard deviation
BMC	Bone mineral content
BMD	Bone mineral density
BMI	Body mass index
BUA	Broadband ultrasound attenuation
CI	Confidence interval
СТх	Carboxy-terminal cross linked telopeptide of type I collagen
CV	Coefficient of variance
dB	Decibel
DEXA	Dual energy X-ray absorptiometry
DOES	Dubbo Osteoporosis Epidemiology Study
DPA	Dual energy photon absorptiometry
Dpy	Deoxypyridinoline
EPIDOS	Epidemiology of Osteoporosis Study
EPOS	European Prospective Osteoporosis Study
EVOS	European Vertebral Osteoporosis Study
FREE	Falls and fracture Risk Epidemiology in the Elderly Study
FRI	Fracture Risk Index
LoA	Limit of agreement

LSC	Least significant criterion
MHz	Megahertz
NH&MRC	National Health and Medical Research Council (Australia)
NOF	Neck of femur
NTx	Amino-terminal cross linked telopeptide of type I collagen
OC	Osteocalcin
OR	Odds ratio
РТН	Parathyroid hormone
Pyr	Pyridinoline
QCT	Quantitative computed tomography
QOL	Quality of life
QUS	Quantitative ultrasound
RCS	Resident classification score
ROI	Region of interest
RR	Relative risk
SD	Standard deviation
SE(M)	Standard error (measurement)
SMMSE	Standard Mini-Mental Status Examination
SPA	Single energy photon absorptiometry
SPSS	Statistical package
SXA	Single energy x-ray absorptiometry
VOS	Velocity of sound
wCV	Within-subjects coefficient of variability
WHO	World Health Organization
wSD	Within-subjects standard deviation

CHAPTER 1

INTRODUCTION AND BACKGROUND

INTRODUCTION AND BACKGROUND

1.1 Introduction

Osteoporosis, a disease characterized by skeletal fragility and an ensuing increased risk of minimal trauma fracture, is a major public health problem which is growing as our population ages. The direct costs alone of treating osteoporotic fracture in Australia have been estimated at \$1.9 billion in 2001 (1). Hip fractures are the most costly to treat and associated with a significant increase in mortality, but any osteoporotic fracture brings with it a significant morbidity and impairment of quality of life (2;3).

The elderly have a high incidence of both involutional osteoporosis and of falling. The combination results in a population who are at particular risk of osteoporotic fracture. The group who are likely to be at highest risk, those elderly individuals who are unable to live independently due to disease or frailty, have not been extensively studied to determine which factors in particular contribute to fracture risk. The purpose of this thesis is to measure the incidence of both osteoporosis and falling in the frail institutionalized elderly, and describe potential predictors of osteoporotic fracture in this high risk group.

1.2 Assessment of Bone

Bone is important as both a solid structure for support and movement, and a dynamic organ involved in calcium homeostasis. Measurement of the both the strength and the activity of bone are important in health and disease. Static measures of bone density are used in clinical practice as a surrogate for bone strength, due to the relative inaccessibility of bone in vivo. Dynamic bone turnover can be quantified by measuring serum markers of bone turnover.

1.2.1 Quantitative Assessment

Bone mineral density (BMD) first became measurable in the 1960s with the introduction of single photon absorptiometry (SPA) (4). Steady improvements in technology have seen the development of single x-ray absorptiometry (SXA), dual photon absorptiometry (DPA) and more recently dual-energy x-ray absorptiometry (DEXA), considered the gold-standard for BMD assessment. Alongside absorptiometry techniques, other imaging modalities used in the assessment of osteoporosis include quantitative computed tomography (QCT) which can assess volumetric BMD, and quantitative ultrasound (QUS) of bone which is thought to incorporate connectivity and porosity in addition to pure density.

1.2.1.1 Dual-energy X-ray Absorptiometry (DEXA)

Dual-energy x-ray absorptiometry is the most commonly used technique for the assessment of osteoporosis. Two incident beams at different energies are passed through a tissue from an x-ray source, and resultant photon attenuation is measured. Differential absorption and scattering of the incident photons in different tissues allow differentiation between bone and soft tissue. The technique is used to measure areal BMD in g/cm², the integral mass of bone mineral with a standard chemical composition (hydroxyapatite) in the measurement region of interest (ROI) divided by the projected area (5).

Early versions of this technique include single-energy x-ray absorptiometry (SXA), which used only one incident beam and thus required correction for the thickness of soft tissue in the ROI. Single- and dual-energy photon absorptiometry (SPA, DPA) used a radionuclide source of photon emissions, which had a lower photon flux and larger diameter source, resulting in much longer scanning times, poorer image definition and poorer precision than DEXA today.

Current precision errors for DEXA are quoted between 1 and 1.5%, and patient radiation is relatively low (10 μ SV at the lumbar spine and 2 μ SV at the hip) (6). Limitations of the technique include problems associated with incident beam hardening and the stability of the polyenergetic spectra, which can result in deterioration in measurement accuracy. Variation in the soft tissue composition of the ROI can also impair accuracy. As calculations are based on bone mineral being uniquely made up of hydroxyapatite (Ca₁₀(PO₄)₆OH₂), problems arise when this assumption is not met, for example with fluoride or more recently strontium therapy. The atomic number of strontium is 38, compared to 20 for calcium, and so when it is incorporated into the bone mineral a disproportionate rise in BMD is seen as measured by DEXA (7).

A working party of the World Health Organization (WHO) has defined osteoporosis in postmenopausal women as a BMD more than 2.5 standard deviations below the young normal mean BMD, (8;9) based on DEXA measurements at the hip. This allows for comparison of fracture risk between different DEXA machines, and simplifies interpretation of scan results in this population. There is some controversy as to whether this classification is generalizable across other populations (10) or different methods of BMD measurement (11); nevertheless the nomenclature has been widely accepted by the clinical community as the standard method for reporting imaging of osteoporosis.

Dual-energy x-ray absorptiometry of the hip and spine has been well proven to predict osteoporotic fracture in post-menopausal women (12-14). Measurement of other sites is less predictive and not used in clinical practice (15;16). There are few studies of DEXA and fracture prediction in the very old. A large prospective study of 7598 independent women over the age of 75 years (17) showed an increase in the relative risk of hip fracture of 1.9 for each standard deviation decrease in BMD at the femoral neck. There are no studies of DEXA and fracture in the institutionalized elderly; Chandler et al (18) showed that low BMD as measured by SXA at the radius is a significant independent risk factor for osteoporotic fracture, particularly in those who were transfer independent as opposed to transfer dependent.

1.2.1.2 Quantitative Ultrasound (QUS)

Quantitative ultrasound (QUS) is a more recently developed technique for assessing bone strength. It uses non-ionizing ultrasound waves at a frequency of 0.1-1MHz which are passed through tissue by means of transducers. As they pass, some of the energy is lost as the beam is attenuated causing a change in the resultant sound intensity measured at the receiving transducer. Any sound waves will lose energy exponentially with distance, and so the waves are measured in the logarithmic scaled decibels (dB) to allow a linear relationship between dB and distance travelled. Bone ultrasound attenuation (BUA) is the measure of this energy loss across bone, and is given as dB/MHz. The interface between air and soft tissue causes pronounced attenuation, and so in practice air needs to be removed from the ROI. Early QUS machines used water baths to achieve this; dry systems use gel applied to the limb and transducers.

Velocity of sound (VOS) is measured as the speed at which ultrasound waves traverse the tissue, in metres per second (m/s). At the calcaneus, this is the average of the velocity across soft tissues and the velocity across bone, which in practice is similar to velocity across bone alone. This is dependent on transducer separation.

It is thought that QUS is more related to the so called 'quality' of the bone than just bone density, as measured by DEXA (19-22). The more bony trabeculae present in a given area, the greater the scattering of sound waves, suggesting BUA is related to bone structure. Velocity of sound also reflects the elastic modulus of the tissue. Whether these factors give an improved estimate of bone strength in clinical practice is not yet clear.

Advantages of QUS over other methods of bone assessment include its portability, ease of use, low cost and absence of ionizing radiation. This makes it an ideal tool for individuals who are less mobile, such as the institutionalized elderly. Disadvantages lie in a slightly poorer precision than DEXA, error ranging from 1-5% for BUA and 0.3-1.0% for VOS across different machines (23). Collagen diseases such as EhlersDanlos syndrome and systemic sclerosis are thought to alter the relationship between bone density and ultrasound properties (24), which might limit the usefulness of the technique in certain disease states.

There are a number of QUS machines in use both in research and in clinical practice today. Most measure QUS at the calcaneus, as a surrogate for the clinically important vertebral bodies or neck of femur; the calcaneus is a weight bearing bone, has a high percentage of trabecular bone, and has a similar pattern of bone loss to the vertebral bodies. Earlier water bath systems, including the Achilles+ (Lunar), UBA575 (Walker Sonix) and DTU-one (Osteometer), have largely been superseded by more portable dry systems. The CUBA Clinical (McCue) was the primary machine used to measure bone density in the following research, and correlates well with the Lunar Achilles+ machine (25) Transducers are 19mm in diameter and spring-loaded to give constant pressure over the patient's heel, and are silicone-capped to accommodate heels of different sizes and shapes. It is easily portable at 10kg, and easy to operate. Calibration is by electronic and water-filled phantoms. Measures of BUA and VOS are calculated. Other dry systems in use include the Sahara (Hologic Inc.), which in addition to BUA and VOS calculates a quantitative ultrasound index (QUI), relating measures back to a young adult normal value of 100. QUS is also used at other sites, including the proximal phalanges, the tibia (primarily measuring cortical bone) and multi-site machines including the Omnisense (Sunlight Ultrasound Technologies).

The DEXA-derived T-score is not necessarily applicable for other methods of assessing bone strength. When the principle of expressing raw data as standard deviations from the young normal mean is extended into QUS measurements, classifying patients as being 'osteoporotic' with a T-score less than -2.5 results in vastly different proportions of disease as is seen with DEXA (11). Faulkner et al showed that in 60 year old postmenopausal women, the prevalence of T-score-defined osteoporosis ranges from 3% when measured by QUS at the heel to 50% as measured by spinal QCT, suggesting that the use of WHO T-score thresholds to define fracture risk should be limited to the DEXA measures they were derived from.

Quantitative ultrasound of the calcaneus has been shown to predict fracture-defined (26;27) and DEXA-defined (28;29) osteoporosis. A number of studies have shown that BUA is significantly higher in men than age-matched women (29-31), although this relationship is not seen as clearly with VOS (31). A cross-sectional study of healthy Japanese shows lower rates of bone density loss as measured by QUS compared to DEXA (32), decreasing by 0.6% per year in both males and females over the age of 20. In women, DEXA at the hip and spine was better than QUS at discriminating between subjects with and without vertebral fractures (33).

Quantitative ultrasound has been well demonstrated to correlate with DEXA measures of BMD at the calcaneus (34;35). It has also been to correlate well with BMD at the spine and hip (35-37), and is associated with similar risk factors for osteoporosis (38;39). Normative data in different populations has confirmed that BUA and VOS decrease with age (40-42), although there were few individuals included over the age of 85 years. The clinical importance of QUS however lies in its ability to predict osteoporotic fracture. Case control and cross-sectional studies have shown that QUS of the calcaneus is able to differentiate between fractured and non-fractured populations in women (43-45) and in men (27;46). The relationship exists for wrist (47;48), hip (49;50) and vertebral (51-53) fractures. There are fewer prospective studies. Huang et al (54) showed that low QUS was significantly associated with increased risk for hip, for non-spine and for any fracture in 590 post-menopausal healthy women over an average follow-up period of 2.7 years.

The best prospective studies have been done in elderly populations, as there is a higher incidence of fracture allowing shorter follow-up periods. A large prospective study of 5662 independently-living elderly French women, mean age 80.4 years (55), showed that both BUA and VOS were significantly associated with incident hip fractures over 2 years, independent of femoral neck BMD as measured by DEXA. Similar results were seen in a 2-year prospective study of 6189 healthy women over the age of 65 years in the United States (56). QUS also appears useful in the frail elderly; Porter et al (57) showed in an elderly, institutionalised population that BUA was lower in those suffering hip fracture than those who did not fracture. More recently, a study of 710 residents of nursing homes or apartments for the elderly (58) showed QUS to be significantly associated with the risk of incident hip fracture over one year follow-up.

With good prospective evidence for the utility of QUS for predicting hip fracture in the institutionalized elderly, this portable and safe method of assessing fracture risk is ideal for studies such as the FREE study presented here.

1.2.1.3 Quantitative computed tomography (QCT)

Conventional computed tomography techniques can also be used to quantify BMD using appropriate software. The cross-sectional tomographic image allows a true volumetric BMD to be calculated in g/cm³, as opposed to the areal value given by DEXA, giving a conceptually more meaningful density measure. The technique has the highest diagnostic sensitivity of available assessment methods for disease and age-related bone loss (59), and has been shown to be an independent predictor of prevalent vertebral fracture in cross-sectional studies (60). However relatively high patient exposure to radiation (6), a precision error at the lumbar spine of 2-4% (5) and high installation and running costs prohibit the use of QCT clinically. Peripheral QCT of the wrist is still used frequently in research, as radiation is much lower, it selectively measures trabecular bone and allows visualization of the bone micro-architecture.

1.2.2 Biochemical markers

Bone is a living tissue, which is constantly modeling and remodeling both to maximize its function as a supportive structure and to contribute to calcium homeostasis. Biochemical measures of calcium metabolism are generally normal in osteopenia and osteoporosis, available assays being not sufficiently sensitive to detect the small changes which may theoretically be present. Serum calcium, phosphate and intact parathyroid hormone must nevertheless be measured in osteoporosis to exclude the presence of other disease states which may be impacting on bone, such as hyperparathyroidism, myeloma, metastatic disease, malnutrition, osteomalacia or phosphate wasting diseases. Vitamin D levels are the exception; as well as indicating malabsorption, nutritional deficiency and osteomalacia, 25-OH-vitamin D is important to incorporate calcium into the bone matrix. Hypovitaminosis D leads to loss of bone mineral and osteoporosis.

Bone formation and resorption markers, shown in Table 1.1, give a picture of what is happening in bone at the cellular level.

Table 1.1: Markers of bone turnover

Bone Formation Markers (serum)				
Bone-specific alkaline phosphatase (bALP)	Osteoblast membrane enzyme			
Osteocalcin (OC)	Bone matrix protein			
Collagen type 1 propeptides (pro-N, pro-C)	Collagen precursors			

Bone Resorption Markers (serum and urine)

Collagen cross-links:	Collagen breakdown products
Total or free puridinalines (Dur Dou)	
Total or free pyridinolines (Pyr, Dpy)	
Telopeptides (NTx, CTx)	
Hydroxyproline, Hydroxylysine glycosides	Collagen breakdown products
Tartrate-resistant acid phosphatase (TRAP)	Osteoclast enzyme

Alkaline phosphatase (ALP) is a cell membrane-associated enzyme expressed by liver, bone, kidney and placenta. The bone-specific isoenzyme (bALP) forms approximately 40% of the total serum ALP, and is derived from osteoblasts and osteoblast precursors. Immunoassays are specific, with an intra-individual coefficient of variance (CV) of less than 10%. There remains a small cross-reactivity with other isoforms of ALP (liver, placenta and intestine). Osteocalcin is a polypeptide expressed by osteoblasts as they actively deposit bone, and is under 1,25-OH-vitamin D control. Serum levels represent a spillover of osteoblastic synthetic activity. It is measured by immunoassay.

The most commonly measured markers of bone resorption are the collagen crosslinks. These are formed in fibrils, and have no means of metabolism once collagen is degraded and so can be measured unaltered in the urine by high-performance liquid chromatography (HPLC). The two forms are pyridinoline (Pyr) and deoxypyridinoline (Dpy). As bone is the largest reservoir for type 1 collagen in the body and turns over more rapidly than other sources, the majority of the cross-links found in the urine are from bone. There is a higher proportion of Dpy in bone than in other tissues, making it logically more specific for bone degradation.

Collagen telopeptide assays are also useful, targeting the two intermolecular sites of Pyr cross-linking in type 1 collagen: amino-terminal (NTx) and carboxy-terminal (CTx) telopeptides. The remaining markers listed in Table 1 will not be discussed, as they are less bone specific and their role in research and clinical practice is less well defined (61).

There is some biological variation in bone turnover markers. There is a distinct circadian rhythm, with levels being lowest in the morning. There are seasonal differences, which may be related to seasonal variation in 25-OH-vitamin D levels.

Changes in bone markers are strongly associated with BMD changes (62-66). A relationship to fracture is also emerging. Garnero et al (67) showed in the large EPIDOS study that elevated CTx and free Dpy levels were associated with a doubling in the risk of hip fracture, but formation markers bALP and OC were not predictive of fracture. Vertebral fracture has also been associated with higher bone turnover markers (CTx and OC) (68). It has been suggested that markers should be interpreted in conjunction with BMD measures rather than in isolation (69), and although bone turnover markers are at present largely a research tool, there may be a role for their use in clinical practice (70).

1.3 Bone and Gender

It is interesting that in most walks of life society is pushing toward gender equality, in the workplace, in opportunity and in basic human rights. However, when it comes to bone disease, osteoporosis in men and women is often treated as the same disease entity, with the same outcomes and many of the same treatment options. The WHO guidelines for grading osteoporosis are based on normative data from women; there are no corresponding well validated measurement scales for men. Gender differences are evident in bone acquisition in childhood, growth and morphometry, peak bone mass, bone loss with age and fracture rates, and so bone disease in the two sexes must be considered in light of these differences.

1.3.1 Bone in childhood

In infancy (age 0 – 18 months), absolute bone mineral content (BMC) has been shown to be the same in males and females (71), but once weight and age are taken into account, males have statistically higher bone mineral density (BMD) and BMC (p=0.02). Total body BMC has also been shown to be significantly higher in prepubertal males than females (72). Nguyen et al measured BMD and BMC in 94 males and 92 females ranging in age from 6 to 36 years, and found no difference between measures at any site before puberty (73). A Tasmanian cohort of 8-year-old children observed boys to have significantly higher hip BMD, and girls higher lumbar spine BMD, once body size had been taken into account (74). At puberty, girls had higher BMD measures at the pelvis, and higher BMD and BMC at the lumbar spine. After puberty, males generally had higher measures at all sites than their age-matched counterparts (73;75). Rates of BMD increase at the lumbar spine are fastest in early childhood (age 0-3 years) and adolescence (76). Changes in cortical bone BMD and BMC at the radius show a similar relationship to age and sexual development with females reaching their maximum rate of bone increase earlier than males, generally within 12 months of the onset of menarche (77;78). By the age of 17, females had reached 93% of their expected peak BMC and BMD, whereas males had reached 86% (77;79). Proximal radius BMD measures in a group of 362 children and young adults aged 6 - 23 years by QCT showed no gender differences pre-puberty (78), but after reaching Tanner stage 3 of development, girls had higher BMD at this site.

Peak bone mass, measured both by DEXA and QUS, have been shown to have a strong genetic influence (80), which is independent of gender. Nevertheless, environmental factors still play a major role in determining both peak bone mass in humans and subsequent bone loss with age. Bone acquisition in 8-year-old boys has been shown to be related to participation in sports and muscle strength, whereas in 8-year-old girls BMD is related to winter sunlight exposure (74). An interventional study of physical activity in 12-16 year old children confirmed a beneficial effect on BMC, areal and volumetric BMD at the femoral neck in boys but not in girls (81). Retrospective studies of exercise in older age groups confirm an effect of moderate to high physical activity on bone density at the lumbar spine and femoral neck in men (82;83).

1.3.2 Age-related changes in bone

From such a gender-defined baseline, BMD measured at the hip is subsequently seen to decrease with age in both men and women (84-86). Cross-sectional data suggests a small bone density loss at the hip in pre-menopausal women, and higher rates of loss in women after the menopause and in men (86). When this cohort was re-examined 2 years later, again a low rate of BMD loss was seen in pre-menopausal women, which tripled at the menopause before decreasing to pre-menopausal rates in the elderly. In men, the longitudinal study showed a small constant BMD loss at the hip with age (86).

BMD changes differently dependent on the site of measurement (87). The picture is not so straightforward at the spine, with BMD showing little change with age in men in cross-sectional or longitudinal studies (86;88), probably due to osteoarthritis. Bone mineral density changes in women closely followed those described at the hip in one study (86), decreasing to zero (no significant change) in the elderly, whereas Krall et al (89) showed both sexes to increase BMD at the lumbar spine after the age of 65. Relatively few studies have compared the loss of cancellous bone histomorphometrically with age between genders (90). Age-related changes in microarchitecture, as seen in iliac crest biopsies, include thinning of the trabeculae in men more so than women, and loss of trabecular connectivity more marked in women (91-94).

Cortical bone changes with age, as measured at the radius with QCT, suggest the BMD remains stable in women aged 22-40 years, then declines slowly with a more rapid drop around the age of 55 years (95). Males show a higher peak BMD which occurs in the 20s, then a rapid decline until the gender difference disappears by the age of approximately 60 years. A large, population-based study in Tromsø, Norway (96) looked at BMD at the distal and ultra-distal radius in 3062 men and 4558 women using single-energy x-ray absorptiometry. Bone mineral density decreased by 0.1% per 1 year age increase in both sexes. Over the age of 50, the rate of BMD loss in

males increased to 0.6% per year, and by a greater extent to 1.3%-1.5% in females. There is little available data on the changes in bone density in the very old.

Cross-sectional studies show that QUS parameters decrease with increasing age (97-104), although changes in the frail elderly with age have not been as extensively documented (57). Longitudinal age-related bone changes over one year can be detectable by VOS and stiffness (101) but the decrease seen in BUA did not reach statistical significance. Such longitudinal studies must be interpreted in light of the precision of the instrument used. Schott et al (105) measured QUS in 88 healthy postmenopausal women at baseline and again after 2 years. The study showed a small, statistically significant decrease in all parameters, but the decrease in BUA approximated the long-term reproducibility precision error for BUA. Velocity of sound and stiffness performed better, decreasing by 0.8% and 1.85% respectively over 2 years, at 2.5 to 5 times their precision errors. Studies using QUS to assess bone after treatment for osteoporosis have shown the instrument to be sensitive to change over 2 to 4 years (106-110) but requires a longer time period for sequential measures than DEXA due to higher precision error.

1.3.3 Anatomy and bone strength

It would be nonsensical to assume that bone size is the same between genders. Males have wider bones, seen as a larger cross-sectional area for matched vertebral bodies (111-113), independent of body size (114). This becomes important when the forces imposed on the vertebral bodies are considered; being in general taller and heavier, males impose greater forces on any one vertebral body compared to females. However, the greater male vertebral cross-sectional area results in the load per unit area being similar between males and females as young adults (112). As the skeleton ages, the increase in vertebral cross-sectional area due to periosteal bone growth is more marked in males, whereas age-related endosteal resorption of bone is more marked in women (111;115). This differential change leads to female vertebral bodies supporting a greater relative stress on a smaller relative area of bone (116).

Volumetric BMD measures allow comparison of bone density between genders allowing for skeletal size. A recent study of opposite-sex twins confirmed that there are only small bone density differences between the sexes (117) measured at the L3 vertebra, femoral neck and distal third of the radius, when bone volume, age, genetic and environmental effects are taken into account. Ebbesen et al (118) examined the L3 vertebral body in 51 female and 50 male cadavers, aged from 18-96, and found that although BMD as measured by DEXA and QCT did not vary in absolute value or rate of decline with age between the sexes, the ash weight of L3 (vertebral body bone mass), vertebral body volume and cross-sectional area were significantly lower in females at any age. Vertebral compressive strength decreased at twice the rate of BMD with age. There was no significant gender difference detected in maximal compressive stress (defined as load divided by cross-sectional area), however no corrections were made for body size.

Duan et al (112) use the term 'Fracture Risk Index (FRI)' to quantify the ratio of vertebral body compression load to vertebral body strength, with unity (or less) equating to appropriate biomechanics. Using this definition, they found male FRI increased by 21% with ageing, in contrast to 102% in women, and 9% of elderly men compared to 26% of older women had an FRI greater than unity. This is further

supported by the cadaveric study by Oyster et al (119), who measured vertebral body and phalangeal strength as the force required to crush the bone. It was shown that a significantly higher force was required to fracture the vertebral bodies of men despite no differences in vertebral BMD (measured by DEXA) by gender. Cortical bone was also stronger in men, with twice the force required for fracture than the age-matched female counterparts.

Hip geometry is related to the risk of hip fracture independent of BMD (120;121). The large European Prospective Osteoporosis Study (EPOS) (122) describes the epidemiology of the different measures of hip strength in Europeans over the age of 50 years. A subset of 1617 individuals underwent femoral DEXA, and scans were analyzed for both bone density and geometry. Men have higher cross-sectional moment of inertia (CSMI) and femoral neck cross-sectional area compared to women, but lower compressive and tensile strength measures which together indicate stronger bone. After adjusting for age, weight and height, males required higher forces at the femoral neck to sustain a fracture.

Other physical attributes of bone have been examined to explain the gender differences in fracture rates. Elastic modulus and hardness of the extra-cellular matrix are not different between genders (123). Macintyre et al (124) examined the nondominant radius of 57 males and 88 females, and reported males to have higher connectivity index, lower mean hole size and higher BMD, all reaching statistically significance. The connectivity index decreased with age in both genders, but the rate of change was higher in females (p<0.001). An age-related increase in hole size reached statistical significance in women but not men. All of these differences support an increased fragility in female bones when compared to men.

1.3.4 Bone turnover

Serum levels of bone-related alkaline phosphatase (bALP), N-terminal osteocalcin (OC), and C-terminal type 1 collagen products are tools used primarily in research to assess bone turnover. A cross-sectional study of 20-40 year old men and women (75) showed serum bALP, OC, Dpy and urinary NTx to be statistically higher in males, however all significant differences were lost after correcting for total body BMC, suggesting that bone size is important to consider when assessing gender differences in bone markers in young adults. In females, markers are seen to increase with age after the 4th decade, and more sharply after the 6th decade, corresponding to menopause (125-129), whereas in males most markers are seen to decrease. Thus bone loss in females is related to a biochemical increase in bone turnover, whereas in men the opposite would seem to be true (127). Parathyroid hormone, important in maintaining calcium homeostasis and acting directly on bone, increases with age but has no differential gender effects (130).

1.3.5 Associations with BMD

Bone mineral content and BMD of the calcaneus, measured by a ¹²⁵I photon absorption method, were significantly higher in men than women (131), but similar relationships are seen with smoking, body mass and early physical activity. Body mass index and weight are associated with BMD at all sites in women, but only at weight bearing sites in men (132-134). Lower fat content is related to lower BMD at the femoral neck in women (135;136), although being moderately overweight is seen to be associated with a higher femoral neck BMD in both genders (133). Lean muscle mass is associated with BMD and total body BMC throughout life, similarly in prepubescent males and females, but more important in females after menarche; Ferretti et al (137) showed women to have higher BMD for a given lean body mass. Ratios of total BMC to lean body mass tended to equalize between genders after the menopause. In Japanese subjects, lean body mass is more highly correlated with BMC as measured by DEXA in males, whereas the correlation between BMC and fat content is more marked in females (138)

1.3.6 Fractures

Knowing that female bones are smaller, less dense and more fragile, it is hardly surprising that age-adjusted incidence of any limb fracture is almost 3 times greater in women (139;140) than men. The incidence of most fractures increases with age in men, except for forearm fractures. Sanders et al (140) found fracture rates were highest for hip, then spine, distal forearm and humerus in decreasing order, similarly in men and women. Overall, fracture incidence was 3-4 times higher in women. In the age group 35-55 years, incidence was 65/10,000 person years in women, and 35/10,000 person years in men, and increased significantly with age independent of gender. Hip fracture rates in Oslo over 50 years were 118/10,000 person years in women and 44/10,000 person years in men (141). A large cross-sectional study showed vertebral fractures to be more common in men than women aged 50-59 years, but a slower increase in incidence is seen with age in men and by over 80 years, females predominate (142). This study however had some methodological inconsistencies in fracture ascertainment, the reporting of some traumatic fractures or osteoarthritic change as osteoporotic fracture introducing a spuriously large vertebral

fracture incidence in men. A large 2 year prospective study of 7428 men and 7865 women showed the incidence of any fracture in men aged 15-49 was 2.3 times that in women (143), whereas over 60 years of age women had 2.3 times the number of fractures as men. Peak fracture rates in men occurred in young adult males, the elderly of both genders, and in 40 year old women, particularly with respect to wrist fracture. Mortality after fracture is also higher in men (144;145).

Just as there are differential risk factors for BMD between sexes, there are also differences in fracture associations. BMD is an important risk for vertebral fracture in both sexes (146), however a case-control study of individuals with vertebral fracture did not reveal a significant difference in vertebral body morphometry between men and women (146). Men have been shown to have a higher prevalence of morphometric vertebral fracture (147), but they fracture at a higher BMD than women, such that at any given spinal BMD, men are 3 times more likely to have morphometric fracture. Men also show a higher risk of multiple (>2) vertebral fractures (148) than women. These observed differences are likely to be in part due to the higher incidence of old traumatic fractures in men. Selby et al (149) replicated this gender difference in BMD fracture threshold, with a 50% risk in fracture occurring at a significantly higher BMD in men (0.908 g/cm^3) than women (0.844) g/cm^{3}) (150). Heavy exercise has been shown to be related to vertebral fracture in men alone, OR 1.5-1.7 (151), but again there is some question of whether the crush fractures recorded were indeed true osteoporotic fractures or the result of earlier trauma, which might have resulted from such heavy exercise.

A cross-sectional study comparing individuals with hip fracture to age-matched controls confirms that men with hip fracture have significantly lower BMD at the femoral neck than their male non-fractured counterparts (152), and women with fracture have significantly less cancellous bone at the fracture site, less cortical bone at impact and significantly larger proximal femur dimensions than female controls. The large Mediterranean Osteoporosis Study (MEDOS) (151;153;154) looked at gender differences in hip fracture and quantified the contribution of various risk factors to this difference. Five percent of the gender difference was attributable to sun exposure, 3% to past history of fracture, 4% to body mass index (BMI), 4% to workrelated exercise and 14% to recreational exercise. The effect of calcium did not show a gender predisposition. The effect of alcohol consumption also shows a gender bias, with a smaller ethanol intake being related to hip fracture risk in women than in men (155). In women, 14 - 27 alcoholic beverages per week inferred a relative risk (RR) of hip fracture of 1.44 (1.03-2.03), whereas in men the risk of hip fracture increases at an alcohol intake of over 28 drinks per week, RR 1.75 (1.06-2.89). Current smoking is associated with an increase in the risk of hip fracture in women only, but a significant interaction is not seen between smoking and gender (156). After 5 years, male ex-smokers show a 25% reduction in fracture risk compared to their smoking counterparts, but this protective effect of smoking cessation is not seen in women.

Genetic influences on hip fracture risk are similar in men and women (157;158). Peak height in women and magnitude of height loss with age in men are independent predictors for hip fracture (151). Bone morphometry does not contribute significantly to the difference in risk of hip fracture between the sexes (159).

23

Similar to other sites, distal forearm fracture is more common in women. Populationbased studies have found an incidence of 18 wrist fractures per 10,000/year in women aged over 35 years, compared to 7/10,000/year in men (140); a similar discrepancy is seen in an older cohort of Europeans (incidence 7.3/1000 patient years in women, 1.7/1000 patient years in men) (139). Forearm fracture increases progressively with age in women in this group (139;160-162), whereas the incidence of wrist fracture in men does not increase until a much older age (161-163). History of diabetes mellitus (odds ratio (OR) 0.34, 95% confidence interval (CI) 0.15-0.75) and a history of prior osteoporotic fracture (OR 2.72, 95% CI 1.20 – 6.19) were independently associated with fracture risk in women but not men (164).

1.4 Falls in the Elderly

Falling and the subsequent problems of injury, loss of mobility and loss of independence is a significant public health issue in the elderly. One in every three to four people over the age of 65 years and up to two-fifths of those over 80 will experience a fall in the next year (165-167), and many will suffer multiple falls. Falling in the community significantly increases the risk of requiring institutionalized care (168). Physiological changes of aging including loss of muscle bulk and strength (169), deteriorating peripheral sensation (169), balance (169;170) and vision (169;171) combine with disease states, drug-related effects and environmental hazards to impede stable posture and gait. Risk factors for falling in independent-living elderly people have been well characterized (Table 1.2). Intuitively, it is likely that the institutionalized elderly are at an even higher risk of falling due to poorer general health, increased frailty and increased age.

Rates of falling in institutions for elderly care are significantly higher than amongst community-dwelling elders (167). An early review of falls literature in nursing homes (172) revealed an overall mean incidence of falls of 1.5 falls per bed per year (range, 0.2 to 3.6). Studies of falls risks in the institutionalized elderly largely identify the same factors as those identified in their community-based compatriots, in addition to risks peculiar to institutionalized care.

Fall history remains the strongest risk factor for falling (173), those residents who have previously fallen having more than 3 times the risk for subsequent falls. The very old (aged over 87 years) are significantly more likely to fall than their younger counterparts.

Table 1.2: Risk factors for falling in community-dwelling elderly persons.

Demographic Factors:
Advanced age (166;174)
History of previous falls (174-179)
Limitations in activities of daily living (180)
Balance and Mobility:
Poor standing balance (177;178;181-183),
Impaired gait and mobility (175;184)
Sensory and Neuromuscular Factors:
Altered proprioception (185)
Altered muscle strength (178;179;182;185)
Vision problems (acuity, depth perception) (171;184)
Poor reaction time (169)
Medical Factors:
Parkinson's Disease (175)
Arthritis (175)
Cognitive function (177;186)
Foot problems (177)
Depression (187;188)
Stroke (166;167)
Urinary incontinence (189;190)
Medications:
Central nervous system-acting (177;191;192)
Cardiac drugs (193)
Analgesics (193)
Use of four or more medications (194;195)
Environmental factors:
Home hazards (196;197)
Use of walking aids (177)

Muscle strength, in particular lower limb strength, is important for postural stability. Most studies in nursing homes have assessed qualitative muscle strength and its relationship with falls. A case-control study of two distinct populations of elderly people found fallers had poorer lower limb strength than non-fallers (167). 'Weak hip strength', assessed on physical examination (not described further), was associated with being a faller in nursing home residents, mean age 88 years (OR 8.36, 95%) confidence interval 2.71-25.79) and in community dwelling elders, mean age 69 years (OR 10.33, 95% confidence interval 2.64-40.36). 'Weak knee strength' was only significantly associated with falling in the community (OR 9.90, 95% confidence interval 2.03-48.44). Tinetti et al. (198) evaluated muscle strength by standard physical examination in 79 nursing home residents, and then followed the group prospectively for 3 months. Reduced muscle strength was recorded as 'decreased knee strength' or 'decreased lower extremity strength', but not quantified. The primary outcome measure was 'multiple (2 or more) falls'. Recurrent fallers were more likely to have reduced muscle strength in either category, compared to residents who fell once or not at all (p < 0.0005).

Quantitative measures of quadriceps strength were used in a prospective cohort study comparing falling and muscle strength between ambulatory residents of aged care facilities in Japan and the United States (199). Strength was measured as maximum kilograms lifted at the ankle for a single knee extension, and the average measure for each quadriceps used for calculations. There was a four-fold higher fall incidence in the US compared to the Japanese group. Within the US cohort, increasing muscle strength was shown to be related to lower risk of falling (p=0.04), and in both cohorts the most falls occurred in residents within the lowest quartile of muscle strength. This study adds quantitative evidence to the observation that reduced quadriceps muscle strength contributes to the risk of falling in the elderly.

Of recent interest has been the role of vitamin D, parathyroid hormone (PTH) and calcium homeostasis in balance, muscle strength and falls. It is known that vitamin D deficiency brings with it muscle weakness and abnormal gait (200-202) but most studies of the effect of vitamin D on muscle do not consider associated PTH status. It has been suggested that PTH itself has an effect on muscle strength, with muscle weakness and fatigue being a feature of primary hyperparathyroidism (203;204), and improvements being seen in patients after treatment (205-207). A study of the effect of both vitamin D and PTH levels and their relation to falls in long-term care facilities for the aged addressed this issue (208). Eighty-three hostel residents were recruited. Vitamin D levels were universally low (median 25-hydroxyvitamin D level was 27 mnol/L, interquartile range 18-37 nmol/L), and inversely related to PTH levels. Fallers (ascertained retrospectively, but after biochemical analysis) had a significantly lower 25-OH-vitamin D and higher PTH level than non-fallers. Logistic regression showed PTH levels to be predictive of having a history of falls, but 25-OH-vitamin D levels became non-significant. This was the first study to link PTH and falling, but was limited by small numbers and a retrospective study design.

Anti-depressant use in nursing home patients has been shown to be associated with a higher risk of falling (209-212) which is not dependent on the type of anti-depressant therapy (tricyclic anti-depressants, selective serotonin-reuptake inhibitors or trazodone) (213). This large retrospective study by Thapa et al. reviewed 2428 nursing home residents in Tennessee between 1993 and 1996 and identified new users

of each of the three anti-depressant types at inception. These residents and a group of randomly-selected non-user controls were then followed for a variable period of time, until leaving that institution or ceasing medication for 14 days, whichever occurred first. Falls were recorded as the primary outcome, and a total of 3524 falls were experienced over 1460 person-years of follow-up. Adjusted rate ratio for tricyclic anti-depressant use compared to no anti-depressant was 2.0 (95% confidence interval 1.8-2.2), for SSRI use was 1.8 (1.6-2.0) and for the triazolopyridine derivative trazodone 1.2 (1.0-1.4). The higher fall rates persisted through the first 180 days of drug therapy. The half-life of benzodiazepines had an impact on falls risk within this cohort (214), drugs with a longer elimination half-life (>= 24 hours) incurring a higher risk of falling, adjusted rate ratio 1.73 (95% confidence interval 1.40-2.14) compared to 1.12 (0.94-1.40) for drugs with a half life <12 hours. Shorter half-life drugs did however bring an increased risk of nocturnal falls, consistent with their peak period of action, adjusted rate ratio 2.82 (2.02-3.94). There is no conclusive evidence of a different effect of oxidative versus non-oxidative benzodiazepines on femoral fracture rate (215).

The same group identified the different range of falls risk factors between those residents who were able to ambulate independently and those who are non-ambulatory (216). Male residents who were not able to transfer independently were twice as likely to sustain an injurious fall, and those residents within the non-ambulatory group who were more mobile had a higher risk of falling. In contrast, the ambulatory group showed a strong association between the use of psychotropic agents and falls, and female residents were twice as likely to fall as their male counterparts. Nevertheless, residents who were able to transfer independently remain at a higher

risk of falling than the less mobile (173), odds ratio 1.64 (95% confidence interval 1.52-1.75).

It is not clear if the increased risk of falling in males seen in some studies is a real effect (217), with the only well-controlled study of proposed gender differences showing similar falls rates (218). Lord et al. (219) showed that community-dwelling men perform better than women at tests of muscle strength, visual field dependence, body sway and dynamic balance, all of which could be considered protective against falling. Campbell et al. (178) showed there were different risk factors for falling between the genders, men with stroke, arthritis, impaired gait or greater body sway being at high risk of falling, while women are at higher risk if they are on polypharmacy, psychotropic drugs, have muscle weakness or low systolic blood pressure. It may be that the differential distribution of risk factors does not enable reliable comparisons to be made.

The use of physical restraints in nursing home residents has been a widespread practice in the past, until it was discovered that their use was associated with a higher occurrence of fall-related injury (220). An increased rate of falls and recurrent falls with the use of physical restraints has been described in confused elderly nursing home residents (221;222). An interventional study looking at the effect of a restraint reduction program found that non-serious falls were higher after implementation of the program, but there was no significant effect on injurious falls (221). More recently, a large 2 year prospective study of education for restraint reduction in the nursing home environment has shown a 90% reduction in the use of physical restraint

30

use brought with it a significant decrease in the number of moderate and serious injuries, rate ratio 1.58 (p=0.003) (223).

So is there a difference between a non-serious fall and a fall that causes injury? Tinetti (224) suggests this is the case, finding in a study of 79 nursing home residents with chronic disabilities that injured fallers were over three times more likely to have bilateral lower extremity weakness, were more independent in ambulation and fell less often than uninjured fallers. In a study of risk factors for falling in the community, Lord et al. (225) showed multiple fallers to have weaker quadriceps muscles, poorer tactile sensation, more dependence on visual fields and higher body sway. More research needs to be done to define types of fallers, and who are at the higher risk of injury; this will be addressed further in Section 1.6. Vitamin D replacement alone has not been shown to have a significant effect on the risk of falling (226;227), with the exception of one study of 86 community-dwelling Japanese elders suffering from Parkinson's disease (228). Even in this group, there was a reduction in falls resulting in fracture, but not a significant reduction in falls overall. Two randomized controlled trials have looked at the effect of vitamin D replacement in conjunction with calcium supplementation. Dawson-Hughes et al. (229) reported no effect on any fall parameters with 500mg of elemental calcium and 700IU of vitamin D over 3 years compared to placebo in independently living older people; a more recent short-term randomized controlled trial (230) in long-stay geriatric care however has shown a beneficial effect on of 1200mg of calcium and 800IU of vitamin D the total number of falls over 12 weeks, compared to calcium supplementation alone. It is interesting to note that the institutionalized group was largely vitamin D deficient, 50% of all participants being profoundly deficient and

90% with 25-OH-vitamin D levels below 30 ng/ml (lower limit of normal). This is not surprising as the group was frail, elderly and relatively immobile with reduced exposure to sunlight. Baseline 25-OH-vitamin D levels in the independently-living study were within the normal range. Taken together, these studies would suggest that the effect of vitamin D supplementation on falling is dependent on the recipients being deficient in vitamin D to be significant.

1.5 Fractures in the Elderly

Osteoporotic fracture is well recognized as an increasing public health problem, conferring significant cost both to the individual and the community. The problem is only growing larger as the population ages, due to the increased incidence of fracture with increasing age (231): 50-60% of all women over the age of 65 are predicted to have a fragility fracture in their lifetime, and up to one third of men (232).

Over the age of 60, women fracture about 2.3 times as often as men, with the incidence of all fracture types increasing with age for both genders (143). Hip fracture incidence for women aged 85-89 years is 2.8%, and for men in this age group 1.4%. Vertebral fracture is under-reported, but prospective studies using radiological fracture as the outcome estimate the incidence to be 1.1% in women and 0.6% in men (233). Using data from the EVOS study (234), the prevalence of morphometric vertebral deformity ranges from 7.3% to 19.2% in men, and 3.5% to 16.6% in women depending on the stringency of the criteria used in reading the films.

Nursing home residents are not immune to sustaining fractures, with an incidence of hip fracture in institutionalized care of 1.9% (235), and a 7.8% incidence of any fracture (236), translating into 30% of all admissions to hospital for hip fracture (237).

There are multiple documented risk factors for sustaining a fracture in the elderly population, including medications (238), body mass (239-241) and loss of weight (242), falls (243) and low bone mineral density measured by DEXA (14;17;244;245) or by QUS (18;55;56;246), but few studies compare the relative contributions of these factors.

The burden of disease is seen as increased morbidity and mortality at an individual level, and increased health resource utilization and health expenditure on a population level. There is significant morbidity associated with vertebral fracture (247), causing pain and disability. Hip fracture increases the risk of subsequent fracture (248) and of requiring institutionalized care (237). It frequently results in loss of mobility and independence; even those individuals who return to living in the community have significantly impaired mobility and self-reported quality of life (249). A time trade-off study of the utility associated with hip fracture revealed that maintaining independence in daily life was of utmost importance to elderly women (250), and the loss of this independence was perceived to impact greatly on quality of life. Even wrist fracture has been shown to have an associated loss of physical function (251) Both vertebral (144;252-254) and hip fracture (144;237) are associated with a significant increase in mortality. The hazards ratio for 10-year mortality is 2.3-2.4 in both women and men with prevalent vertebral fracture (253). Mortality following hip fracture is much higher, ranging from 25 to 30% in the first year (255).

Loss of independence also contributes to the community costs of osteoporotic fracture. A one-year longitudinal study of the costs arising from hip fracture in Canada showed that the mean 1-year cost of hip fracture for an individual who is transferred or readmitted to institutionalized care is over double that of a patient who returns to independent living (256), acute hospitalization costs only contributing to 27% of the overall incurred costs. Vertebral fracture is associated with significant hospitalization costs, estimated to account for over 400,000 bed-days and US\$500 million over one year (257).

CHAPTER 2

STUDY DESIGN AND DESCRIPTION OF

ELDERLY POPULATION

2.1 Aims

The Fracture Risk Epidemiology in the Elderly (FREE) study was designed to test the following hypotheses:

- That quantitative ultrasound (QUS) parameters combined with clinical risk factors for osteoporosis and for falling are useful tools to predict osteoporotic fractures in a high risk population, the frail older person.
- 2: That vitamin D deficiency contributes to hip fracture risk through an increased risk of falling.

The study was supported by a project grant from the National Health and Medical Research Council of Australia (NH&MRC), and employed the equivalent of two full time research assistants who were responsible for data collection and maintenance of the study database. Ethics approval was obtained from the Royal North Shore Hospital Ethics Committee.

2.2 Study Design

The FREE study was set up as a large population-based prospective study of risk factors for falls and fractures, including quantitative ultrasound (QUS) measures to assess osteoporosis. Falls and fractures were then recorded prospectively for a period of 2 years. The study was carried out in the Northern Sydney Health Services area, which covers a geographical area of 1301.8 square kilometres and has a population of 47 275 men and 67 817 women aged 65 years and over (at 2001 Census, http://www.ausstats.abs.gov.au).

Recruitment of aged care facilities to participate in the FREE study began in March 1999. A database of all hostels and nursing homes for aged care in the Northern Sydney Health Services Area was compiled from a list provided by the Department of Health and Aged Care, and institutions were randomly assigned to blocks of 10 nursing homes and 5 hostels. Facilities were then approached block by block (approaching equal numbers of nursing home and hostel blocks) to maintain randomization. Initial contact was made by letter, with a follow up phone call to arrange a preliminary visit.

At each participating facility, all eligible residents were visited and invited to participate. Individuals who were totally non-weight bearing (bed bound, bilateral amputees), non-English speaking or under the age of 65 years were not included in the study. Residents gave informed, signed consent or if unable, their next of kin was approached by letter to give proxy consent. Information on past medical history, falls and fracture risks and a proxy quality of life questionnaire was also obtained from carers when a resident's cognitive impairment precluded accurate history taking. These individuals were identified by facility staff. If a resident gave independent consent but was subsequently found to be cognitively impaired (defined as a Mini Mental Score of less than 18), the appropriate carer was then approached prior to further participation in the study. This method of obtaining proxy consent has been shown to be valid in other populations (258), and will be further evaluated in this study. Copies of each signed consent form were provided for each patient's file, along with an information sheet about the FREE study and contact details for study personnel. Examples of these forms are given in Appendix 1.

Recruitment of aged care facilities was completed in February 2003, when the target of more than 2000 enrolled residents was reached. Population data from Dubbo, New South Wales, Australia identified a fracture rate of 5 to 10% per year for any fracture in independently-living elderly men and women, and a hip fracture rate of 3 to 5% (232). Estimates from published trials in nursing homes suggest a hip fracture rate of up to 6% per year (259;260). Over the 2 year follow up period, the cohort would be expected to experience approximately 60 (3% per annum) to 120 (6% per annum) hip fractures (allowing for some attrition due to death in this elderly group).

2.3 Measurements

Clinical risk factors for both falls and for osteoporosis and fractures were assessed in all subjects at interview. Basic demographics including age, sex, country of birth, activities of daily living status (Barthel Index) (261) and co-morbidities (by Resident Classification Scale graded 1 (lowest care) to 8 (highest level of care required)) (262) were recorded. The RCS is assigned by an aged care assessment team (independent of this research study) and is linked to government funding for ongoing aged care. It incorporates assessments of communication, mobility, ability to eat and drink, hygiene, bladder and bowel management, cognition, aggression or emotional instability, danger to self or others, social needs, medications and technical nursing care requirements (Appendix 2), and as such is a tool to define the degree of nursing care and supervision an individual requires on a daily basis.

The following parameters previously shown to be related to falls (Chapter 1.5) were assessed: use of a walking aid, medications (ascertained by medical record review), history of falls in the previous year, history of stroke, Parkinson's disease, continence, joint replacement, and osteoarthritis of the hip or knee. Knee pain (263) and osteoarthritis of the knee (264) have been associated with objective impairment of balance.

Cognitive function was assessed using the Standardised Mini Mental Status Examination (SMMSE) (265). When one or more (up to 7) of the 30 items were not completed due to visual or hearing impairment, a correction calculation was made (as per personal communication, Dr A. Jorm, NHMRC Psychiatric Epidemiology Research Centre):

(total score possible * total number of items completed satisfactorily) (total score possible – total number of items not attempted).

When 8 or more items were missing, the total score was recorded as missing as a refusal is most likely due to inability to respond correctly (266). A score of less than 18 was interpreted as the most appropriate cut-off to define insufficient cognitive function to give informed consent for the purposes of this study (267).

Quality of life (QOL) was measured using the EQ-5D (268), a categorized scoring model of the generic EuroQol instrument (269) (which uses visual analogue scale assessments). The form used for data collection is shown in Appendix 1. The EQ-5D has been shown to be valid and reliable for measuring health-related QOL in a number of different healthy (270-273) and disease populations (274-277), including postmenopausal osteoporosis (278). Choice of health-related quality of life instrument is largely a function of the population to be studied (279). The EQ-5D has been shown to be completed more reliably than the other commonly used QOL instrument, the SF-36, in elderly patients after a stroke (258), and experience with other projects in Northern Sydney has shown the EQ-5D is more easily completed by frail older people than either the Sickness Impact Profile or the Medical Outcome Study Short Form 12 (personal communication, Prof. Ian Cameron). Analyses were conducted both with and without proxy quality of life data in light of systematic differences between individual and proxy reporting (280;281). A disease-specific instrument was not incorporated into the study due to the high number of confounding variables and comorbidities in this frail population.

A validated implicit reviewer assigned severity score was given for each resident (282), rated at the time of the baseline assessment, where scores are given as:

- 1 no symptoms or medical complications
- 2 mild symptoms
- 3 moderate symptoms
- 4 severe symptoms that severely alter or immediately threaten life or prognosis

according to predefined definitions (Appendix 3).

Information relating to osteoporosis and fracture risk was also collected. This included a history of past corticosteroid and hormone replacement therapy (quantified where possible), current or past smoking and history of previous low trauma fracture (after age \geq 50). Self-report of medical conditions (283-285), in particular of previous fracture (286-288), has been shown to be reliable in elderly persons.

Falls risk was tested by assessments of quadriceps strength and static balance, reaction time, visual contrast sensitivity and proprioception according to published methods proven to predict falls and fractures in the elderly (169;245). Quadriceps muscle strength was measured in the sitting position from a standardized chair in the subject's dominant leg using a strain gauge, measured in kilograms. Quadriceps muscle strength, the ability to stand from a seated position and visual contrast discrimination were measured, but will not be discussed further for the purposes of this thesis.

Static balance was measured as the ability of subjects to maintain balance while standing on a firm and a compliant surface, comprising a 150 x 150 x 10 cm foam mat. Subjects were classified into 5 grades

- 1 unable to maintain balance for any period without support on a firm surface
- 2 able to stand for less than but not equal to 30 seconds unsupported
- 3 able to stand for 30 seconds unsupported on firm ground but not on a compliant surface
- 4 able to stand for 30 seconds unsupported on firm ground but less than
 30 seconds on a compliant surface, and
- 5 capable of maintaining balance whilst standing on a firm or compliant surface for 30 second periods without difficulty.

Weight was measured using the same set of scales in those residents able to stand unsupported. For those who were not sufficiently mobile to use the scales, the scales were calibrated to measure '0 kg' with a ramp and empty wheelchair, and then the resident was weighed whilst sitting in that wheelchair to give a net resident weight. If neither of these methods was possible, the most recent weight in the patient's medical notes was recorded. In order to estimate height, the lower leg length from floor to knee (bent at 90 degrees) was measured for subsequent calculations (289).

An assessment of bone mass was made using two quantitative ultrasound (QUS) machines. Broadband ultrasound attenuation (BUA) and velocity of sound (VOS) were measured at the left calcaneus in all participants using the same McCue CUBA Mark II ultrasound machine. In the event of foot deformity, unilateral foot amputation or injury to the left foot, a measurement was made at the right calcaneus and the reason documented. BUA was also measured using a Metra QUS-2 ultrasound machine in a smaller sample of subjects, as this machine only became

available after study commencement. The coefficient of variation from duplicate measurements for BUA in our laboratory is 2.9% for the CUBA machine and 1.3% for the QUS-2. Each QUS measurement was repeated twice in the study population, repositioning the foot between measurements; a third measurement was taken when the difference between the first two BUA measures was greater than 5 dB/MHz. Each machine was calibrated daily using standard operating phantoms for each machine.

Blood was taken for vitamin D and biochemistry analysis in all consenting participants. Due to the 'invasive' nature of venesection, any resident was permitted to refuse blood being drawn without being excluded from the remainder of the study. Biochemistry was analyzed on a Hitachi 917 analyzer in a single laboratory. Serum 25-hydroxyvitamin D (250HD) was measured using the Diasorin 25-OH-D radioimmunoassay (RIA) kit. Sensitivity was measured at 4 nmol/L, intra-assay precision 7.6% and inter-assay precision 9.0%, with a 'normal' laboratory range of 39 – 140 nmol/L.

Baseline serum levels of intact parathyroid hormone (PTH) were determined by a two-site chemiluminescent enzyme-linked immunometric assay on a DPC Immulite 1000 analyser. The assay procedure measures the intact PTH molecule. The assay has a typical intra-assay precision of 5.5%, an inter-assay precision of 7.9% and the laboratory reference range is 23.7 - 66.2pg/mL (2.5 – 7.0 pmol/L).

Serum albumin was measured by BCG colorimetric assay (Roche), with a normal range of 40-50 g/L. The within-run precision (coefficient of variance, CV) given by

the manufacturer was 0.4% and the between-run CV was 1.7%. Inter-assay precision in this laboratory has been calculated as 2.0%. Serum calcium was measured by colorimetric assay (Roche) using p-cresolphthalein and adjusted for circulating albumin levels, with a normal range 2.15-2.55 mmol/ L, manufacturer's within-run CV 0.9% and between-run CV 1.5%, and measured inter-assay CV in this laboratory 2.1%. A modified Jaffé (picric acid) kinetic colorimetric assay was used to measure serum creatinine with a normal range in males of 70-110 μ mol/L and in females 50-90 μ mol/L, manufacturer's precision of 0.7% intra-assay and 2.3% inter-assay and a measured inter-assay CV of 3.0%. Inorganic phosphorus levels were measured by an endpoint method with sample blanking, based on the formation of ammonium phosphomolybdate complex. The normal range in this laboratory is 0.6-1.3 mmol/L and inter-assay precision of 2.5%; manufacturer's within-run CV is 0.9% and between-run CV is 1.4%.

2.4 Follow Up

Once enrolled in the FREE study, residents were followed up on a regular basis to record the number of falls, injuries and fractures that had occurred since recruitment. A fall was defined as 'unintentionally coming to rest on the ground, floor or other lower level, whether or not an injury occurred'. An injury was recorded if it required treatment, medical consultation, investigation or observation or pain relief. Any fracture was confirmed by x-ray, x-ray report or hospital discharge summary. Where a clinical diagnosis of a fracture was made without radiological evidence, it was recorded as 'unconfirmed' and not included in subsequent analyses.

Incident reports were examined at each facility to confirm falls, and radiology reports obtained to confirm each fracture and its site. Residents' medical progress notes were also reviewed to confirm incident reports, to assess any injury that might have been sustained at the time of the fall and to identify any falls that were not captured on incident report. X-ray reports are recognized as being superior to medical record review for ascertainment of fracture incidence (290). All deaths were recorded. Residents who moved facilities were tracked in order to minimize follow up losses.

The first follow up visit took place about 6 weeks after all baseline assessments had been completed for that aged care facility. Visits continued approximately every 2-3 months for 2 years or until death. After 2 years, only hip fractures and deaths were recorded at follow up.

Previous studies have shown osteoporotic fractures to be associated with a decline in QOL (144;168;291) and an increase in overall mortality (144). To assess whether

fractures in frail elderly residents of aged care facilities lead to a decrease in cognitive function, quality of life and increased mortality, each resident sustaining a fracture during the study period was reassessed four months after fracturing, using the SMMSE and EQ-5D instruments. A control subject, matched for age, gender, institution and broad SMMSE function, was also reassessed at that time.

2.5 Population Characteristics

Thirty hostels and 52 nursing homes were included in the study. A further 2 hostels were due to be approached but had closed between randomization and study participation. Four nursing homes were approached but refused to participate, 2 were undergoing rebuilding and unable to participate at the time they were approached, and 5 had closed since randomization. Numbers of residents available for recruitment and the subsequent participation rates are given in Table 2.1.

		Hostel	Nursing home
Number of residents (total)		1751	2685
Excluded	Age < 65 years	33	59
	Bed bound	0	458
	Non-English speaking	23	89
	Other	3	3
	Total	59	701
Total possible		1692	1984
Self consent	Total	1232	562
	Included (% response)	958 (77.8%)	428 (76.2%)
Carers consent	Total	460	1432
	Included (% response)	149 (32.4%)	470 (32.8%)
Participants	Total (% response)	1107 (65.4%)	898 (45.3%)

Table 2.1: Participation of residents from hostels and nursing homes

A total of 2005 residents were recruited for the study and completed the baseline assessments. Of these, 67% (n=1386) had been self-consents, and the remaining 33% (n=619) had proxy consents completed. As anticipated, carer consent was more difficult to obtain due to a number of reasons, including inability to locate the appropriate carer, introduction of privacy laws that restricted access to information about carers, non-response to multiple mail-outs, mistrust of clinical trials and a desire that a relative be not inconvenienced.

Simple demographic details were collected on non-participants for the first year of the study. Those individuals who refused to participate in the study, either personally or through carer advice, were of a similar age to participants (mean age of non-participants 85.5 years compared to mean age of participants 85.2 years, t=0.79, p=0.4). Non-participants had been residing in aged care facilities for slightly less time than participants (mean difference 3.8 months, 95% confidence interval for the difference 0.15 to 7.4 months, t=-2.04, p=0.04). More non-participants fell into RCS categories 1 and 2 and less into 6-8 than participants (χ^2 =116, df 7, p<0.0001) suggesting a lower level of dependency on nursing care.

Baseline demographic characteristics of the study group are given in Tables 2.2 and 2.3. There was no significant difference in the distribution of RCS scores between men and women (χ^2 =11.08, df=7, p=0.14), suggesting that the range of disability was the same for men as for women.

RCS	Male (n=473)	Female (n=1532)
1 (least care required)	37 (7.8%)	163 (10.6%)
2	99 (20.9%)	304 (19.8%)
3	68 (14.4%)	240 (15.7%)
4	21 (4.4%)	52 (3.4%)
5	26 (5.5%)	119 (7.8%)
6	63 (13.3%)	184 (12.0%)
7	37 (7.8%)	163 (10.6%)
8 (most care required)	99 (20.9%)	304 (19.8%)

Table 2.2: Resident classification scores (RCS) for men and women

		Male			Female		
		Hostel	NH*	Total	Hostel	NH	Total
Number		263	210	473	844	688	1532
Age (years), mean (sd)		82.3 (7.42)	83.4 (6.36)	82.8 (7.81)	86.6 (8.24)	86.7 (6.86)	86.6 (6.59)
Implicit review	1 (no symptoms)	20 (7.5%)	1 (0.5%)	21 (4.4%)	44 (5.2%)	2 (0.3%)	46 (3.0%)
	2 (mild symptoms)	102 (38.8%)	28 (13.3%)	130 (27.5%)	313 (37.1%)	83 (12.1%)	396 (25.8%)
	3 (moderate symptoms)	133 (50.6%)	165 (78.6%)	298 (63.0%)	471 (55.8%)	552 (80.2%)	1023 (66.8%)
	4 (severe symptoms)	7 (2.7%)	14 (6.7%)	21 (4.4%)	10 1.2%)	38 (5.5%)	48 (3.1%)
	Not yet determined	1 (0.4%)	2 (0.9%)	3 (0.7%)	6 (0.7%)	13 (1.9%)	19 (1.3%)
Number of fallers in past 12	months (%)	118 (44.9%)	123 (58.6)	241 (51.0%)	381 (45.1%)	411 (59.7)	792 (51.7%)
Number sustaining a low-	Any fracture	56 (21.3%)	61 (29.0%)	117 (24.7%)	409 (48.5%)	372 (54.1%)	781 (51.0%)
trauma fracture since the age of 50 (%)	Hip	12 (4.6%)	25 (11.9%)	37 (7.8%)	108 (12.8%)	184 (26.7%)	292 (19.1%)
	Spine	7 (2.7%)	7 (3.3%)	14 (3.0%)	80 (9.5%)	145 (21.1%)	125 (8.2%)
	Wrist	5 (1.9%)	8 (3.8%)	13 (2.7%)	138 (16.4%)	85 (12.4%)	223 (14.6%)
Smoker (current)		46 (17.5%)	8 (3.8%)	54 (11.4%)	30 (3.6%)	19 (2.8%)	49 (3.2%)
OA knee		78 (30.0%)	46 (21.9%)	124 (26.2%)	332 (39.3%)	196 (28.5%)	528 (34.5%)
Hip replacement		14 (5.3%)	18 (8.6%)	32 (6.8%)	84 (10.0%)	74 (10.8%)	158 (10.3%)
Knee replacement		21 (8.0%)	13 (6.2%)	34 (7.2%)	56 (6.6%)	34 (4.9%)	90 (5.9%)
Incontinence	Day	76 28.9%)	99 (47.1%)	175 (37.0%)	291 (34.5%)	371 (53.9%)	662 (43.2%)
	Nght	110 (41.8%)	119 (56.7%)	229 (48.4%)	336 (39.8%)	429 (62.4%)	765 (47.5%)
HRT	I	N/A	N/A	N/A	55 (6.5%)	23 (3.3%)	78 (5.1%)
Corticosteroids		28 (10.6%)	10 (4.8%)	38 (8.0%)	81 (9.6%)	59 (8.6%)	140 (9.1%)
Parkinson's disease		23 (8.7%)	32 (15.2%)	55 (11.6%)	17 (2.0%)	44 (6.4%)	61 (4.0%)
History of CVA		69 (26.2%)	62 (29.5%)	131 (27.7%)	151 (17.9%)	153 (22.2%)	304 (19.8%

Table 2.3: Baseline characteristics of the FREE cohort

Women were an average of 3.8 years older than their male counterparts (95% confidence interval for the difference 2.99 to 4.55, t=9.51, p<0.0001). Women were more three times more likely than men to have a history of minimal trauma fracture since the age of 50 years, independent of fracture type (Table 2.4). There was no significant difference in the reporting of falling in the previous 12 months between men and women.

Table 2.4: Odds ratios (OR) for demographic measures for females compared to males.

	OR	95% CI	χ^2 statistic	р
Nursing home accommodation	0.98	0.80 - 1.21	0.038	0.9
History of falls	1.02	0.89 - 1.09	0.116	0.75
History of any fracture	3.18	2.52 - 4.02	101.144	<0.0001*
- hip fracture	2.80	1.96 - 4.01	33.051	<0.0001*
- wrist fracture	6.09	3.45 - 10.76	49.24	<0.0001*
- spine fracture	2.91	1.66 - 5.11	15.143	<0.0001*
Smoking	0.26	0.17 - 0.39	49.69	<0.0001*
Incontinence - day	1.36	1.07 – 1.64	6.46	0.01*
Incontinence - night	1.09	0.89 - 1.34	0.605	0.4
Parkinson's disease	0.32	0.22 - 0.46	38.29	<0.0001*
History of CVA	0.65	0.52 - 0.83	12.32	<0.0001*

*statistically significant p<0.05)

Elderly persons in Australia who require assisted accommodation are assigned places in hostels or nursing homes according to the perceived level of care required, measured by RCS. It would therefore be expected that nursing home residents, requiring a higher level of day to day care, would have significantly more comorbidity and poorer levels of function. When compared to hostel residents, individuals living in nursing homes were seen to have approximately a two-fold increase risk of being fallers (OR 1.9, p<0.0001), were more likely to have sustained a hip fracture (OR 2.5, p<0.0001) and more likely to be incontinent of urine or have a history of stroke (Table 2.5). Nursing home residents were significantly less likely to be current smokers, most probably related to immobility (smoking now rendered an outdoor activity in Australian aged care facilities).

Table 2.5: Odds ratios (OR) for demographic measures for nursing home residents
compared to hostel residents.

	OR	95% CI	χ^2 statistic	р
Female gender	1.02	0.83 - 1.26	0.04	0.9
History of falls	1.93	1.61 – 2.31	50.23	<0.0001*
History of any fracture	1.32	1.11 – 1.58	9.44	0.002*
- hip fracture	2.49	1.95 - 3.18	55.80	<0.0001*
- wrist fracture	0.77	0.58 - 1.01	3.31	0.07
- spine fracture	0.72	0.51 - 1.03	2.97	0.09
Smoking	0.42	0.27 - 0.66	15.16	<0.0001*
Incontinence - day	2.34	1.95 – 2.81	83.97	<0.0001*
Incontinence - night	2.50	2.08 - 3.01	98.00	<0.0001*
Parkinson's disease	2.54	1.71 – 3.77	22.88	<0.0001*
History of CVA	1.38	1.11 – 1.71	8.63	0.002*

*statistically significant, p<0.05)

The FREE cohort has been shown to be a heterogeneous population, with wide ranges of comorbidity. Their commonality lies in their advanced age, and a requirement for assisted living which renders them unable to live independently. This thesis examines the health of the frail elderly with respect to bone, specifically their risk of falls and fractures which incur significant morbidity and mortality to an already impaired population.

2.6 Discussion

The major strengths of this study lie in its size, randomization and prospective follow up. To our knowledge this is one of the largest prospective studies of falls and fracture risk that has been undertaken in frail institutionalized elderly population, with previous studies recruiting in the low hundreds (176;292-295). In particular, this is the largest cohort of elderly men studied in aged care facilities for any of the falls or fracture parameters. The study is population-based, with rigorous prospective data collection over a minimum of 2 years. Aged care facilities were randomized at the outset and this randomization was maintained throughout the study with negligible facility refusal or drop-out. Large numbers of both men and women have been recruited. The response rate from residents able to give self-consent was reasonable, and once recruited all participants completed the study with minimal 'missing' data.

The population was comparable in age and gender to previous published studies in the elderly who live in assisted care facilities (58;173;176), allowing generalizability to other groups around the world. This supports the applicability of our results to other frail elderly populations. Non-participants were of a similar age to participants and had lower RCS classifications, so that we can be confident in the assumption that the study includes the more frail population. This does introduce a potential bias in the study group, which must be considered when relating the results of the FREE study in healthier, more independent populations.

All major outcome measures had been previously validated in published studies, to allow meaningful conclusions to be drawn.

Quantitative ultrasound of the calcaneus has been shown to be a predictor of fracture rate in younger, independently living populations (54-56) and in smaller studies of the frail elderly (57;58). This study has measured BUA and VOS in a much larger population, which has not been carried out on such a large scale previously. It is anticipated that this will provide a greater power to predict fracture in a population of elderly persons living in aged care facilities. This is important from a practical viewpoint, as QUS is much more easily performed in a frail immobile population than the 'gold standard' DEXA measurement.

One of the problems encountered in the recruitment of participants for the FREE study was a poor carers' response, resulting in a smaller proportion of cognitively impaired residents (compared to those with adequate cognition to give self-consent) being included in the study. This skews the population towards a higher cognitive function than truly exists in hostel and nursing home accommodations. Such a recruitment bias must be considered when interpreting results, particularly if poor cognitive function contributes to falls and fracture risk (176;296). This systematic bias would result in an underestimate of the true falls and/or fracture risk in assisted care facilities, and reduce the power of the study to detect possible relationships between cognitive function, falls and fractures.

Those individuals who were non-English speaking, bed-bound or bilateral amputees were not included in the analysis, which might impact on the generalizability of our results. There were a total of 112 (2.5%) non-English speaking residents who were not approached for the study; care must be taken in applying the findings of the FREE study to recent immigrants to Australia. Nevertheless, 83 (4.1%) of residents included

in the study were born in non-English speaking countries, allowing a degree of generalizability of results to elderly individuals of different ethnic backgrounds who have lived in Australia for some years.

There were 458 residents (10.3%) in the nursing homes participating in the study who were bed bound and therefore not included; this is unlikely to have affected outcomes as it is deemed that if a resident does not mobilize in any way, the only factor important in future risk of falling is nursing care, which is not the purpose of this study. The remaining 6 exclusions form only 0.1% of the aged care facility population, and so even if they fell in the highest or lowest risk categories, their exclusion would not impact the final results.

Loss to follow up is a major difficulty in this elderly population, due to the high mortality rate in the frail elderly. Cox regression has been carried out to maximize the information collected during the study, allowing all residents who completed baseline assessment to be included in the analyses up until the time of death. The large number of residents recruited for the study will in part offset the expected losses to mortality from all causes.

In summary, this is a large well-constructed population-based prospective study of falls and fracture risks in the frail elderly living in aged care facilities which is expected to yield valuable information to ultimately assist in formulating preventative strategies for this high-risk population. **CHAPTER 3**

VARIANCE OF QUANTITATIVE ULTRASOUND MEASUREMENT IN THE

FRAIL ELDERLY

VARIANCE OF QUANTITATIVE ULTRASOUND MEASUREMENT IN THE FRAIL ELDERLY

3.1 Introduction

Osteoporosis is an increasing community health problem, particularly as the population ages. Bone mineral density (BMD) measured by dual energy X-ray absorptiometry (DEXA) is the current gold standard for the diagnosis of osteoporosis. BMD has been shown to predict fracture risk, with each standard deviation reduction in femoral neck BMD increasing the age-adjusted risk of hip fracture by about two fold (range 2.0 to 3.5) as well as the risk of any atraumatic fracture by two (range 1.7 to 2.4) (14;17;244;245). DEXA is however not always feasible, for example in large population 'screening' and in elderly, less mobile groups. Calcaneal ultrasound is a portable, easy to use and non-invasive method of assessing trabecular bone and has been recently shown to predict risk of osteoporotic fracture in community-dwelling postmenopausal women (55;56). Little is known of the reliability or reproducibility of heel ultrasound techniques in the very elderly and how this may vary with increasing age.

This study aimed to investigate the reliability and reproducibility of two calcaneal ultrasound machines in a population of frail elderly institutionalized subjects, exploring the variability of broadband ultrasound attenuation (BUA) and velocity of sound (VOS) in the very elderly and to assess sensitivity to change with extreme ageing.

3.2 Methods

As this analysis was carried out prior to the completion of patient recruitment, only the first 1587 subjects recruited into the FREE study who had calcaneal ultrasound measurements were included in this analysis, of which 1210 (76.2%) were female.

3.2.1 Measurements

Broadband ultrasound attenuation (BUA) and velocity of sound (VOS) were measured in all participants using the same CUBA Mark II ultrasound instrument (McCue Ultrasonics, London, UK). BUA was also measured using the QUS-2 ultrasound machine (Quidel Corporation, San Diego, USA) in 684 (509 females) of the participants. BUA measurements were taken in a lesser number of subjects with the QUS-2 machine, as the machine only became available for use after study commencement. Assessments were made at the left calcaneus, unless foot deformity or oedema precluded measurement and then the right heel measurement was taken. Two consecutive measurements were made, repositioning the foot for each measurement. Fifty-six consecutively recruited residents of hostels were remeasured using the same technique on the CUBA after an average period of 2.2 years. T-scores were calculated using machine-defined normal ranges.

Patient details including age and medication history were collected, and weight was measured in all participants, using a wheelchair ramp for those patients unable to stand unsupported on the scales. Height was not measured in this population.

Ambient temperature has been shown to affect QUS accuracy and precision of the Achilles (Lunar) water bath machine, giving higher values at lower temperatures (297), but it is not known if this is applicable to dry systems. Air conditioning in hostels and nursing homes kept relatively stable room temperatures throughout our study. Moreover, follow-up measures were taken in the same season as the original measurements to avoid temperature measurement errors longitudinally.

3.2.2 Statistical methods

Analysis of reproducibility

Three sample statistics were computed to assess the reproducibility of each measure: (a) the standard error measurement (SEM, within-subject standard deviation) which was calculated as the square root of the average variance from individual subjects, (b) this SEM was then expressed in relation to the overall mean as the coefficient of variation, and (c) the limit of agreement (LoA) as described by Bland and Altman (298).

Analysis of concordance between CUBA and QUS-2 machines

The concordance between the CUBA and QUS-2 instruments for BUA was addressed by calculating the coefficient of concordance (299) as

$$C = \frac{2Cov(BUA_c, BUA_q)}{s_c^2 + s_q^2 + (BUA_c - BUA_q)^2}.$$
 Cov(BUA_c, BUA_q) is the covariance between the

measurement of BUA by CUBA, and BUA by QUS-2 instrument, s_c^2 and s_q^2 are the between-subjects variance of BUA measurements by the CUBA and QUS-2, respectively, and BUA_c and BUA_q are the respective means of BUA measured by the two instruments. The coefficient *C* is an index of agreement of measurements between two instruments, i.e. the degree to which pairs of measurements fall on the line of identity. Thus, even with a high correlation between the two methods, if their means are different, the value of C is still small, e.g. poor agreement. Limits of agreement between machines were calculated using repeated BUA measures for each of the two instruments (12). The weighted Kappa statistic was used to assess the degree of agreement between the two instruments in various T-score-based classifications.

Longitudinal analysis

Statistical analysis was carried out following the method of Nguyen and Eisman (300). For subjects with two visits, the rates of change in BUA and VOS (CUBA) were expressed in actual units and in percentage terms. In the latter, the difference between the mean measurement at the follow-up visit and the mean baseline measurement was calculated, then divided by the mean baseline measurement, and expressed as a percentage change. Because each measurement (either at baseline or follow-up visit) was subjected to random measurement error, a change was considered "significant" if it exceeds the change that may be expected from random fluctuation with a certain probability. In traditional analysis, the variance of difference between two successive measurements, x_0 and x_1 is equal to the sum of individual variances, i.e., $var(x_0 - x_1) = 2s^2$, where s is an estimate of SEM as mentioned above. Under the assumption of the Normal distribution, $1.96 \times \sqrt{2}s$ (also known as the least significant criterion (LSC)) covers 95% of the random variability of within-subject differences. However, the above formulation is based on the assumption that successive measurements within an individual are independent. This is unlikely to be true, as measurements within an individual are correlated, although the correlation decreases with increasing intervals between measurements. Therefore, a more logical formulation must take this correlation into account. Assuming a simple

autoregressive model, the variance of difference between successive measurements can be shown to be equal to: $var(x_0 - x_1) = 2s^2[(1 - \rho_i) + \rho_i s_a^2 / s^2]$, where ρ_i is the serial correlation between two consecutive measurements; and σ_a^2 , the analytical variance (e.g. its square root, σ_a , is referred to as measurement error) included in the within-subject variance, s^2 . As the analytical variance is small relative to the withinsubject variance, the formula can be reduced to: $var(x_0 - x_1) = 2s^2(1 - \rho_i)$. Therefore, in this study, we used two criteria to assess the longitudinal change in each ultrasound measure

3.3 Results

3.3.1 Demographics

The characteristics of the study population are shown in Table 3.1. Females were on average of 3.94 years older than their male counterparts (95% confidence interval 3.13 to 4.74 years, p <0.001) and weighed 14.4kg less (95% CI 12.8 - 15.9, p <0.001).

Table 3.1: Descriptive statistics of the study population and duplicate ultrasound

 measures, given as mean and standard deviation.

		Female	Male
Number		1210	377
Age (years)		86.26 (6.68)	82.32 (7.78)
Weight (kg)		56.76 (12.52)	71.14 (13.47)
CUBA BUA (dB/MHz)	Measure 1	45.70 (18.13)	71.83 (23.00)
	Measure 2	45.44 (18.31)	71.88 (23.02)
CUBA VOS (m/s)	Measure 1	1558.73 (42.83)	1596.46 (47.60)
	Measure 2	1558.45 (43.52	1596.26 (47.97
QUS-2 BUA (dB/MHz)	Number	509	175
	Measure 1	53.52 (14.88)	73.50 (21.19)
	Measure 2	53.43 (14.81)	73.46 (21.96)

The mean difference between repeated BUA measures on the CUBA machine for the entire cohort was 0.315 dB/MHz, standard deviation 2.388 (t=5.2, p<0.0001). This is not a clinically meaningful difference, but statistically significant due to the large sample size. Similarly the mean difference between repeated BUA measures on the

QUS-2 machine (for the entire cohort) was -0.014 dB/MHz, standard deviation 2.885 (t=-0.1, p=0.9).

3.3.2 Reproducibility

Reproducibility statistics are given in Tables 3.2 and 3.3. Mean CUBA BUA (F=10.93, p<0.0001), mean CUBA VOS (F= 21.14, p<0.0001) and mean QUS BUA (11.92, p<0.0001) all decreased significantly with increasing age. The within-subject SEM for BUA was 1.6 dB/MHz in the CUBA and 2.0 dB/MHz in the QUS-2 and the within-subject coefficient of variation (wCV) for BUA was 3.1% for CUBA and 3.4% for QUS-2, neither value being significantly different between machines. Variation in SEM was independent of mean BUA and of age (r = -0.01, p = 0.78) as seen in figure 3.1. Limits of agreement between duplicate measurements of BUA ranged from –4.4 to 5.0 dB/MHz with the CUBA, and from –5.7 to 5.6 dB/MHz with the QUS-2.

For VOS (CUBA), the within-subject SEM was 5.0 m/s and the CV 0.3%. The within-subject SEM significantly increased as the mean VOS increased (r = 0.11, p < 0.001), but there was no significant correlation between SEM and age (r = 0.01; p = 0.89). The limits of agreement between two measurements ranged from -15 to 16 m/s.

Table 3.2.	Analysis of reproducibility of CUBA BUA, CUBA VOS, and QUS-2
BUA in an	elderly institutionalized population.

	<u>CUBA</u>		
Statistic and parameter	BUA (dB/MHz)	VOS (m/s)	BUA (dB/M
Mean ± SD	51.8 ± 22.4	1567.5 ± 46.9	58.6 ± 18.9
Error variance	2.6	25.2	4.2
"True" variance	378.7	1975.4	354.3
Coefficient of reliability	0.993	0.987	0.988
Coefficient of reliability			
(T scores)	0.10	0.12	0.16
Within-subject SD (SEM)	1.6	5.0	2.0
Coefficient of variation	3.1%	0.3%	3.4%
Limits of agreement	-4.4 to 5.0	-14.8 to 15.8	-5.7 to 5.6
LSC $(1.96 \times \sqrt{2}\sigma_i)$	4.5	13.9	5.7
	(8.6%)	(0.9%)	(9.7%)
$LSC = 1.96 \times \sigma_i \sqrt{2(1-\rho_i)}$	^a 1.3	5.4	1.6
	(2.4%)	(0.3%)	(2.7%)

^a: The correlation between two consecutive measurements was estimated to be 0.92 for BUA measurement and 0.88 for VOS.

LSC - least signfigicant change

Table 3.3. Mean, between-subjects standard deviation (bSD), within-subjectsstandard deviation (wSD) and within-subject coefficient of variation (wCV) stratifiedby age group.

Variable		<80	80-89	90+	All Ages
CUBA BUA	Mean*	59.6	52.2	46.3	51.9
	wSD	1.7	1.8	1.6	1.7
	wCV	2.8	3.4	3.4	3.3
	bSD	22.4	22.0	21.6	22.4
	bCV (%)	37.6	42.1	46.7	
CUBA VOS	Mean*	1579.8	1569.1	1557.4	1567.7
	wSD	5.0	5.9	5.2	5.5
	wCV	0.3	0.4	0.3	0.4
	bSD	46.7	47.6	43.8	46.9
	bCV (%)	3.0	3.0	2.8	
QUS-2 BUA	Mean*	65.2	57.2	55.6	58.5
	wSD	2.0	1.8	2.4	2.0
	wCV	3.1	3.2	4.3	3.5
	bSD	19.3	18.5	18.3	1.5
	bCV (%)	29.6	32.3	32.9	

*statistically significant, p<0.0001).

Figure 3.1a: Difference between duplicate measures for BUA in women measured by CUBA





Figure 3.1b: Difference between duplicate measures for BUA in women measured by QUS-2



3.3.3 Concordance between CUBA and QUS-2

BUA was measured using both machines in 684 subjects (Table 3.4). The correlation coefficient for BUA between the two instruments was 0.87 (p < 0.001). On average, BUA measured by the QUS-2 was significantly higher (p < 0.0001) than that measured by the CUBA by approximately 1.9 dB/MHz (standard deviation of the differences corrected for repeated measures 9.547 dB/MHz). QUS-2 BUA measures may range from 16.5 dB/MHz below CUBA BUA measurement to 20.3 dB/MHz above CUBA BUA (95% confidence interval).

Table 3.4. Co	omparison of BUA	(dB/MHz) and	T-scores measur	ed by CUBA and
QUS-2 instru	ments			

	CUBA	QUS-2	Difference (95% CI)
Males			
Number of subjects	175	175	
BUA (dB/MHz)	73.1±20.7	73.7 ± 21.4	0.6 (-1.0 to 2.2)
T-scores	-2.1 ± 1.2	-1.2 ± 1.6	0.9 (0.8 to 1.0)
Females			
Number of subjects	509	509	
BUA (dB/MHz)	51.1 ± 15.6	53.4 ± 14.8	2.3 (1.2 to 3.4)
T-scores	-2.3 ± 0.9	-2.8 ± 1.1	-0.4 (-0.38 to -0.50)

When stratified by gender, the difference remains significant in females (average difference 2.3 dB/MHz, 95% LoA –15 to 19 dB/MHz) but not seen in males (average difference 0.6 dB/MHz, 95% LoA –21 to 23dB/MHz). Nonetheless, when BUA is expressed relative to the young normal mean, the QUS-2 T-scores were significantly higher than the CUBA T-scores in males; the reverse was observed in females (Table 3.4). As a result, there was a poor agreement between the two instruments in the classification of subjects based on T-scores (Table 3.5).

	Males	Females
Cuba BUA T-scores		
Less than -2.5	74 (42.3%)	269 (52.9%)
Between -1 and -2.5	74 (42.3%)	184 (36.2%)
More than or equal to -1	27 (15.4%)	55 (10.8%)
US-2 BUA T-scores		
Less than -2.5	47 (26.9%)	331 (65.0%)
Between -1 and -2.5	49 (28.0%)	130 (25.5%)
More than or equal to -1	79 (45.1%)	48 (9.5%)
greement between		
vo instruments (Kappa)	0.44 ± 0.04	0.62 ± 0.03

Table 3.5. T-score classifications from BUA by CUBA and QUS-2 instruments

Product-moment correlation coefficient for BUA: 0.88 (p < 0.0001), and T-score BUA: 0.76 (p < 0.0001).

For example, in males, 42% had CUBA T-scores less than -2.5, while the same classification was found in 27% by QUS-2 T-scores. In contrast, more females are seen with QUS-2 T-scores less than -2.5 (65%) than with CUBA T-scores (53%). The Kappa coefficients for males and females were 0.44 ± 0.04 and 0.62 ± 0.03 , respectively.

3.3.4 Longitudinal Measurements

There were 56 residents with longitudinal measurements. During the 2.2 years (range: 2.0 to 2.3 yrs) of follow-up, the mean \pm SD change in BUA (measured by CUBA) was -1.9 \pm 8.3 dB/MHz (p = 0.09, not statistically significant), but percentage change from baseline was statistically significant at -5.2 \pm 16.5 % (p = 0.02). For VOS, the change was not statistically significant either in absolute units (0.8 \pm 21.5 m/s; p = 0.79) or in percentage terms (0.1 \pm 1.3 %; p = 0.76). Changes in QUS measures were not related to calcium, vitamin D supplementation or to bisphosphonate use in this small study. Eight of the 10 residents measured at 2 years who were on bone preserving medications or commenced these over the study period decreased in BUA (range -1.33 to -17.73 dB/MHz), one resident being maintained on calcium and vitamin D during the study increased by 29.92 dB/MHz.


As expected, the correlation between baseline and follow-up measurements was statistically significant for both BUA (r = 0.92; p < 0.001) and VOS (r = 0.85; p < 0.001). Using the criteria of least significant change with the serial correlation being

taken into account ($LSC = 1.96 \times \sigma_i \sqrt{2(1 - \rho_i)}$), any change in BUA of more than 1.3 dB/MHz and in VOS of more than 5.8 m/s, in either direction, can be considered statistically significant. Accordingly, 28% and approximately 60% of the changes in BUA were classified as "decreased" and "increased", respectively, while the remainder (12%) were classified as "unchanged". Similarly, 46% of the changes in VOS were regarded as a "significant increase" and 54% were regarded as "significant decrease in *both* BUA and VOS, while 18% had significant increase in both measurements (Figure 3.2).

	Increased	Decreased	Unchanged
Using $LSC = 1.96$	$5 \times \sigma_i \sqrt{2(1-\rho_i)} as$	criteria	
BUA (n; %)	16	34	7
	(28.1)	(59.7)	(12.3)
VOS (n; %)	26	31	0
	(45.6)	(54.4)	(0)
Using $LSC = 1.96$	$5 imes \sqrt{2} \sigma_i$ as criter	ia	
BUA (n; %)	9	21	27
	(15.8)	(36.8)	(47.7)
VOS (n; %)	12	13	32
	(21.1)	(22.8)	(56.1)

Table 3.6: Changes in BUA and VOS (N=56), measured by CUBA.

Figure 3.2: Percentage of individuals with significant change in BUA and VOS over 2.2 years using CUBA (N=56)



- a: Using $(LSC = 1.96 \times \sqrt{2}\sigma_i)$ as criteria for significant change
- b: Using $(LSC = 1.96 \times \sigma_i \sqrt{2(1 \rho_i)})$ as criteria for significant change

3.4 Discussion

The interpretation of any diagnostic tool requires a good understanding of the accuracy and reliability inherent in that test. This study has described the precision and reproducibility of two calcaneal ultrasound instruments in an elderly population, in order to estimate the prevalence of osteoporosis (Chapter 4) and assess bone loss over time in this group. Coefficients of variation for BUA for the CUBA ultrasound instrument (3.1%) and the QUS-2 instrument (3.4%) compare well with other machines which measure BUA, but are higher than for VOS or for BMD measured by DEXA (301). VOS appeared more sensitive to change than BUA with the LSC for BUA being 2.4% compared to 0.3% for VOS measurement, but longitudinally VOS was not seen to change in our population, with equivalent numbers of patients increasing as decreasing VOS. This has important implications for study design; choice of sample size and the most appropriate bone measurement will depend on the magnitude of change expected for each parameter measured and the power required in addition to machine precision. Our data suggests age has little influence on the precision and within subject variance of calcaneal ultrasound even in the very elderly, making it a useful tool for epidemiological studies of bone loss in an older population.

Statistically, the correlation in BUA between CUBA and QUS-2 instruments was high (r = 0.87, p<0.0001), however, there was a poor concordance between the two instruments. It is interesting to observe that BUA measured by CUBA was significantly higher than that measured by QUS-2, although the difference was found in women but not in men, probably due to the smaller sample size in the latter group. It is not clear why there was a systematic difference in BUA between the instruments, given the fact that they have virtually identical degree of precision. Nevertheless, the

discordance between the instruments raises the question of accuracy of BUA measurement.

Results of the present study also suggest that if QUS is used for the diagnosis of osteoporosis (or low bone mass) then standardisation across instruments must be addressed first. Between the two instruments evaluated in this study, there was a high discordance in the diagnosis of low bone mass. If the criteria of T-scores < -2.5 was used as a definition of "low bone mass" then QUS-2 was more likely to classify a women as "low bone mass" (65%) compared to CUBA (53%). On the other hand, in men, the prevalence of low bone mass was significantly higher in CUBA T-scores (42%) than in QUS-2 (27%). This suggests that the two instruments use systematically different referent databases, or different algorithms to compute the Tscores. Each instrument is programmed with its own specific normal range, the CUBA using a British normal population and the QUS-2 a US population; inherent differences between these groups may contribute to the different T-score classifications seen. It would seem from our comparative data for two quantitative ultrasound instruments, and the variable precision, reproducibility and sensitivity to change reported by other investigators using other machines, that the use of QUS for the assessment of osteoporosis is instrument specific and should not be generalised between models. Thuy et al (302) recently showed poor agreement between data-based and machine-based Tscore classifications (κ =0.65) in a large Vietnamese population-based study using the QUS-2 instrument. The use of T-scores as a means of classifying osteoporosis was initially described in post-menopausal women using DEXA BMD values at the hip (8) and is not necessarily generalizable to other measurement modalities. Machine-derived

75

T-scores for QUS must certainly be interpreted with caution, and can be a trap for the unwary when QUS is used in the clinical setting.

Over 2 yrs of follow-up more than 50% of subjects had significant changes in BUA. This change appears quite sensitive, because with BMD one might expect that almost 50% would experience no change. Individual changes could not be attributed to the use of bone preserving medications. Despite being a more precise and reproducible measure, VOS was not shown to change significantly over 2 years in this population, limiting its use in this setting. Although the sample size was small compared to the cross-sectional study, it should be sufficient to detect a difference in VOS which was more precise and reproducible than BUA. It is possible that VOS is measuring a different property of bone to BUA, which changes at a different rate with time in the elderly. This does not explain the discrepancies between the FREE longitudinal study of QUS and previous studies showing significant longitudinal changes in VOS. It should be noted that a different ultrasound method (the Achilles system) was used in the Schott study (105), and we have shown in our correlation analysis between the CUBA and QUS-2 methods there are quite large differences in absolute terms between different ultrasound machines. Moreover the sample size of both studies is modest. Importantly, although VOS or SOS is quite precise, it is acknowledged to be much less sensitive to change than BUA and the % differences in change between the two studies (0.1% vs 0.8%) are not large. The only identifiable selection bias in this sub-sample of 56 residents was the institution; as sequential patients who were enrolled in the FREE study were followed up at 2 years, all had lived in one of only 3 hostels at enrolment. Their baseline QUS values however did not differ from the FREE cohort as a whole. The FREE study is older and frailer than previous studies, and we see in

Chapter 4 that QUS does not change rapidly in the very old; it is possible that there is a differential rate of change between BUA and VOS in very old age. Certainly BUA is more informative for longitudinal studies in this population.

It is difficult to determine whether the statistically significant change in BUA of 5% from baseline is clinically important without prospective fracture data. Changes in DEXA measurements at both hip and spine of 5% confer a clinically important change in fracture risk, but this is not generalizable to other modes of bone density assessment. It does not fulfill the 'one standard deviation' drop in BUA which Bauer et al showed to correspond to a doubling in fracture risk (56). The effect was diluted by some residents increasing BUA over time. There was no identifiable factor which may have caused this effect, and it is likely that it is an example of regression to the mean.

Only a few previous studies have examined changes in QUS longitudinally. Schott et al (105) followed 88 post-menopausal women (mean age 63 years) over 2 years, showing a decrease in BUA of 1% (p=0.02) and a decrease in VOS of 0.8% (p=0.0001). This change in BUA approximated the long-term precision value, while the change in VOS was more meaningful at 5 times the precision for BUA. The derived parameter of stiffness decreased by 1.85%, approximating 2.5 times the precision, giving intermediate sensitivity to change. A larger cross-sectional study of 270 female nursing home residents (mean age 85 years) incorporated longitudinal data for 60 women after 1 year (101). This shorter study failed to show a significant decrease in BUA, whereas VOS reached statistical significance decreasing by 0.3% (p=0.001). The coefficient of variation for this study was 1.8% for BUA and 0.3% for

VOS. Other groups have shown QUS to be sensitive to change in high risk populations, including otherwise healthy patients with acute spinal injury (303), prolonged bed rest (304), and in children with rheumatic diseases (305). Daly et al (306) used QUS to look at bone changes with exercise, measuring calcaneal BUA and VOS over an 18-month period in elite male gymnasts and matched normoactive controls. Over 18 months, BUA increased significantly in the gymnasts from baseline whereas VOS did not increase in either group.

In an interventional clinical trial (108) BMD in postmenopausal osteoporotic women treated with alendronate increased by 5.0%, compared with a rather modest increase in VOS (0.7%) and BUA (1.4%) after two years. Change in BMD was poorly correlated with change in QUS. The minimum significant difference between two measurements was 0.8% for VOS and 5.6% for BUA, consistent with our findings. Among the QUS parameters, the authors concluded the calculated value of 'stiffness' showed the greatest longitudinal sensitivity, which was only slightly lower than BMD. Another study of the effect of anti-resorptive therapy on QUS parameters over 2 years in osteoporotic women (107) concluded that the minimum significant difference in BUA was 3 times greater than that for BMD. Sahota et al (106) showed a similar relationship between QUS and BMD over 4 years therapy with hormone replacement therapy in 60 early post-menopausal women. Therapy with calcitonin (109) gave a significant improvement in VOS over 2 years but no improvement in BUA. Conversely, BUA was sensitive to increases with vitamin D and calcium supplementation (110) over a similar time period in elderly institutionalized women who were vitamin D deficient, whereas VOS was seen to decrease. Studies of osteoporosis therapies must be interpreted with caution, as the effect of the

intervention may differ between therapies and therefore cannot be attributed to the method of measurement alone.

In conclusion, QUS measurements were highly reliable and sensitive to longitudinal change. However, the accuracy of BUA measurement could be problematic, and as a result, its use in the diagnosis of osteoporosis remains contentious.

CHAPTER 4

PREVALENCE OF OSTEOPOROSIS

IN THE FRAIL ELDERLY

PREVALENCE OF OSTEOPOROSIS IN THE FRAIL ELDERLY

4.1 Introduction

Osteoporosis, a disease characterised by skeletal fragility, is a major public health problem. Hip fractures are the most costly osteoporotic fracture to treat and are due to a fall or injury in over 90% of cases (307). Nursing home and hostel residents account for approximately 40% of total hip fractures (308) and as populations age, health care costs in this frail elderly group will increase substantially.

Skeletal bone quantity can be assessed by measurement of bone mineral density (BMD), usually by dual energy x-ray absorptiometry. BMD is considered the best predictor for risk of fracture, however its routine use in the frail and institutionalised older person is often impractical. Recently quantitative ultrasound (QUS), a less expensive and more portable technology, has been shown to predict risk of osteoporotic fracture in older women, independently of BMD (55;56) and may have particular use in a high risk elderly population. However there have been few studies of gender differences in QUS.

The aims of this study are as follows:

- To describe bone density as measured by QUS of the calcaneus in a large population of frail elderly residents of hostels and nursing homes
- (2) To identify differences in QUS between men and women
- (3) To show whether QUS changes with age similarly in men and women
- (4) To show whether QUS changes with age similarly in this frail elderly population as in previously studied groups

(5) To describe the relationship of QUS to risk factors for falling in this population.

4.2 Methods

All subjects enrolled in the FREE study (described in detail in Chapter 2) up until the end of June 2001 were included in this analysis. Broadband ultrasound attenuation (BUA) and velocity of sound (VOS) were measured at the left calcaneus in all participants using the same McCue CUBA Mark II ultrasound machine, and BUA measured in a smaller number using the Metra QUS-2 ultrasound machine. Each QUS measurement was repeated twice in the study population and the average of the two values used in the analysis.

The following clinical risk factors were assessed in all subjects at interview: age, gender, weight, height, activities of daily living status (Barthel Index) (261), cognitive function (Standardised Mini Mental Status Examination or SMMSE) (265), co-morbidities (by Resident Classification Scale graded 1 (lowest care) to 8 (highest level of care required)) (262), medications, falls risk and history of previous low trauma fracture (after age \geq 50). Falls risk was tested by assessments of quadriceps strength and static balance, according to published methods proven to predict falls and fractures in the elderly (169;245), explained in detail in Chapter 2.

QUS measurements in the elderly population were expressed as T-scores using the BUA and VOS means and standard deviations from a healthy control population. Controls comprised healthy twin pairs (a total of 264 males and 936 females) aged 18-85 years as previously reported (37). Only one twin from each identical pair, randomly selected, was included. Individuals with a history of underlying bone disease, past or present corticosteroid use, serious concomitant disease and weight over 100kg were excluded. T-scores were calculated using the mean and standard deviation compared to the control subjects aged 20-30 years, defined as representing 'young normals'.

Statistical analyses were carried out using the SPSS 11.0 statistical software package. Independent sample student's t-test was use to compare means between continuous variables, and Pearson's chi-squared statistic calculated to test differences between proportions. One-way analysis of variance (ANOVA) was used to evaluate univariate associations for continuous variables. Multiple linear regression was performed using a stepwise backward method. Comparison of regression lines was carried out using the method described by Armitage (309).

4.3 Results

By June 2001, 1193 residents from 52 randomly selected aged care institutions had been enrolled representing a 50% participation rate for eligible residents. This participation rate is only moderate, but expected and reflects the difficulty in obtaining informed consent from proxy informants in this population. Residents who required proxy consent were less likely to participate in the study than those able to give independent consent (27% vs 68%, p<0.0001). Responders and non-responders were identical with respect to mean age, sex distribution and time of residence in institutionalized care (mean differences not statistically significant, p>0.09).

The study sample comprised 294 men and 899 women. The men were on average 5 years younger than the women (mean \pm SE age 81.2 \pm 7 and 86.7 \pm 6 vs years respectively, p < 0.001). Summary findings according to sex and residential status (nursing home or hostel) are shown in Table 4.1.


Mean (SD)	Men		Wom	ien
Institution Type	Nursing Home	Hostel	Nursing Home	Hostel
Number of residents	124	170	460	439
Age (years)	82.5 (8.8)	81.5 (7.5)	86.2 (7.1)	81.5 (6.4)
Weight (kg)	67.3 (12.1)	73.9 (14.2)	55.7 (12.4)	58.8 (12.8)
SMMSE (score, 0-30)	15.9 (9.7)	24.4 (5.7)	15.3 (10)	24.1 (6.3)
EQ-5D (score, 7-18)	11.9 (2.9)	9.4 (2.1)	12.1 (2.8)	9.8 (2.3)
Quads Strength (kg)	21.7 (9.4)	27.3 (8.5)	15.9 (5.5)	15.9 (6.2)
Static Balance (score, 1-5)*	2.1 (1.2)	4.1 (1.1)	1.9 (1.1)	3.9 (1.2)
BUA CUBA (dB/MHz)	65.6 (24.1)	73.6 (20.0)	41.2 (18.8)	49.2 (16.9)
BUA QUS-2 (dB/MHz)	70.6 (22.8)	74.6 (20.1)	51.6 (15.6)	53.3 (14.8)
Prevalence of prior fracture	19.0%	15.7%	36.9%	35.7%
Length of stay (years)	2.41 (2.17)	3.9 (3.9)	2.42 (2.63)	4.44 (5.39)
25-OH-vitamin D (nmol/L)	21.4 (1.9)	26.2 (2.2)	16.9 (1.3)	17.3 (0.9)

Table 4.1. Characteristics of hostel and nursing home populations, divided by gender

*Static balance was measured on a scale of 1 to 5, where 1 represents 'not able to stand unsupported' and 5 represents 'good balance (able to stand for 30 seconds on an uneven surface unsupported)'.

In nursing home residents, men were younger (p < 0.002), heavier (p < 0.001), and had higher BUA and VOS (p<0.0001) than women (Table 4.1). There were no significant differences in length of stay or cognitive status. Similarly in hostel residents, men were younger (p < 0.001), heavier (p < 0.001), had better static balance (p=0.013) and higher BUA and VOS (P<0.0001) than women. There were no significant differences in SMMSE, but length of stay was longer in women than men in hostels (p < 0.051). Residents of nursing homes had not been institutionalized for as long as those in hostel accommodation (p<0.001), but no gender difference was seen. Gender remained significantly associated with BUA (p<0.0001), VOS (p<0.0001) and 25(OH)D levels (p=0.005) after adjusting for age and institution type. A history of previous low trauma fracture after age 50 was more common in women than men (p < 0.001) but did not differ by institution. Overall fracture prevalence was 32%; sites of fracture, some of which were multiple in certain subjects, were hip (11.7%), wrist (2.9%), humerus (2.7%), and other (16.0%). Mean BUA was 44 (± 18) dB/MHz and mean VOS 1553 (+39) m/s in subjects with a history of previous fracture compared to BUA 56 (+ 22) dB/MHz and VOS 1578 (+52) m/s in those without (p < 0.001 for both comparisons). Those individuals reporting previous hip fracture had significantly lower BUA measures than individuals reporting other fractures (mean difference 8.3 dB/MHz, p < 0.001) but VOS did not reach a statistically significant difference. Use of medications for osteoporosis was infrequent (bisphosphonates 2.3%, raloxifene 0.7%, hormone replacement therapy 2.8%, vitamin D 4% and calcium supplements 13.7%). The above results were unaffected if these subjects were excluded.


BUA measured by the CUBA was highly correlated with BUA measured by the QUS-2 ($R^2 = 0.74$, p<0.001), the latter reading on average 10% higher. There was no significant decline in BUA or VOS measured by either machine with age in men however in women BUA declined by 2-8-4.7% per decade (p<0.0001) and VOS by 1% per decade (p< 0.001) (Figure 4.1).



(a) women and (b) men

BUA = 85-0.46*Age; R² = 0.029





When compared to a healthy population of volunteers in Sydney aged 18-80 years (Figure 4.2), the data suggest that although the sex difference in QUS measures is

maintained into very old age, only minor loss occurs at the calcaneal site in BUA and VOS with very old age in both men and women. These results were obtained from a cohort measured over 10 years before, using a different software version for the ultrasound machine so it was decided not do a formal statistical comparison.

Figure 4.2: BUA changes with age, in a 'normal' healthy population of men and women and the institutionalized elderly.






Figure 4.3 shows the data from this elderly population in comparison to an aged cohort of female subjects measured as part of the Dubbo Osteoporosis Epidemiology Study (310) with the same CUBA machine. The DOES population comprises 452 subjects (age range 58-92 years). Similarly to the young normal population, comparison of the linear regression of age vs BUA for each group (DOES n = 452, FREE n = 1197) suggests a significant difference in intercept (mean \pm SE 126.4 \pm 8.9



• Dubbo Osteoporosis Epidemiology Study • FREE population

Figure 4.3 (b): BUA in institutionalized elderly women, compared with a cohort of independent older women taken from the DOES study (310), related to age, shown with separate regression lines.



vs 91.4 \pm 6.7, p<0.05) but not for the slope of the regression lines (mean slope \pm SE – 0.90 \pm 0.13 vs -0.531 \pm 0.078, p=0.1) suggesting a similar pattern of bone loss with age and considerable generalizability of the BUA data across different populations.

Compared to a young normal population sample measured in Sydney using a McCue CUBA II (37), the mean BUA T score at age 85 was -1.35 in men compared to -2.33 in women and at age 95 -1.72 versus -2.55 for men and women respectively. For VOS the mean T score at age 85 in men was -1.67 compared to -3.32 in women and at age 95 it was -1.95 versus -3.54 for men and women respectively. Compared to 186 healthy women aged 65-85 years living in the community, mean BUA was 1.2 standard deviations lower in the institutionalized females. Similarly compared to 37 healthy men aged 65-85 years, mean BUA was 0.8 standard deviations lower in the institutionalized males. BUA and VOS were significantly inversely related to Resident Classification Scale, a measure of comorbidities (p < 0.0001 respectively).

Univariate analyses showed there were significant correlations between BUA and VOS and body weight (p<0.0001) in both sexes. There were significant correlations in women but not men between BUA, VOS and quadriceps muscle strength (p < 0.0009 and 0.02 respectively). In multiple regression analysis, weight was the only significant predictor of BUA in men (Table 4.2). However in women, significant predictors of BUA included weight and history of previous fracture, age and type of institution (Table 4.2). When both men and women were included in the multivariate model, significant predictors of BUA values included gender, institution type, weight

and history of fracture, which together account for 38% of the variability of BUA.

Age did not contribute significantly to the model.

Table 4.2. Independent predictors of BUA in men and women.

Males

Model variables	β estimate	Standard error	t value	p value
Intercept	38.32	6.81		
Weight	0.46	0.09	4.90	< 0.0001

Females

Model variables	β estimate	Standard error	t value	p value
Intercept	40.99	9.68		
Age	-0.22	0.09	-2.47	0.014
History of fracture	-5.52	1.12	-4.94	< 0.0001
Weight	0.45	0.049	9.19	< 0.0001
Institution type	-6.89	1.19	-5.77	<0.0001

Pooled population

Model variables	β estimate	Standard error	t value	p value
Intercept	53.02	5.22		
Gender	-17.11	1.44	11.90	<0.0001
Institution type	-5.57	1.13	-4.94	< 0.0001
History of fracture	-5.34	1.07	-4.98	< 0.0001
Weight	0.47	0.043	10.86	<0.0001

4.4 Discussion

This study measured sex differences in QUS parameters in a frail elderly institutionalized population of men and women. It was carried out on the first 1000 residents enrolled in the FREE study, in late 2000 and 2001. Past history of minimal trauma fracture in this group was low compared to the total population at the end of the study (Chapter 2), but otherwise the subgroup is representative of the whole study. There are no clear reasons for this phenomenon; there were no notable changes in interviewers who might have elicited a more definitive response from residents. Standards of care in nursing homes and hostels are constantly under scrutiny with quality assurance programs. It is possible that the quality of medical record reporting had improved over the period of the study, resulting in a larger proportion of previous fractures being reported. Personal recall of previous fracture is thought to be reliable in the elderly (286;288), but it may be that the discrepancies seen are a reflection of inherent measurement error due to recall bias. The lower prevalence of reported previous low trauma fracture in this subgroup does not impact on the results of the ensuing study of QUS and gender differences in the frail elderly.

As expected, the prevalence of low bone density as measured by quantitative ultrasound was high in the elderly FREE cohort. Women had significantly lower bone density than men of a similar age, an expected finding as the female population is by definition post-menopausal and has therefore undergone a period of accelerated bone loss as a response to withdrawal of physiologic oestrogen. At advanced age, the rates of bone loss were seen to be similar between men and women suggesting hormonal changes have a less important role in this population. Although rates of BMD loss measured at the femoral neck have been reported to increase significantly in both sexes with advancing age (310), there have been few studies of other skeletal sites and most studies include only small numbers of very elderly men. This study suggests only minor loss occurs at the calcaneal site in BUA and VOS with advanced age in either sex. However one limitation of this study is its cross-sectional nature which allows only inferences about rates of loss. Another explanation would be that increased mortality associated with low bone mass has lead to a survivor bias for 'healthier' elderly subjects who are losing at a slower rate. This seems unlikely given the subjects were all 'institutionalized' and so were frail elderly. Of interest, the sex difference in QUS parameters is maintained into very old age.

A number of studies have examined QUS in older women, mainly in relation to its value as a predictor of osteoporotic fracture risk (55;56). Studies in elderly women have identified age, body composition and physical activity as determinants of QUS parameters (40;45), but there have been few studies of QUS in men, or indeed elderly men. One study compared QUS in 169 normal men and 210 normal women aged 20-80 years (41). BUA was higher in men than women at all ages but showed little decline in either sex after age 60. Another study of 224 men attending an osteoporosis clinic identified significant correlations between weight (and body mass index) and BUA, but no age related decline in QUS measures was observed, except in VOS in 76 fracture cases (27). In both these studies only a small number of subjects were aged greater than 70 years. Pluijm et al (58) measured BUA in 132 men and 578 women aged \geq 70 but did not report determinants of QUS or sex differences. Truscott (40) examined age-related trends in QUS in 2087 women aged from 16 to 93. No

94

decline in QUS measures was observed between the age bands 75-79 and 80-84, however there were a total of only 20 subjects in these groups.

When compared to an independently living population aged over 65 years (taken from the Dubbo study (310)), the frail elderly group were seen to lose bone at a similar rate. However, the baseline bone density was lower in the FREE study as seen by the different intercepts of regression lines for the two groups. This is logical, as the Dubbo group are by definition healthier and more mobile than those of the FREE study who require supportive care with activities of daily living, and it could be assumed that independently living individuals spend more time actively weight bearing (recognized as important to maintain bone density). Interestingly this has an effect on baseline but not rate of bone loss. It might be concluded that other factors, such as age and hormonal changes, are more important than exercise and weight bearing for maintaining bone density from any baseline.

There are currently few published studies of the prevalence of osteoporosis in the frail institutionalized elderly. Intuitively it would be expected that BMD and QUS should be low in this aged, frail population, but the practicalities of measuring BMD in this relatively frail group have precluded large studies. A recent North American study (18) of 1427 elderly female nursing home residents using distal radius BMD measured by single energy x-ray absorptiometry reported that 82% were osteoporotic (T-score < -2.5) and 54% fell under a T-score of -3.5.

It is also important to consider the meaning of a T-score derived from calcaneal ultrasound in terms of fracture risk. Faulkner et al (11) illustrated the differential classification of an individual as osteoporotic or not using T-scores derived from different techniques of bone mass estimation, including DXA, QCT and QUS at different skeletal sites. It is likely that a QUS-derived T-score of -2.5 is in fact related to a much higher fracture rate than a hip DXA-derived T-score of -2.5; one cannot use the two derivations interchangeably in the estimation of the prevalence of osteoporosis.

Prospective information on fracture incidence will be important to define appropriate cut-offs for QUS measures as they pertain to fracture risk in the frail institutionalized elderly population. This information will then allow more definitive recommendations about the usefulness of BUA and/or QUS in assessing osteoporosis in the frail elderly.

CHAPTER 5

DISTRIBUTION, DETERMINANTS AND ASSOCIATIONS OF 25-OH-VITAMIN D LEVELS IN AGED CARE FACILITIES

DISTRIBUTION, DETERMINANTS AND ASSOCIATIONS OF 25-OH-VITAMIN D LEVELS IN AGED CARE FACILITIES

5.1 Introduction

Vitamin D deficiency is a silent epidemic, affecting men and women both in Australia (311-313) and around the globe (311;314-317). Its prevalence increases in older populations as mobility and general health decline (318-320), related to inadequate sunlight exposure, poor diet and increasing renal insufficiency (321;322). Institutionalized elderly persons are at particular risk for low 25-OH-vitamin D levels (323-327), but there is less known about the relationships between vitamin D, gender, season and biochemical measures of calcium homeostasis in this frail elderly population.

The aims of this study are as follows:

1: To describe the distribution of serum 25-OH-vitamin D levels in elderly men and women living in residential aged care facilities (hostels and nursing homes) in northern Sydney.

2: To determine if serum 25-OH-vitamin D levels differ with age, gender, residential status or season.

3: To describe the association between 25-OH-vitamin D levels and other bone-related biochemical parameters, including serum creatinine (as a surrogate for renal function), serum parathyroid hormone (PTH), corrected serum calcium and serum albumin levels. What is the independent influence of age, gender and institution on these associations?

4: To define hypovitaminosis D by relating 25-OH-vitamin D levels to subclinical secondary hyperparathyroidism.

5: To identify the key independent predictors of serum 25-OH-vitamin D level (continuous) and hypovitaminosis D (categorical) in the frail elderly.

6: To estimate the prevalence of vitamin D supplementation in this population, including the use of vitamin D supplements, analogues, combinations of these or other bone active medications.

7: To identify the determinants of the use of vitamin D supplementation. Is the use of vitamin D supplements associated with age, gender or residential status? Does this supplementation impart a benefit on circulating 25-OH-vitamin D levels?

This chapter will describe the distribution and determinants of serum 25-OH-vitamin D levels in an elderly population living in aged care facilities, and identify the significant biochemical interrelationships with serum 25-OH-vitamin D in this population.

5.2 Methods

5.2.1 Subjects

Each resident enrolled in the FREE study (a full description of the FREE study population methods is given in Chapter 2) was asked to give a blood sample for 25-OH-vitamin D and other biochemical analyses. Sixty-three percent of the cohort (268 men and 1000 women) agreed. Remaining study participants or their carers declined venesection and a small percentage of venesections were technically overly difficult and abandoned after one failed attempt. Carer consent was obtained where appropriate. Current medications were recorded from interview and verified by medical records and medication charts, including prescription drugs and over-thecounter vitamin preparations. Individuals with biochemical primary hyperparathyroidism (15 residents with an elevated parathyroid hormone and associated hypercalcaemia) or major renal insufficiency alone (6 individuals with a serum creatinine >300 μ mol/L) were excluded from the analyses, resulting in a final study population of 979 women and 264 men.

5.2.2 Biochemical measurements

Serum 25-hydroxyvitamin D (25OHD), intact parathyroid hormone (PTH), calcium (adjusted for circulating albumin levels), creatinine and inorganic phosphorus levels were measured in 1268 residents using methods described in Chapter 2. Creatinine clearance was calculated using the Cockcroft-Gault formula. For males:

Creatinine clearance (ml/min) = $\frac{(140 - age)*weight}{72*serum creatinine (mg/100ml)}$

The same calculation is used in women, adjusted by x 0.85.

5.2.3 Statistical analysis

Analyses were carried out using the SPSS 11.0 statistical software package. Independent sample student's t-test was use to compare means between continuous variables (log transformed if required to confer normality), and Pearson's chi-squared statistic calculated to test differences between proportions. Vitamin D levels were treated both as a continuous variable, and divided into categories to define 'hypovitaminosis D'. One-way analysis of variance (ANOVA) was used to evaluate univariate associations for continuous variables. Logistic and multiple linear regressions were performed using a stepwise backward method.

5.3 Results

5.3.1 Baseline demographics

Nine hundred and seventy-nine women and 264 men were included in the analysis. The women were on average 3.7 years older than the men (p<0.0001) and were less mobile, being more likely to require a walking aid (usually a frame). Demographics of the group are shown in Table 5. 1.

Table 5.1: Demographics of institutionalized elderly cohort with biochemical measures.

		Ger	nder	Test	
		Male	Female	statistic	Significance
Age (mean)		83.10	86.76	t=7.2097	p<0.0001
Institution	Hostel	136 (48.5%)	560 (57.2%)	χ ² =2.73	p=0.11
	Nursing Home	128 (51.5%)	419 (42.8%)		
Mobility	Walks unaided	94 (35.9%)	287 (29.6%)	χ ² =11.65	p=0.009
	Uses a stick	65 (25.0%)	213 (22.0%)		
	Uses a frame	59 (22.5%)	324 (33.4%)		
	Wheelchair	44 (16.4%)	146 (15.0%)		
On Vitamin D		16 (6.1%)	115 (11.7%)	χ ² =7.13	p=0.007
supplements					

Descriptive statistics for biochemical variables are shown in Table 5.2. Fourteen individuals exhibiting biochemical primary hyperparathyroidism were excluded from analysis. This group had serum PTH levels ranging from 72.3 pg/mL to 602 pg/mL (median 108 pg/mL) and corrected calcium levels between 2.57 mmol/L and 3.08

mmol/L (median 2.66 mmol/L). Five individual with PTH levels above 400 pg/mL but no available calcium measures were also excluded – this did not significantly change any of the calculations. Finally, six individuals with renal insufficiency (creatinine values greater than 300 µmol/L) associated with elevated PTH but normal corrected calcium levels were also excluded.

+/- standard error of the mean (range), median.						
	Hostel		Nursing	Home		
	Male	Female	Male	Female		

Table 5.2: Biochemical measures grouped by gender and institution, given as mean

	Но	stel	Nursing Home		
	Male	Female	Male	Female	
25-OH-vitamin D	35.8 +/- 1.48	29.8 +/- 0.64	29.5 +/- 1.59	25.7 +/- 0.80	
(nmol/L)	(5 - 92), 34	(2 – 104), 27	(5 – 103), 26	(2 – 101), 21	
PTH (pg/mL)	57.5 +/- 3.58	72.8 +/- 2.04	65.2 +/- 3.39	71.2 +/- 2.44	
	(12.6 – 238.0), 45.5	(4.6 – 337.0), 61.2	(12.2 – 209.0), 57.3	(5.6 – 309.0), 55.8	
Corr. Calcium	2.34 +/- 0.017	2.36 +/- 0.009	2.34 +/- 0.018	2.36 +/- 0.010	
(mmol/L)	(1.96 - 2.91), 2.33	(1.87 – 2.73), 2.35	(2.01 – 2.60), 2.36	(1.86 – 2.68), 2.37	
Inorg. phosphorus	1.15 +/- 0.021	1.25 +/- 0.012	1.13 +/- 0.020	1.20 +/- 0.012	
(mmol/L)	(0.75 - 1.66), 1.14	(0.77 – 2.11), 1.25	(0.76 – 1.67), 1.11	(0.59 – 1.77), 1.21	
Albumin (g/L)	42.0 +/- 0.38	41.7 +/- 0.23	40.1 +/- 0.44	39.3 +/- 0.22	
	(32 – 49), 42.0	(30 – 58), 42.0	(32 – 50), 40	(26 – 47), 39	
Creatinine	70.0 +/- 21.4	44.7 +/- 18.6	57.4 +/- 21.3	44.0 +/- 18.4	
clearance	(33.2 – 131.1), 58.0	(16.1 – 140.8), 54.0	(22.9 – 124.4), 41.4	(11.8 – 147.3),	
(µmol/L)				40.6	

Nine hundred and sixty-two residents (77.4%) had low 25-OH-vitamin D levels when compared to the laboratory normal range (39 - 140 nmol/L). Serum parathyroid hormone was low in 103 residents (8.3%) and high in 508 individuals (40.9%) (normal range 23.7 – 66.2 pg/mL). Hypocalcaemia was seen in 38 residents (7.2%) and hypercalcaemia in 22 (4.2%), (normal range 2.15 - 2.55 mmol/L). Inorganic phosphorus was elevated in 136 (25.7%) individuals (normal range 0.6 - 1.3 mmol/L). Albumin was low in 181 (34.2%) and elevated in two individuals (normal range 40 - 50 g/L). One hundred and twenty men (94.5%) had a reduced creatinine clearance (normal range in males 97 - 137 ml/min), and 399 women (97.6%) had a creatinine clearance which fell below the normal range for females (88 – 128 ml/min).

5.3.2 Distribution of 25-OH-vitamin D levels

Table 5.3 shows the prevalence of 'hypovitaminosis D' in men and women, using different cut-off values previously described in the literature concerning 25-OH-vitamin D in elderly populations (322;328;329). The majority of both men and women had low 25-OH-vitamin D levels independent of which cut-off value is used to define 'hypovitaminosis D'.

Gender	25-OH-vitamin	Hypovitaminosis	Mean age	Mean vit D	Mean PTH
	D cut-off	D (number, %)	(years)	(nmol/L)	(pg/mL)
Male	Vit D < 25	109 (42.9%)	83.31	17.7	72.9
	Vit D < 30	139 (54.4%)	83.12	19.9	68.5
	Vit D < 39	187 (75.4%)	83.71	23.7	65.9
	Vit D < 50	217 (85.3%)	83.34	26.0	63.1
	Vit D 25-50	108 (42.4%)	83.38	34.5	53.3
	Low quartile < 18	46 (15.5%)	83.87	12.9	70.9
	Low tertile < 20	62 (24.6%)	83.67	14.4	71.4
	Low quintile < 17	38 (13.9%)	82.29	12.0	68.3
Female	Vit D < 25	525 (57.2%)	87.00	17.0	78.2
	Vit D < 30	643 (70.2%)	86.86	19.0	76.7
	Vit D < 39	732 (84.7%)	86.92	21.6	76.8
	Vit D < 50	884 (94.2%)	86.81	24.3	73.5
	Vit D 25-50	359 (37.0%)	86.60	35.0	66.6
	Low quartile < 18	267 (25.6%)	86.65	12.8	83.6
	Low tertile < 20	336 (36.9%)	86.63	14.0	80.5
	Low quintile < 17	232 (21.2%)	86.72	12.2	80.7

Table 5.3: Classification of residents with low 25-OH-vitamin D levels

5.3.3 Determinants of serum 25-OH-vitamin D levels

Serum 25-OH-vitamin D and PTH were not normally distributed (Figure 5.1), and so have been log transformed to allow comparisons as continuous variables. Serum 25-OH-vitamin D decreased with increasing age in the total cohort (t = -2.71, p = 0.007),

but did not reach statistical significance in either males or females alone. Serum 25-OH-vitamin D levels were higher in hostel accommodation than in nursing home residents in both males (mean difference 6.24 nmol/L, 95% confidence interval 1.94 - 10.94, p = 0.005) and females (mean difference 4.10 nmol/L, 95% confidence interval 2.13 - 6.08, p<0.0001). Males had significantly higher 25-OH-vitamin D levels than women, independent of institution. Serum 25-OH-vitamin D levels were significantly associated with the use of a walking aid (F=10.73, df=3, p<0.0001), being highest in those residents able to walk independently or with a stick, and lowest in those who use a wheelchair. There was no association between current smoking and 25-OH-vitamin D (mean 25-OH-vitamin D levels in smokers 27.4 nmol/L compared to 27.2 nmol/L in non-smokers, t=0.214, p=0.9).




Figure 5.1: Distribution of serum 25-OH-vitamin D and PTH for males and females.

There were a similar proportion of men and women venesected each season, shown in Table 5.4 (χ^2 =4.73, df=3, p=0.19).

	Gender				
Season	Male	Female	Total		
Spring	71 (26.9%)	264 (27.0%)	335		
Summer	73 (27.7%)	231 (23.5%)	304		
Autumn	42 (15.9%)	210 (21.5%)	252		
Winter	78 (29.5%)	274 (28.0%)	352		
Total	264	979	1243		

 Table 5.4:
 Number of residents recruited each season for men and women.

When those individuals taking vitamin D supplements are excluded from the analysis (n=131), 25-OH-vitamin D levels were significantly associated with season in both males (F=4.76, df=3, p=0.003) and females (F=6.46, df=3, p<0.0001). Levels were lowest in winter in both sexes (Table 5.5), but males had an earlier peak in spring/summer compared to an autumn peak in women (Figure 5.2).

	Gender		Mean	95% Confidence	
Season	Male	Female	Difference	Interval	Significance
Spring	34.91 (2.017)	24.84 (0.848)	10.07	5.72 - 14.42	p<0.0001*
Summer	34.77 (2.259)	26.80 (0.932)	7.97	3.11 - 12.82	p=0.002*
Autumn	28.33 (1.879)	29.21 (1.071)	-0.89	-5.21 - 3.43	p=0.7
Winter	26.55 (1.815)	23.93 (0.799)	2.61	-1.37 - 6.55	p=0.2

 Table 5.5:
 Mean 25-OH-vitamin D level (standard error of the mean) for each season

*statistically significant difference between males and females, p<0.05.




Figure 5.2: Changes in 25-OH-vitamin D levels with season in males and females.

5.3.4 Serum 25-OH-vitamin D and associated biochemical parameters

Males had significantly higher creatinine levels and lower PTH and phosphate levels than women, independent of institution (p<0.05). Using the laboratory-derived upper limit of normal for creatinine clearance, 120 men (94.5%, normal range in males 97 – 137 ml/min), and 399 women (97.6%, normal range 88 – 128 ml/min) had a degree of renal insufficiency. There was no statistically significant difference in PTH levels between institution type in either gender (p>0.1). Serum albumin was lower in nursing homes than in hostels in both males and females (p<0.001), and phosphate levels were higher in hostels than nursing homes in females (p=0.02) but not significantly different between institutions in males (p>0.5). Creatinine clearance was not significantly different between institutions in either males (p=0.4) or females (p=0.7). Albumin decreased with increasing age in both males and females (p<0.05), and PTH increased with age (p<0.005). Creatinine clearance decreased with age in both males and females (p<0.0001), and calcium and phosphate were not related to age in either gender (p>0.2).

Parathyroid hormone levels increased as 25-OH-vitamin D decreased in both men and women (Figure 5.3) as would be expected biologically. There was not a clear level of serum 25-OH-vitamin D below which a rise of PTH is seen (secondary hyperparathyroidism) which might support a definition of hypovitaminosis D. Log transformation of 25-OH-vitamin D, iPTH or both does not reveal any point of inflection or level of 25-OH-vitamin D above which iPTH levels might increase exponentially.





Figure 5.3: Parathyroid hormone levels related to 25-OH-vitamin D.

However when 25-OH-vitamin D was split into quintiles, mean PTH levels were seen to rise between the third and fourth quintiles (Figure 5.4), representing a mean 25-OH-vitamin D level of 29 nmol/L, supporting the use of either 25 or 30 nmol/L as an appropriate cut-off for defining hypovitaminosis D in this frail elderly population.

Figure 5.4: Mean PTH values relating to 25-OH-vitamin D quintiles.



Examination of each of the quintiles for 25-OH-vitamin D does not reveal the serum levels to be normally distributed within each quintile. When the median iPTH values are plotted against 25-OH-vitamin D quintiles, a lower cut-off level of 22 nmol/L is seen (Figure 5.5).




Figure 5.5: Median PTH values relating to 25-OH-vitamin D quintiles

Univariate analysis corrected for age showed 25-OH-vitamin D levels to be significantly and inversely associated with PTH in men (t= -3.630, p<0.0001) and in women (t= -3.718, p<0.0001). In females, 25-OH-vitamin D was associated with creatinine clearance (t= -2.901, p=0.004) and with albumin (t=2.429, p=0.016) but there was no association observed in the smaller sample of males (p>0.2). Serum 25-OH-vitamin D was not associated with calcium or phosphate in either gender (p>0.15).

5.3.5 Independent predictors of 25-OH-vitamin D levels

All significant univariate parameters were entered into a multiple regression model for men and women separately (Table 5.6). With multiple regression modeling, lower vitamin D levels in men were associated only with higher PTH and the use of a walking aid. Lower creatinine clearance, higher PTH, more advanced age and the use of a walking aid were significant independent predictors of low vitamin D levels in females. When the total cohort is examined, female gender, reduced mobility, higher PTH, increasing age, cooler season and lower creatinine clearance were independently related to lower 25-OH-D levels.

(Constant) Ln PTH Walking aid	4.264 -0.178	0.217	19.653	< 0.0001
	-0.178	0.053		
Walking aid		0.055	-3.331	0.001
-	-0.116	0.029	-3.951	<0.0001
(Constant)	5.112	0.545	9.381	< 0.0001
Age	-0.011	0.005	-2.056	0.04
Ln PTH	-0.149	0.053	-2.812	0.005
Cr. clearance	0.007	0.002	-3.261	< 0.0001
Walking aid	-0.086	0.028	-3.045	0.003
(Constant)	5.090	0.479	10.625	< 0.0001
Age	-0.009	0.005	-2.000	0.046
Gender	-0.151	0.064	-2.367	0.018
Ln PTH	-0.141	0.046	-3.064	0.002
Cr clearance	-0.006	0.002	-3.386	0.001
Walking aid	-0.100	0.024	-4.158	<0.0001
Season	0.052	0.024	2.205	0.028
	Constant) Age En PTH Cr. clearance Walking aid Constant) Age Gender En PTH Cr clearance Walking aid	Constant) 5.112 Age -0.011 In PTH -0.149 Cr. clearance 0.007 Walking aid -0.086 Constant) 5.090 Age -0.009 Gender -0.151 In PTH -0.141 Cr clearance -0.006 Walking aid -0.100 Season 0.052	Constant) 5.112 0.545 Age -0.011 0.005 Ln PTH -0.149 0.053 Cr. clearance 0.007 0.002 Walking aid -0.086 0.028 Constant) 5.090 0.479 Age -0.009 0.005 Gender -0.151 0.064 Ln PTH -0.141 0.046 Cr clearance -0.006 0.002 Walking aid -0.100 0.024 Season 0.052 0.024	Constant) 5.112 0.545 9.381 Age -0.011 0.005 -2.056 Ln PTH -0.149 0.053 -2.812 Cr. clearance 0.007 0.002 -3.261 Walking aid -0.086 0.028 -3.045 Constant) 5.090 0.479 10.625 Age -0.009 0.005 -2.000 Gender -0.151 0.064 -2.367 Ln PTH -0.141 0.046 -3.064 Cr clearance -0.006 0.002 -3.386 Walking aid -0.100 0.024 -4.158 Season 0.052 0.024 2.205

 Table 5.6:
 Independent determinants of serum 25-OH-vitamin D

Excluded variables: In males, season, age; in females, season, institution, albumin; in total cohort, institution, albumin.

Independent predictors of hypovitaminosis D, defined as a serum 25-OH-vitamin D level less than or equal to 30 nmol/L in this population, are shown in Table 5.7. As for the continuous variable 25-OH-vitamin D, hypovitaminosis D is predicted by higher serum PTH values, use of a walking aid and lower creatinine clearance. In men, use of a walking aid, PTH and creatinine clearance remain independently associated with vitamin D deficiency. Season is an important predictor of hypovitaminosis D in women, along with use of a walking aid and PTH levels.

 Table 5.7: Independent determinants of hypovitaminosis D (25-OH-vitamin D <= 30 nmol/L)</th>

	Variable	Wald	df	Signif.	Likelihood	95% confidence
		statistic		(p)	ratio (LR)	interval for the LR
Males	Walking aid	3.35	1	0.07	1.02	N/A
$(R^2=0.18)$	ln(PTH)	5.30	1	0.002	2.32	1.13 – 4.74
	Cr clearance	3.35	1	0.07	1.02	1.00 - 1.05
Females	Walking aid	19.48	3	0.0001	N/A	N/A
$(R^2=0.10)$	Season	12.47	3	0.006	N/A	N/A
	ln(PTH)	5.37	1	0.02	1.67	1.08 - 2.57
Whole	Gender	9.25	1	0.002	0.47	0.28 - 0.76
cohort	Walking aid	28.39	3	0.0001	N/A	N/A
$(R^2=0.11)$	ln(PTH)	11.39	1	0.001	1.87	1.30 - 2.68
	Cr clearance	5.77	1	0.02	1.02	1.00 – 1.03

Excluded variables: In males, season, age, institution; in females, age, institution, creatinine clearance, albumin.

5.3.6 Vitamin D Supplementation

The use of vitamin D supplementation and other bone-active medications was low overall in this population (Table 5.8). There was not a statistically significant difference between men and women in the use of mixed vitamin supplements containing either vitamin D or calcium; however the use of specific bone-targeted compounds including calcium alone, vitamin D analogues (calcitriol) and bisphosphonates was more frequently seen in females than males. A small number of individuals were taking both vitamin D supplements and analogues.

Table 5.8:	Frequency of bor	ne-active medication	on use among mer	n and women living
in aged car	e facilities.			

Medication	Male	Female	Test statistic	Significance
Vitamin D analogue	8 (3.0%)	91 (9.3%)	$\chi^2 = 11.113$	p<0.0001*
Vitamin supplements containing D	12 (4.5%)	76 (9.6%)	χ ² =3.272	p=0.079
Any vitamin D preparation	16 (6.1%)	115 (11.7%)	χ ² =7.13	p=0.007*
Calcium	13 (4.9%)	94 (9.6%)	χ ² =5.782	p=0.018*
Vitamin supplements with calcium	21 (8.0%)	117 (12.0%)	$\chi^2 = 3.365$	p=0.077
Hormone therapy	1 (0.4%)	33 (3.4%)	χ ² =6.696	p=0.005*
Bisphosphonates	3 (1.1%)	57 (5.8%)	χ ² =9.938	p=0.001*

*statistically significant difference, p<0.05

Supplementation was equally distributed between hostel and nursing home accommodation (χ^2 =3.85, p=0.05). Residents taking vitamin D preparations were much more likely to be taking concomitant calcium supplements (χ^2 =235.0, p<0.0001). Age did not appear to influence the use of vitamin D compounds (Table

5.9). Mean 25-OH-vitamin D levels were 45.2 nmol/L in the supplementation group, which was significantly higher than 27.2 nmol/L for the group without supplementation (mean difference 18.0 nmol/L, 95% CI for the difference 14.0-22.0, p<0.0001), although still below the 50 nmol/L cut-off recommended for vitamin D sufficiency by many authors. This significant difference is maintained in men and women (Table 5.9).

Table 5.9: Comparison of residents on vitamin D supplementation at baseline with

 those not receiving supplements (mean (sd)).

	Male		Female	
	Vit D Suppl	No Suppl	Vit D Suppl	No Suppl
N	16 (6.1%)	248 (93.9%)	115 (11.7%)	864 (88.3%)
Age (years)	81.16 (7.246)	83.23 (7.727)	86.85 (5.174)	86.75 (6.518)
25-OH-vitamin D (ng/mL)	55.56 (24.147)	31.28 (16.505)*	43.71 (21.944)	25.99 (13.388)*
PTH (pg/mL)	46.24 (32.771)	62.23 (43.574)	50.36 (45.016)	74.98 (48.686)*
Corrected calcium (mmol/L)	2.379 (0.0946)	2.335 (0.1407)	2.440 (0.1267)	2.349 (0.1330)*
Creatinine (µmol/L)	119.30 (39.721)	100.81 (26.539)	90.05 (28.984)	89.34 (28.040)
Albumin (g/L)	40.6 (3.44)	40.3 (1.32)	41.1 (3.45)	40.8 (2.96)

*significantly different to supplemented group, p<0.001

Corrected calcium levels were significantly higher in those individuals receiving vitamin D supplements (mean difference 0.082, 95% CI 0.045-0.119). Mean PTH levels were significantly lower in the supplementation group, 49.9 pg/mL compared to 72.1 pg/mL in those not receiving supplements, mean difference 22.3 pg/mL (95% CI 14.3-30.3, p<0.0001). When men and women are considered separately, corrected calcium remains higher and PTH lower in the supplementation group although this

does not reach statistical significance in males (p=0.33 and p=0.12 respectively), likely due to smaller numbers of male participants in the study. Albumin levels were similar between supplemented and non-supplemented residents (males, t=0.29, p=0.77; females, t=0.55, p=0.58).

5.4 Discussion

Sydney is situated at 33 degrees 55' south of the Equator, with average temperatures in summer of 23 degrees Celsius and 17 degrees Celsius in winter. At this latitude, similar to Buenos Aires, there is sufficient sunlight to provide adequate UV exposure for vitamin D synthesis in the skin of healthy individuals all year round (330), with an average of 340 sunny days each year. Despite this, serum 25-OH-vitamin D levels were almost universally low in this cohort of frail elderly people living in residential aged care facilities in northern Sydney. Similarly, a cross-sectional study of 99 elderly men and women conducted in Melbourne, Australia (324) showed that 52% of residents of the aged care facilities studied had 25-OH-vitamin D levels below the normal range, compared to 77% in this more northerly group. Equally low levels of vitamin D in aged care facilities have been described elsewhere (323;326;327), which would suggest that elderly persons living in such facilities are not benefiting from sunlight-derived vitamin D.

Males had higher 25-OH-vitamin D levels than women independent of season or institution, as has been shown previously (331). Residents in hostel accommodation had higher 25-OH-vitamin D and higher albumin levels than their counterparts in nursing homes. If one considers that serum albumin is a reasonable measure of general nutrition and morbidity (332-334), it is logical to infer that differences in vitamin D status between institutions are likely to reflect the poorer mobility and general health of nursing home residents rather than inadequate nutrition. Nevertheless, nutritional intake may confound the relationship between 25-OHvitamin D and residential status, when you consider that poor nutritional intake impacts on general health and mobility. It might also be that poor general health was the primary event, and subsequent reduction in dietary intake is seen. It would be interesting to examine the relationship of 25-OH-vitamin D and residential status when variables such as dietary intake are also considered.

Sunlight exposure is important for maintaining adequate 25-OH-vitamin D levels. Melin et al showed prospectively that seasonal variation of 25-OH-vitamin D levels requires three or more hours per week of sunlight exposure during summer months in Scandinavia (335), and that 25-OH-vitamin D levels measured in autumn reflect this exposure during summer. Individuals living in aged care facilities are less mobile than those living independently, and therefore less likely to get outdoors, contributing to an increased prevalence in hypovitaminosis D (323;326) which is not compensated by dietary intake (327). It has previously been shown that seasonal variation of 25-OH-vitamin D levels is less marked in residents of aged care facilities than in independently-living elderly people (325;336). Nevertheless, we have seen a distinct seasonal variation of 25-OH-vitamin D levels in this institutionalized group, with lowest levels occurring in winter as has been shown in previous cross-sectional (337-340) and longitudinal studies (341) in independently-living elderly populations. Serum 25-OH-vitamin D peaks during the spring/summer in men and in autumn in women; this difference might be explained by the poorer mobility of the female group, resulting in fewer trips outside and hence a slower accumulation of sunlight exposure over the warmer months.

Age-related changes in biochemistry were seen as expected, with increasing age conferring lower albumin and 25-OH-vitamin D and higher PTH levels in both genders. Creatinine clearance was seen to decrease with age in both males and

females as expected. Overall, biochemical changes in this frail elderly group did not behave differently to expected changes in any older population. It was of concern to note that although general nutrition was good in two thirds of the group, reflected by normal albumin values, 35% of residents of hostels and nursing homes have suboptimal albumin levels. This proportion of nutritional deficiency warrants further investigation and appropriate intervetional program to improve general nutrition.

Serum 25-OH-vitamin D levels were inversely associated with PTH levels as expected, reflecting secondary hyperparathyroidism occurring in response to low vitamin D stores. Creatinine clearance (as a surrogate for renal function) was an important determinant of 25-OH-vitamin D in women but not in men; it is possible that this was due to insufficient numbers of male participants to confer statistical significance. It is also possible that creatinine clearance is in part measuring general debility, which then confers poorer mobility, reduced sunlight exposure and subsequently lower vitamin D acquisition in women than in men.

There has been much debate about the most appropriate definition of hypovitaminosis D, with cut-off values previously quoted from as low as 8-10 nmol/L (342) up to 50-80 nmol/L (314;322;343;344). Differences in 25-OH-vitamin D assays might account for some of this variability (345;346), and so associating low 25-OH-vitamin D levels with other parameters such as PTH is important to assess clinical significance. The most commonly used cut-off is below 30 nmol/L (315;318;329;339;347;348), on the basis that at this level of 25-OH-vitamin D a mild increase in serum parathyroid hormone is seen in response to subclinical hypovitaminosis D (329) and so has clinical significance. Our laboratory uses a standard cut-off of 39 nmol/L, being the

lower end of the 'normal' range in healthy adults. This study however supports the use of 22-30 nmol/L as a cut-off for defining hypovitaminosis D in the frail elderly living in aged care facilities. Nevertheless, thinking of 25-OH-vitamin D on a continuum instead of in discrete terms may be more appropriate when assessing a patient's risk for comorbidities such as osteoporosis, fracture and general health.

This study showed creatinine clearance and PTH are both independent predictors of 25-OH-vitamin D levels in females, those individuals with lower creatinine clearance and higher PTH values having significantly lower 25-OH-vitamin D levels. Creatinine clearance was not an independent predictor of vitamin D in males, again most likely due to insufficient numbers of males in this study. This relationship may well be confounded by muscle bulk in men, with higher baseline creatinine levels in men obscuring any effect of renal insufficiency on 25-OH-vitamin D levels. Mobility, as measured by the use of a walking aid, was also an independent predictor of 25-OH-vitamin D; both men and women who were independently mobile or used a stick had higher 25-OH-vitamin D levels than those using a frame or wheelchair. If this is interpreted as a surrogate for sunlight exposure, then those individuals who are healthier (higher creatinine clearance and higher albumin in women, hostel accommodation in men) and more mobile are more likely to obtain vitamin D from direct sunlight, and less likely to develop secondary hyperparathyroidism.

Vitamin D supplementation was infrequent in the FREE study, introducing a potential for selection bias. Nevertheless, vitamin D supplementation was seen to appropriately increase serum 25-OH-vitamin D levels in this elderly group, with a mean difference in levels between residents on supplements and those not taking vitamin D of

18mnol/L, bringing those individuals on treatment just above most cut-offs for hypovitaminosis D. It has been suggested that the elderly require higher doses of vitamin D to overcome hyperparathyroidism related to increasing creatinine levels (349). With a well recognized relationship between low 25-OH-vitamin D levels and a higher risk of osteoporosis and ensuing fracture, the low rate of vitamin D supplementation seen in this frail elderly population was disappointing. Only 10.5% of residents (n=131) were receiving vitamin D supplementation or vitamin D analogues, leaving 60% (n=741) of the total cohort with untreated low 25-OH-vitamin D levels (<30 nmol/L). Despite taking a wide variety of preparations and doses for unspecified indications, those on supplements had higher 25-OH-vitamin D levels and lower PTH levels suggesting appropriate biochemical responses to such therapy. Vitamin D supplementation has been shown to improve bone mineral density (350-352), muscle strength (227;230) and falling rates (230), and is a simple and economical intervention which might improve quality of life in this frail population. Its underutilization in frail elderly persons suggests an important lack of awareness and education which needs to be addressed.

Elderly institutionalized men and women have a high prevalence of hypovitaminosis D which is not adequately addressed by appropriate replacement therapy, and may have important implications for bone and muscle health. The relationship between low 25-OH-vitamin D, falls and fractures is an important area for future study in this high-risk elderly population. **CHAPTER 6**

VITAMIN D AND OSTEOPOROSIS

6.1 Introduction

Vitamin D is important for maintaining calcium homeostasis and in the mineralization of healthy bone. Ageing results in impairment of vitamin D metabolism, with reduced capacity to generate pre-vitamin D in the skin in response to ultraviolet light exposure (353) combining with poor oral intake resulting in low circulating 25-OH-vitamin D levels. The ability to convert 25-OH-vitamin D to its active form, 1,25-OH-vitamin D, in the kidney (regulated by PTH) is also reduced (354;355). Calcium absorption from the intestine is modulated by circulating vitamin D metabolites (356), and with ageing the normal absorption of approximately 30% of dietary calcium is reduced, initiating the ionized calcium-driven feedback loop which results in normalization of serum calcium by way of compensatory secondary hyperparathyroidism, increased renal tubular reabsorption of calcium and increased bone resorption. A low calcium/phosphorus product is also seen, which causes an increase in the deposition of unmineralized osteoid.

It has been shown in Chapter 5 that there is widespread vitamin D insufficiency and secondary hyperparathyroidism in the frail elderly living in aged care facilities. But does this abnormal calcium homeostasis translate into lower bone mass and an increased risk of fracture? Cross-sectional studies have shown BMD is lower at the hip and lumbar spine in younger vitamin D deficient populations (347;357;358), and interventional studies of calcium and vitamin D supplementation have shown reduced bone loss (229) or significant BMD increases at both sites (351;352;359-361). Fracture incidence has also been shown to be significantly reduced by vitamin D supplementation (260;350;362;363).

There is less known about the relationship between 25-OH-vitamin D levels and QUS measures, which may be of more relevance to frail populations.

The aims of this study are as follows:

1: To confirm the distribution of BUA and VOS as measured by quantitative ultrasound of the calcaneus in a subpopulation of elderly men and women living in residential aged care facilities (hostels and nursing homes) in northern Sydney.

2: To establish the effect of vitamin D supplementation on BUA and VOS.

3: To describe the association between QUS measures and bone-related biochemical parameters, including serum 25-OH-vitamin D levels and serum parathyroid hormone (PTH).

4: To define hypovitaminosis D by relating 25-OH-vitamin D levels to BUA and VOS measures.

5: To identify the key independent predictors of BUA and VOS in the frail elderly.

This large cross-sectional study aims to show the relationship between serum 25-OHvitamin D levels and bone mass as measured by QUS in frail elderly men and women, and to describe the independent predictors of low BUA and VOS.

6.2 Methods

6.2.1 Subjects

The population described in Chapter 5 has been used for this analysis, comprising 979 women and 264 men enrolled in the FREE study between 1999 and 2002. Individuals with biochemical primary hyperparathyroidism or significant renal failure were excluded (Chapter 5.2).

6.2.2 Measurements

Serum 25-OH-vitamin D analysis and quantitative ultrasound of the left calcaneus were performed as described in Chapter 2.

6.2.3 Statistical analysis

For describing the basic population demographics and the effect of vitamin D supplementation on QUS measures, Pearson's chi-squared statistic was used for comparing frequencies between categorical variables, and student's independent t-test for comparing the means of continuous variables. Univariate analysis for BUA and VOS was performed using one-way ANOVA for continuous variables, log transforming 25-OH-vitamin D to fulfill assumptions of normality. Associations with categorical variables were carried out using independent sample t-tests for binomial variables and one-way ANOVA for multinomial variables. Multivariate analysis was carried out including all significant univariate variables, using multiple linear regression.

6.3 Results

6.3.1 Baseline demographics

Demographics for the group, including age, QUS values and mean serum 25-OHvitamin D levels are shown in Table 6.1.

Table 6.1: Mean Vitamin D levels and QUS values for men and women in hostels

 and nursing homes

	M	lale	Fen	nale
	Hostel	Nursing home	Hostel	Nursing home
	(N=136)	(N=128)	(N=560)	(N=419)
Age (years),	82.5 (7.5),	83.7 (7.9),	86.8 (6.1),	86.8 (6.8),
mean (sd), range	65.2 - 100.7	65.6 - 101.7	65.6 - 100.7	65.1 – 101.3
BUA (dB/MHz),	77.4 (22.0),	69.5 (24.8),	50.5 (17.1	41.8 (17.9),
mean (sd), range	16.5 – 135.3	18.8 - 138.2	13.9 – 119.8	12.2 – 117.9
VOS (m/s),	1608.8 (45.6),	1590.8 (51.6),	1567.7 (42.5),	1551.4 (42.1),
mean (sd), range	1473.5 – 1738.5	1472.5 – 1748.5	1484.0 - 1733.5	1459.5 - 1721.0
25-OH-vitamin D (nmol/L),	36 (17),	30 (18),	30 (15),	26 (16),
mean (sd), range	5 - 92	5 - 103	2 - 104	2 - 101
25-OH-vitamin D < 30 nmol/L	57 (41.9%)	82 (64.1%)	329 (58.8%)	314 (74.9%)
(number, %)				

6.3.2 Demographic determinants of BUA and VOS

Females were older than males (mean difference 3.7 years, 95% confidence interval (CI) for the difference 2.7 - 4.7 years, t-7.10, p<0.0001). Serum 25-OH-vitamin D was an average of 4.7 nmol/L (95% CI for the mean difference 2.3 - 7.1 nmol/L)

higher in men than women (t=3.86, p<0.0001). BUA was significantly higher in men (mean difference 27.1 dB/MHz, 95% CI 23.5 – 30.8, t=14.79, p<0.0001) than women, as was VOS (males an average of 39.8 m/s higher than women, 95% CI for the difference 32.8 - 46.9 m/s, t=11.06, p<0.0001). Quantitative ultrasound measures were significantly lower in individuals with primary hyperparathyroidism (mean BUA 39.1 dB/MHz, p=0.02 and mean VOS 1544.6 m/s, p<0.05). There were significantly more women with hypovitaminosis D (25-OH-vitamin D <30 nmol/L) than men, χ^2 =15.13, p<0.0001).

Resident mobility was assessed by their use of a walking aid (Table 6.2). Individuals were ranked on their minimum requirements; a resident classified as using a walking frame was unable to walk only using a stick. Those individuals classified as 'unable to walk unassisted' could mobilize with the assistance of nursing staff, but were unable to walk independently using a walking frame.

Table 6.2: 1	Number (%) of men a	and women classified	by mobility.
---------------------	---------------------	----------------------	--------------

Mobility	Men (n=262)	Women (n=970)	Total group (n=1232)
No walking aid	94 (35.9%)	287 (29.6%)	381 (30.9%)
Walking stick	65 (24.8%)	213 (22.0%)	278 (22.6%)
Walking frame	59 (22.5%)	324 (33.4%)	383 (31.1%)
Unable to walk	44 (16.8%)	146 (15.1%)	190 (15.4%)

6.3.3 Vitamin D supplementation

Those individuals receiving vitamin D supplementation at the time of enrolment into the study have been retained in the analysis. There is no statistically significant difference in BUA between these people and those not receiving supplements (supplemented group BUA=48.3 dB/MHz compared to 52.4 dB/MHz in the nonsupplemented group, mean difference 4.1 dB/MHz, 95% confidence interval for the difference -0.7 to 8.9 dB/MHz, p=0.1). For the whole group, VOS is significantly lower in the supplemented group (1558.9 m/s compared to 1569.2 m/s in the nonsupplemented, mean difference 10.3 m/s, 95% CI 0.1 to 20.6, p=0.047). When the cohort is split by gender (Table 6.3), vitamin D supplementation has no effect on either BUA or VOS measures in men or women.

Table 6.3: Comparison of residents on vitamin D supplementation at baseline with those not receiving supplements.

	Male		Female		
	Vit D Suppl	No Suppl	Vit D Suppl	No Suppl	
Ν	16 (6%)	248 (94%)	115 (11.5%)	864 (88.5%)	
Age (mean (sd))	81.2 (7.3)	83.2 (7.7)	86.9 (5.2)	86.8 (6.5)	
BUA (dB/MHz)	85.6 (26.1)	72.4 (23.4)	42.6 (15.6)	46.5 (18.3)	
VOS (m/s)	1598.3 (39.3)	1599.2 (50.1)	1552.9 (39.1)	1560.2 (43.2)	
25-OH-vitamin D	55.6 (24.2)	31.3 (16.5)*	43.7 (21.9)	26.0 (13.5)*	

*significantly different to supplemented group, p<0.05

6.3.4 Quantitative ultrasound and 25-OH-vitamin D

Figures 6.1 to 6.3 show the linear relationship between 25-OH-vitamin D and QUS parameters. In order to examine these relationships more closely, it was necessary to

log-transform 25-OH-vitamin D to fulfil assumptions of normality (as shown in Chapter 5) for univariate analysis. In men, BUA was not significantly related to log transformed 25-OH-vitamin D (F=3.13, p=0.08) or lnPTH (F=1.88, p=0.2). In women, mean BUA was significantly associated with lnPTH (F=4.30, p=0.04) but not with log transformed 25-OH-vitamin D (F=1.51, p=0.2). Mean VOS was not associated with 25-OH-vitamin D or PTH in men (25-OH-vitamin D, F=3.09, p=0.08; PTH F=1.49, p=0.2) or in women (25-OH-vitamin D, F=1.69, p=0.2; PTH F=2.69, p=0.2).




Figure 6.1: Mean BUA and VOS vs 25-OH-vitamin D in men





Figure 6.2: Mean BUA and VOS vs 25-OH-vitamin D in women

(Intercept=45.6, slope=0.02, R²=0.0004)






Figure 6.3: Mean BUA and VOS vs 25-OH-vitamin D in the whole cohort

6.3.5 Univariate associations with BUA and VOS

Those variables shown to be related to BUA and VOS in Chapter 4 were also included for analysis. Univariate analysis of mean BUA in men showed a significant

relationship between BUA and weight (F=23.23, p<0.0001), but BUA was not related to age (F=0.25, p=0.6). In women, mean BUA was significantly associated with age (F=21.45, p<0.0001) and weight (F=102.47, p<0.0001). Mean VOS in men did not reveal statistically significant associations with either age (F=0.009, p=0.9) or weight (F=1.32, p=0.3). In women, mean VOS was significantly associated with both age (F=11.87, p=0.001) and weight (F=8.53, p=0.004). Serum albumin was not associated with BUA in men (F=-0.093, p=0.93) or with VOS in men (F=-0.516, p=0.61). Similarly in women, there was no association between albumin and BUA (F= β 0.572, p=0.57) or between albumin and VOS (F=-0.202, p=0.84)

Interestingly, when men and women are considered together, univariate analysis reveals that mean BUA is significantly associated with age (F=39.95, p<0.0001), weight (F=266.74, p<0.0001), lnPTH (F=14.21, p<0.0001) and 25-OH-vitamin D (F=10.95, p=0.001), and mean VOS also shows statistically significant univariate associations with age (F=24.43, p<0.0001), weight (F=51.77, p<0.0001), lnPTH (F=9.10, p=0.003) and vitamin D (8.77, p=0.003). Serum albumin was not associated with BUA (F=0.393, p=0.70) or with VOS (F=0.053, p=0.96) in the whole group.

Both QUS measures and 25-OH-vitamin D levels decrease with decreasing mobility as measured by the use of walking aids in men and women (table 6.4). In men, QUS variables are significantly higher in more mobile individuals (for BUA, F=4.64, p=0.004, and for VOS F= 4.51, p<0.0001), and serum 25-OH-vitamin D levels are higher (F=3.51, p=0.02). Similarly significant relationships are seen in women, with higher mobility inferring higher BUA (F=22.6, p<0.0001), VOS (F=21.7, p<0.0001) and serum 25-OH-vitamin D (F=7.7, p<0.0001). These relationships are shown pictorially in Figures 6.4 and 6.5.

Table 6.4: Mean QUS and 25-OH-vitamin D levels for each category of mobility inmen and women.

	Mobility	Men (mean (sd))	Women (mean (sd))
BUA (dB/MHz)	Independent	79.8 (23.3)	52.5 (16.2)
	Stick	74.9 (23.1)	48.6 (18.3)
	Frame	69.6 (25.6)	42.9 (17.6)
	Unable to walk	62.8 (19.4)	38.0 (17.5)
VOS (m/s)	Independent	1615.1 (48.1)	1576.1 (41.2)
	Stick	1597.3 (48.8)	1561.2 (45.4)
	Frame	1591.2 (53.2)	1550.4 (38.9)
	Unable to walk	1581.2 (41.9)	1544.1 (42.9)
25-OH-vitamin D	Independent	36.0 (18.2)	29.2 (15.3)
(nmol/L)	Stick	35.0 (16.2)	30.4 (16.0)
	Frame	29.9 (19.6)	28.3 (16.3)
	Unable to walk	26.8 (16.5)	22.7 (13.9)




Figure 6.4: Box and whisker plots* of BUA and VOS related to level of mobility in the whole cohort.

*Boxes represent the interquartile range around the median




Figure 6.5: Box and whisker plots* of 25-OH-vitamin D levels related to level of mobility in the whole cohort.

Mobility

*Boxes represent the interquartile range around the median

Both BUA and VOS were significantly lower in nursing home residents than in hostel dwellers. In men, mean BUA in nursing homes was 69.53 dB/MHz, compared to 77.40 dB/MHz in hostels (mean difference 7.87 dB/MHz, 95% CI 1.20 – 14.53, p=0.02), and mean VOS in nursing homes was 1590.79 m/s compared to 1608.77 m/s in hostels (mean difference 18.00 m/s, 95% CI 4.14 – 31.83 m/s, p=0.01). In women, mean BUA in nursing homes was 41.77 dB/MHz, compared to 50.50 dB/MHz in hostels (mean difference 8.74 dB/MHz, 95% CI 6.18 – 11.29, p<0.0001), and mean VOS in nursing homes was 1551.39 m/s compared to 1537.74 m/s in hostels (mean difference 16.35 m/s, 95% CI 10.17 – 22.53 m/s, p<0.0001).

6.3.6 Hypovitaminosis D and QUS measures

In order to establish the existence of a distinct 25-OH-vitamin D level below which BUA and VOS are significantly low, QUS parameters were plotted against quintiles of vitamin D (Figure 6.6). Low levels of vitamin D are seen to correspond to low levels of both BUA and VOS, but the cut-off value is approximately 22 nmol/L, lower than that identified by secondary hyperparathyroidism (Chapter 5).




Figure 6.6: Mean BUA and VOS values relating to 25-OH-vitamin D quintiles

When men and women are considered separately, hypovitaminosis D (defined at two different cut-off levels, 25-OH-vitamin D <30 nmol/L as was related to secondary hyperparathyroidism in Chapter 5, and 25-OH-vitamin D <22 nmol/L, as is suggested by Figure 6) was not related to either BUA or VOS (tables 6.5, 6.6) in men. A cut-off of 30 nmol/L was also too high to show a significant relationship between 25-OH-vitamin D and QUS in women; using the lower cut-off of 22 nmol/L revealed significantly lower BUA in these vitamin D-deficient women, but no significant differences in VOS. When the group was combined, both QUS variables are significantly lower in individuals with either definition of hypovitaminosis D.

Table 6.5: Unadjusted relationship between hypovitaminosis D and QUS variables(where the mean difference is the magnitude of QUS in 25-OH-vitamin D repleteresidents less QUS in residents with 25-OH-vitamin D < 30 nmol/L)</td>

		Mean	95% CI	t	р
		difference			
BUA (dB/MHz)	Men	4.32	-2.52 - 11.16	1.25	0.2
	Women	1.94	-0.89 - 4.78	1.35	0.2
	Total cohort	4.43	1.38 - 7.49	2.85	0.005*
VOS (m/s)	Men	9.47	-4.78 - 23.72	1.31	0.2
	Women	5.46	-1.31 - 12.23	1.58	0.1
	Total cohort	9.22	2.47 - 15.96	2.79	0.005*

*statistically significant at p<0.05

Table 6.6: Unadjusted relationship between hypovitaminosis D and QUS variables(where the mean difference is the magnitude of QUS in 25-OH-vitamin D repleteresidents less QUS in residents with 25-OH-vitamin D < 22 nmol/L)</td>

		Mean	95% CI	t	р
		difference			
BUA (dB/MHz)	Men	3.49	-3.52 - 10.50	0.98	0.3
	Women	3.08	0.45 - 5.71	2.30	0.02*
	Total cohort	4.87	1.98 – 7.77	3.30	0.001*
VOS (m/s)	Men	9.44	-5.13 - 24.00	1.28	0.2
	Women	5.65	-0.65 - 11.95	1.77	0.08
	Total cohort	8.91	2.75 - 15.07	2.84	0.005*

*statistically significant at p<0.05

6.3.7 Independent predictors of BUA and VOS

Results for multivariate analysis for BUA and VOS are shown in Table 6.7. All variables which were significantly associated with the QUS parameters (age, gender, institution type, mobility, lnPTH, ln25-OH-vitamin D, weight) were included in the analysis. Only mobility, lnPTH, weight and gender remain independently related to BUA in this elderly population, accounting for 41% of the variability of BUA. VOS is predicted by mobility, weight and gender alone, comprising 21% of the variability of VOS. 25-OH-vitamin D becomes nonsignificant in both models. When a binomial variable of 'hypovitaminosis D' is substituted for the continuous log transformed 25-OH-vitamin D variable, hypovitaminosis D is not an independent predictor of either BUA or QUS for either a 22 or 30nmol/L cut-off (p>0.8), data not shown.

	Variable	β	t	р
BUA	Mobility	-0.239	-9.032	0.0001
$R^2 = 0.41$	lnPTH	-0.058	-2.177	0.03
	Weight	0.315	10.904	0.0001
	Gender (female)	-0.366	-12.613	0.0001
VOS	Mobility	-0.259	-8.461	0.0001
$R^2 = 0.21$	Weight	0.093	2.775	0.006
	Gender (female)	-0.318	-9.500	0.0001

Table 6.7: Multivariate analysis of BUA and VOS

Excluded from model: For BUA: Age, Institution, InvitD. For VOS: Age, Institution, InvitD, InPTH.

When men and women are considered separately, slightly different parameters are independent predictors for QUS (Tables 6.8, 6.9), but 25-OH-vitamin D levels remain unrelated to BUA and VOS.

	Variable	β	t	р
Men	Mobility	-0.265	-3.819	0.0001
$R^2 = 0.19$	Age	0.215	3.061	0.003
	Weight	0.345	4.980	0.0001
Women	Mobility	-0.261	-6.792	0.0001
$R^2 = 0.23$	Weight	0.336	9.714	0.0001
	Institution	-0.083	-2.138	0.03
	lnPTH	-0.069	-2.030	0.04

Table 6.8: Multivariate analysis of BUA (dB/MHz) in men and women

Excluded from model: For men: Institution, InvitD, InPTH. For women: Age, InvitD.

Table 6.9: Multivariate analysis of VOS (m/s) in men and women

	Variable	β	t	р
Men	Mobility	-0.245	-3.358	0.001
$R^2 = 0.06$				
Women	Mobility	-0.284	-7.682	0.0001
$R^2 = 0.10$	Weight	0.086	2.293	0.02
	Age	-0.075	-1.985	0.05

Excluded from model: For men: Age, Institution, InvitD, InPTH, weight. For women: Institution, InvitD, InPTH.

6.4 Discussion

Vitamin D deficiency was extremely common in both elderly men and women living in institutions in Sydney, but serum 25-OH-vitamin D was not independently associated with BUA or VOS in this study. Elderly populations have previously been reported as having low 25-OH-vitamin D levels (327;339) at similar latitudes in Europe and the US, and a case-control study of elderly men in Virginia showed 25-OH-vitamin D levels to be significantly lower in men residing in nursing homes than living independently (326). There is some observational evidence suggesting that 25-OH-vitamin D levels relate to bone mass. In a group of postmenopausal women undergoing screening for osteoporosis, those with subclinical vitamin D deficiency were found to have lower vertebral bone mass as measured by quantitative computed tomography than those with normal 25-OH-vitamin D levels (364). A Spanish group has also recently shown a correlation between 25-OH-vitamin D levels and BMD at the lumbar spine and femoral neck in otherwise healthy postmenopausal women (358).

However the relationship between QUS measures and 25-OH-vitamin D is not as clear. A cross-sectional study of elderly institutionalized men and women showed 25-OH-vitamin D levels to be associated with both BUA and VOS, largely through the effect of secondary hyperparathyroidism (365). Calcium and vitamin D supplementation in this group was shown to have a significant beneficial effect on BUA over 2 years compared to placebo, but no effect on VOS (110). An observational study of perimenopausal women did not reveal a similar association between vitamin D supplementation and QUS measures in this younger cohort (366). In the FREE study, cross-sectional analysis did not reveal a relationship between vitamin D supplementation and BUA or VOS, despite significantly higher 25-OH- vitamin D levels in the supplemented group. This may be related to the small numbers of patients receiving vitamin D supplements, since the trend in women was for those on supplements to have lower BUA and VOS measures (despite not reaching statistical significance). It may be that these individuals were preferentially identified as having a high risk of osteoporosis and thus self-selected to have lower QUS readings.

Serum 25-OH-vitamin D was univariately related to BUA in this group of frail elderly men and women. When vitamin D status was considered as 'hypovitaminosis D' at a cut-off of <30nmol/L or <22nmol/L, those individuals classified as being vitamin D deficient had significantly lower BUA and VOS measures. This suggests that a 25-OH-vitamin D level less than 30nmol/L may have an important effect on bone health (as there is on the development of secondary hyperparathyroidism, seen in Chapter 5). Prospective studies of QUS as a predictor of fracture have shown that a decrease of 1 standard deviation in BUA is associated with an approximate doubling of hip fracture risk (54;56;58); the clinical significance of the smaller differences in BUA shown in this study is not clear but prospective fracture analysis will address this issue. When VOS is considered, a distinct cut-off to define 'hypovitaminosis D' is not so clear, and seems to be higher as seen in Figure 6.6. However, the previous studies of prospective fracture support the lower cut-off seen with BUA (22-30nmol/L). Prospective fracture data from the FREE study will help to more clearly elucidate whether BUA or VOS is more important for fracture risk in this frail elderly cohort.

It makes sense to assess 25-OH-vitamin D levels in relation to serum PTH levels, as this will reflect changes over a shorter time frame. Quantitative ultrasound is measuring the outcome of longer periods of vitamin D deficiency. Nevertheless, this has important clinical implications, with univariate analysis suggesting that vitamin D supplementation in the elderly should be considered even in mild cases of suboptimal serum 25-OH-vitamin D, and widens the population at risk of impaired bone metabolism. If those patients with low 25-OH-vitamin D levels can be identified before longstanding bone changes have occurred, then any intervention is likely to be more successful in ultimate fracture prevention.

Independent predictors of BUA include mobility, weight, gender and PTH but not 25-OH-vitamin D. However, when mobility is not included in the analysis, 25-OHvitamin D remains in the model as a significant independent predictor. It has been shown that 25-OH-vitamin D levels are strongly linked to mobility, which suggests that 25-OH-vitamin D is either an intervening variable that plays a role in determining mobility perhaps by improving muscle strength (and hence is lost in a multivariate analysis), or that mobility is a strong determinant of vitamin D status independent of bone health. Weight bearing is important in maintaining BMD, and hence it is logical to assume that the relationship between mobility and QUS measures might be solely due to the amount of time a resident spends standing or walking. The picture is complicated by vitamin D, as an individual who is more mobile is more likely to get outside into the sunlight and have adequate UV light exposure to synthesize vitamin D. Conversely, if it is assumed that vitamin D plays a role in maintaining muscle strength (200;202), it may be that those individuals with higher 25-OH-vitamin D levels are going to be more mobile independent of sunlight exposure. It is likely that 25-OH-vitamin D, mobility and QUS are closely entwined in elderly persons; the relationship between 25-OH-vitamin D and muscle strength relating to falls will be addressed in Chapter 7.

Parathyroid hormone levels remain an independent predictor of BUA but not VOS in multivariate analysis. It has been shown previously in case control studies that primary hyperparathyroidism is associated with low BUA measures (367-369), and that surgical correction results in significant improvement of BUA but not VOS (367). This is supported by the lower BUA seen in individuals with biochemical primary hyperparathyroidism in this elderly cohort. When secondary hyperparathyroidism is considered, Krieg et al (365) in a smaller study of 264 women and 103 men living in aged care facilities found that VOS was also related to PTH levels, but did not correct for mobility. It was suggested that the relationship between secondary hyperparathyroidism and QUS is likely to be mediated by low 25-OH-vitamin D levels, but the bone changes reflect the effect of the PTH levels directly rather than vitamin D deficiency. Continuous exposure to PTH at a bone level results in mobilization of calcium from the bone surface (370) and to an increase in the number and activity of osteoclasts (371), and secondary hyperparathyroidism might therefore predispose to changes which might be detectable by QUS.

Vitamin D and PTH play a role in determining BUA and VOS at the calcaneus in the frail elderly, however how much is a direct action on bone and how much is via improved general health and mobility is not clear. Nevertheless, this study gives further evidence to support vitamin D supplementation in D-deficient elderly persons.

CHAPTER 7

VITAMIN D AND PTH AND THEIR RELATIONSHIP TO FALLING IN THE ELDERLY

7.1 Introduction

Falling is a serious health problem in the elderly, bring with it significant morbidity and mortality. There are a myriad of factors associated with an increased risk of falling in this group, including increasing age, balance, muscle strength and cognitive function (Chapter 1.4). The elderly are at particularly high risk, with 33% of community dwellers over the age of 65 having at least one fall each year (176;177;372). The risk is higher in the institutionalized elderly (293;294;373), and it is estimated that 6-7% of fallers sustain a fracture as a direct result of falling (177;374).

Vitamin D levels are low in residents of aged care facilities (326;327) as seen in Chapter 5, due to inadequate sunlight exposure, impaired ability to convert vitamin D in the skin and poor diet. Vitamin D insufficiency has also been related to poorer muscle strength (200;375), which could logically increase the risk of falling in this group. Cross-sectional studies have not consistently shown a higher incidence of falls in vitamin D deficiency (176;376-378). There is one large prospective study of time to first fall in women living in residential care (379) which shows that lower 25-OHvitamin D levels are related to a shorter time to falling.

Parathyroid hormone has also been shown to be indirectly associated with muscle strength, with secondary hyperparathyroidism patients exhibiting muscle weakness (380). One study of falls in the elderly describes an independent association of higher PTH levels with an increased falls risk (OR 5.6 for each natural logarithmic unit of PTH) (208). This study aims to:

(1) Prospectively measure the incidence of falls in a population of elderly residents of aged care facilities in Sydney, Australia

(2) Describe the demographic factors associated with falling in aged care facilities

(3) Describe the association between serum 25-OH-vitamin D levels and prospective falls in the elderly

(4) Describe the association between serum PTH levels and prospective falls in the elderly

(5) Establish the independent factors, both physical and biochemical, associated with prospective falls in the institutionalized elderly. Do 25-OH-vitamin D and PTH both contribute to falls risk in this population?

7.2 Methods

7.2.1 Subjects

The first 1010 women and 270 men enrolled in the FREE study to reach one year of prospective follow up have been included in this analysis. All falls and deaths were recorded over the follow up period, as described in Chapter 2.

7.2.2 Measurements

Measurements of physical function and biochemistry are described in full in Chapter 2. Potential risk factors for falling in an elderly population, including age, weight, type of accommodation, urinary incontinence, cognitive function and overall health as assessed by both RCS and implicit review were recorded. Two different methods of assessing balance were examined: the 30 second stand test on both firm and pliable surfaces, and the use of a walking aid in daily life as a surrogate for balance on mobilization.

7.2.3 Statistical analysis

Baseline characteristics were examined with relation to falls status using independent t-tests for continuous variables and chi-square tests for categorical variables. Risk to first fall was calculated using proportional cox regression.

7.3 Results

7.3.1 Distribution of falls


The number of falls occurring in the study population at 6 and 12 months is shown in Table 7.1. There was no significant difference in the incidence of falls at 6 months $(\chi^2=16.78, df=15, p=0.3)$ or at 12 months $(\chi^2=31.65, df=23, p=0.1)$ between men and women. In the first 6 months of follow-up, there were more falls in hostels than in nursing homes $(\chi^2=26.96, p=0.03)$ but there was no significant difference between institution type after 12 months follow up $(\chi^2=32.90, p=0.08)$.

	At 6 r	nonths	At 12	months
	Men	Women	Men	Women
	(N=270)	(N=1010)	(N=270)	(N=1000)
No falls	172 (64%)	628 (62%)	142 (53%)	492 (49%)
1 fall	48 (18%)	203 (20%)	51 (19%)	224 (22%)
2 falls	14 (5%)	73 (7%)	21 (8%)	98 (10%)
3 falls	16 (6%)	39 (4%)	11 (4%)	64 (6%)
4 falls	4 (2%)	20 (2%)	6 (2%)	34 (3%)
5 falls	6 (2%)	20 (2%)	9 (3%)	17 (2%)
6 falls	2 (0.7%)	7 (0.7%)	7 (3%)	24 (2%)
7 falls	1 (0.4%)	9 (1%)	3 (1%)	11 (1%)
8 falls	1 (0.4%)	2 (0.2%)	2 (0.7%)	13 (1%)
9 falls	3 (1%)	3 (0.3%)	2 (0.7%)	6 (0.6%)
>=10 falls	3 (1%)	6 (0.6%)	16 (6%)	27 (3%)

Table 7.1: Incidence of falls at 6 and 12 months in men and won
--

7.3.2 Serum 25-OH-vitamin D and balance

At baseline, univariate analysis showed serum 25-OH-vitamin D was significantly related to balance (F=14.09, df 4, p<0.0001) and to implicit review score (F=4.31, df 3, p=0.005). Those individuals who performed badly in either score were seen to have lower 25-OH-vitamin D levels than those with better balance performance (Figure 7.1) or higher implicit review scores (Figure 7.2). Balance was measured on both a firm and compliant surfaces, as described in Chapter 2.3, with 1 being an inability to stand for any length of time unaided on a firm surface and 5 being able to stand on a pliable surface for 30 seconds unsupported.






Table 7.2: Independent determinants of balance

	χ^2 statistic	P value
25-OH-vitamin D < 30nmol/L	67.12	<0.0001
Age	32.65	<0.0001
Weight	27.73	<0.0001
Gender	11.86	0.02
Incontinence	91.00	<0.0001

Logistic regression modeling showed that the relationships between hypovitaminosis D and balance (Table 7.2) and between low 25-OH-vitamin D and implicit review (Table 7.3) were independent of age, weight, gender or presence of urinary incontinence .

Figure 7.2: Relationship between mean 25-OH-vitamin D levels and function, as graded by the implicit review score.




 Table 7.3:
 Independent determinants of implicit review score

	χ^2 statistic	P value
25-OH-vitamin D < 30nmol/L	22.53	<0.0001
Age	5.05	0.17
Weight	19.88	<0.0001
Gender	4.59	0.21
Incontinence	51.90	<0.0001

Characteristics	No falls in 6 months	One or more falls	P value		
Age (years)	85.5 +/- 6.9	86.7 +/- 6.7	0.004*		
Female sex (%)	628 (78.5%)	382 (79.6%)	0.67		
Hostel accommodation (%)	483 (60.4%)	235 (49.0%)	<0.0001*		
Walking aid (%)					
No walking aid	252 (31.5%)	137 (28.5%)			
Walking stick	194 (24.3%)	92 (19.2%)			
Walking frame	216 (27.0%)	184 (38.3%)			
Unable to walk	134 (16.8%)	60 (12.5%)			
Balance			<0.0001*		
Unable to stand unassisted	154 (66.4%)	78 (33.6%)			
Stands for <30 seconds	61 (47.7%)	67 (52.3%)			
Stands 30 seconds on firm surface	99 (45.8%)	117 (54.2%)			
Stands <30 seconds on soft surface	186 (60.2%)	123 (39.8%)			
Stands 30 seconds on soft surface	297 (76.3%)	92 (23.7%)			
Weight (kg)	60.6 +/- 13.7	57.9 +/- 13.0	0.001*		
BUA (dB/MHz)	51.7 +/- 22.1	50.45 +/- 22.7	0.44		
VOS (m/s)	1566.0 +/- 45.7	1567.3 +/- 47.6	0.62		
25-OH-vitamin D (median) (nmol/L)	29.6 +/- 17.0 (26)	27.7 +/- 15.2 (24)	0.04*		
PTH (median) (pg/mL)	75.1 +/- 71.9 (57.8)	79.5 +/- 67.6 (59.5)	0.28		
SMMSE	21.6 +/- 8.1	19.2 +/- 8.7	<0.0001*		
Resident classification			<0.0001*		
RCS 1	72 (9.0%)	48 (10.0%)			
RCS 2	142 (17.8%)	113 (23.5%)			
RCS 3	96 (12.0%)	95 (19.8%)			
RCS 4	26 (3.3%)	19 (4.0%)			
RCS 5	64 (8.0%)	40 (8.3%)			
RCS 6	115 (14.4%)	50 (10.4%)			
RCS 7	203 (25.4%)	87 (18.1%)			
RCS 8	73 (9.1%)	19 (4.0%)			
Implicit review			<0.0001*		
No symptoms	32 (80%)	8 (20%)			
Mild symptoms	255 (72.4%)	97 (27.6%)			
Moderate and severe symptoms	505 (57.7%)	361 (42.3%)			
Incontinence	352 (68.3%)	163 (31.7%)	<0.0001*		

Table 7.4: Baseline characteristics and status with respect to falls at 6 months

*statistically significant p<0.05.

Characteristics	No falls in 12 months	One or more falls	P value		
Age (years)	85.4 +/- 6.8	86.5 +/- 6.8	0.005*		
Female sex (%)	492 (77.6%)	518 (80.2%)	0.27		
Hostel accommodation (%)	388 (61.2%)	330 (51.1%)	<0.0001*		
Walking aid (%)					
No walking aid	203 (32.1%)	184 (28.9%)			
Walking stick	152 (24.1%)	134 (21.0%)			
Walking frame	162 (25.6%)	238 (37.4%)			
Unable to walk	113 (17.9%)	81 (12.7%)			
Balance			<0.0001*		
Unable to stand unassisted	130 (56.0%)	102 (44%)			
Stands for <30 seconds	45 (35.2%)	83 (64.8%)			
Stands 30 seconds on firm surface	70 (32.4%)	146 (67.6%)			
Stands <30 seconds on soft surface	137 (44.3%)	172 (55.7%)			
Stands 30 seconds on soft surface	250 (64.3%)	139 (35.7%)			
Weight (kg)	60.3 +/- 13.8	59.0 +/- 13.2	0.10		
BUA (dB/MHz)	51.4 +/- 22.2	52.0 +/- 23.4	0.10		
VOS (m/s)	1565.0 +/- 45.1	1568.0 +/- 47.6	0.26		
25-OH-vitamin D (median) (nmol/L)	30.2 +/- 17.0 (26)	28.0 +/- 15.6 (24)	0.006*		
PTH (median) (pg/mL)	73.3 +/- 70.0 (56.9)	80.1 +/- 70.6 (59.5)	0.09		
SMMSE	21.8 +/- 8.1	19.6 +/- 8.5	<0.0001*		
Resident classification			<0.0001*		
RCS 1	55 (8.8%)	65 (10.2%)			
RCS 2	112 (17.9%)	143 (22.5%)			
RCS 3	74 (11.8%)	117 (18.4%)			
RCS 4	21 (3.3%)	24 (3.8%)			
RCS 5	46 (7.3%)	58 (9.1%)			
RCS 6	91 (14.5%)	74 (11.7%)			
RCS 7	172 (27.4%)	118 (18.6%)			
RCS 8	56 (8.9%)	36 (5.7%)			
Implicit review			<0.0001*		
No symptoms	32 (80.0%)	8 (20.0%)			
Mild symptoms	210 (59.7%)	142 (40.3%)			
Moderate symptoms	386 (44.1%)	489 (55.9%)			
Incontinence	294 (57.1%)	221 (42.9%)	<0.0001*		

Table 7.5: Baseline characteristics and status with respect to falls at 12 months

*statistically significant p<0.05


Tables 7.4 and 7.5 show the baseline characteristics of those residents who suffered one or more falls over 6 and 12 months respectively, compared to residents who did not fall over the same period. Fallers were seen to be older, more likely to use a walking aid, of lower weight and requiring more nursing care (as measured by resident classification score, RCS) than non-fallers. Fallers had lower serum 25-OHvitamin D levels than non-fallers, but PTH levels were not significantly different between the groups although there was a trend at 12 months. Similar relationships were seen at both 6 and 12 months, and so further analyses have been limited to 12 month data. Time to first fall was calculated, as it gives more useful information than simple comparisons between fallers and non-fallers.

7.3.3 Survival analysis

The median time from baseline assessment to first fall was 274 days (range 244 to 304 days). One hundred and eighty-seven residents (14.6%) died over the first 12 months of follow-up. The incidence of a first fall decreased over time as shown in Figure 7.3, censoring residents at the time of their first fall, at death or at the end of follow-up. Serum 25(OH)D distribution was close to normal for this subgroup and hence not log transformed; PTH was log transformed to preserve analysis assumptions.

When time to first fall was considered (Table 7.6), similar relationships were seen with baseline demographic variables as were seen with fall status.

Figure 7.3: Survival curve for time to first fall.



	LR	Wald statistic	P value
Age	1.02	15.28	<0.0001*
Institution type	0.71	19.20	<0.0001*
Walking aid	N/A	33.15	<0.0001*
Balance	N/A	27.77	<0.0001*
SMMSE	0.98	34.43	<0.0001*
RCS	N/A	43.38	<0.0001*
Implicit review	N/A	10.26	0.016*
Weight	0.99	9.40	0.002*
Incontinence	1.37	6.21	0.013*
25-OH-vitamin D	1.00	3.55	0.06
РТН	1.00	4.99	0.025*

Table 7.6: Univariate associations with time to first fall

*statistically significant at p<0.05.

Cox regression analysis results for the entire cohort are shown in Table 7.7. Independent predictors for falling included increasing age (p = 0.001), poor balance (p=0.001), incontinence (p=0.06) and implicit review (p=0.10). Whether treated as a continuous or a categorical variable (using cutoffs for hypovitaminosis D as described in Chapter 5), serum 25-OH-vitamin D was not significantly related to falling.

	Hazards ratio (HR)	95% Confidence Interval		P value
		Lower	Upper	
Age	1.03	1.01	1.05	0.001
Incontinence	1.28	0.99	1.65	0.058
Balance	N/A	N/A	N/A	0.001
Implicit review	N/A	N/A	N/A	0.10

Table 7.7: Survival analysis for time to first fall

Excluded variables: 25-OH-vitamin D, InPTH, SMMSE

Because it has previously been shown there may be different risk factors for falls in the population according to whether a subject can stand or not (381), a subgroup analysis was performed in more healthy individuals, ie. those aged < 90 years, relatively independent (RCS >=5) and able to walk. Univariate analysis in this subgroup of 455 residents showed 25-OH-vitamin D was related to time to first fall (HR 0.99, 95% CI 0.98-1.00, p=0.002). However in multivariate analysis that included age, balance, continence status and cognitive function, 25-OH-vitamin D was no longer significant (Table 7.8). In univariate analysis, serum PTH (log transformed) was also related to time to first fall (HR 1.43, 95% CI 1.20-1.70, p<0.0001). In multivariate analysis, PTH remained significant (HR 1.52, CI 1.211.91, p<0.0001) as shown in Table 7.8. However PTH was not independently related to falls in elderly residents over the age of 90 years (p=0.9).

Table 7.8: Survival analysis for time to first fall in residents aged <90 years with</th>RCS >=5 and able to walk.

	Hazards ratio (HR)	95% Confidence Interval		P value
		Lower	Upper	
lnPTH	1.52	1.21	1.91	<0.0001
SMMSE	0.97	0.94	1.00	0.047
Incontinence	1.51	1.11	2.05	0.009
Balance	N/A	N/A	N/A	< 0.0001
Implicit review	N/A	N/A	N/A	0.017

Excluded variables: 25-OH-vitamin D

7.4 Discussion

This study showed that 50% of residents in aged care facilities in Sydney will have one or more falls over a year, most of which occurred in the first 6 months. Fallers were more likely to live in hostels than nursing homes, were slightly older, lighter, more cognitively impaired, more likely to be incontinent and had poorer balance and general health scores than non-fallers, consistent with current falls literature (Chapter 1.4). Fallers had significantly lower 25-OH-vitamin D levels than non-fallers (median 25-OH-vitamin D level for fallers 24 nmol/L, compared to 26 nmol/L for residents who did not fall). Parathyroid hormone levels showed a trend towards being higher in fallers although this did not reach statistical significance.

It is considered that low 25-OH-vitamin D levels increase fracture risk (382) and that supplementation reduces the incidence of fractures in the elderly (383;384), but perhaps this effect may not be solely through changes in bone mass such as those seen in Chapter 6. Vitamin D deficiency may lead to reduced muscle strength through loss of type 2 muscle fibres (385), and hence an increased risk of falling. Cross-sectional studies relate lower 25-OH-vitamin D levels to reduced quadriceps and grip strength in elderly people (386;387). Bischoff et al (200) examined the relationship between muscle strength and vitamin D deficiency in ambulatory elderly subjects (103 women, 216 men, mean age for women 74, for men 77 years). Twelve percent of women and 18% of men had 25-OH-vitamin D values below the normal range (< 30 nmol/L). Strength of leg extension showed a modest, but significant, positive correlation with 25-OH-vitamin D in male but not female subjects. A study of both independent living and hospitalized people over 70 years of age (377) showed serum 25-OH-vitamin D

levels were significantly lower in those with poorer handgrip strength and stairclimbing ability.

Interventional studies have shown varied results. A small study in elderly women (378) showed that 6 months of vitamin D replacement improved extensor muscle strength and walking distance over 2 minutes in vitamin D-deficient (25-OH-vitamin D <20 nmol/L) older women. Measures of functional ability and activities of daily living in the frail elderly have been shown to improve in uncontrolled studies of vitamin D supplementation (388;389). Studies in healthier, non-vitamin D deficient populations have failed to show a significant improvement in quadriceps muscle function after vitamin D supplementation (390;391). A comprehensive review of the relationship between 25-OH-vitamin D and muscle function published in 2002 (375) concluded that there is a threshold effect, with vitamin D deficiency being associated with reduced muscle strength but the relationship not persisting in 25-OH-vitamin D replete individuals.

Few studies have prospectively looked at the effect of 25-OH-vitamin D levels on falling. Supplementation with oral calcium and vitamin D has been recently shown to reduce the incidence of fallers and falls by approximately 50% (230;392). Our study showed that balance was related to serum 25-OH-vitamin D, and that fallers had significantly lower 25-OH-vitamin D levels than non-fallers in residents of aged care facilities. This supports findings of a retrospective study by Stein et al (208) in a group of 83 ambulatory residents of hostels and nursing homes in Melbourne, Australia. The mean age was 84 years and the median 25-OH-vitamin D level was 27 (18-37) nmol/L, similar to the participants of the FREE study (393). Fallers had a median 25-OH-vitamin D level of 22 nmol/L, compared to 29 nmol/L for non-fallers

(p=0.019). Parathyroid hormone levels were also shown to be significantly associated with falling (OR=5.6, 95% CI 1.7-18.5 per logarithmic unit of PTH), independent of 25-OH-vitamin D levels.

The only large prospective study of vitamin D and falling in residential care facilities published to date has been an extension of this Melbourne cohort (379), measuring 25-OH-vitamin D and prospective falls in 1261 ambulatory women living in hostels and nursing homes. Higher serum 25-OH-vitamin D levels were independently associated with a lower risk for falling (p<0.01). Parathyroid hormone levels were not reported. When only residents of nursing homes were examined (who as a group had a higher incidence of falls than the hostel residents), the relationship was not seen. The authors postulate that the relationship might be obscured in this population by the substantially higher falls rate in this group. Our study did not show a relationship between vitamin D and falls risk by cox regression analysis, either in the population as a whole or in any subgroup analysis.

There are a number of explanations for the discrepancy seen between the two studies. Flicker et al found a relationship between 25-OH-vitamin D and time to first fall after correcting for psychotropic medication use, the presence of wandering behaviour and previous Colles fracture which were not examined in our group. Likewise, the Melbourne group did not report any association between vitamin D and mobility which was seen in the FREE study. It is likely that the use of different cofactors has impacted the different relationships seen in the two studies. If we accept earlier findings of differing risk factors for falling in the elderly dependent on frailty (381), our results are more in keeping with the Melbourne group. When the frailest elderly (aged >90 years or unable to stand unassisted) were excluded from the analysis, a higher serum PTH was independently associated with increased falls risk in Sydney. It is plausible that the relationship between 25-OH-vitamin D and time to first fall in the more 'healthy' elderly population is mediated by PTH. It is possible that iPTH may be a more sensitive marker of vitamin D deficiency than serum 25-OH-vitamin D. Another explanation for the apparent discrepancy between the study of Flicker et al and the present study is that the assay used for PTH in the present study was in fact a more sensitive marker of vitamin D deficiency than the serum 25OH D assay employed. This merits further studies to tease out the proportional roles of 25-OHvitamin D and PTH in falling in the elderly.

High levels of PTH have previously been linked to muscle weakness (380), which was reversible after treatment hyperparathyroidism and independent of vitamin D status. It may be that the effect we have seen on falling in this elderly population is reflecting a direct action of PTH on muscle, however serum 25-OH-vitamin D levels are known to be tightly linked to PTH and it might be that secondary hyperparathyroidism is in fact an intermediate variable in the relationship between hypovitaminosis D and falling. The absence of a relationship between 25-OH-vitamin D and falling in this subgroup analysis does not support the theory that both 25-OHvitamin D and PTH act independently on muscle function and falling; more studies are required to further define the relationship.

A similar collinearity problem was encountered with balance, general health and 25-OH-vitamin D levels. Residents with lower 25-OH-vitamin D levels had significantly poorer balance and worse implicit review scores. It is likely that the relationship between balance and 25-OH-vitamin D is mediated in part by the action of 25-OHvitamin D on muscle strength and function. Serum 25-OH-vitamin D levels may also be an indirect measure of general health, reflecting mobility, nutrition and comorbidities which may all contribute to falls risk.

In conclusion, hypovitaminosis D was not shown to be independently predictive of falling in the frail institutionalized elderly, but may be indirectly related to falls risk through secondary hyperparathyroidism. Falling in this population is a common problem, with multiple interacting factors contributing to increase an elderly person's risk which still need to be adequately defined.

CHAPTER 8

SUMMARY AND FUTURE DIRECTIONS

SUMMARY AND FUTURE DIRECTIONS

8.1 Introduction

In order to implement preventive health strategies, the first step is to fully understand the size of the problem. Osteoporotic fracture is an important issue both in terms of human morbidity and mortality, and with regard to the health care dollar. Costs associated with fracture increase substantially with age (394), just as the incidence of fracture also increases. Those elderly individuals who are too frail to live independently are at high risk of falling, of involutional osteoporosis and hence of osteoporotic fracture. This thesis aimed to describe the magnitude of the problem of osteoporotic fracture in the frail institutionalized elderly, by assessing risk factors for falls and fractures in a large population of elderly residents of hostels and nursing homes in Sydney, Australia. It is hoped that this will provide a valuable baseline from which interventional studies can be designed to identify the most appropriate, effective strategies for fracture prevention in the frail elderly.

8.2 Summary and implications of each study

This thesis described a range of features related to falls and fractures in the frail elderly. The study was unique in being a large, prospective study of risk factors in a group of frail elderly people in residential care facilities, a group who have not been previously extensively studied. The key findings of this thesis are summarized as follows:

(1) Quantitative ultrasound is a reliable tool for assessing bone density in the frail elderly, as both precision and sensitivity to longitudinal change were independent of age (Chapter 3).

Measurement of BUA and VOS was reliable, least significant change for BUA being 2.4% and for VOS 0.3% using the CUBA McCue instrument and 2.7% for BUA using the Metra QUS-2. Comparison of the instruments showed the QUS-2 measures BUA higher than the CUBA by 1.9 dB/MHz (+/- 9.55 dB/MHz). Moreover classification of individuals as osteoporotic by machine-derived T-score was different between instruments (k=0.44 in men and k=0.62 in women). In women, BUA was observed to decrease by 5.2% over 2 years without significant change in VOS, suggesting BUA is more useful in longitudinal studies.

Ultrasound measurements at the calcaneus are reliable and sensitive to longitudinal change even in the extreme elderly, which makes it a useful tool in frail, immobile populations. Measurements are however instrument-specific, and should be interpreted with caution in the assessment of fracture risk. It is now widely accepted that the same machine should be used for progress studies of DEXA measurements; this study extends the recommendation to the use of QUS machines for measuring bone. The same QUS machine should be used for baseline and follow-up measurements in longitudinal studies to ensure meaningful conclusions.

(2) There is a high prevalence of low bone density as measured by QUS in the frail elderly (Chapter 4).

Broadband ultrasound attenuation and VOS were higher in men than women by approximately 30% and 2% respectively, and this difference was maintained at all ages. There was no significant decline in BUA or VOS with age in men; however, for women BUA declined by 2.8-4.7% per decade and VOS by 1% per decade. Mean BUA T-scores were -1.55 and -2.48 at age 90 years in men and women.

This study confirms that both men and women of advanced age are at significant risk of low bone density. In a group recognized to have a high incidence of falls, this is likely to be an important risk factor for future fracture. Although limited by its crosssectional design, this study suggests only minor bone loss occurs at the calcaneal site with very old age in either sex. Treatment must be aimed not merely at preventing bone loss in this group, but at improving bone density in order to reduce potential fracture risk.

(3) The prevalence of low 25-OH-vitamin D is high in this population and under-treated (Chapter 5).

Over 77% of the population had low serum 25-OH-vitamin D levels compared to the laboratory reference range, and almost 41% had secondary hyperparathyroidism. A

25-OH-vitamin D cut-off of 30 nmol/L predicted secondary hyperparathyroidism. Independent predictors of lower serum 25-OH-vitamin D levels included female gender, increasing age, elevated PTH levels, cooler season, decreased mobility and lower creatinine clearance. Independent predictors of hypovitaminosis D (25-OHvitamin D < 30 nmol/L) included female gender, reduced mobility higher PTH levels, and a lower creatinine clearance. Only 6% of men and 12% of women were on vitamin D supplementation. Vitamin D supplementation conferred significantly higher serum 25-OH-vitamin D and lower PTH levels, but this was not apparently related to nutritional status as albumin levels were not different between the groups.

Although Sydney has adequate sunlight hours all year long this population was not being exposed on a regular basis. Vitamin D supplementation is a simple and cheap intervention strategy that could be undertaken in a frail elderly population to improve calcium metabolism, particularly in less mobile individuals.

(4) The relationship between QUS and 25-OH-vitamin D levels was significant only in elderly people who were vitamin D deficient, and secondary hyperparathyroidism was an independent predictor of BUA (Chapter 6).

Mean 25-OH-vitamin D levels ranged from 26 nmol/L in female residents of nursing homes to 36 nmol/L in male residents of hostels. Women were more likely to be low in vitamin D than men. There was no difference in BUA between residents on vitamin D supplementation and those not receiving therapy. Univariate associations with BUA and VOS included age, weight, serum 25-OH-vitamin D and PTH levels. Quantitative ultrasound measures were seen to change significantly at a 25-OHvitamin D cut-off of 22 nmol/L, lower than that seen to correspond to secondary hyperparathyroidism in Chapter 5. Multivariate analysis showed BUA to beassociated with mobility, gender, weight and PTH but not with serum 25-OH-vitaminD. Velocity of sound was not independently related to 25-OH-vitamin D or to PTH.

This study gives further evidence to support vitamin D supplementation in D-deficient elderly persons. Whether it is the improvement in 25-OH-vitamin D levels which is most important to maintain BUA or the suppression of secondary hyperparathyroidism is not clear. The relevance of PTH as an independent factor in bone density in the elderly requires further exploration.

(5) Falls are common in aged care facilities but not related to 25-OH-vitamin D(Chapter 7).

The incidence of falls in aged care facilities was 50% over 12 months. Serum 25-OHvitamin D levels were related to the incidence of falls in univariate analysis. Fallers were significantly older, lighter, required more walking aids, more nursing care and had lower serum 25-OH-vitamin D levels than non-fallers over 12 months. The median time to first fall was 274 days. Falling was independently associated with increasing age, poorer balance, incontinence and poorer implicit health grade but not 25-OH-vitamin D or PTH. When subgroup analyses were performed, falling was associated with higher PTH levels in residents under 90 years of age who could walk but not in older residents.

It may be that falling in the frail elderly is related to secondary hyperparathyroidism, which obscured a possible link with vitamin D deficiency. It is important that the mechanism of PTH action in muscle is more fully described in the setting of muscle

8.3 Future directions

This thesis has described the characteristics of elderly residents of aged care facilities in the northern Sydney area, with respect to risk factors for falls and fractures. We know that they are osteoporotic, which confers an increased risk of low-trauma fracture. We know they have a high incidence of falling, which also predisposes to fracture. The predictive role of serum 25-OH-vitamin D is less clear; this population is overwhelmingly deficient in 25-OH-vitamin D but this thesis does not show a straightforward relationship between vitamin D and either falling or QUS measures. The complex interplay between serum 25-OH-vitamin D and PTH and their relationship to objective measures of falls risks needs further investigation.

Analysis of muscle strength and quantitative balance will also be important to quantify the contribution of falls to the prospective incidence of fracture. Ultimately it is important that we are able to define discrete risk factors for osteoporotic fracture, which might be modified by targeted interventions to reduce the incidence of fracture in the frail institutionalized elderly and thereby improve not just morbidity and mortality, but quality of life. A trend is emerging that implicates vitamin D as important in both muscle and bone strength – prospective studies need to confirm this relationship to allow this relatively cheap and simple intervention to be introduced into everyday practice.

The most important issue to be addressed is which, if any, of the demographic and physiologic characteristics of this group predict fracture. The FREE study is measuring prospective fracture incidence at all sites, and so the baseline characteristics of the population described in this thesis will be examined for any relationship to fracture. This will further define the role of both vitamin D and PTH in falls and fractures.

Reference List

- Sambrook PN, Seeman E, Phillips SR, Ebeling PR, Osteoporosis A, National Prescribing Service. Preventing osteoporosis: outcomes of the Australian Fracture Prevention Summit. Med J Aust 2002; 176 Suppl:S1-16.
- (2) Melton LJ, III, Gabriel SE, Crowson CS, Tosteson AN, Johnell O, Kanis JA. Costequivalence of different osteoporotic fractures. Osteoporos Int 2003; 14(5):383-388.
- (3) Suh TT, Lyles KW. Osteoporosis considerations in the frail elderly. Curr Opin Rheumatol 2003; 15(4):481-486.
- (4) Cameron JR, Sorenson J. Measurement of bone mineral in-vivo: an improved method. Science 1963; 142:230-236.
- (5) Blake GM, Wahner HW, Fogelman I. Technical principles of x-ray absorptiometry. The Evaluation of Osteoporosis: Dual energy x-ray absorptiometry and ultrasound in clinical practice. Martin Dunitz Ltd, 1999: 45-72.
- (6) Kalender WA. Effective dose values in bone mineral measurements by photon absorptiometry and computed tomography. Osteoporos Int 1992; 2(2):82-87.
- (7) Meunier PJ, Slosman DO, Delmas PD, Sebert JL, Albanese C, Tandi ML et al. Strontium ranelate as a treatment of vertebral osteoporosis. J Bone Miner Res 1997; 12(Suppl):S129.
- (8) Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. Report of a WHO Study Group. World Health Organization Technical Report Series 1994; 843:1-129.
- (9) Kanis JA, Melton LJ, III, Christiansen C, Johnston CC, Khaltaev N. The diagnosis of osteoporosis. J Bone Miner Res 1994; 9(8):1137-1141.
- (10) Faulkner KG, Orwoll E. Implications in the use of T-scores for the diagnosis of osteoporosis in men. J Clin Densitom 2002; 5(1):87-93.
- (11) Faulkner KG, von Stetten E, Miller P. Discordance in patient classification using Tscores. J Clin Densitom 1999; 2(3):343-350.
- (12) Cummings SR, Bates D, Black DM. Clinical use of bone densitometry: scientific review. JAMA 2002; 288(15):1889-1897.
- (13) Cummings SR, Black DM, Nevitt MC, Browner W, Cauley J, Ensrud K et al. Bone density at various sites for prediction of hip fractures. The Study of Osteoporotic Fractures Research Group. Lancet 1993; 341(8837):72-75.
- (14) Marshall D, Johnell O, Wedel H. Meta-analysis of how well measures of bone mineral density predict occurrence of osteoporotic fractures. BMJ 1996; 312:1254-1259.
- (15) Abrahamsen B, Hansen TB, Jensen LB, Hermann AP, Eiken P. Site of osteodensitometry in perimenopausal women: correlation and limits of agreement between anatomic regions. J Bone Miner Res 1997; 12(9):1471-1479.

- (16) O'Gradaigh D, Debiram I, Love S, Richards HK, Compston JE. A prospective study of discordance in diagnosis of osteoporosis using spine and proximal femur bone densitometry. Osteoporos Int 2003; 14(1):13-18.
- (17) Schott AM, Cormier C, Hans D, Favier F, Hausherr E, Dargent-Molina P et al. How hip and whole-body bone mineral density predict hip fracture in elderly women: the EPIDOS Prospective Study. Osteoporos Int 1998; 8(3):247-254.
- (18) Chandler JM, Zimmerman SI, Girman CJ. Low bone mineral density and risk of fracture in white female nursing home residents. JAMA 2000; 284:972-977.
- (19) Hans D, Arlot ME, Schott AM, Roux JP, Kotzki PO, Meunier PJ. Do ultrasound measurements on the os calcis reflect more the bone microarchitecture than the bone mass?: a two-dimensional histomorphometric study. Bone 1995; 16(3):295-300.
- (20) Gluer CC, Wu CY, Genant HK. Broadband ultrasound attenuation signals depend on trabecular orientation: an in vitro study. Osteoporos Int 1993; 3(4):185-191.
- (21) Bouxsein ML, Radloff SE. Quantitative ultrasound of the calcaneus reflects the mechanical properties of calcaneal trabecular bone. J Bone Miner Res 1997; 12(5):839-846.
- (22) Lee SC, Coan BS, Bouxsein ML. Tibial ultrasound velocity measured in situ predicts the material properties of tibial cortical bone. Bone 1997; 21(1):119-125.
- (23) Blake GM, Wahner HW, Fogelman I. Commercial Ultrasound Instruments. The Evaluation of Osteoporosis: Dual energy x-ray absorptiometry and ultrasound in clinical practice. Martin Dunitz Ltd, 1999: 127-146.
- (24) Cheng S, Tylavsky F, Rho JY, Carbone L. Do collagen diseases have an abnormal effect on ultrasound assessment? J Bone Miner Res 1997;(Suppl.):S175.
- (25) Pocock NA. Quantitative diagnostic methods in osteoporosis: a review. Australas Radiol 1998; 42(4):327-334.
- (26) Agren M, Karellas A, Leahey D, Marks S, Baran D. Ultrasound attenuation of the calcaneus: a sensitive and specific discriminator of osteopenia in postmenopausal women. Calcif Tissue Int 1991; 48(4):240-244.
- (27) Pluskiewicz W, Drozdzowska B. Ultrasound measurements at the calcaneus in men: differences between healthy and fractured persons and the influence of age and anthropometric features on ultrasound parameters. Osteoporos Int 1999; 10(1):47-51.
- (28) Adler RA, Funkhouser HL, Holt CM. Utility of heel ultrasound bone density in men. J Clin Densitom 2001; 4(3):225-230.
- (29) Pluskiewicz W, Drozdzowska B. Ultrasonic measurement of the calcaneus in Polish normal and osteoporotic women and men. Bone 1999; 24(6):611-617.
- (30) Karlsson MK, Duan Y, Ahlborg H, Obrant KJ, Johnell O, Seeman E. Age, gender, and fragility fractures are associated with differences in quantitative ultrasound independent of bone mineral density. Bone 2001; 28(1):118-122.

- (31) Minnisola S, Tirafili C, Rosso R, Costa G, Mazzouli G. Ultrasonic parameters of bone in young normals: effect of age and sex. Eur J Clin Invest 1996; 26(11):947-950.
- (32) Takeda N, Miyake M, Kita S, Tomomitsu T, Fukunaga M. Sex and age patterns of quantitative ultrasound densitometry of the calcaneus in normal Japanese subjects. Calcif Tissue Int 1996; 59(2):84-88.
- (33) Stewart A, Torgerson DJ, Reid DM. Prediction of fractures in perimenopausal women: a comparison of dual energy x ray absorptiometry and broadband ultrasound attenuation. Ann Rheum Dis 1996; 55(2):140-142.
- (34) Kang C, Speller R. Comparison of ultrasound and dual energy X-ray absorptiometry measurements in the calcaneus. Br J Radiol 1998; 71(848):861-867.
- (35) Graafmans WC, von Lingen A, Ooms ME. Ultrasound measurements in the calcaneus: Precision and its relation with bone mineral density of the heel, hip and lumbar spine. Bone 1996; 19(2):97-100.
- (36) Knapp KM, Blake GM, Spector TD, Fogelman I. Multisite quantitative ultrasound: precision, age- and menopause-related changes, fracture discrimination, and Tscore equivalence with dual-energy X-ray absorptiometry. Osteoporos Int 2001; 12(6):456-464.
- (37) Naganathan V, March L, Hunter D, Pocock NA, Markovey J, Sambrook PN. Quantitative heel ultrasound as a predictor for osteoporosis. Med J Aust 1999; 171(6):297-300.
- (38) Cheng S, Fan B, Wang L, Fuerst T, Lian M, Njeh C et al. Factors affecting broadband ultrasound attenuation results of the calcaneus using a gel-coupled quantitative ultrasound scanning system. Osteoporos Int 1999; 10(6):495-504.
- (39) Frost ML, Blake GM, Fogelman I. Quantitative ultrasound and bone mineral density are equally strongly associated with risk factors for osteoporosis. J Bone Miner Res 2001; 16(2):406-416.
- (40) Truscott JG. Reference data for ultrasonic bone measurement: variation with age in 2087 Caucasian women aged 16-93 years. Br J Radiol 1997; 70(838):1010-1016.
- (41) Langton CM, Langton DK. Male and female normative data for ultrasound measurement of the calcaneus within the UK adult population. Br J Radiol 1997; 70(834):580-585.
- (42) Sosa M, Saavedra P, Munoz-Torres M, Alegre J, Gomez C, Gonzalez-Macias J et al. Quantitative ultrasound calcaneus measurements: normative data and precision in the Spanish population. Osteoporos Int 2002; 13(6):487-492.
- (43) Cheng S, Tylavsky F, Carbone L. Utility of ultrasound to assess risk of fracture. J Am Geriatr Soc 1997; 45(11):1382-1394.
- (44) Njeh CF, Boivin CM, Langton CM. The role of ultrasound in the assessment of osteoporosis: a review. Osteoporos Int 1997; 7(1):7-22.
- (45) Gregg EW, Kriska AM, Salamone LM, Roberts MM, Anderson SJ, Ferrell RE et al. The epidemiology of quantitative ultrasound: a review of the relationships with bone mass, osteoporosis and fracture risk. Osteoporos Int 1997; 7(2):89-99.

- (47) Porter RW, Johnson K, McCutchan JD. Wrist fracture, heel bone density and thoracic kyphosis: a case control study. Bone 1990; 11(3):211-214.
- (48) Kroger H, Jurvelin J, Arnala I, Penttila K, Rask A, Vainio P et al. Ultrasound attenuation of the calcaneus in normal subjects and in patients with wrist fracture. Acta Orthopaed Scand 1995; 66(1):47-52.
- (49) Schott AM, Weill-Engerer S, Hans D, Duboeuf F, Delmas PD, Meunier PJ. Ultrasound discriminates patients with hip fracture equally well as dual energy Xray absorptiometry and independently of bone mineral density. J Bone Miner Res 1995; 10(2):243-249.
- (50) Hadji P, Hars O, Gorke K, Emons G, Schulz KD. Quantitative ultrasound of the os calcis in postmenopausal women with spine and hip fracture. J Clin Densitom 2000; 3(3):233-239.
- (51) Kung AW, Luk KD, Chu LW, Tang GW. Quantitative ultrasound and symptomatic vertebral fracture risk in Chinese women. Osteoporos Int 1999; 10(6):456-461.
- (52) Mikhail MB, Flaster E, Aloia JF. Stiffness in discrimination of patients with vertebral fractures. Osteoporos Int 1999; 9(1):24-28.
- (53) Gonnelli S, Cepollaro C, Agnusdei D, Palmieri R, Rossi S, Gennari C. Diagnositc value of ultrasound analysis and bone densitometry as predictors of vertebral deformity in postmenopausal women. Osteoporos Int 1995; 5:413-418.
- (54) Huang C, Ross PD, Yates AJ, Walker RE, Imose K, Emi K et al. Prediction of fracture risk by radiographic absorptiometry and quantitative ultrasound: a prospective study. Calcif Tissue Int 1998; 63(5):380-384.
- (55) Hans D, Dargent-Molina P, Schott AM. Ultrasonographic heel measurements to predict hip fracture in elderly women: the EPIDOS prospective study. Lancet 1996; 348:511-514.
- (56) Bauer DC, Gluer CC, Cauley JA, Vogt TM, Ensrud KE, Genant HK et al. Broadband ultrasound attenuation predicts fractures strongly and independently of densitometry in older women. A prospective study. Study of Osteoporotic Fractures Research Group. Arch Intern Med 1997; 157(6):629-634.
- (57) Porter RW, Miller CG, Grainger D, Palmer SB. Prediction of hip fracture in elderly women: a prospective study. BMJ 1990; 301(6753):638-641.
- (58) Pluijm SM, Graafmans WC, Bouter LM, Lips P. Ultrasound measurements for the prediction of osteoporotic fractures in elderly people. Osteoporos Int 1999; 9(6):550-556.
- (59) Grampp S, Genant HK, Mathur A, Lang P, Jergas M, Takada M et al. Comparisons of noninvasive bone mineral measurements in assessing age-related loss, fracture discrimination, and diagnostic classification. J Bone Miner Res 1997; 12(5):697-711.
- (60) Rehman Q, Lang T, Modin G, Lane NE. Quantitative computed tomography of the lumbar spine, not dual x-ray absorptiometry, is an independent predictor of

prevalent vertebral fractures in postmenopausal women with osteopenia receiving long-term glucocorticoid and hormone-replacement therapy. Arthritis Rheum 2002; 46(5):1292-1297.

- (61) Eyre DR. Bone biomarkers as tools in osteoporosis management. Spine 1997; 22(24S):17S-24S.
- (62) Garnero P, Sornay-Rendu E, Chapuy MC, Delmas PD. Increased bone turnover in late postmenopausal women is a major determinant of osteoporosis. J Bone Miner Res 1996; 11(3):337-349.
- (63) Melton LJ, III, Khosla S, Atkinson EJ, O'fallon WM, Riggs BL. Relationship of bone turnover to bone density and fractures. J Bone Miner Res 1997; 12:1083-1091.
- (64) Rosen CJ, Chesnut CH, III, Mallinak NJS. The predictive value of bichemical markers of bone turnover for bone mineral density in early postmenopausal women treated with hormone replacement or calcium supplementation. J Clin Endocrinol Metab 1997; 82:1904-1910.
- (65) Schneider DL, Barrett-Connor EL. Urinary N-telopeptide levels discriminate normal, osteopenic and osteoporotic bone mineral density. Arch Intern Med 1997; 157:1241-1245.
- (66) Ross P, Knowlton W. Rapid bone loss is associated with increased levels of biochemical markers. J Bone Miner Res 1998; 13:297-302.
- (67) Garnero P, Hausherr E, Chapuy MC, Marcelli C, Grandjean H, Muller C et al. Markers of bone resorption predict hip fracture in elderly women: the EPIDOS Prospective Study. J Bone Miner Res 1996; 11(10):1531-1538.
- (68) Ravn P, Rix M, Andreassen H, Clemmesen B, Bidstrup M, Gunnes M. High bone turnover is associated with low bone mass and spinal fracture in postmenopausal women. Calcif Tissue Int 1997; 60:255-260.
- (69) Souberbielle J-C, Cormier C, Kindermans C. Bone markers in clinical practice. Curr Opin Rheumatol 1999; 11(4):312.
- (70) Ebeling PR, Akesson K. Role of biochemical markers in the management of osteoporosis. Best Pract Res Clin Rheum 2001; 15(3):385-400.
- (71) Rupich RC, Specker BL, Lieuw AF, Ho M. Gender and race differences in bone mass during infancy. Calcif Tissue Int 1996; 58(6):395-397.
- (72) Horlick M, Thornton J, Wang J, Levine LS, Fedun B, Pierson RN, Jr. Bone mineral in prepubertal children: gender and ethnicity. J Bone Miner Res 2000; 15(7):1393-1397.
- (73) Nguyen TV, Maynard LM, Towne B, Roche AF, Wisemandle W, Li J et al. Sex differences in bone mass acquisition during growth: the Fels Longitudinal Study. J Clin Densitom 2001; 4(2):147-157.
- (74) Jones G, Dwyer T. Bone mass in prepubertal children: gender differences and the role of physical activity and sunlight exposure. J Clin Endocrinol Metab 1998; 83(12):4274-4279.

- (75) Henry YM, Eastell R. Ethnic and gender differences in bone mineral density and bone turnover in young adults: effect of bone size. Osteoporos Int 2000; 11(6):512-517.
- (76) del Rio L, Carrascosa A, Pons F, Gusinye M, Yeste D, Domenech FM. Bone mineral density of the lumbar spine in white Mediterranean Spanish children and adolescents: changes related to age, sex, and puberty. Pediatr Res 1994; 35(3):362-366.
- (77) Magarey AM, Boulton TJ, Chatterton BE, Schultz C, Nordin BE, Cockington RA. Bone growth from 11 to 17 years: relationship to growth, gender and changes with pubertal status including timing of menarche. Acta Paediatrica 1999; 88(2):139-146.
- (78) Schoenau E, Neu CM, Rauch F, Manz F. Gender-specific pubertal changes in volumetric cortical bone mineral density at the proximal radius. Bone 2002; 31(1):110-113.
- (79) Matsukura T, Kagamimori S, Yamagami T, Nishino H, Iki M, Kajita E et al. Reference data of forearm bone mineral density in healthy Japanese male and female subjects in the second decade based on calendar age and puberty onset: Japanese Population Based Osteoporosis (JPOS) study. Osteoporos Int 2000; 11(10):858-865.
- (80) Naganathan V, Macgregor A, Snieder H, Nguyen T, Spector T, Sambrook P. Gender differences in the genetic factors responsible for variation in bone density and ultrasound. J Bone Miner Res 2002; 17(4):725-733.
- (81) Sundberg M, Gardsell P, Johnell O, Karlsson MK, Ornstein E, Sandstedt B et al. Peripubertal moderate exercise increases bone mass in boys but not in girls: a population-based intervention study. Osteoporos Int 2001; 12(3):230-238.
- (82) Florindo AA, Latorre MR, Jaime PC, Tanaka T, Pippa MG, Zerbini CA. Past and present habitual physical activity and its relationship with bone mineral density in men aged 50 years and older in Brazil. J Gerontology Series A-Biological Sciences & Medical Sciences 2002; 57(10):M654-M657.
- (83) Neville CE, Murray LJ, Boreham CA, Gallagher AM, Twisk J, Robson PJ et al. Relationship between physical activity and bone mineral status in young adults: the Northern Ireland Young Hearts Project. Bone 2002; 30(5):792-798.
- (84) Burger H, van Daele PL, Algra D, van den Ouweland FA, Grobbee DE, Hofman A et al. The association between age and bone mineral density in men and women aged 55 years and over: the Rotterdam Study. Bone & Mineral 1994; 25(1):1-13.
- (85) Tsai KS, Cheng WC, Sanchez TV, Chen CK, Chieng PU, Yang RS. Bone densitometry of proximal femur in Chinese subjects: gender differences in bone mass and bone areas. Bone 1997; 20(4):365-369.
- (86) Warming L, Hassager C, Christiansen C. Changes in bone mineral density with age in men and women: a longitudinal study. Osteoporos Int 2002; 13(2):105-112.
- (87) Clarke BL, Ebeling PR, Jones JD, Wahner HW, O'fallon WM, Riggs BL et al. Predictors of bone mineral density in aging healthy men varies by skeletal site. Calcif Tissue Int 2002; 70(3):137-145.

- (88) Blunt BA, Klauber MR, Barrett-Connor EL, Edelstein SL. Sex differences in bone mineral density in 1653 men and women in the sixth through tenth decades of life: the Rancho Bernardo Study. J Bone Miner Res 1994; 9(9):1333-1338.
- (89) Krall EA, Dawson-Hughes B, Hirst K, Gallagher JC, Sherman SS, Dalsky G. Bone mineral density and biochemical markers of bone turnover in healthy elderly men and women. J Gerontology Series A-Biological Sciences & Medical Sciences 1997; 52(2):M61-M67.
- (90) Chavassieux P, Meunier PJ. Histomorphometric approach of bone loss in men. Calcif Tissue Int 2001; 69(4):209-213.
- (91) Compston JE, Mellish RWE, Croucher P, Newcombe R, Garrahan NJ. Structural mechanisms of trabecular bone loss in man. Bone Miner 1989; 6:339-350.
- (92) Croucher PI, Mellish RWE, Vedi S, Garrahan NJ, Compston JE. The relationship between resorption depth and mean interstitial bone thickness: age-related changes in man. Calcif Tissue Int 1989; 45:15-19.
- (93) Mellish RWE, Garrahan NJ, Compston JE. Age-related changes in trabecular width and spacing in human iliac crest biopsies. Bone Miner 1989; 6:331-338.
- (94) Aaron JE, Makins NB, Sagreiya K. The microanatomy of trabecular bone loss in normal aging men and women. Clin Orthop Rel Res 1987; 215:260-271.
- (95) Rogucka E, Bielicki T, Welon Z, Medras M, Susanne C. Variation in bone mineral density in adults in Poland: age and sex differences. Ann Hum Biol 2000; 27(2):139-148.
- (96) Berntsen GK, Fonnebo V, Tollan A, Sogaard AJ, Magnus JH. Forearm bone mineral density by age in 7,620 men and women: the Tromso study, a populationbased study. Am J Epidemiol 2001; 153(5):465-473.
- (97) Waud CE, Lew R, Baran DT. The relationship between ultrasound and densitometric measurements of bone mass at the calcaneus in women. Calcif Tissue Int 1992; 51(6):415-418.
- (98) van Daele PL, Burger H, Algra D, Hofman A, Grobbee DE, Birkenhager JC et al. Age-associated changes in ultrasound measurements of the calcaneus in men and women: the Rotterdam Study. J Bone Miner Res 1994; 9(11):1751-1757.
- (99) Palacios S, Menendez C, Calderon J, Rubio S. Spine and femur density and broadband ultrasound attenuation of the calcaneus in normal Spanish women. Calcif Tissue Int 1993; 52(2):99-102.
- (100) Herd RJ, Ramalingham T, Ryan PJ, Fogelman I, Blake GM. Measurements of broadband ultrasonic attenuation in the calcaneus in premenopausal and postmenopausal women. Osteoporos Int 1992; 2(5):247-251.
- (101) Krieg MA, Thiebaud D, Burckhardt P. Quantitative ultrasound of bone in institutionalized elderly women: a cross-sectional and longitudinal study. Osteoporos Int 1996; 6(3):189-195.
- (102) Damilakis JE, Dretakis E, Gourtsoyiannis NC. Ultrasound attenuation of the calcaneus in the female population: normative data. Calcif Tissue Int 1992; 51(3):180-183.

- (103) Yamazaki K, Kushida K, Ohmura A, Sano M, Inoue T. Ultrasound bone densitometry of the os calcis in Japanese women. Osteoporos Int 1994; 4(4):220-225.
- (104) Schott AM, Hans D, Sornay-Rendu E, Delmas PD, Meunier PJ. Ultrasound measurements on os calcis: precision and age-related changes in a normal female population. Osteoporos Int 1993; 3(5):249-254.
- (105) Schott AM, Hans D, Garnero P, Sornay-Rendu E, Delmas PD, Meunier PJ. Agerelated changes in Os calcis ultrasonic indices: a 2-year prospective study. Osteoporos Int 1995; 5(6):478-483.
- (106) Sahota O, San P, Cawte SA, Pearson D, Hosking DJ. A comparison of the longitudinal changes in quantitative ultrasound with dual-energy X-ray absorptiometry: the four-year effects of hormone replacement therapy. Osteoporos Int 2000; 11(1):52-58.
- (107) Frost ML, Blake GM, Fogelman. Changes in QUS and BMD measurements with antiresorptive therapy: a two-year longitudinal study. Calcif Tissue Int 2001; 69(3):138-146.
- (108) Gonnelli S, Cepollaro C, Montagnani A, Martini S, Gennari L, Mangeri M et al. Heel ultrasonography in monitoring alendronate therapy: a four-year longitudinal study. Osteoporos Int 2002; 13(5):415-421.
- (109) Gonnelli S, Cepollaro C, Pondrelli C, Martini S, Rossi S, Gennari C. Ultrasound parameters in osteoporotic patients treated with salmon calcitonin: a longitudinal study. Osteoporos Int 1996; 6(4):303-307.
- (110) Krieg MA, Jacquet AF, Bremgartner M, Cuttelod S, Thiebaud D, Burckhardt P. Effect of supplementation with vitamin D3 and calcium on quantitative ultrasound of bone in elderly institutionalized women: a longitudinal study. Osteoporos Int 1999; 9(6):483-488.
- (111) Duan Y, Turner CH, Kim BT, Seeman E. Sexual dimorphism in vertebral fragility is more the result of gender differences in age-related bone gain than bone loss. J Bone Miner Res 2001; 16(12):2267-2275.
- (112) Duan Y, Seeman E, Turner CH. The biomechanical basis of vertebral body fragility in men and women. J Bone Miner Res 2001; 16(12):2276-2283.
- (113) Arfai K, Pitukcheewanont PD, Goran MI, Tavare CJ, Heller L, Gilsanz V. Bone, muscle, and fat: sex-related differences in prepubertal children. Radiology 2002; 224(2):338-344.
- (114) Looker AC, Beck TJ, Orwoll ES. Does body size account for gender differences in femur bone density and geometry? J Bone Miner Res 2001; 16(7):1291-1299.
- (115) Seeman E. During aging, men lose less bone than women because they gain more periosteal bone, not because they resorb less endosteal bone. Calcif Tissue Int 2001; 69(4):205-208.
- (116) Gilsanz V, Boechat MI, Gilsanz R, Loro ML, Roe TF, Goodman WG. Gender differences in vertebral sizes in adults: biomechanical implications. Radiology 1994; 190(3):678-682.

- (117) Naganathan V, Sambrook P. Gender differences in volumetric bone density: a study of opposite-sex twins. Osteoporos Int 2003; 14(7):564-569.
- (118) Ebbesen EN, Thomsen JS, Beck-Nielsen H, Nepper-Rasmussen HJ, Mosekilde L. Age- and gender-related differences in vertebral bone mass, density, and strength. J Bone Miner Res 1999; 14(8):1394-1403.
- (119) Oyster N. Sex differences in cancellous and cortical bone strength, bone mineral content and bone density. Age & Ageing 1992; 21(5):353-356.
- (120) Faulkner KG, Cummings SR, Black D, Palermo L, Gluer CC, Genant HK. Simple measurement of femoral geometry predicts hip fracture: the Study of Osteoporotic Fractures. J Bone Miner Res 1993; 8:1211-1217.
- (121) Peacock M, Turner CH, Liu G, Manatunga AK, Timmerman L, Johnston CCJr. Better discrimination of hip fracture using bone density, geometry and architecture. Osteoporos Int 1995; 5:167-173.
- (122) Crabtree N, Lunt M, Holt G, Kroger H, Burger H, Grazio S et al. Hip geometry, bone mineral distribution, and bone strength in European men and women: the EPOS study. Bone 2000; 27(1):151-159.
- (123) Hoffler CE, Moore KE, Kozloff K, Zysset PK, Goldstein SA. Age, gender, and bone lamellae elastic moduli. J Ortho Res 2000; 18(3):432-437.
- (124) MacIntyre NJ, Adachi JD, Webber CE. Gender differences in normal agedependent patterns of radial bone structure and density: a cross-sectional study using peripheral quantitative computed tomography. J Clin Densitom 1999; 2(2):163-173.
- (125) Chailurkit LO, Pongchaiyakul C, Charoenkiatkul S, Kosulwat V, Rojroongwasinkul N, Rajatanavin R. Different mechanism of bone loss in ageing women and men in Khon Kaen Province. J Med Assoc Thailand 2001; 84(8):1175-1182.
- (126) Chailurkit LO, Piaseu N, Rajatanavin R. Influence of normal ageing on mechanism of bone loss in women and men in Bangkok. J Med Assoc Thailand 2002; 85(8):915-921.
- (127) Midtby M, Magnus JH, Joakimsen RM. The Tromso Study: a population-based study on the variation in bone formation markers with age, gender, anthropometry and season in both men and women. Osteoporos Int 2001; 12(10):835-843.
- (128) Tsai KS, Pan WH, Hsu SH, Cheng WC, Chen CK, Chieng PU et al. Sexual differences in bone markers and bone mineral density of normal Chinese. Calcif Tissue Int 1996; 59(6):454-460.
- (129) Ebeling PR, Atley LM, Guthrie JR, Burger HG, Dennerstein L, Hopper JL et al. Bone turnover markers and bone density across the menopausal transition. J Clin Endocrinol Metab 1996; 81(9):3366-3371.
- (130) Haden ST, Brown EM, Hurwitz S, Scott J, El Hajj FG. The effects of age and gender on parathyroid hormone dynamics. Clin Endocrinol 2000; 52(3):329-338.
- (131) Cheng S, Suominen H, Heikkinen E. Bone mineral density in relation to anthropometric properties, physical activity and smoking in 75-year-old men and women. Aging (Milano) 1993; 5(1):55-62.

- (132) Felson DT, Zhang Y, Hannan MT, Anderson JJ. Effects of weight and body mass index on bone mineral density in men and women: the Framingham study. J Bone Miner Res 1993; 8(5):567-573.
- (133) Kirchengast S, Knogler W, Hauser G. Protective effect of moderate overweight on bone density of the hip joint in elderly and old Austrians. Anthropologischer Anzeiger 2002; 60(2):187-197.
- (134) Guthrie JR, Ebeling PR, Hopper JL, Barrett-Connor E, Dennerstein L, Dudley EC et al. A prospective study of bone loss in menopausal Australian-born women. Osteoporos Int 1998; 8(3):282-290.
- (135) Kirchengast S, Peterson B, Hauser G, Knogler W. Body composition characteristics are associated with the bone density of the proximal femur end in middle- and old-aged women and men. Maturitas 2001; 39(2):133-145.
- (136) Taaffe DR, Cauley JA, Danielson M, Nevitt MC, Lang TF, Bauer DC et al. Race and sex effects on the association between muscle strength, soft tissue, and bone mineral density in healthy elders: the Health, Aging, and Body Composition Study. J Bone Miner Res 2001; 16(7):1343-1352.
- (137) Ferretti JL, Capozza RF, Cointry GR, Garcia SL, Plotkin H, Alvarez Filgueira ML et al. Gender-related differences in the relationship between densitometric values of whole-body bone mineral content and lean body mass in humans between 2 and 87 years of age. Bone 1998; 22(6):683-690.
- (138) Tsunenari T, Tsutsumi M, Ohno K, Yamamoto Y, Kawakatsu M, Shimogaki K et al. Age- and gender-related changes in body composition in Japanese subjects. J Bone Miner Res 1993; 8(4):397-402.
- (139) Ismail AA, Pye SR, Cockerill WC, Lunt M, Silman AJ, Reeve J et al. Incidence of limb fracture across Europe: results from the European Prospective Osteoporosis Study (EPOS). Osteoporos Int 2002; 13(7):565-571.
- (140) Sanders KM, Seeman E, Ugoni AM, Pasco JA, Martin TJ, Skoric B et al. Age- and gender-specific rate of fractures in Australia: a population-based study. Osteoporos Int 1999; 10(3):240-247.
- (141) Lofthus CM, Osnes EK, Falch JA, Kaastad TS, Kristiansen IS, Nordsletten L et al. Epidemiology of hip fractures in Oslo, Norway. Bone 2001; 29(5):413-418.
- (142) Jackson SA, Tenenhouse A, Robertson L. Vertebral fracture definition from population-based data: preliminary results from the Canadian Multicenter Osteoporosis Study (CaMos). Osteoporos Int 2000; 11(8):680-687.
- (143) Singer BR, McLauchlan GJ, Robinson CM, Christie J. Epidemiology of fractures in 15,000 adults: the influence of age and gender. J Bone Joint Surg - Brit 1998; 80(2):243-248.
- (144) Center JR, Nguyen TV, Schneider D, Sambrook PN, Eisman JA. Mortality after all major types of osteoporotic fracture in men and women: an observational study. Lancet 1999; 353(9156):878-882.
- (145) Trombetti A, Herrmann F, Hoffmeyer P, Schurch MA, Bonjour JP, Rizzoli R. Survival and potential years of life lost after hip fracture in men and age-matched women. Osteoporos Int 2002; 13(9):731-737.

- (146) Lunt M, Felsenberg D, Reeve J, Benevolenskaya L, Cannata J, Dequeker J et al. Bone density variation and its effects on risk of vertebral deformity in men and women studied in thirteen European centers: the EVOS Study. J Bone Miner Res 1997; 12(11):1883-1894.
- (147) Kudlacek S, Schneider B, Resch H, Freudenthaler O, Willvonseder R. Gender differences in fracture risk and bone mineral density. Maturitas 2000; 36(3):173-180.
- (148) Ismail AA, O'Neill TW, Cooper C, Silman AJ. Risk factors for vertebral deformities in men: relationship to number of vertebral deformities. European Vertebral Osteoporosis Study Group. J Bone Miner Res 2000; 15(2):278-283.
- (149) Selby PL, Davies M, Adams JE. Do men and women fracture bones at similar bone densities? Osteoporos Int 2000; 11(2):153-157.
- (150) Hordon LD, Raisi M, Aaron JE, Paxton SK, Beneton M, Kanis JA. Trabecular architecture in women and men of similar bone mass with and without vertebral fracture: I. Two-dimensional histology. Bone 2000; 27(2):271-276.
- (151) Silman AJ, O'Neill TW, Cooper C, Kanis J, Felsenberg D. Influence of physical activity on vertebral deformity in men and women: results from the European Vertebral Osteoporosis Study. J Bone Miner Res 1997; 12(5):813-819.
- (152) Cody DD, Divine GW, Nahigian K, Kleerekoper M. Bone density distribution and gender dominate femoral neck fracture risk predictors. Skelet Radiol 2000; 29(3):151-161.
- (153) Johnell O, Gullberg B, Kanis JA, Allander E, Elffors L, Dequeker J et al. Risk factors for hip fracture in European women: the MEDOS Study. Mediterranean Osteoporosis Study. J Bone Miner Res 1995; 10(11):1802-1815.
- (154) Kanis J, Johnell O, Gullberg B, Allander E, Elffors L, Ranstam J et al. Risk factors for hip fracture in men from southern Europe: the MEDOS study. Mediterranean Osteoporosis Study. Osteoporos Int 1999; 9(1):45-54.
- (155) Hoidrup S, Gronbaek M, Gottschau A, Lauritzen JB, Schroll M. Alcohol intake, beverage preference, and risk of hip fracture in men and women. Copenhagen Centre for Prospective Population Studies. Am J Epidemiol 1999; 149(11):993-1001.
- (156) Hoidrup S, Prescott E, Sorensen TI, Gottschau A, Lauritzen JB, Schroll M et al. Tobacco smoking and risk of hip fracture in men and women. Int J Epidemiol 2000; 29(2):253-259.
- (157) Kannus P, Palvanen M, Kaprio J, Parkkari F, Koskenvuo M. Genetic factors and osteoporotic fractures in elderly people: prospective 25 year follow up of a nationwide cohort of elderly Finnish twins. BMJ 1999; 319:1334-1337.
- (158) Langdahl BL, Ralston SH, Grant SF, Eriksen EF. An Sp1 binding site polymorphism in the COLIA1 gene predicts osteoporotic fractures in both men and women. J Bone Miner Res 1998; 13(9):1384-1389.
- (159) Center JR, Nguyen TV, Pocock NA, Noakes KA, Kelly PJ, Eisman JA et al. Femoral neck axis length, height loss and risk of hip fracture in males and females. Osteoporos Int 1998; 8(1):75-81.

- (160) van Staa TP, Dennison EM, Leufkens HG, Cooper C. Epidemiology of fractures in England and Wales. Bone 2001; 29:517-522.
- (161) O'Neill TW, Cooper C, Finn JD, Lunt M, Purdie DW, Reid DM et al. Incidence of distal forearm fracture in British men and women. Osteoporos Int 2001; 12:555-558.
- (162) Bengner U, Johnell O. Increaseing incidence of forearm fractures: a comparison of epidemiological patterns 25 years apart. Acta Orthopaed Scand 2003; 56:158-160.
- (163) Melton LJ, III, Amadio PC, Crowson CS, O'fallon WM. Long term trends in the incidence of disteal forearm fractures. Osteoporos Int 1998; 8:341-348.
- (164) Melton LJ, III, Achenbach SJ, O'fallon WM, Khosla S. Secondary osteoporosis and the risk of distal forearm fractures in men and women. Bone 2002; 31(1):119-125.
- (165) Campbell A, Reinken J, Martinez GS. Falls in old age: a study of frequency and related clinical factors. Age & Ageing 1981; 10:264-270.
- (166) Prudham D, Evans JG. Factors associated with falls in the elderly: a community study. Age & Ageing 1981; 10(3):141-146.
- (167) Robbins AS, Rubenstein LZ, Josephson KR, Schulman BL, Osterweil D, Fine G. Predictors of falls among elderly people. Results of two population-based studies. Arch Intern Med 1989; 149(7):1628-1633.
- (168) Tinetti ME, Williams CS. Falls, injuries due to falls, and the risk of admission to a nursing home. N Engl J Med 1997; 337(18):1279-1284.
- (169) Lord SR, Ward JA, Williams P, Anstey KJ. Physiological factors associated with falls in older community-dwelling women. J Am Geriatr Soc 1994; 42(10):1110-1117.
- (170) Baloh RW, Corona S, Jacobson KM, Enrietto JA, Bell T. A prospective study of posturography in normal older people. J Am Geriatr Soc 1998; 46(4):438-443.
- (171) Lord SR, Clark RD, Webster IW. Visual acuity and contrast sensitivity in relation to falls in an elderly population. Age & Ageing 1991; 20(3):175-181.
- (172) Rubenstein LZ, Josephson KR, Robbins AS. Falls in the nursing home. Ann Int Med 1994; 121(6):442-451.
- (173) Kiely DK, Kiel DP, Burrows AB, Lipsitz LA. Identifying nursing home residents at risk for falling. J Am Geriatr Soc 1998; 46(5):551-555.
- (174) Maciorowski L, Munro B, Dietrick-Gallagher M, McNew C, Sheppard-Hinkel I, Wanich C. A review of the patient fall literature. J Nurs Qual Ass 1988.
- (175) Nevitt MC, Cummings SR, Kidd S, Black D. Risk factors for recurrent nonsyncopal falls. A prospective study. JAMA 1989; 261(18):2663-2668.
- (176) Graafmans WC, Ooms ME, Hofstee HM, Bezemer PD, Bouter LM, Lips P. Falls in the elderly: a prospective study of risk factors and risk profiles. Am J Epidemiol 1996; 143(11):1129-1136.
- (177) Tinetti ME, Speechley M, Ginter SF. Risk factors for falls among elderly persons living in the community. N Engl J Med 1988; 319(26):1701-1707.

- (178) Campbell AJ, Borrie MJ, Spears GF. Risk factors for falls in a community-based prospective study of people 70 years and older. J Gerontol 1989; 44(4):M112-M117.
- (179) Davis JW, Ross PD, Nevitt MC, Wasnich RD. Risk factors for falls and for serious injuries on falling among older Japanese women in Hawaii. J Am Geriatr Soc 1999; 47(7):792-798.
- (180) Lord SR, Ward JA, Williams P, Anstey KJ. An epidemiological study of falls in older community-dwelling women: the Randwick falls and fractures study. Aust J Pub Health 1993; 17(3):240-245.
- (181) Kerber KA, Enrietto JA, Jacobson KM, Baloh RW. Disequilibrium in older people: a prospective study. Neurology 1998; 51(2):574-580.
- (182) Lord SR, McLean D, Stathers G. Physiological factors associated with injurious falls in older people living in the community. Gerontology 1992; 38(6):338-346.
- (183) Lord SR, Sambrook PN, Gilbert C, Kelly PJ, Nguyen T, Webster IW et al. Postural stability, falls and fractures in the elderly: results from the Dubbo Osteoporosis Epidemiology Study. Med J Aust 688; 160(11):684-685.
- (184) Dargent-Molina P, Favier F, Grandjean H, Baudoin C, Schott AM, Hausherr E et al. Fall-related factors and risk of hip fracture: the EPIDOS prospective study. Lancet 1996; 348(9021):145-149.
- (185) Lord SR, Rogers MW, Howland A, Fitzpatrick R. Lateral stability, sensorimotor function and falls in older people. J Am Geriatr Soc 1999; 47(9):1077-1081.
- (186) Kippenbrock T, Soja ME. Preventing falls in the elderly: interviewing patients who have fallen. Researchers attempt to identify fall-risk factors from the patients' point of view. Geriatr Nurs 1993; 14(4):205-209.
- (187) Biderman A, Cwikel J, Fried AV, Galinsky D. Depression and falls among community dwelling elderly people: a search for common risk factors. J Epidemiol Comm Health 2002; 56(8):631-636.
- (188) Gostynski M, Ajdacic-Gross V, Heusser-Gretler R, Gutzwiller F, Michel JP, Herrmann F. Dementia, depression and activity of daily living as risk factors for falls in elderly patients. Sozial- und Praventivmedizin 2001; 46(2):123-130.
- (189) Tinetti ME, Inouye SK, Gill TM, Doucette JT. Shared risk factors for falls, incontinence, and functional dependence. Unifying the approach to geriatric syndromes. JAMA 1995; 273(17):1348-1353.
- (190) Brown JS, Vittinghoff E, Wyman JF, Stone KL, Nevitt MC, Ensrud KE et al. Urinary incontinence: does it increase risk for falls and fractures? Study of Osteoporotic Fractures Research Group. J Am Geriatr Soc 2000; 48(7):721-725.
- (191) Ensrud KE, Blackwell TL, Mangione CM, Bowman PJ, Whooley MA, Bauer DC et al. Central nervous system-active medications and risk for falls in older women. J Am Geriatr Soc 2002; 50(10):1629-1637.
- (192) Leipzig RM, Cumming RG, Tinetti ME. Drugs and falls in older people: a systematic review and meta-analysis: I. Psychotropic drugs. J Am Geriatr Soc 1999; 47(1):30-39.

- (193) Leipzig RM, Cumming RG, Tinetti ME. Drugs and falls in older people: a systematic review and meta-analysis: II. Cardiac and analgesic drugs. J Am Geriatr Soc 1999; 47(1):40-50.
- (194) Lim KD, Ng KC, Ng SK, Ng LL. Falls amongst institutionalised psycho-geriatric patients. Singap Med J 2001; 42(10):466-472.
- (195) Neutel CI, Perry S, Maxwell C. Medication use and risk of falls. Pharmacoepidemiol Drug Safety 2002; 11(2):97-104.
- (196) Northridge ME, Nevitt MC, Kelsey JL. Non-syncopal falls in the elderly in relation to home environments. Osteoporos Int 1996; 6(3):249-255.
- (197) Connell BR, Wolf SL. Environmental factors influencing falls in older individuals. Arch Phys Med Rehab 1997; 78:179-186.
- (198) Tinetti ME, Williams TF, Mayewski R. Falls risk index for elderly patients based on number of chronic disabilities. Am J Med 1996; 80(3):429-434.
- (199) Lipsitz LA, Nakajima I, Gagnon M, Hirayama T, Connelly CM, Izumo H et al. Muscle strength and fall rates among residents of Japanese and American nursing homes: an International Cross-Cultural Study. J Am Geriatr Soc 1994; 42(9):953-959.
- (200) Bischoff HA, Stahelin HB, Urscheler N, Ehrsam R, Vonthein R, Perrig-Chiello P et al. Muscle strength in the elderly: its relation to vitamin D metabolites. Arch Phys Med Rehab 1999; 80(1):54-58.
- (201) Glerup H, Mikkelsen K, Poulsen L, Hass E, Overbeck S, Andersen H. Hypovitaminosis D myopathy without biochemical signs of osteomalacic bone involvement. Calcif Tissue Int 2000; 66(6):419-424.
- (202) Pfeifer M, Begerow B, Minne HW. Vitamin D and muscle function. Osteoporos Int 2002; 13(3):187-194.
- (203) Baczynski R, Massry SG, Magott M, el Belbessi S, Kohan Rl, Brautbar N. Effect of parathyroid hormone on energy metabolism of skeletal muscle. Kidney Int 1985.
- (204) Joborn C, Rastad J, Stalberg E, Akerstrom G, Ljunghall S. Muscle function in patients with primary hyperparathyroidism. Muscle Nerve 1989.
- (205) Kristoffersson A, Bostrom A, Soderberg T. Muscle strength is improved after parathyroidectomy in patients with primary hyperparathyroidism. Br J Surg 1992; 79(2):165-168.
- (206) Hedman I, Grimby G, Tisell LE. Improvement of muscle strength after treatment for hyperparathyroidism. Acta Chir Scand 1984; 150(7):521-524.
- (207) Wersall-Robertson E, Hamberger B, Ehren H, Eriksson E, Granberg PO. Increase in muscular strength following surgery for primary hyperparathyroidism. Acta Med Scand 1986; 220(3):233-235.
- (208) Stein MS, Wark JD, Scherer SC, Walton SL, Chick P, Di Carlantonio M et al. Falls relate to vitamin D and parathyroid hormone in an Australian nursing home and hostel. J Am Geriatr Soc 1999; 47(10):1195-1201.

- (209) Lipsitz LA, Jonsson PV, Kelley MM, Koestner JS. Causes and correlates of recurrent falls in ambulatory frail elderly. J Gerontol 1991; 46(4):M114-M122.
- (210) Ruthazer R, Lipsitz LA. Antidepressants and falls among elderly people in long-term care. Am J Pub Health 1993; 83(5):746-749.
- (211) Thapa PB, Gideon P, Fought RL, Ray WA. Psychotropic drugs and the risk of recurrent falls in ambulatory nursing home residents. Am J Epidemiol 1995; 142:202-211.
- (212) Myers AH, Baker SP, Van Natta ML, Abbey H, Robinson EG. Risk factors associated with falls and injuries among elderly institutionalized persons. Am J Epidemiol 1991; 133(11):1179-1190.
- (213) Thapa PB, Gideon P, Cost TW, Milam AB, Ray WA. Antidepressants and the risk of falls among nursing home residents. N Engl J Med 1998; 339(13):875-882.
- (214) Ray WA, Thapa PB, Gideon P. Benzodiazepines and the risk of falls in nursing home residents. J Am Geriatr Soc 2000; 48(6):682-685.
- (215) Sgadari A, Lapane KL, Mor V, Landi F, Bernabei R, Gambassi G. Oxidative and nonoxidative benzodiazepines and the risk of femur fracture. J Clin Psychopharmacol 2000; 20(2):234-239.
- (216) Thapa PB, Brockman KG, Gideon P, Fought RL, Ray WA. Injurious falls in nonambulatory nursing home residents: a comparative study of circumstances, incidence, and risk factors. J Am Geriatr Soc 1996; 44(3):273-278.
- (217) Rawsky E. Review of the literature on falls among the elderly. J Nurs Scholar 1998; 30(1):47-52.
- (218) Morse J, Tylko S, Dixon H. Characteristics of the fall prone patient. The Gerontologist 1987; 27(4):516-522.
- (219) Lord SR, Castell S. Physical activity program for older persons: effect on balance, strength, neuromuscular control, and reaction time. Arch Phys Med Rehab 1994; 75:648-652.
- (220) Tinetti ME, Liu WL, Ginter SF. Mechanical restraint use and fall-related injuries among residents of skilled nursing facilities. Ann Int Med 1992; 116:369-374.
- (221) Ejaz F, Jones J, Rose M. Falls among nursing home residents: an examination of incident reports before and after restraint reduction programs. J Am Geriatr Soc 1994; 42:960-964.
- (222) Capezuti E, Evans L, Strumpf N, Maislin G. Physical restraint use and falls in nursing home residents. J Am Geriatr Soc 1996; 44(6):627-633.
- (223) Neufeld RR, Libow LS, Foley WJ, Dunbar JM, Cohen C, Breuer B. Restraint reduction reduces serious injuries among nursing home residents. Journal of the American Geriatrics Society 1999; 47(10):1202-1207.
- (224) Tinetti ME. Factors associated with serious injury during falls by ambulatory nursing home residents. Journal of the American Geriatrics Society 2003; 35:644-648.

- (225) Lord SR, Lloyd DG, Li SK. Sensori-motor function, gait patterns and falls in community-dwelling women. Age & Ageing 1996; 25(4):292-299.
- (226) Pfeifer M, Begerow B, Minne HW, Abrams C, Nachtigall D, Hansen C. Effects of a short-term vitamin D and calcium supplementation on body sway and secondary hyperparathyroidism in elderly women. J Bone Miner Res 2000; 15(6):1113-1118.
- (227) Latham NK, Anderson CS, Lee A, Bennett DA, Moseley A, Cameron ID. A randomized, controlled trial of quadriceps resistance exercise and vitamin D in frail older people: the frailty interventions trial in elderly subjects (FITNESS). J Am Geriatr Soc 2003; 51:291-299.
- (228) Sato Y, Manabe S, Kuno H, Oizumi K. Amelioration of osteopenia and hypovitaminosis D by 1alpha-hydroxyvitamin D3 in elderly patients with Parkinson's disease. J Neurol Neurosurg Psychiatr 1999; 66(1):64-68.
- (229) Dawson-Hughes B, Harris SS, Krall EA, Dallal GE. Effect of calcium and vitamin D supplementation on bone density in men and women 65 years of age or older. N Engl J Med 1997; 337(10):670-676.
- (230) Bischoff HA, Stahelin HB, Dick W, Akos R, Knecht M, Salis C et al. Effects of vitamin D and calcium supplementation on falls: a randomized controlled trial. J Bone Miner Res 2003; 18(2):343-351.
- (231) Sanders KM, Nicholson GC, Ugoni AM, Pasco JA, Seeman E, Kotowicz MA. Health burden of hip and other fractures in Australia beyond 2000: Projections based on the Geelong Osteoporosis Study. Med J Aust 1999; 170:467-470.
- (232) Jones G, Nguyen T, Sambrook PN, Kelly PJ, Gilbert C, Eisman JA. Symptomatic fracture incidence in elderly men and women: the Dubbo Osteoporosis Epidemiology Study (DOES). Osteoporos Int 1994; 4(5):277-282.
- (233) Incidence of vertebral fracture in europe: results from the European Prospective Osteoporosis Study (EPOS). J Bone Miner Res 2002; 17(4):716-724.
- (234) Leidig-Bruckner G, Limberg B, Felsenberg D, Bruckner T, Holder S, Kather A et al. Sex difference in the validity of vertebral deformities as an index of prevalent vertebral osteoporotic fractures: a population survey of older men and women. Osteoporos Int 2000; 11(2):102-119.
- (235) Maggio D, Ubaldi E, Simonelli G, Cenci S, Pedone C, Cherubini A. Hip fracture in nursing homes: An Italian study of prevalence, latency, risk factors, and impact on mobility. Calcif Tissue Int 2001; 68:337-341.
- (236) Visentin P, Ciravegna R, Uscello L, Molaschi M, Fabris F. Site-specific relative risk of fractures in the institutionalized elderly. Gerontology 1995; 41(5):273-279.
- (237) Diamond TH, Thornley SW, Sekel R, Smerdely P. Hip fracture in elderly men: prognostic factors and outcomes. Med J Aust 1997; 167(8):412-415.
- (238) Ensrud KE, Blackwell T, Mangione CM, Bowman PJ, Bauer DC, Schwartz A et al. Central nervous system active medications and risk for fractures in older women. Arch Intern Med 2003; 163(8):949-957.
- (239) Bernstein J, Grisso JA, Kaplan FS. Body mass and fracture risk. A study of 330 patients. Clin Orthop Rel Res 1999;(364):227-230.

- (240) Ensrud KE, Lipschutz RC, Cauley JA, Seeley D, Nevitt MC, Scott J et al. Body size and hip fracture risk in older women: a prospective study. Study of Osteoporotic Fractures Research Group. Am J Med 1997; 103(4):274-280.
- (241) Cummings SR, Nevitt MC, Browner WS, Stone K, Fox KM, Ensrud KE et al. Risk factors for hip fracture in white women. Study of Osteoporotic Fractures Research Group. N Engl J Med 1995; 332(12):767-773.
- (242) Ensrud KE, Cauley J, Lipschutz R, Cummings SR. Weight change and fractures in older women. Study of Osteoporotic Fractures Research Group. Arch Intern Med 1997; 157(8):857-863.
- (243) Cali CM, Kiel DP. An epidemiologic study of fall-related fractures among institutionalized older people. J Am Geriatr Soc 1995; 43(12):1336-1340.
- (244) Hui SL, Slemenda CW, Johnston CCJr. Age and bone mass as predictors of fracture in a prospective study. J Clin Invest 1987; 81:1804-1809.
- (245) Nguyen T, Sambrook P, Kelly P, Jones G, Lord SL, Freund J et al. Prediction of osteoporotic fractures by postural instability and bone density. BMJ 1993; 307:1111-1115.
- (246) Bauer DC, Gluer CC, Genant HK, Stone K. Quantitative ultrasound and vertebral fracture in postmenopausal women. Fracture Intervention Trial Research Group. J Bone Miner Res 1995; 10(3):353-358.
- (247) Papaioannou A, Adachi JD, Parkinson W, Stephenson G, Bedard M. Lengthy hospitalization associated with vertevral fractures despite control for comorbid conditions. Osteoporos Int 2001; 12:870-874.
- (248) Colon-Emeric CS, Sloane R, Hawkes WG, Magaziner J, Zimmerman SI, Pieper CF et al. The risk of subsequent fractures in community-dwelling men and male veterans with hip fracture. Am J Med 2000; 109(4):324-326.
- (249) Hall SE, Williams JA, Senior JA, Goldswain PRT, Criddle RA. Hip fracture outcomes: quality of life and functional status in older adults living in the community. A N Z J Med 2000; 30:327-332.
- (250) Salkeld G, Cameron ID, Cumming RG, Easter S, Seymour J, Kurrle SE et al. Quality of life related to fear of falling and hip fracture in older women: a time trade off study. BMJ 2000; 320(7231):341-346.
- (251) Scaf-Klomp W, van Sonderen E, Sanderman R, Ormel J, Kempen GIJM. Recovery of physical function after limb injuries in independent older people living at home. Age & Ageing 2001; 30:213-219.
- (252) Ismail AA, O'Neill TW, Cooper C, Finn JD, Bhalla AK, Cannata JB et al. Mortality associated with vertebral deformity in men and women: results from the European Prospective Osteoporosis Study (EPOS). Osteoporos Int 1998; 8(3):291-297.
- (253) Hasserius R, Karlsson MK, Nilsson BE, Redlund-Johnell I, Johnell O. Prevalent vertebral deformities predict increased mortality and increased fracture rate in both men and women: A 10-year population-based study of 598 individuals from the Swedish cohort in the European Vertebral Osteoporosis Study. Osteoporos Int 2002; 14:61-68.

- (254) Ensrud KE, Thompson DE, Cauley JA, Nevitt MC, Kado DM, Hochberg MC et al. Prevalent vertebral deformities predict mortality and hospitalization in older women with low bone mass. Fracture Intervention Trial Research Group. J Am Geriatr Soc 2000; 48(3):241-249.
- (255) Hip fracture outcomes in people age 50 and over background paper. OTA-BP-H-120. 1994. U.S.Congress, Office of Technology Assessment, U.S.Government Printing Office.
- (256) Wiktorowicz ME, Goeree R, Papaioannou A, Adachi JD, Papadimitropoulos E. Economic implications of hip fracture: health service use, institutional care and cost in Canada. Osteoporos Int 2001; 12(4):271-278.
- (257) Gelbach SH, Burge RT, Puleo E, Klar J. Hospital care of osteoporosis-related vertebral fractures. Osteoporos Int 2003; 14:53-60.
- (258) Dorman PJ, Waddell F, Slattery J, Dennis M, Sandercock P. Are proxy assessments of health status after stroke with the EuroQol questionnaire feasible, accurate, and unbiased? Stroke 1997; 28(10):1883-1887.
- (259) Lauritzen JB. Hip fractures. Epidemiology, risk factors, falls, energy absorption, hip protectors, and prevention. Dan Med Bull 1997; 44(2):155-168.
- (260) Chapuy MC, Arlot ME, Duboeuf F, Brun J, Crouzet B, Arnaud S et al. Vitamin D3 and calcium to prevent hip fractures in elderly women. N Engl J Med 1992; 327:1637-1642.
- (261) Granger CV, Albrecht GL, Hamilton BB. Outcome of comprehensive medical rehabilitation: measurement by PULSES profile and the Barthel Index. Arch Phys Med Rehab 1979; 60(4):145-154.
- (262) Residential Care Manual. Commonwealth of Australia, 1998.
- (263) Jadelis K, Miller ME, Ettinger WHJ, Messier SP. Strength, balance, and the modifying effects of obesity and knee pain: Results from the Observational Arthritis Study in Seniors (OASIS). J Am Geriatr Soc 2001; 49(7):884-891.
- (264) Birmingham TB, Kramer JF, Kirkley A, Inglis JT, Spaulding SJ, Vandervoort AA. Association among neuromuscular and anatomic measures for patients with knee osteoarthritis. Arch Phys Med Rehab 2001; 82:1115-1118.
- (265) Molloy DW, Alemayehu E, Roberts R. Reliability of a Standardized Mini-Mental State Examination compared with the traditional Mini-Mental State Examination. Am J Psychiatr 1991; 148(1):102-105.
- (266) Fillenbaum GG, George LK, Blazer DG. Scoring nonresponse on the Mini-Mental State Examination. Psychol Med 1988; 18(4):1021-1025.
- (267) Forsell Y, Fratiglioni L, Grut M, Viitanen M, Winblad B. Clinical staging of dementia in a population survey: comparison of DSM-III-R and the Washington University Clinical Dementia Rating Scale. Acta Psychiat Scand 1992; 86(1):49-54.
- (268) Rabin R, de Charro F. EQ-5D: a measure of health status from the EuroQol Group. Ann Med 2001; 33(5):337-343.
- (269) EuroQol--a new facility for the measurement of health-related quality of life. The EuroQol Group. Health Policy 1990; 16(3):199-208.

- (270) Johnson JA, Ohinmaa A, Murti B, Sintonen H, Coons SJ. Comparison of Finnish and U.S.-based visual analog scale valuations of the EQ-5D measure. Medical Decision Making 2000; 20(3):281-289.
- (271) Johnson JA, Pickard AS. Comparison of the EQ-5D and SF-12 health surveys in a general population survey in Alberta, Canada. Medical Care 2000; 38(1):115-121.
- (272) Johnson JA, Coons SJ, Ergo A, Szava-Kovats G. Valuation of EuroQOL (EQ-5D) health states in an adult US sample. Pharmacoeconomics 1998; 13(4):421-433.
- (273) Johnson JA, Coons SJ. Comparison of the EQ-5D and SF-12 in an adult US sample. Quality of Life Research 1998; 7(2):155-166.
- (274) Fransen M, Edmonds J. Reliability and validity of the EuroQol in patients with osteoarthritis of the knee. Rheumatology 1999; 38(9):807-813.
- (275) Hurst NP, Kind P, Ruta D, Hunter M, Stubbings A. Measuring health-related quality of life in rheumatoid arthritis: validity, responsiveness and reliability of EuroQol (EQ-5D). Br J Rheumatol 1997; 36(5):551-559.
- (276) Schrag A, Selai C, Jahanshahi M, Quinn NP. The EQ-5D--a generic quality of life measure-is a useful instrument to measure quality of life in patients with Parkinson's disease. J Neurol Neurosurg Psychiatr 2000; 69(1):67-73.
- (277) Siderowf AD, Werner RM. The EQ-5D--a generic quality of life measure--is a useful instrument to measure quality of life in patients with Parkinson's disease. J Neurol Neurosurg Psychiatr 2001; 70(6):817.
- (278) Oleksik A, Lips P, Dawson A, Minshall ME, Shen W, Cooper C et al. Healthrelated quality of life in postmenopausal women with low BMD with or without prevalent vertebral fractures. J Bone Miner Res 2000; 15(7):1384-1392.
- (279) Coons SJ, Rao S, Keininger DL, Hays RD. A comparative review of generic quality-of-life instruments. Pharmacoeconomics 2000; 17(1):13-35.
- (280) Coucill W, Bryan S, Bentham P, Buckley A, Laight A. EQ-5D in patients with dementia: an investigation of inter-rater agreement. Medical Care 2001; 39(8):760-771.
- (281) Quine S, Helby L, Cameron I, Lyle D. Carer burden after proximal femoral fracture. Disabil Rehab 1994; 16(4):191-197.
- (282) Holtzman J, Lurie N. Causes of increasing mortality in a nursing home population. J Am Geriatr Soc 1996; 44:258-264.
- (283) Bush TL, Miller SR, Golder AL, Hale WE. Self-report and medical record report agreement of selected medical conditions in the elderly. Am J Pub Health 1989; 79:1554-1556.
- (284) Colditz GA, Martin P, Stampfer MJ, Willett WC, Sampson L, Rosner B et al. Validation of questionnaire information on risk factors and disease outcomes in a prospective cohort study of women. Am J Epidemiol 1986; 123(5):894-900.
- (285) Harlow SD, Linet MS. Agreement between questionnaire data and medical records. Am J Epidemiol 1989; 129(2):233-248.

- (286) Paganini-Hill A, Chao A. Accuracy of recall of hip fracture, heart attack, and cancer: A comparison of postal survey data and medical records. Am J Epidemiol 1993; 138(2):101-106.
- (287) Jonsson B, Gardsell P, Johnell O, Redlund-Johnell I, Sernbo I. Remembering fractures: fracture registration and proband recall in southern Sweden. J Epidemiol Comm Health 1994; 48(5):489-490.
- (288) Nevitt MC, Cummings SR, Browner WS, Seeley DG, Cauley JA, Vogt TM et al. The accuracy of self-report of fractures in elderly women: Evidence from a prospective study. Am J Epidemiol 1992; 125:490-499.
- (289) Han TS, Lean MEJ. Lower leg length as an index of stature in adults. Int J Obesity 1996; 20:21-27.
- (290) Pasco JA, Henry MJ, Gaudry TM, Nicholson GC, Kotowicz MA. Identification of incident fractures: the Geelong Osteoporosis Study. A N Z J Med 1999; 29(2):203-206.
- (291) Borgquist L, Nilsson LT, Lindelow G, Wiklund I, Thorngren KG. Perceived health in hip-fracture patients: a prospective follow-up of 100 patients. Age & Ageing 1992; 21(2):109-116.
- (292) Bueno-Cavanillas A, Padilla-Ruiz F, Jimenez-Moleon JJ, Peinado-Alonso CA, Galvez-Vargas R. Risk factors in falls among the elderly according to extrinsic and intrinsic precipitating causes. Eur J Epidemiol 2000; 16(9):849-859.
- (293) Bueno-Cavanillas A, Padilla RF, Peinado AC, Espigares GM, Galvez VR. Risk factors associated with falls in institutionalized elderly population. A prospective cohort study. Medicina Clinica 1999; 112(1):10-15.
- (294) Nurmi I, Sihvonen M, Kataja M, Luthje P. Falls among institutionalized elderly--a prospective study in four institutions in Finland. Scandinavian Journal of Caring Sciences 1996; 10(4):212-220.
- (295) Visentin P, Ciravegna R, Uscello L, Corcelli F, Fabris F. Longitudinal study of fractures in institutionalized elderly. Minerva Medica 1996; 87(5):195-200.
- (296) van Schoor NM, Smit JH, Pluijm SM, Jonker C, Lips P. Different cognitive functions in relation to falls among older persons. Immediate memory as an independent risk factor for falls. J Clin Epidemiol 2002; 55(9):855-862.
- (297) Iki M, Kajita E, Mitamura S, Nishino H, Yamagami T, Nagahama N. Precision of quantitative ultrasound measurement of the heel bone and effects of ambient temperature on the parameters. Osteoporos Int 1999; 10(6):462-467.
- (298) Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. Lancet 1986; 1(8476):307-310.
- (299) Lin LI. A concordance correlation coefficient to evaluate reproducibility. Biometrics 1989; 45(1):255-268.
- (300) Nguyen TV, Eisman JA. Assessment of significant change in BMD: a new approach. J Bone Miner Res 2000; 15(2):369-372.

- (301) Blake GM, Wahner HW, Fogelman I. Commercial ultrasound instruments. The evaluation of osteoporosis: dual energy x-ray absorptiometry and ultrasound in clinical practice. 2 ed. Martin Dunitz Ltd, 1999.
- (302) Thuy VT, Chau TT, Cong ND, De DV, Nguyen TV. Assessment of low bone mass in Vietnamese: comparison of QUS calcaneal ultrasonometer and data-derived Tscores. J Bone Miner Metab 2003; 21(2):114-119.
- (303) Warden SJ, Bennell KL, Matthews B, Brown DJ, McMeeken JM, Wark JD et al. Quantitative ultrasound assessment of acute bone loss following spinal cord injury: a longitudinal pilot study. Osteoporos Int 2002; 13(7):586-592.
- (304) Laugier P, Novikov V, Elmann-Larsen B, Berger G. Quantitative ultrasound imaging of the calcaneus: precision and variations during a 120-day bed rest. Calcif Tissue Int 2000; 66:16-21.
- (305) Falcini F, Bindi G, Simonini G, Stagi S, Galluzzi F, Masi L et al. Bone status evaluation with calcaneal ultrasound in children with chronic rheumatic diseases. A one year followup study. J Rheumatol 2003; 30:179-84.
- (306) Daly RM, Rich PA, Klein R, Bass S. Effects of high-impact exercise on ultrasonic and biochemical indices of skeletal status: A prospective study in young male gymnasts. J Bone Miner Res 1999; 14(7):1222-1230.
- (307) Norton R, Campbell AJ, Lee-Joe T, Robinson E, Butler M. Circumstances of falls resulting in hip fractures among older people. J Am Geriatr Soc 1997; 45(9):1108-1112.
- (308) March LM, Cameron ID, Cumming RG, Chamberlain AC, Schwarz JM, Brnabic AJ et al. Mortality and morbidity after hip fracture: can evidence based clinical pathways make a difference? J Rheumatol 2000; 27(9):2227-2231.
- (309) Armitage P. Statistical Methods in Medical Research. Blackwell, 1971.
- (310) Jones G, Nguyen T, Sambrook P, Kelly PJ, Eisman JA. Progressive loss of bone in the femoral neck in elderly people: longitudinal findings from the Dubbo osteoporosis epidemiology study. BMJ 1994; 309(6956):691-695.
- (311) Grover SR, Morle R. Vitamin D deficiency in veiled or dark-skinned pregnant women. Med J Aust 2001; 175:251-252.
- (312) Skull SA, Ngeow JY, Biggs BA, Street A, Ebeling PR. Vitamin D deficiency is common and unrecognized among recently arrived adult immigrants from The Horn of Africa. Int Med J 2003; 33(1-2):47-51.
- (313) Nowson CA, Margerison C. Vitamin D intake and vitamin D status of Australians. Med J Aust 2002; 177(3):149-152.
- (314) Chapuy MC, Preziosi P, Maamer M, Arnaud S, Galan P, Hercberg S et al. Prevalence of vitamin D insufficiency in an adult normal population. Osteoporos Int 1997; 7(5):439-443.
- (315) Kudlacek S, Schneider B, Peterlik M, Leb G, Klaushofer K, Weber K et al. Assessment of vitamin D and calcium status in healthy adult Austrians. Eur J Clin Invest 2003; 33(4):323-331.

- (316) Harris SS, Soteriades E, Coolidge JA, Mudgal S, Dawson-Hughes B. Vitamin D insufficiency and hyperparathyroidism in a low income, multiracial, elderly population. J Clin Endocrinol Metab 2000; 85(11):4125-4130.
- (317) Goswami R, Gupta N, Goswami D, Marwaha RK, Tandon N, Kochupillai N. Prevalence and significance of low 25-hydroxyvitamin D concentrations in healthy subjects in Delhi. Am J Clin Nutrit 2000; 72:472-475.
- (318) Thomas MK, Lloyd-Jones DM, Thadhani RI, Shaw AC, Deraska DJ, Kitch BT et al. Hypovitaminosis D in medical inpatients. N Engl J Med 1998; 338(12):777-783.
- (319) Kauppinen-Makelin R, Tahtela R, Loyttyniemi E, Karkkainen J, Valimaki MJ. A high prevalence of hypovitaminosis D in Finnish medical in- and outpatients. J Intern Med 2001; 249(6):559-563.
- (320) Gloth FM, Gundberg CM, Hollis BW, Haddad JG, Tobin JD. Vitamin D deficiency in homebound elderly persons. JAMA 1995; 274(21):1683-1686.
- (321) Inderjeeth CA, Nicklason F, Al Lahham Y, Greenaway TM, Jones G, Parameswaran VV et al. Vitamin D deficiency and secondary hyperparathyroidism: clinical and biochemical associations in older non-institutionalised Southern Tasmanians. A N Z J Med 2000; 30(2):209-214.
- (322) Gallagher JC, Kinyamu HK, Fowler SE, Dawson-Hughes B, Dalsky GP, Sherman SS. Calciotropic hormones and bone markers in the elderly. J Bone Miner Res 1998; 13(3):475-482.
- (323) Corless D, Gupta SP, Sattar DA, Switala S, Boucher BJ. Vitamin D status of residents of an old people's home and long-stay patients. Gerontology 1979; 25:350-355.
- (324) Stein MS, Scherer SC, Walton SL, Gilbert RE, Ebeling PR, Flicker L et al. Risk factors for secondary hyperparathyroidism in a nursing home population. Clin Endocrinol 1996; 44(4):375-383.
- (325) Vir SC, Love HG. Nutritional status of institutionalized and noninstitutionalized aged in Belfast, Northern Ireland. Am J Clin Nutrit 1979;1934-1937.
- (326) McMurtry CT, Young SE, Downs RW, Adler RA. Mild vitamin D deficiency and secondary hyperparathyroidism in nursing home patients receiving adequate dietary vitamin D. J Am Geriatr Soc 1992; 40(4):343-347.
- (327) Webb AR, Pilbeam C, Hanafin N, Holick MF. An evaluation of the relative contributions of exposure to sunlight and of diet to the circulating concentrations of 25-hydroxyvitamin D in an elderly nursing home population in Boston. Am J Clin Nutrit 1990; 51(6):1075-1081.
- (328) Glendenning P, Vasikaran SD. Vitamin D status and redefining serum PTH reference range in the elderly. J Clin Endocrinol Metab 2002; 87(2):946-947.
- (329) Souberbielle J-C, Cormier C, Kindermans C, Gao P, Cantor T, Forette F et al. Vitamin D status and redefining serum parathyroid hormone reference range in the elderly. J Clin Endocrinol Metab 2001; 86(7):3086-3090.
- (330) Holick MF. McCollum Award Lecture, 1994: vitamin D--new horizons for the 21st century. Am J Clin Nutrit 1994; 60(4):619-630.

- (331) Jacques PF, Felson DT, Tucker KL, Mahnken B, Wilson PWF. Plasma 25hydroxyvitamin D and its determinants in an elderly population sample. Am J Clin Nutrit 1997; 66:929-936.
- (332) Morley JE, Glick Z, Rubenstein LZ. Geriatric nutrition: a comprehensive review. New York: Raven Press, 1995.
- (333) Selberg O, Sel S. The adjunctive value of routine biochemistry in nutritional assessment of hospitalized patients. Clin Nutrit 2001; 20(6):477-485.
- (334) Seiler WO. Clinical pictures of malnutrition in ill elderly subjects. Nutrition 2001; 17(6):496-498.
- (335) Melin A, Wilske J, Ringertz H, Saaf M. Seasonal variations in serum levels of 25hydroxyvitamin D and parathyroid hormone but no detectable change in femoral neck bone density in an older population with regular outdoor exposure. J Am Geriatr Soc 2001; 49(9):1190-1196.
- (336) Lamberg-Allardt C. Vitamin D intake, sunlight exposure and 25-hydroxyvitamin D levels in the elderly during one year. Ann Nutr Metab 1984; 28:144-150.
- (337) Rapuri PB, Kinyamu HK, Gallagher JC, Haynatzka V. Seasonal changes in calciotropic hormones, bone markers, and bone mineral density in elderly women. J Clin Endocrinol Metab 2002; 87(5):2024-2032.
- (338) Chapuy MC, Schott AM, Garnero P, Hans D, Delmas PD, Meunier PJ. Healthy elderly French women living at home have secondary hyperparathyroidism and high bone turnover in winter. EPIDOS Study Group. J Clin Endocrinol Metab 1996; 81(3):1129-1133.
- (339) van der Wielen RP, Lowik MR, van den BH, de Groot LC, Haller J, Moreiras O et al. Serum vitamin D concentrations among elderly people in Europe. Lancet 1995; 346(8969):207-210.
- (340) Dawson-Hughes B, Harris SS, Dallal GE. Plasma calcidiol, season, and serum parathyroid hormone concentrations in healthy elderly men and women. Am J Clin Nutrit 1997; 65:67-71.
- (341) Perry HM, Horowitz M, Morley JE, Patrick P, Vellas B, Baumgartner R et al. Longitudinal changes in serum 23-hydroxyvitamin D in older people. Metabolism 1999; 48(8):1028-1032.
- (342) Preece M, Tomlinson S, Ribot CA, Pietrek J, Korn HT, Davies DM. Studies on human vitamin D deficiency in man. QJM 1975; 44:5575-5589.
- (343) Lips P. Problems with direct 25-hydroxyvitamin D assays, and the target amount of vitamin D nutrition desirable for patients with osteoporosis (letter). Osteoporos Int 2000; 11:636.
- (344) Glendenning P, Vasikaran SD. Vitamin D status and redefining serum PTH reference range in the elderly (letter). J Clin Endocrinol Metab 2002; 87(2):946-947.
- (345) Vieth R. Problems with direct 25-Hydroxyvitamin D assays, and the target amount of vitamin D nutrition desirable for patients with osteoporosis. Osteoporos Int 2000; 11:635-636.

- 11 1 1 1²⁵1 1 DIA d 1 C
- (346) Hollis BW. Comparison of commercially available ¹²⁵I-based RIA methods for the determination of circulating 25-hydroxyvitamin D. Clinical Chemistry 2000; 46(10):1657-1661.
- (347) Sahota O, Masud T, San P, Hosking DJ. Vitamin D insufficiency increases bone turnover markers and enhances bone loss at the hip in patients with established vertebral osteoporosis. Clin Endocrinol 1999; 51(2):217-221.
- (348) Brot C, Jorgensen N, Madsen OR, Jensen LB, Sorensen OH. Relationships between bone mineral density, serum vitamin D metabolites and calcium:phosphorus intake in healthy perimenopausal women. J Intern Med 1999; 245:509-516.
- (349) Vieth R, Ladak Y, Walfish PG. Age-related changes in the 25-hydroxyvitamin D versus parathyroid hormone relationship suggest a different reason why older adults require more vitamin D. J Clin Endocrinol Metab 2003; 88(1):185-191.
- (350) Dawson-Hughes B, Harris SS, Krall EA, Dallal GE. Effect of calcium and vitamin D supplementation on bone density in men and women 65 years of age or older. New England Journal of Medicine 1997; 337(10):670-676.
- (351) Ooms ME, Roos JC, Bezemer PD, van der Vijgh WJ, Bouter LM, Lips P. Prevention of bone loss by vitamin D supplementation in elderly women: a randomized double-blind trial. J Clin Endocrinol Metab 1995; 80(4):1052-1058.
- (352) Baeksgaard L, Andersen KP, Hyldstrup L. Calcium and vitamin D supplementation increases spinal BMD in healthy, postmenopausal women. Osteoporos Int 1998; 8:255-260.
- (353) Holick MF, Matsuoka LY, Wortsman J. Age, vitamin D, and solar ultraviolet. Lancet 1989; 2(8671):1104-1105.
- (354) Slovik DM, Adams JS, Neer RM, Holick MF, Potts JT, Jr. Deficient production of 1,25-dihydroxyvitamin D in elderly osteoporotic patients. N Engl J Med 1981; 305(7):372-374.
- (355) Riggs BL, Hamstra A, DeLuca HF. Assessment of 25-hydroxyvitamin D 1 alphahydroxylase reserve in postmenopausal osteoporosis by administration of parathyroid extract. J Clin Endocrinol Metab 1981; 53(4):833-835.
- (356) Devine A, Wilson SG, Dick IM, Prince RL. Effects of vitamin D metabolites on intestinal calcium absorption and bone turnover in elderly women. Am J Clin Nutrit 2002; 75:283-288.
- (357) Lips P, Duong T, Oleksik A, Black D, Cummings S, Cox D et al. A global study of vitamin D status and parathyroid function in postmenopausal women with osteoporosis: baseline data from the multiple outcomes of raloxifene evaluation clinical trial. J Clin Endocrinol Metab 2001; 86(3):1212-1221.
- (358) Mezquita-Raya P, Munoz-Torres M, Luna JD, Luna V, Lopez-Rodriguez F, Torres-Vela E et al. Relation between vitamin D insufficiency, bone density, and bone metabolism in healthy postmenopausal women. J Bone Miner Res 2001; 16(8):1408-1415.
- (359) Hunter D, Major P, Arden N, Swaminathan R, Andrew T, MacGregor AJ et al. A randomized controlled trial of vitamin D supplementation on preventing

postmenopausal bone loss and modifying bone metabolism using identical twin pairs. J Bone Miner Res 2000; 15(11):2276-2283.

- (360) Peacock M, Liu G, Carey M, McClintock R, Ambrosius W, Hui S et al. Effect of calcium or 25OH Vitamin D3 dietary supplementation on bone loss at the hip in men and women over the age of 60. J Clin Endocrinol Metab 2000; 85:3011-3019.
- (361) Grados F, Brazier M, Kamel S, Duver S, Heurtebize N, Maamer M et al. Effects on bone mineral density of calcium and vitamin D supplementation in elderly women with vitamin D deficiency. Joint, Bone, Spine 2003; 70:203-208.
- (362) Chapuy MC, Arlot ME, Delmas PD, Meunier PJ. Effect of calcium and cholecalciferol treatment for three years on hip fractures in elderly women. BMJ 1994; 308(6936):1081-1082.
- (363) Lips P, Graafmans WC, Ooms ME, Bezemer PD, Bouter LM. Vitamin D supplementation and fracture incidence in elderly persons: a randomized, placebocontroled clinical trial. Ann Int Med 1996; 124(4):400-406.
- (364) Villareal DT, Civitelli R, Chines A, Avioli LV. Subclinical vitamin D deficiency in postmenopausal women with low vertebral bone mass. J Clin Endocrinol Metab 1991; 72(3):628-634.
- (365) Krieg MA, Cornuz J, Jacquet AF, Thiebaud D, Burckhardt P. Influence of anthropometric parameters and biochemical markers of bone metabolism on quantitative ultrasound of bone in the institutionalized elderly. Osteoporos Int 1998; 8(2):115-120.
- (366) Gregg EW, Kriska AM, Salamone LM, Wolf RL, Roberts MM, Ferrell RE et al. Correlates of quantitative ultrasound in the Women's Healthy Lifestyle Project. Osteoporos Int 1999; 10(5):416-424.
- (367) Ingle BM, Thomas WE, Eastell R. Differential effects of primary hyperparathyroidism on ultrasound properties of bone. Osteoporos Int 2002; 13(7):572-578.
- (368) Montagnani A, Gonnelli S, Cepollaro C, Bruni D, Franci MB, Lucani B et al. Graphic trace analysis of ultrasound at the phalanges may differentiate between subjects with primary hyperparathyroidism and with osteoporosis: a pilot study. Osteoporos Int 2002; 13(3):222-227.
- (369) Cortet B, Cortet C, Blanckaert F, Racadot A, d'Herbomez M, Marchandise X et al. Bone ultrasonometry and turnover markers in primary hyperparathyroidism. Calcif Tissue Int 2000; 66(1):11-15.
- (370) Talmage RV, Elliott JR. Removal of calcium from bone as influenced by the parathyroids. Endocrinology 1958; 62:717-722.
- (371) Mundy GR, Roodman GD. Osteoclast ontogeny and function. In: Meunier PW, editor. Bone and mineral research. Amsterdam: Elsevier, 1987: 209-280.
- (372) Blake AJ, Morgan K, Bendall MJ, Dallosso H, Ebrahim SB, Arie TH et al. Falls by elderly people at home: prevalence and associated factors. Age & Ageing 1988; 17(6):365-372.
- (373) Niino N, Nakamura K. Circumstances and factors related to falls in the institutionalized elderly. Japan J Geriatr 1996; 33(1):12-16.

- (374) Speechley M, Tinetti M. Falls and injuries in frail and vigorous community elderly persons. J Am Geriatr Soc 1991; 39(1):46-52.
- (375) Janssen HC, Samson MM, Verhaar HJ. Vitamin D deficiency, muscle function, and falls in elderly people. Am J Clin Nutrit 2002; 75(4):611-615.
- (376) Pfeifer M, Begerow B, Minne HW, Schlotthauer T, Pospeschill M, Scholz M et al. Vitamin D status, trunk muscle strength, body sway, falls, and fractures among 237 postmenopausal women with osteoporosis. Exp Clin Endocrinol Diabetes 2001; 109(2):87-92.
- (377) Mowe M, Haug E, Bohmer T. Low serum calcidiol concentration in older adults with reduced muscular function. J Am Geriatr Soc 1999; 47(2):220-226.
- (378) Verhaar HJ, Samson MM, Jansen PA, de Vreede PL, Manten JW, Duursma SA. Muscle strength, functional mobility and vitamin D in older women. Aging (Milano) 2000; 12(6):455-460.
- (379) Flicker L, Mead K, MacInnis RJ, Nowson C, Scherer S, Stein MS et al. Serum vitamin D and falls in older women in residential care in Australia. J Am Geriatr Soc 2003; 51:1533-1538.
- (380) Chou F-F, Lee C-H, Chen J-B. General weakness as an indication for parathyroid surgery in patients with secondary hyperparathyroidism. Arch Surg 1999; 134(10):1108-1111.
- (381) Lord SR, March LM, Cameron ID, Cumming RG, Schwarz J, Zochling J et al. Differing risk factors for falls in nursing home and intermediate-care residents who can and cannot stand unaided. J Am Geriatr Soc 2003; 51:1645-1650.
- (382) Weatherall M. A meta-analysis of 25 hydroxyvitamin D in older people with fracture of the proximal femur. N Z Med J 2000; 113(1108):137-140.
- (383) Heikinheimo RJ, Inkovaara JA, Harju EJ, Haavisto MV, Kaarela RH, Kataja JM et al. Annual injection of vitamin D and fractures of aged bones. Calcif Tissue Int 1992; 51(2):105-110.
- (384) Chapuy MC, Pamphile R, Paris E, Kempf C, Schlichting M, Arnaud S et al. Combined calcium and vitamin D3 supplementation in elderly women: confirmation of reversal of secondary hyperparathyroidism and hip fracture risk: the Decalyos II study. Osteoporos Int 2002; 13(3):257-264.
- (385) Russell JA. Osteomalacic myopathy. Muscle Nerve 1994; 17:578-580.
- (386) Mets T. Calcium, vitamin D and hip fractures. Incidence of falls may have decreased. BMJ 1994; 309:193.
- (387) Glennon DG, Inderjeeth CA, Petta A, Reberger C, Tapper J, Sondestrom J et al. Vitamin D and muscle strength in older women with previous fractures. Int Med J 2002; 33(Suppl.):A24.
- (388) Sorensen OH, Lund BI, Saltin B. Myopathy in bone loss of ageing: improvement by treatment with 1-alpha-hydroxycholecalciferol and calcium. Clin Sci 1979; 56:157-161.

- (389) Gloth FM3, Smith CE, Hollis BW, Tobin JD. functional improvement with vitamin D replenishment in a cohort of frail, vitamin D-deficient older people. J Am Geriatr Soc 1995; 43:1269-1271.
- (390) Grady D, Halloran B, Cummings S, Leveille S, Wells L, Black D et al. 1,25dihydroxyvitamin D3 and muscle strength in the elderly: a randomized controlled trial. J Clin Endocrinol Metab 1991; 73:1111-1117.
- (391) Boonen S, Lysens R, Verbeke G, Joosten E, Dejaeger E, Pelemans W et al. Relationship between age-associated endocrine deficiencies and muscle function in elderly women: a cross-sectional study. Age & Ageing 1998; 27:449-454.
- (392) Dukas L, Bischoff HA, Lindpaintner LS, Schacht E, Birkner-Binder D, Thalmann B et al. Alfacalcidol reduces the number of fallers and falls in a communitydwelling elderly population with a minimum calcium intake above 500mg daily. Results from a randomised double-blind, placebo-controlled trial. Calcif Tissue Int 2003; 72(4):371.
- (393) Sambrook PN, Cameron ID, Cumming RG, Lord SR, Schwarz JM, Trube A et al. Vitamin D deficiency is common in frail institutionalised older people in northern Sydney. Med J Aust 2002; 176(11):560.
- (394) Kanis JA, Johnell O, Oden A, De Laet C, Oglesby A, Jonsson B. Intervention thresholds for osteoporosis. Bone 2002; 31(1):26-31.

APPENDIX 1

PATIENT INFORMATION SHEET PATIENT INFORMED CONSENT FORM PROXY INFORMATION AND CONSENT FORM DATA COLLECTION SHEET