Integrated study of group B streptococcus and human ureaplasmas – the paradigm shifts

Fanrong Kong

A thesis submitted to the University of Sydney in fulfilment of the requirement for the degree of Doctor of Philosophy.

Department of Medicine

University of Sydney

Declaration

This thesis is less than 100, 000 words in length and contains no material that has been accepted for the award of any other degree or diploma in any other universities.

I declare that the author, with the following exception, conducted all the practical work incorporated into this thesis:

- All the New Zealand GBS strains were provided by Dr Diana Martin (ESR, New Zealand) and GBS conventional serotyping was mainly performed by Sonia Gowan (ESR, New Zealand). Fourteen selected isolates were serotyped by Abbie Weisner (PHLS, United Kingdom).
- Ureaplasma and mycoplasma reference strain culture was partially performed by Susannah Gordon (CIDM, Westmead Hospital).
- Gregory James (CIDM, Westmead Hospital) provided all the needed molecular equipment, reagents, softwares and some protocols.
- Zhenfang Ma assisted in some ureaplasma PCR and data analysis.
- Mark Wheeler (WMI, Westmead Hospital) performed sequencing.
- Professor Lyn Gilbert (supervisor) supervised the overall project. Assisted in research design, in the analysis and interpretation of data, and made a significant contribution to the project.

Preface

Someone told me "a PhD thesis is just like a **story**". From my viewpoint, the **story** formed the major part of my past six-year life, and they will significantly affect my future research career and even my whole life. I tried to make the story something that would be interesting and useful to be read. On the other hand, I tried to cut its length to save the possible readers precious time.

As a PhD student, besides to **master** or to learn the techniques and methodologies as a **Master** student or a **technician**, to think about the **philosophy** or to explain the findings (including that of the others besides myself) would be also or even more important for the future related research fields. You would find the thesis in some extent (in particular, in section 3) reflected the above considerations.

From the teacher side, the selection of **good** teachers would be very important for a **good** student. I was so lucky to be a student of Professors Lyn Gilbert and Tania Sorrell. I also want to thank so many "informal teachers", I may only know many of them from their excellent publications (I tried my best to include their contributions in my reference sections though I might still have lost many for cutting the volume reason), and these teachers are also highly appreciated. Molecular lab – I located myself there in my past six years, all the staffs there were so kind and the conditions there were so good – the experience there will be kept in my mind as a very memorable life in my personal history.

The two **tiny** microorganisms that I selected, as my study objectives – GBS and ureaplasmas – were also my very **great** teachers! They showed me that they also had some kind of "wisdom", which I tried to expose partial in the thesis (especially in section 3) but far from enough. No doubt, their "wisdom" deserves to be further studied – especially based on their invaluable genome resources and benefit from the new bioinformatic concepts, theory and research tools.

The thesis contains three sections. The first section of general introduction provided the background and some useful concepts of the next two sections. The second section mainly contained those so called "traditional" molecular microbiology study of the two microorganisms. It included mainly the genotyping studies, which included most of my previous publications (also see appendices). My past six year publications (in PDF format) had been used as appendices and were put at the end of the thesis. Their inner relationship was also given at the beginning of the appendices. So only selected parts (after modification and reorganization) that were believed to be important to explain the outline of the story were put into the body of the thesis (mainly in section 2). The third section of the thesis contained some so called "novel" genomic and bioinformatics based study of the two organisms. Because we are facing the dramatic paradigm shift in their post genomic era, the ideas contained in the study may not be absolutely correct; even for the correct ones, some of them may still look a little bit ugly. But at least I would like to use them to show the potential values for doing this kind of "novel" study.

It was not easy to be a PhD student, especially considering the era of knowledge explosion, and the so many paradigm shifts. In this case, I really feel that what I had done was far from enough. However, to get the PhD or to pass the **milestone** is my long-term dream (even from my early childhood). So could I dare to use this thesis to have a **try**?

If the thesis can be seen as a **very tiny** drop of but **useful** water comparing with the endless sea of the **truth**, I will be more than happy!

Acknowledgments

The work described in this thesis was completed at:

Molecular Laboratory, Reference Laboratory Section, Centre for Infectious Diseases and Microbiology (CIDM), Institute of Clinical Pathology and Medical Research (ICPMR), Westmead Hospital, Westmead, NSW 2145, Australia. All the staffs in Molecular Laboratory and Reference Laboratory Section are highly appreciated.

The project was partially supported by Australian Postgraduate Awards 2000 (APA 2000) and Westmead Millennium Institute (WMI) Initiating Grant.

This thesis couldn't become true without:

Professor Lyn Gilbert, a great and very knowledgeable person, my highly respected supervisor – I was so lucky to be her student. Her excellent supervision, patient editing, and many other supports made my past six years productive and enjoyable. Professor Tania Sorrell, a very kind person, my associate supervisor – she always prepared to give her kind help to me when I needed.

I thank Mark Wheeler for his precious help in sequencing and Glenys Connor for some GBS data collection. Dr Hui Wang, in the third year of my PhD, while I was training him, gave me a great deal of help, was much appreciated.

I would like to thank all my family members; their continuous encouragements and supports made me feel full of energy. Zhenfang Ma, my deeply beloved wife, my another half, the most valuable **gift** I could get from my God – she gave me the greatest and invaluable supports, any words can't express my thanks to her – I **love** you! At the final stage of my PhD thesis writing, my lovely son (Alan, Xiangyu Kong) joined with us – welcome you my little honey and hope you can make my future **story** more exciting!

Abstract

Group B streptococcus (GBS, S. agalactiae) and human ureaplasmas (U. parvum and *U. urealyticum*) are two clinically and phylogenetically related, potential perinatal pathogens. Their relationships between genotypes and pathogenesis of GBS and ureaplasma infection were still not well understood, one of the reason is that both of them are still short of a very practical genotyping system. In the study, to solve the above problem we developed genotyping systems for the organisms (the second section). For human ureaplasmas, based on four genes/gene clusters (rRNA gene clusters, the elongation factor Tu genes, urease gene complexes and multiple banded antigen genes), we designed many primer pairs suitable for developing species identification assays for the two newly established human ureaplasma species (U. parvum and U. urealyticum). Further, based on the heterogeneity of ureaplasma multiple banded antigen gene (which contains species- and serovarspecific regions), we developed genotyping methods for each ureaplasma species. For GBS, based on three sets of molecular markers (capsular polysaccharide synthesis gene clusters, surface protein antigen genes and mobile genetic elements), we developed a genotyping system. The primary evaluation of the genotyping systems showed that the genotyping systems were practical alternative assays for the conventional serotyping and they will be useful to further explore the relationships between genotypes and pathogenesis of GBS and ureaplasma infection. In the study, we introduced novel data and tools into GBS and ureaplasma studies especially from genomic- and bioinformatics-based molecular microbiology

(the third section). For two newly established human ureaplasma species, based on the *U. parvum* serovar-3 genome, and using the above four important genes/gene clusters, we exposed some interesting problems in the understanding of new ureaplasma taxonomy especially in the post genomic era. For GBS, we studied the two published full genomes and exposed some new problems or possible future new research fields. In particular we found the two finished and one ongoing GBS genomes were all non-typical and suggest that future genomic project had better have genetic population structure viewpoint. Finally, we suggested that integrated studies of the two potential or conditional perinatal pathogens, from the viewpoint of evolution, would provide a new understanding angle of the pathogenesis of the two organisms. Studies suggested that during coevolution, human ureaplasmas (especially *U. parvum*) became friendlier than their ancestors to their human host (by losing most of its virulence genes); however, GBS tried to increase its invasive abilities (by getting more virulence genes) to fight against the human host attack.

Table of Contents

TITLE PAGE	
DECLARATION	Il
PREFACE	II
ACKNOWLEDGMENTS	V
ABSTRACT	V
TABLE OF CONTENTS	VIII
LIST OF FIGURES	X
LIST OF TABLES.	Xl
ABBREVIATIONS	XIII
SECTION 1 GENERAL INTRODUCTION	1
A. BRIEF INTRODUCTION	1
B. WHY GBS AND UREAPLASMAS?	2
B.1. GBS AND UREAPLASMAS	2
B.2. CLINICAL ASPECTS	2
B.3. VIRULENCE AND PATHOGENESIS	9
B.4. THE ROLE OF ANTIBODY	12
B.5. PREVENTION EFFORTS.	14
C. WHY GENOTYPING?	15
C.1. SEVERAL CONCEPTS OR GLOSSARY	15
C.2. GENOTYPING OF GBS AND UREAPLASMAS	16
C.3. MOLECULAR EPIDEMIOLOGY OF GSB AND UREAPLASMAS	18
C.4. INTEGRATED STUDY OF INFECTION, POPULATION GENETICS AND EVOLUTION.	19
D. WHY GENOMICS AND COMPARATIVE GENOMICS?	21
D.1. SOME USEFUL CONCEPTS OR GLOSSARY	22
D.2. UREAPLASMA AND GBS GENOMIC STUDIES	26

E. THESIS ARRANGEMENT	31
F. HYPOTHESIS TO BE TESTED BY THE STUDY	31
SECTION 2 GENOTYPING SYSTEMS	33
UREAPLASMAS	33
CHAPTER 1 UREAPLASMA SPECIES IDENTIFICATION	33
CHAPTER 2 UREAPLASMA GENOTYPING	44
GBS	65
CHAPTER 3 GBS MOLECULAR SEROTYPE IDENTIFICATION	65
CHAPTER 4 GBS PROTEIN GENE PROFILES	91
CHAPTER 5 GBS MOBILE GENETIC ELEMENTS	113
SECTION 3 GENOMIC- AND BIOINFORMATICS-BASED STUDIES	134
CHAPTER 6 POSTGENOMIC TAXONOMY OF HUMAN UREAPLASMAS	134
CHAPTER 7 IN SILICO GBS COMPARATIVE GENOMICS	152
CHAPTER 8 GBS AND UREAPLASMA EVOLUTIONARY "WISDOM"	179
REFERENCES	202
ADDENDICES	225

List of Figures

Figure 2.1. Multiple sequence alignment of the 5'-end and upstream of <i>mba</i>
Figure 3.1. Molecular serotype identification based on the sequence heterogeneity of the 790-br fragment at the 3'-end of <i>cpsE-cpsF</i> -and the 5'-end of <i>cpsG</i>
Figure 5.1. Genetic relationship of 194 invasive Australasian GBS strains (or 56 genotypes)
Figure 8.1. The branching order of the main groups within <i>Bacteria</i> based on conserved indelegresent in various proteins
Figure 8.2. Supertrees of 45 species constructed with 730 trees

List of Tables

Table 1.1. Primers targeting ureaplasma 3 different genes/regions
Table 2.1. Primers for ureaplasma genotyping
Table 2.2. Specificity and expected lengths of ureaplasma genotyping primer pairs60
Table 3.1. Reference strains used in GBS molecular serotyping study
Table 3.2. Primers used in GBS molecular serotyping study
Table 3.3. The heterogeneity of 8 GBS serotypes in the regions of the 3'-end of <i>cpsD</i> and the 5'-end of <i>cpsE</i>
Table 3.4. Comparison of the results of conventional serotyping (CS) and molecular serotype identification (MS)/subtyping of 206 clinical GBS isolates
Table 3.5. Algorithm for GBS molecular serotype (MS) identification by PCR and sequencing87
Table 4.1. Primers used in GBS protein gene profiling study96
Table 4.2. Specificity and expected lengths of amplicons of GBS protein gene profiling primer pairs
Table 4.3. Genetic groups and subgroups of GBS bac based on amplicon length and sequence heterogeneity
Table 4.4. The relationship between GBS protein gene profiles and capsular polysaccharide synthesis (<i>cps</i>) gene molecular sero/subtypes
Table 5.1. Primers used in GBS mobile genetic element study
Table 5.2. Distribution of mobile genetic elements among 194 invasive GBS isolates123
Table 5.3. Relationship between genotypes of invasive GBS isolates and patients' age-groups
Table 6.1. Primers used for sequencing 3 ureaplasma different genes/gene clusters
Table 6.2. Comparison of inter-species, intra-species and inter-copy heterogeneity in rRNA gene complexes of <i>U. parvum</i> and <i>U. urealyticum</i>
Table 6.3. Comparison of interspecies heterogeneity of DNA and amino acid sequences of the urease gene clusters of <i>U. parvum</i> and <i>U. urealyticum</i>
Table 7.1. General features of NEM316 and 2603 V/R genomes
Table 7.2. SmaI restriction map of NEM316 and 2603 V/R genomes
Table 7.3. NEM316 and 2603 V/R selected "virulence"-related proteins
Table 7.4. The cell wall protein heterogeneity of NEM316 and 2603 V/R

Table 7.5. Possible genomic islands in NEM316 and 2603 V/R genomes	175
Table 8.1. General features of GBS and ureaplasma genomes	183
Table 8.2. Some similar charizations of GBS, ureaplasma related organisms that support the confederative evolution from <i>Streptococcus</i> to <i>Mollicutes</i>	
Table 8.3. Sequence comparison of human ureaplasma and GBS major surface protein antigens	

Abbreviations

ABI Applied Biosystems
Alp2 C alpha-like 2 antigen

alp2 C alpha-like 2 antigen gene

Alp3 C alpha-like 3 antigen

alp3 C alpha-like 3 antigen gene

Alp4 C alpha-like 4 antigen

alp4 C alpha-like 4 antigen gene

Alp5 C alpha-like 4 antigen

alp5 C alpha-like 4 antigen geneAmiC Amidase family protein

ANGIS Australian National Genomic Information Service

ATCC American Type Culture Collection (Manassas, VA, USA)

Bac C beta antigen or IgA binding protein

bac C beta antigen or IgA binding protein gene

Bca C alpha antigen

bca C alpha antigen gene

BPS group B protective surface protein

bp base pair

BspA A cell surface associated leucine-rich repeat protein involved in adhesion to

fibronectin and fibrinogen

CAMP (discovered by) Christie, Atkins, and Munch-Petersen

CDC Centers for Disease Control and Prevention

CDS Coding sequences

CGH Comparative genome hybridization

CI Confidence interval

CLD Chronic lung disease (of prematurity)

Clp Clp ATPase family of molecular chaperones

CNS Central nervus system
CbpD Choline binding protein D

CpdB Cyclo-nucleotide phosphodiesterase

CPS Capsular polysaccharide

cps Capsular polysaccharide synthesis (gene cluster)

CS Conventional serotyping/serotype

CSF Cerebrospinal fluid

DNA Deoxyribonucleic acid

dNTP Deoxynucleotide triphosphate

EaeH EaeH of *Escherichia coli* O157:H7

EF-TU Elongation factor Tu

EOD Early onset disease (infection occurring within the first week of life)

FbsA A fibrinogen receptor from group B streptococcus

G+C Guanine plus Cytosine

GBS Group B streptococcus or group B streptococci or S. agalactiae

g Gram

GIs genomic islands

Human ureaplasmas – *U. parvum* and *U. urealyticum*

h Hour

Hsa (antigen that recognition of) sialic acid-containing host receptors

IAP Intrapartum antibiotic prophylaxis

Indels Insertations and deletions
IS Insertion sequence(s)
Lmb Laminin-binding protein

lmb Laminin-binding protein gene

LOD Late onset disease (infection occurring between 1 week and 2 to 3 months of age)

LSU rRNA Large submit rRNA

l Liter/LitreM Molar

MBA Multiple banded antigen

mba Multiple banded antigen genemge Mobile genetic element(s)

min Minute

ml Microlitre/Microliter

MLEE Mutiple locus enzyme electrophoresis

MLST Multiple locus sequence typing

MS Molecular serotype(s)

NanA Sialic acid lyase (catalyzes the hydrolysis of sialic acid into pyruvate and N-

acetylmannosamine)

NCTC National Culture Type Collection (Colindale, UK)

No. Number(s)

NGU Non-gonococcal urethritis

nra Encode a response regulator (no response to atmospheric conditions)

OR Odds ratio

ORF Open reading frame(s)

Pas The surface protein antigen I/II of Streptococcus intermedius

PCR Polymerase chain reaction

PFBP Streptococcus pyogenes fibronectin-binding protein

PFGE Pulsed-field gel electrophoresis

pgp Protein gene profile(s)PIs Pathogenicity islands

PrtS Serine proteinase, subtilase family
PspC Pneumococcal surface protein C

PulA Alkaline amylopullulanase

R28 Streptococcus pyogenes surface protein R28
R5 (or BPS protein) – group B protective surface protein

RAPD Random amplified polymorphic DNA

RFLP Restriction length fragment polymorphisms

rgg Encode a response regulator

Rib Rib antigen

rib Rib antigen geneRNA Ribonucleic acid

rof Encode a response regulator

s Second

ScpB Serine protease and C5a peptidase

Sec10 Surface exclusion protein

SpaA Streptococcal protein antigen A of *Streptococcus sobrinus*

Sip Surface immunogenic protein
SrpA Periplasmic linker protein

Srt Sortase

SSH Suppressive subtractive hybridization

Ssp5 Agglutinin receptor sst Serosubtype(s)

SSU rRNA Small subunit rRNA

Ta Annealing temperature
Tm Melting temperature

tuf Elongation factor Tu gene

U Unit

UAB University of Alabama at Birmingham

UP Ureaplasma parvum

 ${\it Ureaplasmas} - {\it U. parvum}$ and ${\it U. urealyticum}$

UU Ureaplasma urealyticum

Ureaplasma broth – 10B broth (Shepard, 1970)

WebANGIS - Website of Australian National Genomic Information Service

 $(\underline{http://www1.angis.org.au/pbin/WebANGIS/wrapper.pl})$

YbgE Putative branched-chain aminotransferase
YfkN 2`,3`-cyclic-nucleotide 2`-phosphodiesterase