

This is an Accepted Manuscript of an article published in [*Epidemiology and Infection*], available online: <http://dx.doi.org/10.1017/S0950268810001135> (paywalled)

**Please cite as:**

Tenzin, Dhand, N. K., Dorjee, J., & Ward, M. P. (2011). Re-emergence of rabies in dogs and other domestic animals in eastern Bhutan, 2005-2007. *Epidemiol Infect*, 139(2), 220-225. Available online at: <http://dx.doi.org/10.1017/S0950268810001135> (paywalled)

## Re-emergence of rabies in dogs and other domestic animals in eastern Bhutan, 2005–2007

Tenzin<sup>1,2</sup>, Navneet K Dhand<sup>1</sup>, Jambay Dorjee<sup>3</sup> and Michael P Ward<sup>1</sup>

<sup>1</sup>Faculty of Veterinary Science, The University of Sydney, Camden, NSW 2570, Australia;

<sup>2</sup>Regional Livestock Development Centre, Department of Livestock, Gelephu, Bhutan;

<sup>3</sup>Renewable Natural Resource Research Centre, Bumthang, Bhutan

*Epidemiology and Infection* (2010); 139: 220 – 225.

Available online at: <http://dx.doi.org/10.1017/S0950268810001135> (paywalled)

### Summary

We report a major outbreak of rabies in dogs and other domestic animals that occurred in eastern Bhutan between May 2005 and November 2007. The outbreak peaked in February 2006 and subsided by the end of April 2006 with sporadic cases reported until November 2007. Rabies affected 18 of the 40 sub-districts in the three eastern districts of Bhutan. There were reportedly one human and 256 domestic animal fatalities. The outbreak affected cattle (n=141, 55%), dogs (n=106, 41%), horses (n=7, 3%) and cats (n=2, 1%). Rabies was primarily diagnosed by clinical signs but 36 cases were confirmed by fluorescent antibody test of brain samples. High densities and movements of free-roaming dogs might have been responsible for the rapid spread and persistence of the infection for a longer period than expected in dogs in eastern Bhutan.

**Key words:** Eastern Bhutan, epidemic, rabies outbreak.

Rabies is an acute viral disease transmitted by the bite of a rabid animal. Each year, about 55,000 people die of rabies in Asia and Africa, mainly because of endemic canine rabies (Knobel et al., 2005). In Asia, a large number of human rabies deaths occur in India, and a national multi-centre epidemiological survey has estimated an annual burden of about 20 000 human deaths due to rabies in India (Sudarshan et al., 2007). Rabies has also been described as a re-emerging problem in China, with about 2000 human deaths reported annually (Sudarshan et al., 2007; Si et al., 2008; Wu et al., 2009). Bhutan is a small Himalayan kingdom situated in South Asia in between India and China. Until the early 1990s, rabies in dogs and other domestic animals was reported from most parts of Bhutan. A national rabies control

programme—vaccination of dogs (both free-roaming and pet dogs), implemented in 1992—controlled or eliminated canine rabies from the interior part of the country (Owoyele 1992; MoA, 2007). Since then, canine rabies has been endemic only in areas of southern Bhutan that share the porous border with India. Domestic dogs are the main reservoir of rabies in Bhutan, with spill over infection to other domestic animals, especially cattle. However, the reported incidence of rabies in humans is low and only sporadic cases occur in Bhutan (MoH, 2008)

Rabies is a notifiable disease in Bhutan and it is mandatory to report even suspected cases (RGoB, 2000). A thorough investigation is undertaken in the event of a suspected outbreak in animals. A fluorescent antibody test (FAT) – which is recommended by both the WHO and World Animal Health Organization (OIE) – is used to confirm rabies in animals (Dean et al., 1996; McElhinney et al., 2008; Fooks et al., 2009). The epidemiological data are collected and stored in the Veterinary Information System (VIS) database. As a control programme, vaccination and sterilization of dogs are conducted annually in the country focusing mostly on towns that have high, free-roaming dog densities. However, the programme coverage has been low (<20%) because of limited financial resources and the high population turnover in the free-roaming dog population (MoA, 2007).

In this report, we describe an outbreak of rabies in dogs and other domestic animals that occurred in the three eastern districts (Tashiyangtse, Trashigang, Mongar) of Bhutan (Figure 4.1a). Since the early 1990s, the villages and towns in these districts had been free from canine rabies and no wildlife-mediated rabies had ever been reported or confirmed in this region or elsewhere in Bhutan. However, between 2005 and 2007, rabies cases were reported in the region and therefore an outbreak was declared.

On 2 May 2005, the first case suggestive of rabies was reported in a cow at Gongza village, Toetsho subdistrict within Tashiyangtse district, located adjacent to the villages of Tawang district, Arunachal Pradesh state, India (see Figure 1b). Earlier, a suspected rabid, stray dog was sighted around a cattle herd but could not be traced subsequently. Following this incident, dog-bite cases in cattle and horses were reported in the adjacent villages of the index case. By September 2005, the disease apparently spread into the urban centres and villages of Trashigang district, located about 30 km to the south of the index case (see Figure 1) and then further spread into the villages and towns of five sub-districts of Mongar district, located next to Trashigang. There was a chain of spread from Tashiyangtse to Trashigang and then to Mongar districts. Epidemiological investigation and the typical clinical signs manifested by affected animals led to suspicion of rabies which was later confirmed by laboratory testing. Of the 51 brain samples (40 cattle, 11 dog) collected from the carcasses in the affected villages, 36 (70%) samples (30 cattle, 6 dog) were confirmed to be rabies virus positive by FAT. Samples from all suspected cases were not collected due to logistics and to avoid accidental rabies virus exposure of the people handling carcasses during sample collection. When one or more samples from the infected areas were confirmed to be positive, other suspected cases showing similar clinical signs were diagnosed as rabies. Observations suggested an incursion of rabies virus and subsequent local diffusion through the susceptible dog population, with spill-over over particularly to cattle.

The epidemiological data – day, month and year of occurrence; species of animals affected (cattle, horses, dogs, cats); and location (village, sub-district, district) of outbreaks were acquired. An epidemic curve was constructed for the whole study region by counting the total number of cases reported per month between May 2005 and November 2007, with the number of cases on the y-axis and the month on the x-axis. The cumulative incidence of rabies was calculated as the proportion of the population at risk that were diagnosed as rabies between 2005 and 2007. The animal census data of 2005 was used as the denominator to calculate the incidence risk for cattle, horses and cats. Since accurate dog

population data for the subdistricts of the three districts were not available, we estimated the dog population based on the mean subdistrict human population using ratios proposed and validated by [Knobel et al. \(2005\)](#) in Asia, namely 9.5 humans to one dog (95% CI 4.5:1–14.6:1).




Figure 1: Map of South East Asia with an administrative map of Bhutan and the three eastern districts shaded; (a) inset. The monthly report and spread of rabies cases in each sub-district of eastern Bhutan between May 2005 and November 2007 is shown. (b) May 2005 to October 2005 [the index case (●) is indicated]. (c) November 2005 to April 2006. (d) May 2006 to October 2006. (e) November 2006 to April 2007. (f) May 2007 to November 2007 (the last case was reported in November 2007).

These authors calculated the ratios based on the available published reports of field surveys conducted in many Asian countries. The estimated dog population was then used to calculate the cumulative incidence of rabies in dogs. A Geographical Information System (ArcGIS 9.3. ESRI, CA, USA) was used to map and visualize the monthly reports and the spread of rabies in eastern Bhutan.

A total of 256 animal deaths [141 cattle, 106 dogs (38 pets, 68 free-roaming), seven horses and two cats] were reported between May 2005 and November 2007. Table 1 shows the cumulative incidence of rabies in cattle, dogs, horses and cats in the three districts. The animal pattern showed a much higher incidence risk in dogs compared to other species of animals.

The yearly species-specific incidence risk of rabies showed higher risk in all species of animals during 2006 than in 2005 and 2007 (Table 2) indicating increased transmission of infection during 2006. This is also evident from the epidemic curve (Figure 2).

Table 1: Cumulative incidence of rabies from May 2005 to November 2007, stratified by species and district, eastern Bhutan.

District	Species	Cases	Population	Incidence* (95% CI)
Mongar	Cattle	40	28158	14.2 (10.4–19.3)
	Dog	48	3902	123 (92.9–162.7)
	Horse	3	3121	9.6 (3.32–28.2)
Tashiyangtse	Cattle	28	12122	23.1 (15.9–33.3)
	Dog	27	1876	143.9 (99.1–208.5)
	Horse	2	1837	10.8 (2.9–39.6)
	Cat	2	1156	17.3 (4.7–62.8)
Trashigang	Cattle	73	37552	19.4 (15.4–24.4)
	Dog	31	5383	57.5 (40.6–81.6)
	Horse	2	3399	5.8 (1.61–21.4)

CI: Confidence interval

\*Cumulative incidence, cases per 10,000 population at-risk

Table 2: Cumulative incidence of rabies between May 2005 and November 2007, stratified by year and species in the eastern region of Bhutan.

Period	Species	Cases	Incidence* (95% CI)
May–December 2005	Cattle	20	2.5 (1.6–3.9)
	Dog	10	8.9 (4.8–16.4)
	Horse	2	2.3 (0.6–8.7)

January–December 2006	Cattle	118	15.1 (12.6–18.1)
	Dog	94	84.2 (68.8–102.9)
	Horse	5	5.9 (2.5–13.9)
	Cat	2	3.2 (0.8–11.8)
January–November 2007	Cattle	3	0.3 (0.1–1.1)
	Dog	2	1.7 (0.4–6.5)

CI: Confidence interval

\* Cumulative incidence, cases per 10000 population at-risk




Figure 2: Epidemic curve of rabies cases in dogs and cattle in eastern Bhutan reported between May 2005 and November 2007. Cases in horses (n=7) and cats (n=2) are not shown because of the low numbers. ■ Cattle; □ free-roaming dogs; ▒, pet dogs.

Only one human death was reported during this epidemic, which accounted for 0.94 deaths/100 000 populations at-risk in the three districts. About 900 people were potentially exposed to rabies during the epidemic and they received 5–6 doses (Essen regimen) of rabies vaccination from the hospital. The maximum exposures were reported during 2006. The majority of these people consumed dairy products from a suspected rabid cow or handled carcasses of animals that died of rabies. However, it is not known how many of these people had actually been bitten by confirmed rabid dogs.

Figure 1 shows the monthly reports and spread of rabies in eastern Bhutan between May 2005 and November 2007. There was a rapid spread between January and April 2006 and a large number of sub-districts were affected during this period (Figure 4.1c). Some sub-districts reported persistence of infection for a longer period of time than other sub-districts. The rabies outbreak was eventually confirmed in 18 (45%) of the 40 sub-districts affecting 6/8, 7/15 and 5/17 sub-districts in Tashiyangtse, Trashigang and Mongar districts, respectively.

The proportion of sub-districts that reported rabies from May 2005 to December 2005, January 2006 to December 2006, and January 2007 to November 2007 were 15%, 35% and 7%, respectively. Both dog and cattle rabies was reported in all the sub-districts and almost at the same time, suggesting the role of dog bite in the transmission of disease to cattle (Figure 2). The source of the disease could not be

established, although it is likely to have spread from across the border, since a rabies outbreak in dogs had been reported in Tawang during this period (NS Tamang, personal communication).

The movement of infected free-roaming dogs is believed to have been responsible for the spread of infection in the region. The presence of a road network and the clustered settlement of villages and towns within the region could have facilitated the inter-mixing of dogs between the infected and non-infected areas resulting in rapid spread and persistence of infection. However, landscape features such as mountains and rivers may have been physical barriers to the further movement of dogs and prevented spread into other areas in the region. For example some villages/sub-districts, although located close to infected villages, did not report rabies because they are separated by a major river.

A series of measures were taken to control the outbreak. A public awareness programme was conducted during 2005 and 2006, when the outbreak was ongoing. Consultative meetings and discussions were held with farmers and village leaders to manage the outbreak. About 5000 dogs (both pets and free-roaming dogs) in the region were vaccinated against rabies. In addition, about 900 free-roaming dogs were caught and impounded in 12 temporary shelters constructed in the three districts. Impounding was done during March and April, 2006 in an attempt to prevent the spread of disease by the movement of free-roaming dogs. They were kept for about 4 months in the shelters and were released in the vicinity of capture after vaccination and sterilization. Although impounding is not a recommended strategy for the control of rabies, it was used during this outbreak as an alternative measure to mass culling of dogs, in response to the religious (Buddhist) sentiment of people in the community. No rabies cases were reported or confirmed in the dogs in the shelters. Further studies are necessary to determine the beneficial or deleterious effects of impounding dogs during an outbreak. Although mass culling of free-roaming dogs was not implemented during this outbreak, rabid and some in-contact dogs were eliminated after people realized the seriousness of the disease (many people had lost their valuable animals because of rabies). Such restricted culling may have removed the reservoir of infection resulting in fade-out of the epidemic. A similar major rabies epidemic that occurred in Chukha district in the south western part of Bhutan between January and July 2008 was controlled by immediate culling of in-contact free-roaming dogs in the outbreak areas. This outbreak affected three sub-districts and resulted in the death of 42 cattle, 52 dogs and three horses. Another rabies outbreak that occurred during 2008 in Samtse district – which is located adjacent to Chukha district (Figure 1a) – was also controlled by immediate culling of in-contact free-roaming dogs in the outbreak areas.

The experiences and the review of this epidemic suggest either the lack of a clear rabies outbreak control strategy for a major outbreak or the failure to implement the strategy in eastern Bhutan. Effective control of any disease depends on clear policy guidelines and their implementation, with legal support as well as effective communication. Therefore, control of any rabies outbreak in the future should be aimed at eliminating the reservoir of infection by immediate culling of in-contact and unvaccinated free-roaming dogs in the outbreak areas. An awareness programme on the public health importance of rabies is also necessary to promote good community participation in the control programme. As a preventive measure, sustained sterilization and vaccination of dogs with >70% coverage would induce herd immunity and eliminate rabies in canines (WHO, 2005). In addition, dog population management should be targeted towards habitat control by proper waste management and promoting a sense of responsible dog ownership in the community. Blanket mass culling or removal of dogs may not be a practical solution since it will be neither acceptable to the community (as evidenced from this outbreak) nor scientifically justifiable. We also recommend that the collection of good quality epidemiological data such as estimates of the dog population, combined with laboratory surveillance, would provide accurate information for the planning and management of rabies in Bhutan. In this

report, the risk estimates in dogs could be biased due to errors in estimating the population at risk in the region or inclusion of non-rabies cases in the data. Therefore, attempts should be made to collect all suspected samples in any future outbreak for laboratory confirmation of the cases. In addition, since no genetic characterization was done during this outbreak, it is also important to study phylogenetic relationships in order to understand rabies virus variants circulating in Bhutan.

The analyses presented in this report are descriptive and provide a basis for further analytical investigations to explain the evolution of the epidemic and generate hypotheses for the spread of rabies in eastern Bhutan.

### **Acknowledgements**

We thank the livestock field staff of Trashiyangtse, Trashigang and Mongar districts and the staff of the Regional Veterinary Laboratory, Khaling for their support during the time of investigation and for the field data.

**Declaration of interest:** None.

### **References**

1. Knobel DL, et al. Re-evaluating the burden of rabies in Africa and Asia. *Bulletin of the World Health Organization* 2005; 83: 360–368.
2. Sudarshan MK, et al. Assessing the burden of human rabies in India: results of a national multi-center epidemiological survey. *International Journal of Infectious Diseases* 2007; 11: 29–35.
3. Si H, et al. Rabies trend in China (1990–2007) and post-exposure prophylaxis in the Guangdong province. *BMC Infectious Diseases* 2008; 8: 113.
4. Wu X, et al. Reemerging rabies and lack of systemic surveillance in People’s Republic of China. *Emerging Infectious Diseases* 2009; 15: 1159–1164.
5. Owoyele GD. Rabies outbreak in Thimphu: Case Report. *Bhutan Journal of Animal Husbandry* 1992; 13: 36–39.
6. Rabies Prevention and Control Project for Bhutan 2007. (<http://www.moa.gov.bt/ncah/project/projects.php?id=3&pname=WHO%20Rabies%20Project>). Accessed 28 December 2009.
7. Government of Bhutan. Annual Health Bulletin 2008. Ministry of Health, Government of Bhutan, Thimphu.
8. Government of Bhutan. The Livestock Act of Bhutan 2000. The Royal Government of Bhutan, Ministry of Agriculture, Bhutan.
9. Dean DJ, Ableseth MK, Atanasiu P. The fluorescent antibody test. In: Meslin FX, Kaplan MM, Koprowski H, eds. *Laboratory Techniques in Rabies*, 4th edn. Geneva: World Health Organization, 1996, pp. 88–95.
10. McElhinney LM, Fooks AR, Radford AD. Diagnostic tools for the detection of rabies virus. *European Journal of Companion Animal Practice* 2008; 18:224–231.
11. Fooks AR, et al. Emerging technologies for the detection of rabies virus: challenges and hopes in the 21<sup>st</sup> century. *PLOS Neglected Tropical Diseases* 2009; 3:e530.
12. WHO. WHO Expert Consultation on Rabies: first report (WHO technical report series No. 931), 2004, Geneva, Switzerland.
- 13.