

## Effects of $\text{La}^{3+}$ on ATPase Activities of Plasma Membrane Vesicles Isolated from *Casuarina Equisetifolia* Seedlings under Acid Rain Stress\*

Li Yuhong (李裕红)<sup>1,2</sup>, Yan Chongling (严重玲)<sup>1\*</sup>, Liu Jingchun (刘景春)<sup>1</sup>, Chen Yinghua (陈英华)<sup>1</sup>, Hu Jun (胡俊)<sup>1</sup>, Xue Bo (薛博)<sup>1</sup>

(1. School of Life Sciences, Xiamen University, Xiamen 361005, China; 2. Department of Biology, Quanzhou Normal College, Quanzhou 362000, China)

**Abstract:** The effects of  $\text{La}^{3+}$  on the growth and the ATPases activities of plasma membrane (PM) vesicles isolated from *Casuarina equisetifolia* seedlings under artificial acid rain (pH 4.5) stress were studied. The results show that the height, length of roots, fresh weight and PM  $\text{H}^+$ -ATPase activities of *Casuarina equisetifolia* seedlings increase by the treatments of soaking seeds in  $\text{LaCl}_3$  solutions with lower concentrations, and those can reach their peak values by treating with  $200 \text{ mg L}^{-1} \text{La}^{3+}$ . However, in comparison with the CK, those are inhibited by the higher  $\text{La}^{3+}$  concentrations; PM  $\text{Ca}^{2+}$ -ATPase activity is inhibited with the treatments of  $\text{La}^{3+}$ . The results also reveal that the  $\text{H}^+$ -ATPase activity and the growth of cell enlarge have a remarkable positive correlation, and  $\text{La}^{3+}$  activating  $\text{H}^+$ -ATPase can facilitate plant growth.  $\text{La}^{3+}$  also can alleviate cytosolic acidification of plant under acid rain stress and indirectly maintain the stability of intracellular environment. In order to resistant to acid rain and accelerate the growth of *Casuarina equisetifolia*, the suitable range of  $\text{La}^{3+}$  concentrations to soak seeds for 8 h is  $50 \sim 200 \text{ mg L}^{-1}$ .

**Key words:** ecology; PM ATPase; lanthanum; *Casuarina equisetifolia*; acid rain stress; rare earths

**CLC number:** O614.33      **Document code:** A      **Article ID:** 1002 - 0721(2003)06 - 0675 - 05

As a main kind of membrane transporter, plasma membrane ATPases facilitate the transport of minerals, sugars, metabolites, and other compounds across the membrane of the cells, only by which could plant cells communicate with environment to transfer substance and information, which is essential for plant growth and cellular homeostasis<sup>[1]</sup>. As the experiment materials, the highly purified plasma membrane vesicles from *Casuarina equisetifolia* seedlings cultivated under artificial acid rain stress, were isolated by aqueous two-phase partitioning methods. The study mainly focused on the effects of  $\text{La}^{3+}$  on the PM ATPases activities, and was expected to show some inherent mechanisms and provide some scientific basis for seeking an approach of utilizing rare earth elements to improve the resistance of *Casuarina* shelter forests to resist acid rain stress.

## 1 Materials and Methods

### 1.1 Plant material

The *Casuarina equisetifolia* seeds were supplied

by the *Casuarina* Seedling Base of ChiHu Forestry Center in HuiAn County, Fujian Province. The average weight of one thousand grains was 1.44 g. After being randomly selected, the seeds were surface sterilized in 0.5%  $\text{KMnO}_4$  (5 min), then washed and filtered three times in sterile distilled water. The sterilized seeds were soaked for 8 h in  $\text{LaCl}_3 \cdot 7\text{H}_2\text{O}$  with the concentrations of 0 (CK, sterile distilled water); 50, 100, 200, 300 and 400  $\text{mg L}^{-1}$  respectively, and 10 g seeds each treatment were weighed, and flushed, then filtered to remove  $\text{La}^{3+}$  from the surfaces. Then the seeds were cultured in the basin filled with sands, and the culture solution is the artificial acid rain with pH value of 4.5 as the same as that of naturally happened in higher frequency in Xiamen region<sup>[2]</sup>. The ions concentration of artificial acid rain was according to Ref. [3]. Every treatment repeated nine times. The basins were kept in the glasshouse. In the period of culture, the basins were taken out of the solution to aerate for 1 h every nightfall, and the pH value of artificial acid rain was kept invariable. The culture temperature was  $(22 \pm 2)$ . Two weeks later, at that

\* Received date: 2003 - 01 - 30; revised date: 2003 - 05 - 10

Foundation item: Project supported by National Natural Science Foundation (30170190); Key Natural Science Foundation of Fujian Province (D0120001) and Scientific Research Foundation of Quanzhou Normal College (2002LI04)

Biography: Li Yuhong (1969 - ), Female, Doctor candidate, Lecturer

\* Corresponding author (E-mail: ycl@xmu.edu.cn)

time the plants were about 5 ~ 6 cm height, the seedlings were taken to experiment.

## 1.2 Determination

Determination of the plant growth and biomass: 10 seedlings were randomly taken from each treatment group, then the plant height and the length of main root were measured. The fresh weight of each seedling was weighed by electron scale.

The technique method for preparing plasma membrane vesicles by the method of two - phase partition was referred to that described by Zheng<sup>[4]</sup>.

The PM H<sup>+</sup>-ATPase activity was determined by the method described by Zhang<sup>[5]</sup>: 20  $\mu$ l vesicle membrane preparations was added into 0.5 ml reaction medium consisted of MgSO<sub>4</sub> 3 mmol  $\cdot$  L<sup>-1</sup>, K<sub>2</sub>SO<sub>4</sub> 25 mmol L<sup>-1</sup>, Triton X-100 0.02% (V/V), Tris-Mes 50 mmol L<sup>-1</sup>, pH 6.5, and ATP-Na<sub>2</sub> 3 mmol  $\cdot$  L<sup>-1</sup>. The reactants were incubated in 37 °C for 30 min and the reaction was stopped by 20% (W/V) TCA 200  $\mu$ l. The activities of the enzyme were calculated by the amounts of Pi formed by hydrolysis in the reactions.

PM Ca<sup>2+</sup>-ATPase hydrolysis activity was determined by the method of Li X M<sup>[6]</sup>: 1.1 ml reaction system included imidazole 10 mmol  $\cdot$  L<sup>-1</sup>, MgCl<sub>2</sub> 5 mmol L<sup>-1</sup>, CaCl<sub>2</sub> 50  $\mu$ mol L<sup>-1</sup>, ATP-Na<sub>2</sub> 3 mmol L<sup>-1</sup>, pH 7.0, and 20  $\mu$ l membrane preparations. The enzyme reaction followed the steps of measuring H<sup>+</sup>-ATPase activity. Determination of Pi was according to Ohnishi<sup>[7]</sup>; Protein content was determined by using Coomassie Brilliant Blue G 250 referred to Bradford<sup>[8]</sup>. All the determinations were repeated six times.

## 2 Results and Discussion

### 2.1 Effects of lanthanum on seedling growth

The experiment results (Table 1) indicate that the effects of lanthanum on the growth of *Casuarina equisetifolia* seedlings under acid rain stress with pH value 4.5 is obvious, and the growth including the height, length of roots, and fresh weight of *Casuarina equisetifolia* seedlings is promoted gradually by soaking seeds for 8 h in LaCl<sub>3</sub>  $\cdot$  7H<sub>2</sub>O solution with the increasing

concentrations from 50 to 200 mg  $\cdot$  L<sup>-1</sup>, and in the treatment of 200 mg  $\cdot$  L<sup>-1</sup> La<sup>3+</sup>, all those values reach the highest points. But in the treatment of 300 mg  $\cdot$  L<sup>-1</sup> La<sup>3+</sup>, the effects descend. As to treating with 400 mg  $\cdot$  L<sup>-1</sup> La<sup>3+</sup>, the plant growth is inhibited. The experiment results indicate that rare earth elements have obvious function in alleviating the effects of acid rain on plant biomass; It also suggests that rare earth elements with suitable concentration can promote the growth of plant. Therefore it is feasible that La<sup>3+</sup> could enhance the ability of *Casuarina equisetifolia* to resistant to acid rain stress.

### 2.2 Effects of La<sup>3+</sup> on PM H<sup>+</sup>-ATPase activity

After treating with La<sup>3+</sup> of different concentrations, the changes of PM H<sup>+</sup>-ATPase activity of *Casuarina equisetifolia* seedlings show that lower concentration of La<sup>3+</sup> can stimulate the activity, but higher concentration will inhibit it. The seedling PM H<sup>+</sup>-ATPase activity increases gradually with the increase of La<sup>3+</sup> from 50 to 200 mg  $\cdot$  L<sup>-1</sup>. While in the treatment of 200 mg  $\cdot$  L<sup>-1</sup>, PM H<sup>+</sup>-ATPase activity reaches its peak value, and increases 35.85% compared with CK; However, when the concentration is over 200 mg  $\cdot$  L<sup>-1</sup>, the activity of PM H<sup>+</sup>-ATPase is inhibited and lower than that of CK. In the treatments of 300 and 400 mg  $\cdot$  L<sup>-1</sup> La<sup>3+</sup>, the activities are 72.29% and 90.88% of CK respectively.

H<sup>+</sup>-ATPase of the plasma membrane is considered as a master enzyme. The improvement of PM H<sup>+</sup>-ATPase activity would facilitate building up the gradient of proton electrochemical potentials<sup>[9]</sup>, which can provide stronger driving force for inorganic ions, or organic solutes such as sugars and amino acids, thereby the transport of substance and growth of plant are accelerated. Acid rain in China belongs to sulfate type. The transport of sulfate on plant plasma membrane is powered by an electrochemical proton gradient generated by proton pump of ATPase that extrudes protons to the cell exterior. Sulfate is actively cotransported into plant cells along with protons at a stoichiometry of one sulfate to three protons, thus sulfate transporter is able to couple the influx of net protons to the transport of sulfate into the cell<sup>[1]</sup>. Which would influence the stabilization of cytosolic pH (cytosolic pH is remark-

**Table 1** Effect of La<sup>3+</sup> on seeding growth of *Casuarina equisetifolia* under acid rain stress (X  $\pm$ SD)

| La <sup>3+</sup> / (mg L <sup>-1</sup> ) | 0               | 50              | 100             | 200             | 300             | 400             |
|--|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| Height of plant/cm                       | 5.43 $\pm$ 0.49 | 5.66 $\pm$ 0.46 | 5.76 $\pm$ 0.39 | 6.32 $\pm$ 0.79 | 5.54 $\pm$ 0.38 | 5.31 $\pm$ 0.24 |
| Length of roots/cm                       | 3.09 $\pm$ 0.36 | 3.17 $\pm$ 0.39 | 3.22 $\pm$ 0.32 | 3.54 $\pm$ 0.34 | 3.28 $\pm$ 0.41 | 2.85 $\pm$ 0.26 |
| Fresh weight/mg                          | 8.04 $\pm$ 0.1  | 8.36 $\pm$ 0.15 | 9.70 $\pm$ 0.36 | 9.90 $\pm$ 0.65 | 9.30 $\pm$ 0.15 | 7.82 $\pm$ 0.26 |




Fig. 1 Effects of  $\text{La}^{3+}$  on PM  $\text{H}^{+}$ -ATPase activity of *Casuarina equisetifolia* seedlings under acid rain stress (pH 4.5)

ably stable, generally remaining at 7.3 ~ 7.5).  $\text{H}^{+}$ -ATPase has a function in extruding protons to the cell exterior, and the promotion of  $\text{H}^{+}$ -ATPase activity facilitates transporting proton out of cell to prevent acidification of the cytoplasm, which keeps inner cellular environment in normal and stable state when plant is under acid rain stress.

Two aspects are known to the growth of plants. The first is an increase in the number of cells in limited parts in plant body; the second is the enlargement in the size of each cell, and this kind of growth promotion is also caused by the plant hormone, auxin<sup>[9]</sup>. The proton pump is activated by auxin. The inner surface of the cell wall is acidified by the  $\text{H}^{+}$  formed. The acidification of the cell wall activates the proteins of loosening H-bonds within the cell wall and allows turgorgenerated growth<sup>[11]</sup>. Finally, elongation of the plant body is accelerated.

Therefore the activation of  $\text{H}^{+}$ -ATPase directly contributes to the growth of plant. The growth promotion by lanthanum is just due to a similar mechanism to that described above. Table 1 and Fig. 1 indicate that there are coherent trend of responding to lanthanum between plant height and  $\text{H}^{+}$ -ATPase activation. Analyzing the correlation using the two group average data of plant height and  $\text{H}^{+}$ -ATPase activity shows that they have a remarkable positive correlation. It could be supposed that  $\text{La}^{3+}$  activating PM  $\text{H}^{+}$ -ATPase is one of the mechanisms that  $\text{La}^{3+}$  could accelerate plant enlarging growth.

The C terminal of  $\text{H}^{+}$ -ATPase polypeptide in cytoplasm forms an autoinhibitory domain, and the removal of the autoinhibitory domain activates the enzyme considerably<sup>[11]</sup>. The modulation mechanism of autoinhibitory domain in the C-terminal of PM  $\text{H}^{+}$ -ATPase is possibly by the way of binding an effector, thus the C-terminal part of the  $\text{H}^{+}$ -ATPase loses the autoinhibitory function<sup>[10]</sup>. The mechanism of  $\text{La}^{3+}$  modulating  $\text{H}^{+}$ -ATPase activity might be as follows: first,  $\text{La}^{3+}$  as an effector binding the C terminus re-

gion of  $\text{H}^{+}$ -ATPase relieves autoinhibition function, which could increase the express of  $\text{H}^{+}$ -ATPase activation; second,  $\text{La}^{3+}$  predominantly combines phospholipid of plasma membrane<sup>[11]</sup>, and the combination is as an effector to relieve the autoinhibition function of  $\text{H}^{+}$ -ATPase C terminus, therefore the enzyme activation is affected. As to PM  $\text{H}^{+}$ -ATPase activity decreases with  $\text{La}^{3+}$  treatment in higher concentration (300 ~ 400  $\text{mg L}^{-1}$ ), it might attribute to that when the  $\text{La}^{3+}$  concentration is over the concentration limit and become a kind of ion stress, it would inhibit the gene expression of part of  $\text{H}^{+}$ -ATPase.

### 2.3 Effects of $\text{La}^{3+}$ on PM $\text{Ca}^{2+}$ -ATPase activity

The effects of  $\text{La}^{3+}$  on the plasma membrane  $\text{Ca}^{2+}$ -ATPase activities show as Fig. 2 that the activities of PM  $\text{Ca}^{2+}$ -ATPase are inhibited and the activities decrease with  $\text{La}^{3+}$  concentration increasing from 50 to 400  $\text{mg L}^{-1}$ .

As a second messenger,  $\text{Ca}^{2+}$  participates in many intracellular actions, which is essential for conveying intracellular information and modulating cell metastasis. The transport of  $\text{Ca}^{2+}$  across the plasma membrane is finished by three kinds of transport proteins:  $\text{Ca}^{2+}$ -ATPase,  $\text{Ca}^{2+}/\text{H}^{+}$  antiporter and  $\text{Ca}^{2+}$  channel.  $\text{Ca}^{2+}$ -ATPase and  $\text{Ca}^{2+}/\text{H}^{+}$  antiporter are in charge of efflux of  $\text{Ca}^{2+}$  across the plasma membrane, and  $\text{Ca}^{2+}$  channel is in charge of influx of  $\text{Ca}^{2+}$ <sup>[12]</sup>. It is known that, in biology systems, rare earth elements can represent similar physiological action as  $\text{Ca}^{2+}$ , and replace  $\text{Ca}^{2+}$  binding on the plasma membrane surface, and block the efflux and influx of  $\text{Ca}^{2+}$ <sup>[13]</sup>. Several studies indicate that  $\text{La}^{3+}$  inhibits  $\text{Ca}^{2+}$ -ATPase activity<sup>[14]</sup>, and rare earth elements are  $\text{Ca}^{2+}$  antagonist and  $\text{Ca}^{2+}$  channel blocker of cell membrane surface<sup>[15]</sup>. However, the mechanism of action is not clear yet.

PM  $\text{Ca}^{2+}$ -ATPase, acting as specific active transport system of  $\text{Ca}^{2+}$  on plasma membrane of higher plant, can directly hydrolyze ATP and actively transport  $\text{Ca}^{2+}$ , therefore it plays an important role in low-




Fig. 2 Effects of  $\text{La}^{3+}$  on PM  $\text{Ca}^{2+}$ -ATPase activity of *Casuarina equisetifolia* seedlings under acid rain stress (pH 4.5)

ing the  $\text{Ca}^{2+}$  concentration of plasma and maintaining  $\text{Ca}^{2+}$  concentration at a stable level (the cytosolic concentration of free  $\text{Ca}^{2+}$  is about  $0.2 \mu\text{mol L}^{-1}$ ). In plasma membrane,  $\text{Ca}^{2+}$ -ATPase activity can be enhanced by calmodulin<sup>[1]</sup>.

In this experiment, the reasons that  $\text{La}^{3+}$  inhibits the activities of PM  $\text{Ca}^{2+}$ -ATPase may be: Firstly, the rise of PM  $\text{Ca}^{2+}$ -ATPase activities increases its resistance to acid rain stress. One of our experiments showed that compare to *Casuarina equisetifolia* seedlings without the acid rain stress, the seedlings treated with acid rain stress with pH 4.5 have a higher value of PM  $\text{Ca}^{2+}$ -ATPase activities. Owing to using rare earth elements, plants sensitivity to the acid rain is reduced<sup>[3]</sup>, the sensitivity of PM  $\text{Ca}^{2+}$ -ATPase to the acid rain can be reduced, and keep the activities of PM  $\text{Ca}^{2+}$ -ATPase at a relatively low level by substituting  $\text{Ca}^{2+}$  with  $\text{La}^{3+}$ . Secondly, as  $\text{La}^{3+}$  is the antagonist of  $\text{Ca}^{2+}$  on PM surface, the position of  $\text{Ca}^{2+}$  can be occupied by  $\text{La}^{3+}$  competitively, blocking the efflux and influx of  $\text{Ca}^{2+}$  in the cell, and influencing  $\text{Ca}^{2+}$  signal transmission, which hinders the activation of  $\text{Ca}^{2+}$  binding proteins such as calmodulin, and inhibits  $\text{Ca}^{2+}$ -ATPase activity. Thirdly,  $\text{Ca}^{2+}$  has a relatively higher dissociation constant with PM than that of  $\text{La}^{3+}$ , and  $\text{La}^{3+}$  can replace the bond position of  $\text{Ca}^{2+}$  in PM. The electricity of  $\text{La}^{3+}$  is higher than that of  $\text{Ca}^{2+}$ , so this kind of replace may result in the changes of PM electric potential and then indirectly influence the activities of PM  $\text{Ca}^{2+}$ -ATPase. Fourthly, acid rain can bring some damage to the PM lipid, and the stress can affect the structure and function of plasma membrane lipid. Some studies have showed that the  $\text{Ca}^{2+}$ -ATPase activities of the purified plasma membrane without lipid would be inhibited more and more strongly by  $\text{La}^{3+}$  with increasing concentrations<sup>[6]</sup>. It can be supposed that, the decrease of PM  $\text{Ca}^{2+}$ -ATPase activities by  $\text{La}^{3+}$  might be owing to the abnormal change of plasma membrane lipid environment brought by acid rain, and the inhibition of the activities are strengthened with the increasing  $\text{La}^{3+}$  concentrations.

### 3 Conclusions

1. The growth including height, length of roots, fresh weight and PM  $\text{H}^{+}$ -ATPase activities of *Casuarina equisetifolia* seedlings are stimulated and increase by the treatments of soaking seeds for 8 h in  $\text{LaCl}_3$  with lower concentrations, and their peak values all appear with  $200 \text{ mg L}^{-1} \text{La}^{3+}$ . However, those are inhibited by higher  $\text{La}^{3+}$  concentrations. There is remarkable positive correlation between  $\text{H}^{+}$ -ATPase activity and

plant height.  $\text{La}^{3+}$  activates PM  $\text{H}^{+}$ -ATPase is one of the reasons that  $\text{La}^{3+}$  can accelerate the growth of cell enlarging.  $\text{La}^{3+}$  can enhance  $\text{H}^{+}$ -ATPase activity, alleviate cytosolic acidification of plant under acid rain stress and indirectly maintain the stability of intracellular environment.

2.  $\text{La}^{3+}$  inhibits PM  $\text{Ca}^{2+}$ -ATPase activity, and the inhibition is strengthened with the increasing  $\text{La}^{3+}$  concentrations from 50 to  $400 \text{ mg L}^{-1}$ . The replacing of  $\text{Ca}^{2+}$  with  $\text{La}^{3+}$  in PM could weaken the sensitivity of  $\text{Ca}^{2+}$ -ATPase to acid rain stress, and hinder calmodulin activation, and change membrane voltage, which may lower  $\text{Ca}^{2+}$ -ATPase activity. And the decreased activity also is due to the abnormal membrane lipid environment brought by acid rain.

3. In order to resistant to acid rain stress and accelerate the growth of *Casuarina equisetifolia*, the suitable concentration range of  $\text{LaCl}_3$  treatment is  $50 \sim 200 \text{ mg L}^{-1}$ .

### References :

- [1] Bob B. Buchanan, Wilhelm Gruissem, et al. Biochemistry & Molecular Biology of Plants [M]. Beijing: Science Press (by arrangement with the American Society of Plant physiologists, 2000), 2002, 110, 121, 123, 831, 832, 833.
- [2] Chen D Y, Zhuang Z X, Wang X R. Distribution of acid rain and the pH variations during raining process in Xiamen city, China [J]. Research of Environmental Sciences, 1994, 7(6): 26.
- [3] Yan Chongling, Hong Yetang, Lin Peng, et al. Accumulation of rare earth elements in spinach and soil under condition of using REE and acid rain stress [J]. Journal of Rare Earths, 2002, 20(2): 133.
- [4] Zheng H L, Zhao Z Q, Zhang C G, et al. Changes in lipid peroxidation, the redox system and ATPase activities in plasma membranes of rice seedling roots caused by lanthanum chloride [J]. BioMetals, 2000, 13: 157.
- [5] Zhang W H, Liu Y L, Yu B J. Improvement of determination of ATP-Dependent proton pump in tonoplast vesicles from barley roots [J]. J. Plant Physiology Commun., 1997, 33(4): 282.
- [6] Li X M, Ni J Z. Effects of rare earths on the activity of sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase [J]. Chinese Biochemical Journal, 1995, 11(3): 281.
- [7] Ohnishi T, Gall R S, Mayer M L. An improved assay of inorganic phosphate in the presence of extralabile phosphate compounds: Application to the ATPase assay in the presence of phosphocreatine [J]. Anal. Biochem., 1975, 69: 261.
- [8] Bradford M M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding [J]. Anal. Biochem., 1976, 72: 248.
- [9] Pierre Morsomme, Marc Boutry. The plant plasma

- membrane H<sup>+</sup>-ATPase: Structure, function and regulation [J]. *Biochimica et Biophysica Acta.*, 2000, (1465): 1.
- [10] Palmgren M G. Regulation of plant plasma membrane H<sup>+</sup>-ATPase activity [J]. *Physiol. Plant*, 1991, 83: 316.
- [11] Cai J B, Wang P, Huang B X, et al. Speciation of La in cell plasma membrane [J]. *Journal of the Chinese Rare Earth Society (in Chin.)*, 2001, 19(2): 191.
- [12] Douglas S B. Calcium regulation in plant cells and its role in signaling [J]. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 1995, 46: 95.
- [13] Rai M, Yang S Q, McClain C, et al. Kupffer cell depletion by gadolinium chloride enhances liver regeneration after partial hepatectomy in rats [J]. *Am. J. Physiol.*, 1996, 270: G909.
- [14] Peter Gräf, Elmar W. Weiler. ATP-driven Ca<sup>2+</sup> transport in sealed plasma membrane vesicles prepared by aqueous two-phase partitioning from leaves of *Commelina communis* [J]. *Physiol. Plant*, 1989, 75: 469.
- [15] Terry B T, Findlay G P, Tyerman S D. Direct effects of Ca<sup>2+</sup>-channel blockers on plasma membrane cation channels of *Amaranthus tricolor* protoplasts [J]. *J. Exp. Bot.*, 1992, 43(256): 1457.

## Influence of RE on Cohesive Force of Spray Coating from One-shot Self-felt Iron-Based Composite Powder

Ma Chen<sup>1\*</sup>, Li Muqin<sup>1,2</sup>, Yin Ke<sup>1</sup>, Zou Renjie<sup>1</sup> (1. *Biomedicine Material Key Laboratory, Jiamusi University, Jiamusi 154007, China*; 2. *National Key Laboratory of Advanced Welding Production Technology, Harbin Institute of Technology, Harbin 150001, China*)

**Abstract:** The strength of coating layer obtained by one-shot self-felt powder with subsonic oxygen-acetylene spraying technique was discussed. Due to self-heating effect among elements in powders and micro-alloying, the highest strength is gotten when RE contents in the one-shot self-felt powder are 0.9%. The experimental results show that coating layer could be

**Key words:** inorganic materials; self-felt; composite powder; wear-ability; rare earths

purified by adding proper amounts of RE, which make the inclusion change from reticulate pattern to spherical. RE-compound could be formed when the RE acted with other elements, which make the melting point of powders fall down and the wetting ability could be increased, which should be beneficial to density coating layer and the wear-ability could be increased.

(See *J. Chin. RE. Soc. (in Chin.)*, 2003, 21(6): 733 for full text)

## Effects of Rare Earth Elements La and Ce on Ti-Ni Alloy

Zhao Yuechao<sup>1</sup>, Ma Zhuang<sup>1</sup>, Fu Dajun<sup>1</sup>, Liu Ailian<sup>2</sup> (1. *Institute of Machinery Liaoning Technical University, Fuxin 123000, China*; 2. *Materials Institute, Harbin University of Technology, Harbin 150001, China*)

**Abstract:** Rare earth elements Ce and La were added into the near-equal atomic titanium-nickel alloy. Differential scanning calorimetry and X-ray diffraction were applied to study the effect of the Ce and La. The results show that adding a small amount of Ce into

**Key words:** metal materials; TiNi alloy; phase transformation; rare earths

TiNi alloy can improve the phase transformation obviously, but the phases in TiNi alloy do not change. The results show that adding a small amount of Ce or La can improve the properties of titanium-nickel alloy.

(See *J. Chin. RE. Soc. (in Chin.)*, 2003, 21(6): 736 for full text)