

## Rac1-MKK3-p38-MAPKAPK2 Pathway Promotes Urokinase Plasminogen Activator mRNA Stability in Invasive Breast Cancer Cells\*

Received for publication, September 17, 2002, and in revised form, October 3, 2002  
Published, JBC Papers in Press, October 10, 2002, DOI 10.1074/jbc.M209542200

Qiwei Han<sup>‡</sup>, Jay Leng<sup>§</sup>, Dafang Bian<sup>‡</sup>, Chitlada Mahanivong<sup>‡</sup>, Kevin A. Carpenter<sup>§</sup>,  
Zhixing K. Pan<sup>||</sup>, Jiahuai Han<sup>‡</sup>, and Shuang Huang<sup>‡\*\*</sup>

From the Departments of <sup>‡</sup>Immunology and <sup>||</sup>Molecular and Experimental Medicine, The Scripps Research Institute, La Jolla, California 92037 and <sup>§</sup>Chemicon International, Inc., Temecula, California 92590

We reported previously that down-regulating or functionally blocking  $\alpha v$  integrins inhibits endogenous p38 mitogen-activated protein kinase (MAPK) activity and urokinase plasminogen activator (uPA) expression in invasive MDA-MB-231 breast cancer cells whereas engaging  $\alpha v$  integrins with vitronectin activates p38 MAPK and up-regulates uPA expression (Chen, J., Baskerville, C., Han, Q., Pan, Z., and Huang, S. (2001) *J. Biol. Chem.* 276, 47901–47905). Currently, it is not clear what upstream and downstream signaling molecules of p38 MAPK mediate  $\alpha v$  integrin-mediated uPA up-regulation. In the present study, we found that  $\alpha v$  integrin ligation activated small GTPase Rac1 preferentially, and dominant negative Rac1 inhibited  $\alpha v$  integrin-mediated p38 MAPK activation. Using constitutively active MAPK kinases, we found that both constitutively active MKK3 and MKK6 mutants were able to activate p38 MAPK and up-regulate uPA expression, but only dominant negative MKK3 blocked  $\alpha v$  integrin-mediated p38 MAPK activation and uPA up-regulation. These results suggest that MKK3, rather than MKK6, mediates  $\alpha v$  integrin-induced p38 MAPK activation. Among the potential downstream effectors of p38 MAPK, we found that only MAPK-activated protein kinase 2 affects  $\alpha v$  integrin-mediated uPA up-regulation significantly. Finally, using  $\beta$ -globin reporter gene constructs containing uPA mRNA 3'-untranslated region (UTR) and adenosine/uridine-rich elements-deleted 3'-UTR, we demonstrated that p38 MAPK/MAPK-activated protein kinase 2 signaling pathway regulated uPA mRNA stability through a mechanism involving the adenosine/uridine-rich elements sequence in 3'-UTR of uPA mRNA.

Urokinase plasminogen activator (uPA)<sup>1</sup> is overexpressed in various malignancies including breast, ovarian, and colon can-

cer (1–6). Both *in vitro* and *in vivo* studies have demonstrated clearly that the levels of uPA are associated closely with the degree of tumor cell invasion and that uPA plays a key role in tumor progression and metastasis (7–15). When uPA binds to its cell surface receptor, uPA rapidly converts plasminogen into plasmin, which then degrades a variety of extracellular matrix proteins and also initiates the activation of a metalloproteinase cascade (16, 17). In addition, the interaction of uPA with the uPA receptor also mediates a variety of cell responses including adhesion, migration, proliferation, and transcription of specific genes (18–26), and these processes have potential impact in tumor invasion and metastasis.

The p38 MAPKs include four isoforms ( $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\delta$ ) (27, 28). Despite the structural similarity among the members of p38 MAPK family, differences in activation profile and substrate specificity have been observed (28, 29). The p38 MAPKs can be activated by a wide spectrum of stimuli, including cellular stress, proinflammatory cytokines, and growth factors (29, 30). The activation of p38 MAPKs is regulated by upstream MAPKK (MKK3, MKK6, and probably MKK4) via phosphorylation of a TGY phosphorylation site (28, 31). The p38 MAPK effects are carried out by downstream substrates including protein kinases and transcription factors (28, 30). Recently, p38 MAPK signaling pathway has also been demonstrated to play an important role in regulating mRNA stability (32–34).

The concentration of an mRNA is a function of its rates of synthesis and degradation (35). The regulation of mRNA stability is therefore an important means of modulating gene expression. Generally, mRNA stability is controlled by *cis*-acting sequences within 5'- or 3'-untranslated regions (UTRs) or, in some cases, within the coding region (35, 36). The best characterized regulatory elements are the adenosine/uridine-rich elements (ARE) within 3'-UTR of cytokine, growth factor, and proto-oncogene mRNAs, and these elements often contain several copies of the motif AUUUA (36). Attaching ARE within a uridine-rich context results in the instability of otherwise stable reporter mRNA such as  $\beta$ -globin mRNA (37, 38). Furthermore, mRNA stability is regulated by *trans*-acting RNA binding factors that interact with AREs (36), and these include AUF1 (39, 40), HuR (41, 42), and tristetraprolin (32, 43).

Our previous studies showed that engaging  $\alpha v$  integrins with vitronectin (Vn) activates p38 MAPK and up-regulates uPA expression in invasive MDA-MB-231 cells and that the expression/function of  $\alpha v$  integrins is essential for constitutive p38 MAPK activity and uPA expression in invasive cancer cells

\* This work was supported by NCI, National Institutes of Health Grant R01 CA93926, Department of Army Grant OC990129, and California Cancer Research Program Grant 00-00755V-20041. This is Paper IMM-15307 from The Scripps Research Institute. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>‡</sup> Supported by National Institutes of Health Training Grant T32 HL07195.

\*\* To whom correspondence should be addressed: Dept. of Immunology, IMM19, The Scripps Research Inst., 10550 N. Torrey Pines Rd., La Jolla, CA 92037. Tel.: 858-784-9211; Fax: 858-784-8472; E-mail: shuang@scripps.edu.

<sup>1</sup> The abbreviations used are: uPA, urokinase plasminogen activator; MAPK, mitogen-activated protein kinase; MAPKK, MAPK kinase; UTR, untranslated region; ARE, adenosine/uridine-rich element; Vn, vitronectin; mAb, monoclonal antibody; Ad, adenovirus; PBD, Rac1/Cdc42 binding domain; RBD, Rho binding domain; MAPKAPK2,

MAPK-activated protein kinase 2; MKK, MAPK kinase; GST, glutathione S-transferase; MEK, MAPK/extracellular signal-regulated kinase; MSK, mitogen and stress-activated kinase; MNK, MAPK-interacting kinase; pfu, plaque-forming unit; IL, interleukin; PRAK, p38 MAPK-activated kinase.

(44). We and others (45, 46) have also shown that p38 MAPK regulates uPA expression by promoting uPA mRNA stability. However, the upstream and downstream signaling molecules of p38 MAPK involved in Vn/ $\alpha$ v integrin ligation-mediated uPA up-regulation remain to be elucidated. In the present study, we demonstrate that Vn/ $\alpha$ v integrin ligation activates Rac1 preferentially and that the Rac1 activity is important for Vn-induced p38 MAPK activation. We also show that although both constitutively active MKK3 and MKK6 enhance uPA expression, only dominant negative MKK3 blocks Vn/ $\alpha$ v integrin ligation-induced p38 MAPK activation and uPA up-regulation. Among the potential p38 MAPK downstream kinases, only dominant negative MAPKAPK2 inhibits uPA expression and destabilizes uPA mRNA. In the parallel experiments, we find that constitutively active MAPKAPK2 restores uPA expression and prolongs uPA mRNA stability in p38 MAPK-inhibited MDA-MB-231 cells. These findings suggest that a signaling pathway involving Rac1-MKK3-p38-MAPKAPK2 mediates Vn/ $\alpha$ v integrin-mediated uPA up-regulation. In addition, we provide evidence that the ARE-containing 3'-UTR of uPA mRNA is essential for p38 MAPK/MAPKAPK2-regulated uPA mRNA stability.

#### EXPERIMENTAL PROCEDURES

**Materials and Cell Culture**—Polyclonal antibodies to phospho-p38, p38, phospho-MKK3/6, and phospho-MKK4 were purchased from Cell Signaling (Beverly, MA). The polyclonal antibody to uPA was obtained from American Diagnostica (Greenwich, CT). Rac1 monoclonal antibody (mAb), anti-active MAPKAPK2, and MAPKAPK2 polyclonal antibodies were obtained from Upstate Biotechnology (Lake Placid, NY). Cdc42, RhoA mAbs, and MKK3, MKK4, MKK6, myc tag polyclonal antibodies were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). p38 MAPK inhibitor, SB203580, and control 201474 were obtained from Calbiochem (San Diego, CA). MDA-MB-231 cell line was obtained from ATCC (Manassas, VA) and was maintained in Dulbecco's modified Eagle's medium (high glucose) containing 10% fetal calf serum at 37 °C in a humidified atmosphere of 5% CO<sub>2</sub>.

**Recombinant Adenovirus Construction**—To construct recombinant adenovirus (Ad) encoding dominant negative Ras and Rho GTPase, cDNAs for Myc-tagged H-Ras (N17), RhoA (N19), Rac1 (N17), and Cdc42 (N17) cDNAs were cloned into the adenovirus shuttle vector pAd.CI (containing cytomegalovirus promoter). Ad vector containing Myc-tagged wild-type and constitutively active Rac1 (V12) were constructed similarly. To construct Ad vector encoding dominant negative PRAK, MAPKAPK2, MAPKAPK3, MNK1, MSK1 and MSK2, cDNAs of these kinases containing an alanine mutation in lysine residue in the kinase domain were subcloned into pAd.RSV vector. Ad vectors were prepared by cotransfecting these vectors with pJM17 into 293 cells as described previously (47). Construction of the control Ad vector (Ad.RSV) and Ad vectors containing constitutively active or dominant negative MAPKs have been described elsewhere (47).

**Analyzing the Effect of  $\alpha$ v Integrin Ligation on the Activity of Rho GTPase**—Activated Rho GTPase can be detected by analyzing GTP-bound Rho GTPase. Rac1/Cdc42 and RhoA activities were determined by the recently developed PBD and RBD assays, respectively (48, 49). In Rac1/Cdc42 assays, the PBD from the effector protein p21-activated kinase is used as a probe to specifically isolate the active form of Rac1 or Cdc42 (48). In RhoA assays, the RBD from Rhotekin is used as probe to interact specifically with active RhoA (49). To determine the effect of  $\alpha$ v integrin ligation on the activities of Rac1, Cdc42, and RhoA, MDA-MB-231 cells were starved in serum-free medium for 24 h, and 3-cm culture dishes were coated with 2  $\mu$ g/ml of Vn solution at 4 °C. Cells were detached with Versene (Invitrogen), kept in suspension in serum-free medium for 1 h and then plated onto Vn-coated dishes. At varying times, cells were lysed in ice-cold lysis buffer (50 mM Tris, pH 7.5, 100 mM NaCl, 5 mM MgCl<sub>2</sub>, 1 mM EDTA, 0.1 mM EGTA, 1% Nonidet P-40, 5% glycerol, 5 mM NaF, 0.5 mM Na<sub>3</sub>VO<sub>4</sub>, 1 mM dithiothreitol, and protease inhibitor mixture). Cell lysates were incubated with GST-PBD or GST-RBD (20  $\mu$ g) beads at 4 °C for 1 h, and the beads were subsequently washed five times with lysis buffer. Bound Rac1, Cdc42, or RhoA were detected by immunoblotting using specific mAbs, respectively.

**Analyzing the Effect of  $\alpha$ v Integrin Ligation on the Activities of MAPKs**—To determine the effect of  $\alpha$ v integrin ligation on the activities of

MAPKs, MDA-MB-231 cells were starved for 24 h and kept in suspension for 1 h in serum-free medium. Cells were then plated on Vn-coated plates and at varying times lysed in ice-cold radioimmune precipitation assay buffer (phosphate-buffered saline containing 1% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS, 10 mM NaF, and protease inhibitor mixture). Cell lysates were boiled in non-reducing sample buffer and subjected to immunoblotting to detect total and active MKK3/6 or MKK4 with respective antibodies.

**Analysis of p38 MAPK Activity and uPA Expression**—To determine the effect of dominant negative Ras, Rho GTPase, and MAPKs on  $\alpha$ v integrin ligation-induced p38 MAPK activity, MDA-MB-231 cells were infected individually with dominant negative H-Ras, RhoA, Rac1, Cdc42, MEK1, MKK3, MKK4, or MKK6 Ad vectors (100 pfu/cell) for 24 h. After another 24 h of serum starvation, cells were plated on Vn-coated surface for 30 min and subsequently lysed for immunoblotting to detect active p38 MAPK with phospho-p38 polyclonal antibody.

To determine the effect of dominant negative p38 MAPK downstream kinases on Vn-up-regulated uPA expression, MDA-MB-231 cells were infected with dominant negative PRAK, MAPKAPK2, MAPKAPK3, MNK1, and MSK1 or MSK2 Ad vector (100 pfu/cell) for 24 h and then plated on Vn-coated dishes and cultured for 24 h. Cells were lysed, and cell lysates were subjected to immunoblotting to detect the levels of uPA. To determine whether constitutive active MAPKAPK2 was able to rescue SB203580-inhibited uPA expression, MDA-MB-231 cells were infected with Ad vector containing constitutive active PRAK (100 pfu/cell) or MAPKAPK2 (10 and 100 pfu/cell) for 24 h. Cells were then treated with SB203580 (5  $\mu$ M) for 24 h prior to cell lysis and immunoblotting to detect uPA.


**Analyzing the Effect of Constitutively Active Rac1 on MKK3 and MKK6 Activities**—MDA-MB-231 cells were infected with constitutively active or wild-type Rac1 Ad vector (100 pfu/cell) for 48 h and subsequently lysed in radioimmune precipitation assay buffer. Cell lysates were precleared with Gamma-bind beads (Amersham Biosciences) and then immunoprecipitated with either anti-MKK3 or MKK6 polyclonal antibody. The immunoprecipitates were boiled in sample buffer and subjected to immunoblotting with anti-phospho-MKK3/6 polyclonal antibody to detect the levels of active MKK3 or MKK6, respectively.

**Analyzing the Effect of Constitutively Active MKK3 on MAPKAPK2 Activity**—To determine the effect of constitutively active MKK3 on MAPKAPK2 activity, MDA-MB-231 cells were infected with constitutively active MKK3 Ad vector (100 pfu/cell) for 48 h and subsequently lysed for immunoblotting to detect active MAPKAPK2 with anti-phospho-MAPKAPK2 antibody.

To determine whether p38 MAPK is involved in MKK3-induced MAPKAPK2 activation, MDA-MB-231 cells were first infected with constitutively active MKK3 or MKK4 Ad vector (100 pfu/cell) for 24 h and followed by addition of SB203580 or SB202474 (5  $\mu$ M) for another 24 h. Cells were lysed, and the immunoblotting was performed to detect active MAPKAPK2.

**uPA mRNA Reporter Plasmids and mRNA Stability Analysis**—Plasmid pBBB was provided by Dr. Ann-Bin Shyu (University of Texas School of Medicine, Houston, TX). The entire and truncated forms of uPA 3'-UTR were synthesized by reverse transcriptase PCR using total RNA isolated from MDA-MB-231 cells. The PCR products were digested with BamHI or BamHI/BglII and subcloned into the unique BglII site of pBBB. A common 5'-primer, CTCTGAGGATCCCCAGGGAG-GAAACGGCA, corresponding to nucleotides 1367–1396 of human uPA mRNA sequence was used in the amplification of both PCR products. This 5'-primer contains a mutation at position 1375 (G  $\rightarrow$  A) to generate a BamHI site. The 3'-primers used for PCR are as follows. 1) For construct pBBB/3'-uPA (containing the entire 928-bp uPA 3'-UTR), a primer, ATCAGATCTATTTAAAAATTACAAAAATATAAAATAAAAAT-AG, corresponding to nucleotide 2282–2242 of human uPA mRNA sequence was synthesized, which contains mutations at positions 2278, 2277, and 2275 (C  $\rightarrow$  G, T  $\rightarrow$  A, and T  $\rightarrow$  C) to generate a BamHI site. 2) For construct pBBB/3'-uPA $\Delta$ ARE (pBBB containing ARE-deleted 3'-UTR of uPA mRNA), a primer, TAAGGATTCGTCCTCACCACCCAGTGAG, corresponding to nucleotides 2192–2222 was synthesized. This primer includes mutations at positions 2215 and 2213 (G  $\rightarrow$  T and G  $\rightarrow$  C) to generate a BglII site.

To determine the effect of p38 MAPK activity on uPA mRNA stability, MDA-MB-231 cells were cotransfected with pBBB and pBBB/3'-uPA for 24 h using LipofectAMINE 2000 (Invitrogen) in medium containing 10% fetal calf serum. SB203580 (10  $\mu$ M) or control compound SB202474 (10  $\mu$ M) was then added to cells for 1 h. Cells were switched to serum-free medium containing SB203580 or SB202474, and poly(A)<sup>+</sup> RNA was isolated at various time points (1–4 h). The levels of  $\beta$ -globin transcripts were determined by Northern blotting. To determine




**FIG. 1. Rac1 is involved in Vn-induced p38 MAPK activation.** *A*, MDA-MB-231 cells were starved overnight and plated on Vn-coated surfaces for various times. Cells were lysed, and the cell lysates were incubated with GST-PBD or GST-RBD beads at 4 °C. Rac1 and RhoA activities were determined by the amount of PBD-bound Rac1 and RBD-bound RhoA, respectively. The levels of total Rac1 and RhoA were determined by immunoblotting using cell lysates with the respective mAbs as described under “Experimental Procedures.” *B*, MDA-MB-231 cells were infected with dominant negative H-Ras, Rac1, Cdc42, or RhoA-containing Ad vector for 48 h. Cells were plated on Vn-coated surface for 30 min and subsequently lysed for immunoblotting to detect phosphorylated p38 MAPK. The membrane was stripped and first reprobbed for p38 MAPK to ensure equal protein loading and then reprobbed for Myc-tagged dominant negative H-Ras, Rac1, Cdc42, and RhoA expression.

whether ARE in the 3'-UTR of uPA mRNA affects p38 MAPK-regulated uPA mRNA stability, pBBB/3'-uPA $\Delta$ ARE vector was transfected into MDA-MB-231 cells in complete medium for 24 h. Cells were then treated with SB203580 (10  $\mu$ M) for 1 h and then switched to serum-free medium containing SB203580. poly(A)<sup>+</sup> RNA was isolated, and Northern blotting was performed to detect  $\beta$ -globin transcripts.

To determine the effect of constitutively active MAPKAPK2 and PRAK on uPA mRNA stability in p38 MAPK-inhibited condition, MDA-MB-231 cells were infected with control Ad or Ad vector containing constitutive active MAPKAPK2 or PRAK2 for 24 h and followed by transfection with pBBB/3'-uPA for another 24 h. SB203580 was added to cells for 1 h, and cells were then switched to serum-free medium containing SB203580 for 1–4 h followed by poly(A)<sup>+</sup> RNA extraction.

## RESULTS

**Rac1 Is Involved in  $\alpha$  Integrin-induced p38 MAPK Activation**—Our previous studies showed that Vn/ $\alpha$  integrin ligation induced p38 MAPK activation (44). Several recent studies have implicated that Rho GTPases including Rac, Cdc42, and RhoA are involved in integrin-mediated cellular responses (50–54) and that Rac1 and Cdc42 both activate p38 MAPK (55, 56). To determine whether Rho GTPases were involved in  $\alpha$  integrin-mediated p38 MAPK activation, we first examined the effect of Vn/ $\alpha$  integrin ligation on Rac1, Cdc42, and RhoA activities. MDA-MB-231 cells were starved overnight and then suspended in serum-free medium for 1 h. Cells were plated on Vn-coated surfaces for varying times (2 to 60 min) and subsequently lysed for PBD or RBD assay to determine Rac1/Cdc42 and RhoA GTPase activity, respectively. Rac1 was activated by Vn/ $\alpha$  integrin ligation as early as 2 min and remained active in the entire 60 min (Fig. 1A). The activity of Cdc42 was not detected even though Cdc42 protein expression was readily detectable in these cells (data not shown). Interestingly, the activity of Rho A decreased upon Vn/ $\alpha$  integrin ligation (Fig. 1A). These results suggest that Vn/ $\alpha$  integrin ligation may induce Rac1 activation preferentially.




**FIG. 2. Constitutively active MKK3 and MKK6 can up-regulate uPA expression.** *A*, MDA-MB-231 cells were infected with constitutively active MEK1-, MKK3-, MKK4-, or MKK6-containing Ad vector in serum-free condition for 48 h. Cells were lysed, and immunoblotting was performed to detect phosphorylated p38 MAPK and uPA. *B*, MDA-MB-231 cells were infected with constitutively active MKK3- or MKK6-containing Ad vector for 24 h. SB203580 (5  $\mu$ M) was then added to cells for 24 h and subsequently lysed for immunoblotting to detect cell-associated uPA expression. The membrane was stripped and reprobbed for actin to ensure equal protein loading.

We next examined the importance of Rac1 activity in Vn/ $\alpha$  integrin ligation-induced p38 MAPK activation. Dominant negative mutants of H-Ras or Rho GTPases (Rac1, Cdc42, and RhoA) were expressed in MDA-MB-231 cells using recombinant adenovirus for 48 h, and cells were then plated on Vn-coated surfaces for 30 min. Immunoblotting with an antibody specific for phospho-p38 MAPK showed that the expression of dominant negative Rac1 (N17), but not dominant negative Cdc42, RhoA, or H-Ras, inhibited Vn/ $\alpha$  integrin-induced p38 MAPK phosphorylation (Fig. 1B) significantly. These results suggest that Vn/ $\alpha$  integrin ligation may signal through Rac1 to activate p38 MAPK in invasive breast cancer cells.

**MKK3 Mediates Vn/ $\alpha$  Integrin Ligation-induced p38 MAPK Activation and uPA Up-regulation**—MKK3, MKK6, and MKK4 have been reported capable of activating p38 MAPK (28, 31). We thus determined whether these MAPKKs were able to increase uPA expression in invasive breast cancer cells. Constitutively active mutants of MEK1, MKK3, MKK4, and MKK6 were expressed in MDA-MB-231 cells using recombinant adenovirus for 48 h. Total cellular protein was isolated from the cells and analyzed for the levels of uPA expression. Immunoblotting analyses showed that both constitutively active MKK3 and MKK6 activated p38 MAPK and also up-regulated uPA expression significantly (Fig. 2A). In contrast, constitutive active MEK1 or MKK4 exhibited no stimulatory effect in p38 MAPK activity or uPA expression. To determine whether p38 MAPK is indeed required for constitutively active MKK3/6-induced uPA expression, SB203580 (a specific p38 MAPK inhibitor) was added to constitutively active MKK3 or MKK6-expressing MDA-MB-231 cells. SB203580 at 5  $\mu$ M concentration abrogated MKK3/6-induced uPA expression almost completely (Fig. 2B). These results suggest that both MKK3 and MKK6 can up-regulate uPA expression through p38 MAPK.


We next examined the effect of Vn/ $\alpha$  integrin ligation on MKK3/6 activity. Serum-starved MDA-MB-231 cells were plated on Vn-coated surface for varying times (5 min to 1 h), and immunoblotting was then performed to detect the active MKK3/6 and MKK4 (phosphorylated MAPKKs) with the relative antibodies. Vn/ $\alpha$  integrin ligation induced rapid increase in MKK3/6 phosphorylation; in contrast, the levels of MKK4



**FIG. 3. Vn-induced p38 MAPK activation and uPA up-regulation are signaled through MKK3.** *A*, MDA-MB-231 cells were plated on Vn-coated surface for various times. Cells were then lysed, and immunoblotting was performed to detect phosphorylated MKK4 or MKK3/6 (*pMKK4* or *pMKK3/6*). *B*, MDA-MB-231 cells were infected with dominant negative MEK1-, MKK3-, MKK4-, or MKK6-containing Ad vector for 48 h. Cells were then plated on Vn-coated surface for 30 min and followed by immunoblotting to detect phosphorylated p38 MAPK and cell-associated uPA.

phosphorylation were not altered significantly (Fig. 3A). In the subsequent experiment, we expressed dominant negative mutants of MEK1, MKK3, MKK4, or MKK6 in MDA-MB-231 cells and determined their effect on  $\alpha$ v integrin-mediated p38 MAPK activation and uPA up-regulation. Only the dominant negative MKK3 reduced  $\alpha$ v integrin-mediated p38 MAPK activation and uPA up-regulation (Fig. 3B). These results suggest that MKK3, rather than MKK6, mediates Vn/ $\alpha$ v integrin ligation-induced p38 MAPK activation and uPA expression in MDA-MB-231 cells although both MKK3 and MKK6 can up-regulate uPA expression.


**MAPKAPK2 Is Involved in p38 MAPK-regulated uPA Expression**—Up to six kinases (PRAK, MAPKAPK2, MAPKAPK3, MNK1, MSK1, and MSK2) are activated by p38 MAPK (28, 31) and can potentially serve as the p38 MAPK downstream effectors. To determine which of these molecules is involved in p38 MAPK-regulated uPA expression, dominant negative mutants of PRAK, MAPKAPK2, MAPKAPK3, MNK1, MSK1, or MSK2 were expressed in MDA-MB-231 cells using recombinant adenovirus. Cellular protein was extracted from these cells 48 h post-Ad infection and analyzed for the levels of uPA expression. Immunoblotting showed that a dominant negative MAPKAPK2 inhibited ~80% of Vn-up-regulated uPA expression in MDA-MB-231 cells (Fig. 4A). In contrast, only a slight reduction in the levels of uPA protein can be observed in cells expressing dominant negative PRAK, MAPKAPK3, MNK1, MSK1, or MSK2 in comparison with control (Fig. 4A). In a parallel study, we examined whether constitutively active MAPKAPK2 could rescue uPA expression in SB203580-treated (p38 MAPK-inhibited) MDA-MB-231 cells. Constitutively active MAPKAPK2 was expressed in cells using recombinant adenovirus for 24 h and followed by addition of SB203580 (5  $\mu$ M) to the cells for another 48 h. Immunoblotting using anti-uPA antibody showed that SB203580 inhibited uPA expression significantly (Fig. 4B). However, the expression of constitutively active MAPKAPK2 (at  $10^3$  pfu), but not constitutively active PRAK, restored at least 80% of SB203580-decreased uPA level (Fig. 4B). These results suggest that MAPKAPK2 is the main downstream effector of p38 MAPK for regulating uPA expression.



**FIG. 4. MAPKAPK2 is the main p38 MAPK downstream effector for uPA up-regulation.** *A*, MDA-MB-231 cells were infected with dominant negative PRAK-, MAPKAPK2-, MAPKAPK3-, MNK1-, MSK1-, or MSK2-containing Ad vector for 48 h. Cells were lysed, and immunoblotting was performed to detect cell-associated uPA. *B*, MDA-MB-231 cells were infected with constitutively active PRAK (1,000 pfu/cell) or MAPKAPK2 (200 and 1,000 pfu/cell) for 24 h. SB203580 (5  $\mu$ M) was added to cells for 24 h, and cells were then lysed for immunoblotting to detect cell-associated uPA. Membranes were stripped and re probed for actin to ensure equal protein loading.

**Rac1-MKK3-p38 MAPK-MAPKAPK Pathway Exists in Invasive Breast Cancer Cells**—Because Vn-mediated uPA up-regulation is blocked by dominant negative Rac1, MKK3, and MAPKAPK2 (see Figs. 1, 3, and 4), and SB203580-inhibited uPA expression is restored largely by constitutively active MAPKAPK2 (Fig. 4), it suggests strongly that a signaling pathway involving Rac1-MKK3-p38 MAPK-MAPKAPK2 is responsible for Vn-mediated uPA up-regulation in MDA-MB-231 cells. To confirm the presence of this signaling pathway, we first investigated the effect of active Rac1 on MKK3 and MKK6 activities. Wide-type or constitutively active Rac1 was expressed in MDA-MB-231 cells with the aid of adenovirus, and cells were then lysed and subjected immunoprecipitation with anti-MKK3 and -MKK6 antibodies. Immunoblotting on these immunoprecipitates with active MKK3/6 antibody showed that wide-type Rac1 did not alter the level of active MKK3 or MKK6 (Fig. 5A). In contrast, the levels of active MKK3, rather than MKK6, were elevated significantly by the constitutively active Rac1 expression (Fig. 5A). These results suggest that active Rac1 activates MKK3 specifically in MDA-MB-231 cells and are consistent with the results that only dominant negative MKK3 blocks Vn-mediated p38 MAPK activation and uPA up-regulation (Fig. 3).


To determine the effect of active MKK3 on MAPKAPK2 activity, constitutively active MKK3 or MKK4 was introduced into MDA-MB-231 cells using recombinant Ad vector for 2 days. Immunoblotting with phospho-MAPKAPK2-specific antibody showed that the levels of MAPKAPK2 phosphorylation in cells expressing constitutively active MKK4 was very similar to the control (Fig. 5B). In contrast, the expression of constitutively active MKK3 enhanced the levels of MAPKAPK2 phosphorylation significantly (Fig. 5B). To determine whether p38 MAPK participates in MKK3-induced MAPKAPK2 activation, MDA-MB-231 cells were infected with constitutively active MKK3 Ad vector for 24 h, and SB203580 or control compound 202474 (5  $\mu$ M) was then added to cells for another 24 h. Cells




**FIG. 5. Rac1-MKK3-p38 MAPK-MAPKAPK2 pathway is present in MDA-MB-231 cells.** A, MDA-MB-231 cells were infected with control, wild-type, or constitutively active Rac1 Ad vector for 48 h. Cells were lysed, and cell lysates were immunoprecipitated with MKK3 or MKK6 polyclonal antibody. The immunoblotting was then performed with these immunoprecipitates to detect active MKK3 or MKK6 (phospho-MKK3 or phospho-MKK6; *pMKK3* or *pMKK6*). Membranes were stripped and reprobed with MKK3 or MKK6 antibody to ensure equal loading. B, MDA-MB-231 cells were infected with constitutively active MKK3 or MKK4 for 24 h and then switched to medium in the absence or presence of SB203580 or SB202474 for another 24 h. Cells were lysed, and immunoblotting was performed to detect active MAPKAPK2 (phospho-MAPKAPK2; *pMAPKAPK2*) using phospho-MAPKAPK2-specific antibody. The membrane was stripped and reprobed with MAPKAPK2 polyclonal antibody to detect cellular MAPKAPK2 protein.

were lysed, and immunoblotting was performed to determine the levels of MAPKAPK2 phosphorylation. The treatment of SB203580, but not control SB202474, abrogated the constitutively active MKK3-induced MAPKAPK2 activation almost completely (Fig. 5B). These results suggest that p38 MAPK mediates MKK3-induced MAPKAPK2 activation.

**p38 MAPK-regulated uPA mRNA Stability Requires the ARE in 3'-UTR of uPA mRNA**—Highly conserved ARE have been found in 3'-UTR of uPA mRNAs and reported to be involved in p38 MAPK-regulated uPA mRNA stability (46, 57). To determine the importance of uPA 3'-UTR and its ARE in p38 MAPK-regulated uPA mRNA stability, we subcloned uPA 3'-UTR and ARE-deleted 3'-UTR in plasmid pBBB. The pBBB vector contains the rabbit  $\beta$ -globin cDNA under the control of the *c-fos* promoter. Because *c-fos* promoter is a serum-dependent promoter, the presence or absence of serum can be used to control promoter activity, and thus the stability of  $\beta$ -globin mRNA can be monitored without the addition of actinomycin D. Also, by inserting the sequence of interest in pBBB, the effect of this sequence on mRNA stability can be determined by monitoring otherwise stable  $\beta$ -globin mRNA (37, 38). The pBBB vector containing uPA 3'-UTR (pBBB/3'-uPA) was transfected into MDA-MB-231 cells, and the transfected cells were then treated with SB203580 or control compound SB202474 (5  $\mu$ M). poly(A)<sup>+</sup> RNA was isolated at various times (1–4 h), and Northern blotting was conducted to detect the levels of  $\beta$ -globin transcripts. Both  $\beta$ -globin and  $\beta$ -globin/uPA 3'-UTR RNA transcripts were equally stable in cells treated with control SB202474 (Fig. 6). In contrast, SB203580 destabilized  $\beta$ -globin/uPA 3'-UTR RNA transcript significantly whereas  $\beta$ -globin mRNA stability was not affected (Fig. 6). Interestingly, the pBBB vector containing ARE-deleted 3'-UTR (removal of the last 68 bp) (pBBB/3'-uPA $\Delta$ ARE) could no longer destabilize  $\beta$ -globin RNA transcript in SB203580-treated MDA-MB-231 cells (Fig. 6). These results suggested that p38 MAPK pathway



**FIG. 6. p38 MAPK-regulated uPA mRNA stability depends on the ARE in 3'-UTR.** MDA-MB-231 cells were either cotransfected with phospho-BBB and phospho-BBB/3'-uPA or transfected with phospho-BBB/3'-uPA $\Delta$ ARE. Cells were switched to serum-free medium, and poly(A)<sup>+</sup> RNA was isolated at varying times. Northern blotting was performed to detect  $\beta$ -globin transcripts.



**FIG. 7. MAPKAPK2 regulates uPA mRNA stability through an ARE-targeted mechanism.** A, MDA-MB-231 cells were infected with dominant negative PRAK or MAPKAPK2-containing Ad vector for 48 h. Cells were switched to serum-free medium, and poly(A)<sup>+</sup> RNA was isolated at varying times. Northern blotting was performed to detect  $\beta$ -globin transcripts. B, MDA-MB-231 cells were infected with constitutively active PRAK or MAPKAPK2 for 48 h, followed by 1 h of SB203580 (5  $\mu$ M) treatment. Cells were then switched to serum-free medium containing SB203580, and poly(A)<sup>+</sup> RNA was isolated at varying times. Northern blotting was performed to detect  $\beta$ -globin transcripts.

regulates uPA mRNA stability through an ARE-targeted mechanism.

**MAPKAPK2 Affects uPA mRNA Stability through uPA 3'-UTR**—We also investigated the effect of dominant negative MAPKAPK2 and PRAK on  $\beta$ -globin/uPA 3'-UTR transcript stability. MDA-MB-231 cells were transfected with pBBB containing uPA 3'-UTR and followed by a 2-day infection with control Ad vector or Ad vector encoding dominant negative MAPKAPK2 or PRAK. Although the stability of  $\beta$ -globin/uPA 3'-UTR transcript was not affected significantly by the expression of dominant negative PRAK, it was reduced greatly by the expression of dominant negative MAPKAPK2 (Fig. 7A). In further experiments, we examined the ability of constitutively active MAPKAPK2 to restore SB203580-induced uPA mRNA stability. Constitutively active MAPKAPK2 was introduced into  $\beta$ -globin/uPA 3'-UTR RNA transcript-expressing MDA-MB-231 cells prior to SB203580 treatment for various times. Subsequently, poly(A)<sup>+</sup> RNA was isolated, and Northern analysis was performed to detect  $\beta$ -globin transcripts. Although SB203580 reduced  $\beta$ -globin/uPA 3'-UTR RNA transcript stability greatly, the expression of constitutively active MAPKAPK2 prolonged  $\beta$ -globin/uPA 3'-UTR RNA transcript stability significantly in SB203580-treated MDA-MB-231 cells (Fig. 7B). In a parallel control experiment, we also determined the effect of constitutively active PRAK on  $\beta$ -globin/uPA 3'-UTR transcript stability in SB203580-treated cells and found that it was unable to prolong  $\beta$ -globin/uPA 3'-UTR transcript stability (Fig. 7B). These results suggest that MAPKAPK2 is the main p38 MAPK downstream effector for promoting uPA mRNA stability in invasive breast cancer cells.

#### DISCUSSION

Our previous studies have demonstrated that the elevated endogenous p38 MAPK is essential for uPA up-regulation in

invasive breast cancer (58). We have further shown that the elevated p38 MAPK activity requires  $\alpha$ v integrin expression/function, and  $\alpha$ v integrin ligation activates p38 MAPK and up-regulates uPA expression (44). In this report, we identified the upstream and downstream signaling molecules of p38 MAPK that participate in p38 MAPK-regulated uPA mRNA stability. Small GTPase Rac1 and Cdc42 are capable of activating p38 MAPK (55, 56) and are important for many integrin-mediated cellular responses (50–54). We thus investigated whether Rho GTPases including Rac1, Cdc42, and RhoA were involved in  $\alpha$ v integrin-mediated p38 MAPK activation. We found that Vn/ $\alpha$ v integrin ligation activated Rac1 preferentially (Fig. 1A). Further studies also showed that dominant negative Rac1, rather than dominant negative Cdc42 and RhoA, inhibited  $\alpha$ v integrin ligation-induced p38 MAPK activation significantly (Fig. 1B). These findings suggest strongly that  $\alpha$ v integrin signals through Rac1 for p38 MAPK activation. Several recent studies reported that  $\alpha$ 2 $\beta$ 1 integrin ligation can activate p38 MAPK in various cell types (59–61). Furthermore,  $\alpha$ 2 $\beta$ 1-induced p38 MAPK activation requires Cdc42 in human osteosarcoma cells (59) and Rac1 in murine NmuMG cells (60). Therefore, we consider the possibility that Rac1 or Cdc42 as a common signaling molecule for mediating integrin-induced p38 MAPK activation.

The specific upstream activator kinases for p38 MAPK are MKK3 and MKK6. MKK4, a known activator kinase for JNK, has also been implicated in p38 MAPK activation *in vitro* (27, 28). Although MKK3 and MKK6 are 80% homologous to each other and, in many cases, mediate same signals for p38 MAPK activation, they have been reported to exhibit differential involvement in other cellular events. For example, MKK6, rather than MKK3, is required for FasL and c-Abl-induced cell death in Jurkat T and NIH3T3 cells, respectively (47, 62). Also, MKK6 is important for IL-12-induced p38 MAPK activation and STAT4 activation in T and NK cells (63). However, MKK3, rather than MKK6, mediates tumor necrosis factor  $\alpha$  and lipopolysaccharide-induced p38 MAPK activation and cytokine expression in both fibroblast and macrophages (64, 65). We showed that dominant negative MKK3, but not dominant negative MKK6, was capable of inhibiting  $\alpha$ v integrin ligation-induced p38 MAPK activation and uPA up-regulation significantly (Fig. 3), suggesting that integrin-induced p38 MAPK activation may be mediated strictly by MKK3. Our study provides additional evidence for distinct involvement of MKK3 and MKK6 for p38 MAPK-mediated cellular responses.

p38 MAPK signaling pathway is involved in a wide spectrum of cellular functions, and the diversified role of p38 MAPK may be explained by the existence of various p38 MAPK downstream signaling molecules. Multiple kinases and transcription factors have been identified as p38 MAPK substrates and are potentially downstream effectors of p38 MAPK (28, 29). In fact, p38 MAPK/MSK-1 pathway is found to mediate vascular endothelial cell growth factor-induced cAMP-response element-binding protein phosphorylation and activation in human endothelial cells (66), p38 MAPK/MNK1 pathway is essential for stress-induced eukaryotic initiation factor-4E activation (67, 68), and phosphorylation of 5-lipoxygenase appears to require p38 MAPK/MAPKAPK2 and MAPKAP3 pathways (69). Our studies showed that dominant negative MAPKAPK2 inhibited  $\alpha$ v integrin ligation-induced uPA up-regulation specifically (Fig. 4A), and constitutively active MAPKAPK2 rescued uPA expression in p38 MAPK-inhibited cells (Fig. 4B). These results suggest strongly that MAPKAPK2 is the main p38 MAPK downstream effector in regulating uPA expression in invasive breast cancer cells. These studies also further support the

notion that the diversified role of p38 MAPK is determined by the variety of downstream signaling molecules.

In the past several years, growing evidence has revealed clearly that the ARE in 3'-UTR of mRNA can affect mRNA stability significantly (70, 71). The minimum consensus sequence for ARE is UUAUUUAU or UUAUU(A/U)(A/U) (36), and this sequence is present in the 3'-UTR of uPA mRNA (57). Early studies have shown that p38 MAPK pathway regulates uPA mRNA stability in an ARE-targeted mechanism (46), and this finding was further confirmed by our studies (Fig. 6). We demonstrated that the stability of uPA mRNA 3'-UTR-containing  $\beta$ -globin transcript was decreased specifically by dominant negative MAPKAPK2 in MDA-MB-231 cells by (Fig. 7A), and prolonged by, constitutively active MAPKAPK2 in p38 MAPK-inhibited MDA-MB-231 cells (Fig. 7B). These results suggest that MAPKAPK2 is the downstream effector of p38 MAPK for regulating uPA mRNA stability. Several recent studies have also demonstrated the importance of MAPKAPK2 in regulating mRNA stability. Blocking MAPKAPK2 activity was reported to inhibit cytokine/stress-induced IL-8 and IL-8 expression by destabilizing their mRNA (72). In MAPKAPK2-deficient cells, the half-life of IL-2 mRNA was reduced more than 10-fold, and re-expressing MAPKAPK2 in these cells restored IL-2 mRNA stability (73). Furthermore, tristetraprolin, a protein regulating mRNA stability, was found to be a direct substrate of MAPKAPK2 (32). Taken together, it is very likely that MAPKAPK2 may be a common p38 MAPK downstream signaling molecule for regulating ARE-mediated mRNA stability.

In conclusion, we have defined the signaling pathway linking  $\alpha$ v integrin ligation, p38 MAPK activation to uPA up-regulation in invasive breast MDA-MB-231 cells. Because uPA plays a significant role in tumor invasion and progression, the therapeutic approaches may be developed by intercepting the signaling pathway responsible for uPA overexpression.

**Acknowledgments**—We thank Dr. Ulla Knaus for providing the GST-PBD expression construct. We are also grateful to Joan Gausepohl and Kelly White for preparation of this manuscript.

#### REFERENCES

- Xing, R. H., and Rabbani, S. A. (1996) *Int. J. Cancer* **67**, 423–429
- Dubuisson, L., Monvoisin, A., Nielsen, B. S., Bail, B., Bioulac-Sage, P., and Rosenbaum, J. (2000) *J. Pathol.* **190**, 190–195
- Bouchet, C., Spyros, F., Hacène, K., Durcos, L., Bécette, V., and Oglobine, J. (1998) *Br. J. Cancer* **77**, 1495–1501
- Foekens, J. A., Peters, H. A., Look, M. P., Portengen, H., Schmitt, M., Kramer, M. D., Brünner, N., Jänicke, F., Meijer-van Gelder, M. E., Henzen-Logmans, S. C., van Putten, W. L. J., and Klijn, J. G. M. (2000) *Cancer Res.* **60**, 636–643
- Duggan, C., Maguire, T., McDermott, E., O'Higgins, N., Fennelly, J. J., and Duffy, M. J. (1995) *Int. J. Cancer* **61**, 597–600
- Hildenbrand, R., Leitz, M., Magdolen, V., Albrecht, S., Graeff, H., Stutte, H. J., Bleyl, U., and Schmitt, M. (2000) *Histopathology* **36**, 499–504
- Aguirre Ghiso, J. A., Alonso, D. F., Farias, E. F., Gomez, D. E., and de Kier Joffe, E. M. (1999) *Eur. J. Biochem.* **263**, 295–304
- Yu, W., Kim, J., and Ossowski, L. (1997) *J. Cell Biol.* **137**, 767–777
- Kim, J., Yu, W., Kovalski, K., and Ossowski, L. (1998) *Cell* **94**, 353–362
- Aguirre Ghiso, J. A., Kovalski, K., and Ossowski, L. (1999) *J. Cell Biol.* **147**, 89–103
- Farina, A. R., Coppa, A., Tiberio, A., Tacconelli, A., Turco, A., Colletta, G., Gulino, A., and Mackay, A. R. (1998) *Int. J. Cancer* **75**, 721–730
- Jeffers, M., Rong, S., and Vande Woude, G. F. (1996) *Mol. Cell. Biol.* **16**, 1115–1125
- Frandsen, T. L., Holst-Hansen, C., Nielsen, B. S., Christensen, I. J., Nyengaard, J. R., Carmeliet, P., and Brünner, N. (2001) *Cancer Res.* **61**, 532–537
- Gutierrez, L. S., Schulman, A., Brito-Robinson, T., Noria, F., Ploplis, V. A., and Castellino, F. J. (2000) *Cancer Res.* **60**, 5839–5847
- Bugge, T. H., Kombrinck, K. W., Xiao, Q., Holmbäck, K., Daugherty, C. C., Witte, D. P., and Degen, J. L. (1997) *Blood* **90**, 4522–4531
- Blasi, F. (1999) *Thromb. Haemostasis* **82**, 298–304
- Collen, C. (1999) *Thromb. Haemostasis* **82**, 259–270
- Koshelnick, Y., Ehart, M., Stockinger, H., and Binder, B. R. (1999) *Thromb. Haemostasis* **82**, 305–311
- Degryse, B., Resnati, M., Rabbani, S. A., Villa, A., Fazioli, F., and Blasi, F. (1999) *Blood* **94**, 649–662
- Koopman, J. L., Slomp, J., de Bart, A. C. W., and Quax, H. A. (1998) *J. Biol. Chem.* **273**, 33267–33272
- Preissner, K. T., Kanse, S. M., and May, A. E. (2000) *Curr. Opin. Cell Biol.* **12**, 621–628

22. Wei, Y., Yang, X., Liu, Q., Wilkins, J. A., and Chapman, H. A. (1999) *J. Cell Biol.* **144**, 1285–1294
23. Yebra, M., Goretzki, L., Pfeifer, M., and Mueller, B. M. (1999) *Exp. Cell Res.* **250**, 231–240
24. Ossowski, L., and Aguirre-Ghiso, J. A. (2001) *Curr. Opin. Cell Biol.* **12**, 613–620
25. Nguyen, D. H. D., Hussaini, I. M., and Gonias, S. L. (1998) *J. Biol. Chem.* **273**, 8502–8507
26. Yebra, M., Parry, G. C. N., Stromblad, S., Mackman, N., Rosenberg, S., Mueller, B. M., and Cheresch, D. A. (1996) *J. Biol. Chem.* **271**, 29393–29399
27. Nebreda, A. R., and Porras, A. (2000) *Trends Biochem. Sci.* **25**, 257–260
28. Ono, K., and Han, J. (2000) *Cell Signal.* **12**, 1–13
29. New, L., and Han, J. (1999) *Trends Cardiovasc Med.* **8**, 220–229
30. Martin-Blanco, E. (2000) *Bioessays* **22**, 637–645
31. Lee, J. C., Kumar, S., Griswold, D. E., Underwood, D. C., Votta, B. J., and Adams, J. L. (2000) *Immunopharmacology* **47**, 185–201
32. Mahtani, K. R., Brook, M., Dean, J. L. E., Sully, G., Saklatvala, J., and Clark, A. R. (2001) *Mol. Cell. Biol.* **21**, 6461–6469
33. Kontoyiannis, D., Kotlyarov, A., Carballo, E., Alexopoulou, L., Blakeshear, P. J., Gaestel, M., Davis, R., Flavell, R., and Kollias, G. (2001) *EMBO J.* **20**, 3760–3770
34. Vasudevan, S., and Peltz, S. W. (2001) *Mol. Cell* **7**, 1191–1200
35. Mitchell, P., and Tollervey, D. (2001) *Curr. Opin. Genet. Dev.* **10**, 193–198
36. Chen, C.-Y. A., and Shyu, A.-B. (1995) *Trends Biochem. Sci.* **20**, 465–470
37. Loflin, P. T., Chen, C.-Y. A., Xu, N., and Shyu, A.-B. (1999) *Methods* **17**, 11–20
38. Xu, N., Loflin, P., Chen, C.-Y. A., and Shyu, A.-B. (1998) *Nucleic Acids Res.* **26**, 558–565
39. Zhang, W., Wagner, B. J., Ehrenman, K., Schaefer, A. W., DeMaria, C. T., Crater, D., DeHaven, K., Long, L., and Brewer, G. (1993) *Mol. Cell. Biol.* **13**, 7652–7665
40. Buzby, J. S., Brewer, G., and Nugent, D. J. (1999) *J. Biol. Chem.* **274**, 33973–33978
41. Ma, W.-J., Cheng, S., Campbell, C., Wright, A., and Furneaux, H. (1996) *J. Biol. Chem.* **271**, 8144–8151
42. Peng, S. S. Y., and Chen, C.-Y. A. (1998) *EMBO J.* **17**, 3461–3470
43. Carballo, E., Lai, W. S., and Blakeshear, P. J. (1998) *Science* **281**, 1001–1005
44. Chen, J., Baskerville, C., Han, Q., Pan, Z., and Huang, S. (2001) *J. Biol. Chem.* **276**, 47901–47905
45. Huang, S., New, L., Pan, Z., Han, J., and Nemerow, G. R. (2000) *J. Biol. Chem.* **275**, 12266–12272
46. Montero, L., and Nagamine, Y. (1999) *Cancer Res.* **59**, 5286–5293
47. Huang, S., Jiang, Y., Li, Z., Nishida, E., Mathias, P., Lin, S., Ulevitch, R. J., Nemerow, G. R., and Han, J. (1997) *Immunity* **6**, 739–749
48. Benard, V., Bohl, B. P., and Bokoch, G. M. (1999) *J. Biol. Chem.* **274**, 13198–13204
49. Ren, X., Kiosses, W. B., and Schwatz, M. A. (1999) *EMBO J.* **18**, 578–585
50. Price, L. S., Leng, J., Schwartz, M. A., and Bokoch, G. M. (1998) *Mol. Biol. Cell* **9**, 1863–1871
51. Mainiero, F., Soriani, A., Strippoli, R., Jacobelli, J., Gismondi, A., Piccoli, M., Frati, L., and Santoni, A. (2000) *Immunity* **12**, 7–16
52. Bourdoulous, S., Orend, G., MacKenna, D. A., Pasqualini, R., and Ruoslahti, E. (1998) *J. Cell Biol.* **143**, 267–276
53. Bialkowska, K., Kulkarni, S., Du, X., Goll, D. E., Saido, T. C., and Fox, J. E. B. (2000) *J. Cell Biol.* **151**, 685–695
54. Clark, E. A., King, W. G., Brugge, J. S., Symons, M., and Hynes, R. O. (1998) *J. Cell Biol.* **142**, 573–586
55. Philips, A., Roux, P., Coulon, V., Bellanger, J.-M., Vié, A., Vignais, M.-L., and Blanchard, J. M. (2000) *J. Biol. Chem.* **275**, 5911–5917
56. Molnár, A., Theodoras, A. M., Zon, L. I., and Kyriakis, J. M. (1997) *J. Biol. Chem.* **272**, 13229–13235
57. Nanbu, R., Montero, L., D'Orazio, D., and Nagamine, Y. (1997) *Eur. J. Biochem.* **247**, 169–174
58. Chan, S. Y., Empig, C. J., Welte, F. J., Speck, R. F., Schmaljohn, A., Kreisberg, J. F., and Goldsmith, M. A. (2001) *Cell* **106**, 117–126
59. Ivaska, J., Reunanen, H., Westermarck, J., Koivisto, L., Kähari, V.-M., and Heino, J. (1999) *J. Cell Biol.* **147**, 401–415
60. Klekotka, P. A., Santora, S. A., and Zutter, M. M. (2001) *J. Biol. Chem.* **276**, 9503–9511
61. Xu, J., Clark, R. A., and Parks, W. C. (2001) *Biochem. J.* **355**, 437–447
62. Cong, F., and Goff, S. P. (1999) *Proc. Natl. Acad. Sci. U. S. A.* **96**, 13819–13824
63. Visconti, R., Gadina, M., Chiariello, M., Chen, E. H., Stancato, L. F., Gutkind, J. S., and O'Shea, J. J. (2000) *Blood* **96**, 1844–1852
64. Wysk, M., Yang, D. D., Lu, H.-T., Flavell, R. A., and Davis, R. J. (1999) *Proc. Natl. Acad. Sci. U. S. A.* **96**, 3763–3768
65. Lu, H.-T., Yang, D. D., Wysk, M., Gatti, E., Mellman, I., Davis, R. J., and Flavell, R. A. (1999) *EMBO J.* **18**, 1845–1857
66. Mayo, L. D., Kessler, K. M., Pincheira, R., and Warren, R. S. (2001) *J. Biol. Chem.* **276**, 25184–25189
67. Knauf, U., Tschopp, C., and Gram, H. (2001) *Mol. Cell. Biol.* **21**, 5500–5511
68. Waskiewicz, A. J., Flynn, A., Proud, C. G., and Cooper, J. A. (1997) *EMBO J.* **16**, 1909–1920
69. Werz, O., Klemm, J., Samuelsson, B., and Rådmark, O. (2000) *Proc. Natl. Acad. Sci. U. S. A.* **97**, 5261–5266
70. Dreyfuss, G., Kim, V. N., and Kataoka, N. (2002) *Nat. Rev. Mol. Cell. Biol.* **3**, 195–205
71. Guhaniyogi, J., and Brewer, G. (2001) *Gene* **265**, 11–23
72. Winzen, R., Kracht, M., Ritter, B., Wilhelm, A., Chen, C.-Y. A., Shyu, A.-B., Müller, M., Gaestel, M., Resch, K., and Holtmann, H. (1999) *EMBO J.* **18**, 4969–4980
73. Neininger, A., Kontoyiannis, D., Kotlyarov, A., Winzen, R., Eckert, R., Volk, H.-D., Holtmann, H., Kollian, G., and Gaestel, M. (2002) *J. Biol. Chem.* **277**, 3065–3068