

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/227272895>

Limit points of eigenvalues of (di)graphs

Article in Czechoslovak Mathematical Journal · September 2006

DOI: 10.1007/s10587-006-0064-y

CITATIONS

11

READS

29

2 authors:



[Fuji Zhang](#)

Xiamen University

147 PUBLICATIONS 1,650 CITATIONS

[SEE PROFILE](#)



[Zhibo Chen](#)

Pennsylvania State University

45 PUBLICATIONS 272 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Applications of combinatorics [View project](#)

All content following this page was uploaded by [Fuji Zhang](#) on 23 June 2017.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.

Fu Ji Zhang; Zhibo Chen

Limit points of eigenvalues of (di)graphs

Czechoslovak Mathematical Journal, Vol. 56 (2006), No. 3, 895--902

Persistent URL: <http://dml.cz/dmlcz/128114>

Terms of use:

© Institute of Mathematics AS CR, 2006

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

LIMIT POINTS OF EIGENVALUES OF (DI)GRAPHS

FUJI ZHANG, Xiamen, ZHIBO CHEN, McKeesport

(Received March 31, 2004)

Abstract. The study on limit points of eigenvalues of undirected graphs was initiated by A. J. Hoffman in 1972. Now we extend the study to digraphs. We prove:

1. Every real number is a limit point of eigenvalues of graphs. Every complex number is a limit point of eigenvalues of digraphs.
2. For a digraph D , the set of limit points of eigenvalues of iterated subdivision digraphs of D is the unit circle in the complex plane if and only if D has a directed cycle.
3. Every limit point of eigenvalues of a set \mathcal{D} of digraphs (graphs) is a limit point of eigenvalues of a set $\tilde{\mathcal{D}}$ of bipartite digraphs (graphs), where $\tilde{\mathcal{D}}$ consists of the double covers of the members in \mathcal{D} .
4. Every limit point of eigenvalues of a set \mathcal{D} of digraphs is a limit point of eigenvalues of line digraphs of the digraphs in \mathcal{D} .
5. If M is a limit point of the largest eigenvalues of graphs, then $-M$ is a limit point of the smallest eigenvalues of graphs.

Keywords: limit point, eigenvalue of digraph (graph), double cover, subdivision digraph, line digraph

MSC 2000: 05C50, 15A48

1. INTRODUCTION

Since A. J. Hoffman [9] initiated the study of limit points of graph eigenvalues in 1972, many interesting results have been obtained on this topic (see, for example, [2]–[7], [9], [10], [15]). Hoffman [9] studied limit points of the largest eigenvalues of graphs, and he determined all limit points less than $\sqrt{2 + \sqrt{5}}$ (the golden mean) and showed that these limit points constitute an increasing sequence (a_n) with $a_1 = 2$ and $\lim_{n \rightarrow \infty} a_n = \sqrt{2 + \sqrt{5}}$. He also suggested that possibly there exists a real number λ such that every number not less than λ is a limit point of the largest eigenvalues

The first author was supported by NSFC. The second author was supported in part by the RDG grant of the Penn State University.

of graphs. In fact this is true with $\lambda = \sqrt{2 + \sqrt{5}}$, as proved by Shearer [15]. From these works the set of limit points of the largest eigenvalues of graphs has been determined. For $k \geq 2$, the limit points of the k th largest eigenvalues of graphs have been studied for a long time, but they have not been completely determined. Dasong Cao and Hong Yuan [2], [3] showed that for each $k \geq 2$ there is a gap in the set L_k of limit points of the k th largest eigenvalues of graphs, and that $L_k \subseteq L_{k+1}$ for all k . They conjectured in [3] that $\lim_{k \rightarrow \infty} L_k = R$, the set of all real numbers.

In this paper we will first prove that every real number is a limit point of eigenvalues of graphs, and then extend the study to the limit points of eigenvalues of digraphs.

We follow [1] for general graph theoretical terminology. All digraphs in this paper are finite without loops or multiple arcs. Undirected simple graphs are simply called graphs. Let D be a digraph (graph) with adjacency matrix A . The characteristic polynomial of D is $\chi(D, x) = |xI - A|$, where I denotes the identity matrix. The roots (complex numbers) of $\chi(D, x)$ are called the eigenvalues of D . It is well known that all eigenvalues of D must be real numbers when D is a graph.

To state our main results, we need the following

Definition 1. Let \mathcal{D} denote an infinite set of digraphs (graphs). The complex number ζ is said to be a *limit point of eigenvalues of \mathcal{D}* if there is an infinite sequence of distinct complex numbers λ_n , each of which is an eigenvalue of a digraph (graph) in \mathcal{D} , such that $\zeta = \lim_{n \rightarrow \infty} \lambda_n$.

Remarks.

- (i) When \mathcal{D} is the set of all digraphs (graphs), we simply say that ζ is a limit point of digraph (graph) eigenvalues.
- (ii) Every limit point of graph eigenvalues is a limit point of digraph eigenvalues.

Remark (ii) can be justified as follows. For any graph G there corresponds a unique digraph $D(G)$, called its associated digraph, which is obtained when each edge e of G is replaced by two oppositely oriented arcs with the same ends as e . It is obvious that a graph G and its associated digraph $D(G)$ have the same adjacency matrix, and so they have the same eigenvalues. Hence, when we consider eigenvalues, a graph can be seen as a digraph with a symmetric adjacency matrix.

We say that a digraph is bipartite if its underlying graph is bipartite. Note that a bipartite graph and a bipartite digraph both have an adjacency matrix in the block form $\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix}$, where 0 denotes the block with all entries zero. While for bipartite graphs B must equal to the transpose A^T of A , there is no such restriction for bipartite digraphs.

For an n by n matrix A , let \ddot{A} denote the $2n$ by $2n$ matrix $\begin{bmatrix} 0 & A \\ A & 0 \end{bmatrix}$.

Definition 2. Let D be a digraph (graph) with adjacency matrix A . The digraph (graph) with the adjacency matrix \ddot{A} is called the *double cover of D* , denoted by \ddot{D} .

Remarks.

- (i) \ddot{D} is a graph when D is a graph, since \ddot{A} is symmetric when A is symmetric.
- (ii) \ddot{D} is a bipartite digraph (graph) when D is a digraph (graph).

The concept of line digraphs was introduced in Harary and Norman [8]. For a digraph D with vertex set $V(D)$ and arc set $A(D)$, the *line digraph $L(D)$* of D has $A(D)$ as its vertex set; (a, b) is an arc of $L(D)$ if and only if there are vertices u, v, w in D with $a = (u, v)$ and $b = (v, w)$, i.e., the head of a coincides with the tail of b . The subdivision digraph $S(D)$ of D is obtained by inserting a new vertex onto every arc of D ; that is, each arc (u, v) of D is replaced by two new arcs (u, w) and (w, v) where w is a new vertex. The iterated subdivision digraphs are defined inductively by $S^r(D) = S(S^{r-1}(D))$ with $S^0(D) = D$ and $S^1(D) = S(D)$.

We will prove:

1. Every real number is a limit point of eigenvalues of graphs. Every complex number is a limit point of eigenvalues of digraphs.
2. For a digraph D , the set of limit points of eigenvalues of iterated subdivision digraphs of D is the unit circle in the complex plane if and only if D has a directed cycle.
3. Every limit point of eigenvalues of a set \mathcal{D} of digraphs (graphs) is a limit point of eigenvalues of a set $\ddot{\mathcal{D}}$ of bipartite digraphs (graphs), where $\ddot{\mathcal{D}}$ consists of the double covers of the members in \mathcal{D} .
4. Every limit point of eigenvalues of a set \mathcal{D} of digraphs is a limit point of eigenvalues of line digraphs of the digraphs in \mathcal{D} .
5. If M is a limit point of the largest eigenvalues of graphs, then $-M$ is a limit point of the smallest eigenvalues of graphs.

2. LEMMAS

Lemma 1. *Let D be a digraph or graph. If the eigenvalues of D are λ_i ($i = 1, \dots, n$), then the eigenvalues of the double cover \ddot{D} of D are $\pm\lambda_i$ ($i = 1, \dots, n$).*

Proof. Let A be the adjacency matrix of D . Then

$$\chi(\ddot{D}, x) = |xI - \ddot{A}| = \begin{vmatrix} xI & -A \\ -A & xI \end{vmatrix}.$$

It is well known in matrix theory (see, for example, [11, p. 45]) that if M is an invertible matrix then $\begin{vmatrix} M & N \\ P & Q \end{vmatrix} = |M| \cdot |Q - PM^{-1}N|$. So,

$$\begin{aligned} \chi(\ddot{D}, x) &= x^n \left| xI - (-A) \left(\frac{1}{x} I \right) (-A) \right| \\ &= x^n \left| xI - \frac{1}{x} A^2 \right| = |x^2 I - A^2| = |xI - A| \cdot |xI + A| \\ &= (-1)^n |xI - A| \cdot |-xI - A| = (-1)^n \chi(D, x) \chi(D, -x). \end{aligned}$$

This proves Lemma 1. □

Let D_1 and D_2 be digraphs (graphs) with matrices A and B , respectively. The Kronecker product $D_1 \otimes D_2$ is the digraph (graph) with adjacency matrix $A \otimes B$, the Kronecker product of the matrix A with the matrix B . From matrix theory (see, e.g., [14, p. 24]) we know that the set of the eigenvalues of a Kronecker product $A \otimes B$ of square matrices A and B is the same as the set of all possible products $\lambda_A \lambda_B$ where λ_A and λ_B are eigenvalues of A and B , respectively. Since the Kronecker product is associative, we have the following Lemma 2.

Lemma 2. *Let D_i denote digraphs (graphs), $i = 1, 2, \dots, n$. The set of eigenvalues of $D_1 \otimes D_2 \otimes \dots \otimes D_n$ is the same as the set of all possible products $\lambda_1 \lambda_2 \dots \lambda_n$, where λ_i is an eigenvalue of D_i .*

Lemma 3. *If a real number r is a limit point of graph eigenvalues, then*

- i) $-r$ is also a limit point of graph eigenvalues;
- ii) every point on the circle with radius $|r|$ centered at the origin in the complex plane is a limit point of digraph eigenvalues.

Proof. i) If λ is an eigenvalue of a graph D , then by Lemma 1, $-\lambda$ is an eigenvalue of \ddot{D} . This implies i) directly.

ii) From Remark (ii) following Definition 1, r is also a limit point of digraph eigenvalues. Let $r = \lim_{n \rightarrow \infty} \lambda_n$, where λ_n are distinct and each λ_n is an eigenvalue of a digraph G_n . It is well known (see [4, p. 53]) that the eigenvalues of a directed cycle D_n with n vertices are $\exp(i2k\pi/n)$ ($k = 0, 1, \dots, n-1$; $i = \sqrt{-1}$), the n th complex roots of unit. Note that these n complex roots are evenly distributed in the unit circle on the complex plane.

Now, let \mathcal{D} denote the set of all digraphs $G_n \otimes D_n$. By Lemma 2, $\lambda_n \exp(i2k\pi/n)$ ($k = 0, 1, \dots, n-1$) are eigenvalues of $G_n \otimes D_n$. Note that these n complex numbers are evenly distributed on the circle with radius $|\lambda_n|$ centered at the origin in the complex plane, and that $\lim_{n \rightarrow \infty} |\lambda_n| = |r|$. Then it is easily seen that every point on

the circle with radius $|r|$ centered at the origin in the complex plane is a limit point of eigenvalues of \mathcal{D} , and so it is a limit point of digraph eigenvalues. This completes the proof for Lemma 3. \square

Lemma 4. *A digraph D has a nonzero eigenvalue if and only if D has a directed cycle.*

Proof. From the ‘‘Coefficients Theorem for Digraphs’’ (see [4, p. 32]), we know that for a digraph D with n vertices,

$$\chi(D, x) = x^n + a_1x^{n-1} + \dots + a_n \quad \text{with } a_i = \sum_{L \in \mathcal{L}_i} (-1)^{p(L)} \quad (i = 1, 2, \dots, n)$$

where \mathcal{L}_i is the set of all such subdigraphs L of D with exactly i vertices that each component of L is a directed cycle; $p(L)$ denotes the number of components of L .

So, the necessity of the lemma follows immediately. To show the sufficiency, since D has a directed cycle, we consider the directed cycles of D with the shortest length ϱ . Then we see that in $\chi(D, x) = x^n + a_1x^{n-1} + \dots + a_n$, $a_\varrho \neq 0$, and this proves the sufficiency. \square

Lemma 5 (Lin and Zhang [12]; also see [13, p. 19]). *Let D be a digraph with n vertices and m arcs. Then $\chi(L(D), x) = x^{m-n}\chi(D, x)$.*

Lemma 6 (Zhang, Lin, and Meng [16]). *Let D be a digraph with n vertices and m arcs. Then $\chi(S(D), x) = x^{m-n}\chi(D, x^2)$.*

3. MAIN RESULTS

Theorem 1.

- i) *Every real number r is a limit point of graph eigenvalues.*
- ii) *Every complex number ζ is a limit point of digraph eigenvalues.*

Proof. i) It is well known that a cycle C_n with n vertices has eigenvalues $2\cos(2k\pi/n)$, $k = 1, 2, \dots, n$. So, it is obvious that $r = 0$ is a limit point of graph eigenvalues. Then, by Lemma 3 (i), we may assume $r > 0$ without loss of generality. Note that C_{4n+1} has a positive eigenvalue $\lambda_n = 2\cos(2n\pi/(4n+1))$. Obviously $\lambda_n \rightarrow 0$ as $n \rightarrow \infty$. So, for any $0 < \varepsilon < \frac{1}{2}r$, we may take an n such that $\lambda_n < \frac{1}{2}\varepsilon$. Let $N = \lfloor (r - \frac{1}{2}\varepsilon)/\lambda_n \rfloor$. Then $N \leq (r - \frac{1}{2}\varepsilon)/\lambda_n < N+1$. It follows that $N\lambda_n \leq r - \frac{1}{2}\varepsilon < r$ and that $N\lambda_n > r - \frac{1}{2}\varepsilon - \lambda_n > r - \varepsilon$. Thus we have $r - \varepsilon < N\lambda_n < r$. Now let G_ε denote the product graph of N copies of C_{4n+1} . It is well known (see, e.g., [4])

that the eigenvalues of the product graph $\Gamma_1 \times \Gamma_2$ of two graphs Γ_1 and Γ_2 are the sums of eigenvalues of Γ_1 with those of Γ_2 . (Note that $\Gamma_1 \times \Gamma_2$ is denoted differently in [4] as $\Gamma_1 + \Gamma_2$.) Hence, $N\lambda_n$ are eigenvalues of G_ε . Therefore, for any $\varepsilon > 0$ there is a graph G_ε with an eigenvalue in the open interval $(r - \varepsilon, r)$. It clearly implies i).

ii) directly follows from i) and Lemma 3 (ii). □

It should be pointed out that the graphs and digraphs in Theorem 1 can be restricted as regular graphs and regular digraphs, which can be easily seen from the proof of Theorem 1.

Theorem 2. *For a digraph D , the set of limit points of eigenvalues of iterated subdivision digraphs of D is the unit circle in the complex plane if and only if D has a directed cycle.*

Proof. If D has no directed cycles, then all the iterated subdivision digraphs $S^r(D)$ have no directed cycles. Then by Lemma 4, all eigenvalues of $S^r(D)$ are zero so that no limit points exist.

Let D be a digraph with a directed cycle. Then by Lemma 4, D has nonzero eigenvalues. Let α be any one of the nonzero eigenvalues. By Lemma 6, the subdivision digraph $S(D)$ has the two square roots of α among its eigenvalues. Clearly, for any positive integer r , the iterated subdivision digraph $S^r(D)$ has all the 2^r th roots of α among its eigenvalues. Note that in the complex plane the 2^r th roots of α are evenly distributed on the circle centered at the origin with the radius equal to the 2^r th root of the modulus of α . Since $\lim_{r \rightarrow \infty} 2^r = \infty$, $\lim_{r \rightarrow \infty} \sqrt[2^r]{|\alpha|} = 1$, and every nonzero eigenvalue of $S^r(D)$ is a 2^r th root of a nonzero eigenvalue of D , we see that the set of the nonzero eigenvalues of $S^r(D)$ approaches the unit circle in the complex plane as $r \rightarrow \infty$. Hence Theorem 2 follows. □

When we consider the limit points of eigenvalues of a set \mathcal{D} of digraphs (graphs), the next two theorems are helpful.

Theorem 3. *Let \mathcal{D} be an infinite set of digraphs (graphs), and let $\ddot{\mathcal{D}}$ denote the set of the double covers of digraphs (graphs) in \mathcal{D} . Every limit point of eigenvalues of \mathcal{D} is a limit point of eigenvalues of $\ddot{\mathcal{D}}$.*

Proof. It follows from Lemma 1. □

From Theorems 1 and 3, we immediately have

Corollary 1. *Every real number is a limit point of eigenvalues of bipartite graphs. Every complex number is a limit point of eigenvalues of bipartite digraphs.*

For an infinite set \mathcal{D} of digraphs, let $\vec{\mathcal{D}}$ denote the set of all line digraphs $L(D)$ with D in \mathcal{D} . We have

Theorem 4. *Every limit point of eigenvalues of \mathcal{D} is a limit point of eigenvalues of $\vec{\mathcal{D}}$.*

Proof. It follows from Lemma 5. □

From Theorems 1 and 4, we immediately have

Corollary 2. *Every complex number is a limit point of eigenvalues of line digraphs.*

Note that if λ is the largest eigenvalue of a graph G , then by Lemma 1, $-\lambda$ is the smallest eigenvalue of the double cover \check{G} of G . This implies

Theorem 5. *If M is a limit point of the largest eigenvalues of graphs, then $-M$ is a limit point of the smallest eigenvalues of graphs.*

By Theorem 5 we can obtain the following result of [6].

Corollary 3. *Every number in the interval $(-\infty, -\sqrt{2 + \sqrt{5}}]$ is a limit point of the smallest eigenvalues of graphs.*

Proof. It was initiated by Hoffman [9] and proved by Shearer [15] that every number not less than $\sqrt{2 + \sqrt{5}}$ (the golden mean) is a limit point of the largest eigenvalues of graphs. Then Corollary 3 immediately follows from Theorem 5. □

References

- [1] *J. A. Bondy, U. S. R. Murty:* Graph Theory with Applications. Macmillan, London, 1976.
- [2] *D. Cao D, H. Yuan:* Graphs characterized by the second eigenvalue. *J. Graph Theory* 17 (1993), 325–331. Zbl 0786.05055
- [3] *D. Cao, H. Yuan:* The distribution of eigenvalues of graphs. *Linear Algebra Appl.* 216 (1995), 211–224. Zbl 0820.05047
- [4] *D. M. Cvetković, M. Doob, and H. Sachs:* Spectra of Graphs. Academic Press, New York, 1980, third edition Johann Ambrosius Barth Verlag 1995. Zbl 0824.05046
- [5] *D. M. Cvetković, P. Rowlinson:* The largest eigenvalue of a graph: a survey. *Linear and multilinear Algebra* 28 (1990), 3–33. Zbl 0744.05031
- [6] *M. Doob:* The limit points of eigenvalues of graphs. *Linear Algebra Appl.* 114/115 (1989), 659–662. Zbl 0671.05050

- [7] *M. Doob*: Some new results on the limit points of eigenvalues of graphs. *Abstracts Amer. Math. Soc.* 12 (1991), 450.
- [8] *F. Harary, R. Z. Norman*: Some properties of line digraphs. *Rend. Circ. Mat. Palermo* 9 (1960), 161–168.
- [9] *A. J. Hoffman*: On limit points of spectral radii of non-negative symmetric integral matrices. *Graph Theory and its Application. Lecture Notes in Math.* 303 (Y. Alavi et al., eds.). Springer-Verlag, New York, 1972, pp. 165–172.
- [10] *A. J. Hoffman*: On limit points of the least eigenvalue of a graph. *Ars Combinatoria* 3 (1977), 3–14. [Zbl 0445.05067](#)
- [11] *F. R. Gantmacher*: *Theory of Matrices*, Vol. I. AMS Chelsea Publishing, Providence, 1960.
- [12] *G. Lin, F. Zhang*: On the characteristic polynomial of directed line graph and a type of cospectral directed digraph. *KeXue TongBao (Chinese Sci. Bulletin)* 22 (1983), 1348–1350.
- [13] *B. Liu, H. Lai*: *Matrices in Combinatorics and Graph Theory*. Kluwer, Dordrecht, 2000. [Zbl 0958.05003](#)
- [14] *M. Marcus, H. Minc*: *A Survey of Matrix Theory and Matrix Inequalities*. Allyn and Bacon, Boston, 1964. [Zbl 0247.15002](#)
- [15] *J. B. Shearer*: On the distribution of the maximum eigenvalue of graphs. *Linear Algebra and Appl.* 114/115 (1989), 17–20. [Zbl 0672.05059](#)
- [16] *F. Zhang, G. Lin, and J. Meng*: The characteristic polynomials of digraphs formed by some unary operations. *J. Xinjiang Univ.* 4 (1987), 1–6.

Authors' addresses: F u j i Z h a n g, Department of Mathematics, Xiamen University, Xiamen 361005, P.R. China; Z h i b o C h e n, Department of Mathematics, Penn State University, McKeesport, PA 15132, USA.