

**Metallacycles**

International Edition: DOI: 10.1002/anie.201502412

German Edition: DOI: 10.1002/ange.201502412

**Five-Membered Cyclic Metal Carbyne: Synthesis of Osmapentalynes by the Reactions of Osmapentalene with Allene, Alkyne, and Alkene\*\***

Congqing Zhu, Yuhui Yang, Jingjing Wu, Ming Luo, Jinglan Fan, Jun Zhu,\* and Haiping Xia\*\*


**Abstract:** The synthesis of small cyclic metal carbynes is challenging due to the large angle strain associated with the highly distorted nonlinear triple bonds. Herein, we report a general route for the synthesis of five-membered cyclic metal carbyne complexes, osmapentalynes, by the reactions of an osmapentalene derivative with allene, alkyne, and alkene. Experimental observations and theoretical calculations document the aromaticity in the fused five-membered rings of osmapentalynes. The realization of transforming osmapentalene to osmapentalyne through this general route would not only allow further exploration of metallapentalyne chemistry but also show promising applications of this novel aromatic system with broad absorption band and high molar absorption coefficient.

Small cyclic alkynes and metal carbynes have attracted many interests from chemists for their highly strained structures with the nonlinear triple bonds.<sup>[1,2]</sup> Not surprisingly, the first example of a cyclic metal carbyne complex, metalabenzynes (six-membered ring), has not been reported until 2001.<sup>[2a]</sup> In principle, the realization of smaller rings requires more effort to overcome larger ring strain. On the other hand, compared with Hückel aromatic compounds containing  $4n + 2$   $\pi$  electrons, Möbius aromatic ones with  $4n$   $\pi$  electrons have proven much more difficult to synthesize and identify.<sup>[3,4]</sup> So far, the reported Möbius aromatic compounds are limited and most of them have a twisted topology.<sup>[5]</sup>

Recently, we reported the first five-membered cyclic metal carbyne, metallapentalyne.<sup>[6]</sup> Density functional theory (DFT) calculations not only confirmed the larger ring strain (24.3 kcal mol<sup>-1</sup> in osmapentalyne versus 9.6 kcal mol<sup>-1</sup> in osmabenzynes) but also revealed planar Möbius aromaticity in osmapentalyne.<sup>[6,7]</sup> Further study demonstrated that these unique aromatics show aggregation-enhanced near-infrared

photoluminescence<sup>[6]</sup> and the carbyne carbon can be attacked by both electrophile and nucleophile.<sup>[7]</sup> However, only the alkyl propiolates have been proven to be effective substrates for the synthesis of osmapentalynes so far. Thus, developing alternative synthetic approaches is desirable to explore the chemistry of metallapentalyne, as well as planar Möbius aromaticity. Here, we report a general route to osmapentalynes by the reactions of complex **1**<sup>[8]</sup> with allene, alkyne, and alkene.

Complex **1** can be readily prepared and exhibits remarkable stability in the solid state due to  $\sigma$ -aromaticity in the metallacyclopropene unit and  $\pi$ -aromaticity in the fused five-membered rings.<sup>[8]</sup> In addition, metallacyclopropene species are considered as important intermediates in various reactions.<sup>[9]</sup> Therefore, we study the reactions of **1** with a series of unsaturated molecules, such as allene, alkyne, and alkene. As shown in Scheme 1, treatment of complex **1** with excess



**Scheme 1.** Synthesis of osmapentalynes **2**–**4**.


allenylboronic acid pinacol ester in the presence of  $\text{HBF}_4 \cdot \text{H}_2\text{O}$  at  $60^\circ\text{C}$  for 3 days led to the formation of complex **2**, which was isolated in 60% yield by column chromatography. A plausible mechanism for the formation of **2** was proposed in Scheme S1 in the Supporting Information (SI).

The structure of complex **2** has been verified by X-ray diffraction analysis.<sup>[10]</sup> As shown in Figure 1, the metallacycle of **2** is almost planar, which is reflected by the mean deviation (0.0143 Å) from the least-squares plane through the eight atoms ( $\text{Os}1, \text{C}1\cdots\text{C}7$ ). The  $\text{Os}-\text{C}1$  bond length (1.847 Å) is close to that of the first osmapentalyne (1.845 Å).<sup>[6]</sup> The  $\text{C}-\text{C}$  bond distances (1.375–1.411 Å) in the fused five-membered rings are between the lengths of  $\text{C}-\text{C}$  single and double bonds without significant bond-length alternation. The planarity as well as the  $\text{Os}-\text{C}$  and  $\text{C}-\text{C}$  bond distances in the fused five-

[\*] Dr. C. Zhu, Y. Yang, J. Wu, M. Luo, J. Fan, Dr. J. Zhu, Prof. Dr. H. Xia State Key Laboratory of Physical Chemistry of Solid Surfaces and Collaborative Innovation Center of Chemistry for Energy Materials (iChEM), and Department of Chemistry, College of Chemistry and Chemical Engineering, Xiamen University Xiamen 361005 (China) E-mail: jun.zhu@xmu.edu.cn hpxia@xmu.edu.cn Homepage: <http://junzhu.chem8.org/> <http://chem.xmu.edu.cn/group/hpxia/index.htm>

[\*\*] We thank the 973 Program (2012CB821600), the NSFC (21332002, 21174115, and 21172184), the Program for New Century Excellent Talents in University (NCET-13-0511), and the program for Changjiang Scholars and Innovative Research Team in University.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201502412>.




**Figure 1.** Molecular structure of the cation of **2** (thermal ellipsoids set at 50% probability). Phenyl groups in  $\text{PPh}_3$  have been omitted for clarity.

membered ring indicates a delocalized metallabicyclic compound.<sup>[11]</sup>

Complex **2** has also been characterized by high-resolution mass spectrometry (HRMS), NMR spectroscopy, and elemental analysis. The HRMS showed a molecular ion peak at  $m/z = 1155.2817$  with expected isotopic distribution (calculated value for complex **2**  $[\text{C}_{65}\text{H}_{55}\text{ClOsP}_3]^+$  at  $m/z = 1155.2809$ ). In the  $^1\text{H}$  NMR spectrum, the typical H signal of  $\text{Os}-\text{CH}$  disappears, which is consistent with the solid structure. The proton signals in the fused five-membered rings of **2** are observed at the aromatic region (i.e. 9.03 ppm for H5, 7.60 ppm for H6, and 7.28 ppm for H3, confirmed by  $^1\text{H}-^{13}\text{C}$  HSQC). In the  $^{13}\text{C}$  NMR spectrum, the signal of the carbyne carbon appears at 320.0 ppm, which is close to the signals of the reported osmapentalynes (323.8–330.8 ppm).<sup>[6]</sup>

The delocalized structure of **2** can be rationalized by two major resonance structures (Scheme 2): osmapentalyne **2A**



**Scheme 2.** Two major resonance structures of **2**.

(with an  $\text{Os}\equiv\text{C}$  unit) and osmacycloallene **2B** (with an  $\text{Os}=\text{C}=\text{C}$  unit). The structures with  $\text{C}=\text{C}=\text{C}$  unit in five-membered metallacycles, e.g., metallacycloallenes<sup>[12]</sup> and metallacyclocumulenes,<sup>[13]</sup> have been reported experimentally and theoretically. In contrast, the similar structure with a  $\text{M}=\text{C}=\text{C}$  unit in a five-membered ring has only one example reported by us recently.<sup>[14]</sup>


Besides allene, osmapentalene **1** can also react with alkyne. As shown in Scheme 1, treatment of **1** with excess phenylacetylene in the presence of  $\text{HBF}_4\cdot\text{H}_2\text{O}$  at 60 °C for 2 d led to the formation of complex **3**, which was isolated in 65% yield by column chromatography. The structure of **3** was fully supported by NMR spectroscopy, HRMS, elemental analysis, and X-ray diffraction. The structural features of the metallabicyclic in **3** are very similar to those of **2** (Figure S6).

Although these reactions provide a new route for the synthesis of osmapentalynes via osmapentalene, they are relatively slow. To target a more rapid reaction, we carried out density functional theory (DFT) calculations to probe the nature of the reaction. Formally, it is the internal  $\text{sp}^2$ -hybridized carbons of allene and alkyne that attacked the  $\text{sp}^3$ -hybridized carbon atom (C8) of osmapentalene **1** (Fig-

ure S1). As the B(pin) substituent is electron-deficient, the internal carbon of allene should be electrophilic. Thus the C8 atom of osmapentalene **1** is expected to be nucleophilic. Indeed, the calculation on the model complex **1'** (for which  $\text{PH}_3$  was used to replace  $\text{PPh}_3$  in **1**) shows that the Mulliken charge on the C8 atom is  $-0.61$  (Figure S1). The charge on the internal carbon atoms in allene, and in alkyne are  $+0.56$  and  $+1.19$ , respectively, in line with our hypothesis. If the carbon in a substrate becomes more electrophilic, this reaction should become faster. Interestingly, the Mulliken charge of alkene carbon in tetracyanoethylene (TCNE) is  $+2.35$ , indicating that it could be a better substrate.

Therefore, we tried TCNE to react with osmapentalene **1**. Osmapentalyne **4** was produced at room temperature for only 3 h and isolated in 82% yield (Scheme 1). The proton signals of the fused five-membered rings in **4** were located at the aromatic region in the  $^1\text{H}$  NMR spectrum. Specifically, the signals of H3, H5, and H6 were observed at 7.82, 9.55, and 8.11 ppm, respectively. In the  $^{13}\text{C}$  NMR spectrum, the C1 signal (319.7 ppm) was comparable to the signals for osmapentalynes (323.8–330.8 ppm)<sup>[6]</sup> and osmabenzynes (264.9–316.4 ppm).<sup>[2]</sup>

The structure of complex **4** was further confirmed by HRMS, elemental analysis, and single-crystal X-ray diffraction. As shown in Figure 2, it contains an  $\text{Os}\equiv\text{C}$  triple bond in




**Figure 2.** Molecular structure of the cation of **4** (thermal ellipsoids set at 50% probability). Phenyl groups in  $\text{PPh}_3$  have been omitted for clarity.

the planar fused five-membered rings (the mean deviation from the least-square plane is only 0.0214 Å). The length of the  $\text{Os1}-\text{C1}$  (1.795 Å) and the bond angle at the carbyne carbon ( $130.9^\circ$ ) are comparable to those of  $\text{Os}\equiv\text{C}$  triple bonds (1.845 and 1.808 Å) and the bond angles ( $129.5$  and  $131.2^\circ$ ) in osmapentalynes reported earlier.<sup>[6]</sup> The bond lengths of  $\text{C7}-\text{C8}$  (1.512 Å),  $\text{C8}-\text{C9}$  (1.556 Å), and  $\text{C9}-\text{C12}$  (1.572 Å) are obviously carbon–carbon single bonds.

The downfield proton chemical shifts, the planarity of the fused five-membered rings, and the delocalized carbon–carbon bond indicate that the osmapentalynes **2–4** are aromatic. To confirm their aromaticity, we employed the “isomerization stabilization energy” (ISE) method, a convenient tool to evaluate the magnitude of aromaticity<sup>[15]</sup> by DFT calculations. We calculated the ISE values based on the model complexes of **2'**, **3'**, and **4'** for which  $\text{PH}_3$ 's were used to replace the  $\text{PPh}_3$ 's. The computed ISE values of **2'**, **3'**, and **4'** are  $-36.1$ ,  $-37.6$ , and  $-32.6$  kcal mol $^{-1}$  ( $\text{R} = \text{PH}_3^+$ , Scheme S5), respectively, which are more negative than those of the model complexes **2''** ( $-24.1$  kcal mol $^{-1}$ ), **3''** ( $-25.0$  kcal mol $^{-1}$ ), and **4''** ( $-23.9$  kcal mol $^{-1}$ ) in which the

phosphonium substituent was removed ( $R = H$ , Scheme S5), further demonstrating the stabilization effect of the phosphonium substituent.<sup>[16]</sup>

We also performed nucleus-independent chemical shift (NICS) calculations<sup>[17]</sup> on the model complexes of **2'**, **3'**, and **4'**. In general, negative values indicate aromaticity and positive values mean antiaromaticity. Indeed, the computed NICS(1)<sub>zz</sub> for the left and right five-membered rings of complexes **2'** (−15.7 and −16.1 ppm), **3'** (−15.3 and −15.9 ppm), and **4'** (−18.1 and −18.0 ppm) are all negative. These larger negative NICS(1)<sub>zz</sub> values suggest that all these osmapentalynes are aromatic. In addition, the aromaticity in **2'** is further supported by the anisotropy of the induced current density (AICD) analysis.<sup>[18]</sup> As shown in Figure 3, the clockwise current density vectors plotted on the AICD



**Figure 3.** AICD isosurfaces of **2'** by  $\pi$  contribution. Current density vectors are plotted onto the AICD isosurface of 0.035 to indicate diatropic ring currents. The magnetic field vector is orthogonal with respect to the ring plane and points upward (clockwise currents are diatropic).


isosurface indicate that a diatropic ring current in the  $\pi$  system is displayed along the periphery of fused five-membered rings, thus confirming the  $\pi$ -aromaticity in **2'** (see also Figure S2).

Note that the cyano group is one of the strongest electron-withdrawing groups, which also induce stronger electron-accepting effects in a conjugated system. However, the four cyano groups in complex **4** are unconjugated with osmapentalyne unit. Fortunately, treatment of complex **4** with excess triethylamine at room temperature for an hour led to the color change from yellow to green. The conjugated product **5** was isolated as a green solid (Scheme 3). A possible mechanism for the formation of **5** is proposed in Scheme S3 in the SI.

Complex **5** was characterized by HRMS, elemental analysis, <sup>1</sup>H and <sup>31</sup>P NMR spectra. Unfortunately, the poor solubility of complex **5** hampers the characterization by <sup>13</sup>C NMR spectroscopy. Nevertheless, the solid structure of **5**,




**Scheme 3.** Reaction of osmapentalyne **4** with  $\text{NEt}_3$ .



**Figure 4.** Molecular structure of **5** (thermal ellipsoids set at 50% probability). Phenyl groups in  $\text{PPh}_3$  have been omitted for clarity.

which also contains an osmapentalyne unit, was confirmed by single-crystal X-ray diffraction (Figure 4). The characteristic Os1–C1 bond length (1.855 Å) and the Os1–C1–C2 angle (130.0°) are close to those of osmapentalynes reported earlier.<sup>[6]</sup> In addition, the three cyano groups in complex **5** are conjugated with the osmapentalyne unit.

Structurally, the most distinct feature of complex **5** is that all of the cyano groups are conjugated with the osmapentalyne unit compared with complex **4**, which could lead to the significant difference of the UV/Vis absorption spectra. As shown in Figure 5, with the extended conjugation, the lowest-



**Figure 5.** UV/Vis absorption spectra of **4** and **5** measured in  $\text{CH}_2\text{Cl}_2$  at room temperature.

energy absorption band is red-shifted and the molar extinction coefficient is increased. Specifically, the absorption maximum of complex **5** in the low-energy absorption band located at 694 nm ( $\epsilon = 4.68 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ,  $\epsilon$ : molar extinction coefficient), which is red-shifted by 242 nm compared with that of **4** ( $\lambda_{\text{max}} = 452 \text{ nm}$ ,  $\epsilon = 0.84 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ) due to the effective conjugation of multicyno groups with osmapentalyne unit (Figure S3). Remarkably, the molar extinction coefficient of **5** ( $4.68 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ) in the low-energy absorption band is even three times more than that of the classical ruthenium complex **N3** ( $\epsilon = 1.42 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ).<sup>[19]</sup> The UV/Vis absorption maximum of **2** ( $\lambda_{\text{max}} = 510 \text{ nm}$ ,  $\epsilon = 1.73 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ) and **3** ( $\lambda_{\text{max}} = 526 \text{ nm}$ ,  $\epsilon = 1.71 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ) are also recorded under the same conditions which are slightly red-shifted by 75 and 91 nm compared with that of complex **1** ( $\lambda_{\text{max}} = 435 \text{ nm}$ ,  $\epsilon = 0.82 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ; Figure S4). The absorption maximum of complexes **4**, **2**, **3**, and **5** are red-shifted gradually, which is consistent with the decreasing HOMO–LUMO gaps (3.45, 3.08, 2.98, and 2.31 eV for **4**, **2**, **3**, and **5**, respectively) from calculations.



Metallapentalyne, containing a metal–carbon triple bond in five-membered ring, is a unique aromatic system. Here, we developed a general and effective method for the synthesis of osmapentalynes via osmapentalene with a metallacyclopropene unit. Such a metallacyclopropene unit can react with allene, alkyne, and alkene to form osmapentalynes. The experimental data together with the computations (the negative ISE, NICS values, and AICD plot) confirm the aromaticity of these fused five-membered rings. Developing such a general method allows us to realize various metallaromatics with potential applications in photoacoustic imaging and photothermal therapy.<sup>[20]</sup>

**Keywords:** aromaticity · density functional calculations · metal carbynes · osmapentalynes · osmium

**How to cite:** *Angew. Chem. Int. Ed.* **2015**, *54*, 7189–7192  
*Angew. Chem.* **2015**, *127*, 7295–7298

- [1] a) “Cyclic alkynes: preparation and properties”: R. Gleiter, R. Merger in *Modern Acetylene Chemistry*, (Eds.: P. J. Stang, F. Diederich), Wiley-VCH, Weinheim, **1995**; b) G. Wittig, *Angew. Chem. Int. Ed. Engl.* **1962**, *1*, 415; *Angew. Chem.* **1962**, *74*, 479; c) N. Suzuki, M. Nishiura, Y. Wakatsuki, *Science* **2002**, *295*, 660; d) U. Rosenthal, *Angew. Chem. Int. Ed.* **2004**, *43*, 3882; *Angew. Chem.* **2004**, *116*, 3972; e) N. Suzuki, D. Hashizume, *Coord. Chem. Rev.* **2010**, *254*, 1307.
- [2] Selected examples for metallabenzynes: a) T. B. Wen, Z. Y. Zhou, G. Jia, *Angew. Chem. Int. Ed.* **2001**, *40*, 1951; *Angew. Chem.* **2001**, *113*, 2005; b) T. B. Wen, S. M. Ng, W. Y. Hung, Z. Y. Zhou, M. F. Lo, L.-Y. Shek, I. D. Williams, Z. Lin, G. Jia, *J. Am. Chem. Soc.* **2003**, *125*, 884; c) T. B. Wen, W. Y. Hung, H. H. Y. Sung, I. D. Williams, G. Jia, *J. Am. Chem. Soc.* **2005**, *127*, 2856; d) W. Y. Hung, J. Zhu, T. B. Wen, K. P. Yu, H. H. Y. Sung, I. D. Williams, Z. Lin, G. Jia, *J. Am. Chem. Soc.* **2006**, *128*, 13742; e) J. Chen, H. H. Y. Sung, I. D. Williams, Z. Lin, G. Jia, *Angew. Chem. Int. Ed.* **2011**, *50*, 10675; *Angew. Chem.* **2011**, *123*, 10863.
- [3] a) D. P. Craig, N. L. Paddock, *Nature* **1958**, *181*, 1052; b) H. S. Rzepa, *Chem. Rev.* **2005**, *105*, 3697; c) R. Herges, *Chem. Rev.* **2006**, *106*, 4820.
- [4] a) E. Heilbronner, *Tetrahedron Lett.* **1964**, *5*, 1923; b) H. E. Zimmerman, *J. Am. Chem. Soc.* **1966**, *88*, 1564.
- [5] a) D. Ajami, O. Oeckler, A. Simon, R. Herges, *Nature* **2003**, *426*, 819; b) J. K. Park, Z. S. Yoon, M.-C. Yoon, K. S. Kim, S. Mori, J.-Y. Shin, A. Osuka, D. Kim, *J. Am. Chem. Soc.* **2008**, *130*, 1824; c) Z. S. Yoon, A. Osuka, D. Kim, *Nat. Chem.* **2009**, *1*, 113; d) G. R. Schaller, F. Topić, K. Rissanen, Y. Okamoto, J. Shen, R. Herges, *Nat. Chem.* **2014**, *6*, 608.
- [6] C. Zhu, S. Li, M. Luo, X. Zhou, Y. Niu, M.-L. Lin, J. Zhu, Z. Cao, X. Lu, T. Wen, Z. Xie, P. v. R. Schleyer, H. Xia, *Nat. Chem.* **2013**, *5*, 698.
- [7] a) C. Zhu, M. Luo, Q. Zhu, J. Zhu, P. v. R. Schleyer, J. I.-C. Wu, X. Lu, H. Xia, *Nat. Commun.* **2014**, *5*, 3265; b) C. Zhu, Q. Zhu, J. Fan, J. Zhu, X. He, X.-Y. Cao, H. Xia, *Angew. Chem. Int. Ed.* **2014**, *53*, 6232; *Angew. Chem.* **2014**, *126*, 6346.
- [8] C. Zhu, X. Zhou, H. Xing, K. An, J. Zhu, H. Xia, *Angew. Chem. Int. Ed.* **2015**, *54*, 3102; *Angew. Chem.* **2015**, *127*, 3145.
- [9] a) C. P. Casey, J. T. Brady, T. M. Boller, F. Weinhold, R. K. Hayashi, *J. Am. Chem. Soc.* **1998**, *120*, 12500; b) U. Rosenthal, V. V. Burlakov, M. A. Bach, T. Beweries, *Chem. Soc. Rev.* **2007**, *36*, 719; c) S. Zhang, W.-X. Zhang, J. Zhao, Z. Xi, *J. Am. Chem. Soc.* **2010**, *132*, 14042; d) R. Castro-Rodrigo, M. A. Esteruelas, A. M. López, F. López, J. L. Mascareñas, M. Oliván, E. Oñate, L. Saya, L. Villarino, *J. Am. Chem. Soc.* **2010**, *132*, 454; e) A. Collado, M. A. Esteruelas, E. Oñate, *Organometallics* **2011**, *30*, 1930; f) J. Zhao, S. Zhang, W.-X. Zhang, Z. Xi, *Coord. Chem. Rev.* **2014**, *270*, 2.
- [10] CCDC 1041475 (2), 1041476 (3), 1041477 (4), and 1041478 (5) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
- [11] a) J. R. Bleeker, *Chem. Rev.* **2001**, *101*, 1205; b) H.-P. Wu, T. J. R. Weakley, M. M. Haley, *Chem. Eur. J.* **2005**, *11*, 1191; c) C. W. Landorf, M. M. Haley, *Angew. Chem. Int. Ed.* **2006**, *45*, 3914; *Angew. Chem.* **2006**, *118*, 4018; d) V. Jacob, C. W. Landorf, L. N. Zakharov, T. J. R. Weakley, M. M. Haley, *Organometallics* **2009**, *28*, 5183; e) B. J. Frogley, L. J. Wright, *Coord. Chem. Rev.* **2014**, *270*, 151.
- [12] Selected examples for metallacycloallenes, see: a) N. Suzuki, D. Hashizume, H. Koshino, T. Chihara, *Angew. Chem. Int. Ed.* **2008**, *47*, 5198; *Angew. Chem.* **2008**, *120*, 5276; b) J. Ugolotti, G. Dierker, G. Kehr, R. Fröhlich, S. Grimme, G. Erker, *Angew. Chem. Int. Ed.* **2008**, *47*, 2622; *Angew. Chem.* **2008**, *120*, 2662; c) J. Ugolotti, G. Kehr, R. Fröhlich, S. Grimme, G. Erker, *J. Am. Chem. Soc.* **2009**, *131*, 1996; d) K. Kaleta, M. Ruhmann, O. Theilmann, T. Beweries, S. Roy, P. Arndt, A. Villinger, E. D. Jemmis, A. Schulz, U. Rosenthal, *J. Am. Chem. Soc.* **2011**, *133*, 5463; e) G. Bender, G. Kehr, R. Fröhlich, J. L. Petersen, G. Erker, *Chem. Sci.* **2012**, *3*, 3534; f) G. Bender, G. Kehr, C. G. Daniliuc, B. Wibbeling, G. Erker, *Dalton Trans.* **2013**, *42*, 14673.
- [13] Selected examples for metallacyclocumulenes, see: a) U. Rosenthal, A. Ohff, W. Baumann, R. Kempe, A. Tillack, V. V. Burlakov, *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 1605; *Angew. Chem.* **1994**, *106*, 1678; b) P.-M. Pellny, V. V. Burlakov, P. Arndt, W. Baumann, A. Spannenberg, U. Rosenthal, *J. Am. Chem. Soc.* **2000**, *122*, 6317; c) U. Rosenthal, V. V. Burlakov, P. Arndt, W. Baumann, A. Spannenberg, *Organometallics* **2005**, *24*, 456; d) U. Rosenthal, *Angew. Chem. Int. Ed.* **2008**, *47*, 5118; *Angew. Chem.* **2008**, *120*, 5196; e) T. Beweries, U. Rosenthal, *Nat. Chem.* **2013**, *5*, 649; f) S. Roy, U. Rosenthal, E. D. Jemmis, *Acc. Chem. Res.* **2014**, *47*, 2917.
- [14] T. Wang, J. Zhu, F. Han, C. Zhou, H. Chen, H. Zhang, H. Xia, *Angew. Chem. Int. Ed.* **2013**, *52*, 13361; *Angew. Chem.* **2013**, *125*, 13603.
- [15] a) P. v. R. Schleyer, F. Pühlhofer, *Org. Lett.* **2002**, *4*, 2873; b) J. Zhu, K. An, P. v. R. Schleyer, *Org. Lett.* **2013**, *15*, 2442; c) I. Fernández, G. Frenking, G. Merino, *Chem. Soc. Rev.* **2015**, DOI: 10.1039/c5s00004a.
- [16] a) X. Wang, C. Zhu, H. Xia, J. Zhu, *Organometallics* **2014**, *33*, 1845; b) Y. Huang, X. Wang, K. An, J. Fan, J. Zhu, *Dalton Trans.* **2014**, *43*, 7570.
- [17] a) P. v. R. Schleyer, C. Maerker, A. Dransfeld, H. Jiao, N. J. R. v. E. Hommes, *J. Am. Chem. Soc.* **1996**, *118*, 6317; b) Z. Chen, C. S. Wannere, C. Corminboeuf, R. Puchta, P. v. R. Schleyer, *Chem. Rev.* **2005**, *105*, 3842; c) H. Fallah-Bagher-Shaidaei, C. S. Wannere, C. Corminboeuf, R. Puchta, P. v. R. Schleyer, *Org. Lett.* **2006**, *8*, 863.
- [18] a) R. Herges, D. Geuenich, *J. Phys. Chem. A* **2001**, *105*, 3214; b) D. Geuenich, K. Hess, F. Köhler, R. Herges, *Chem. Rev.* **2005**, *105*, 3758.
- [19] M. K. Nazeeruddin, A. Kay, I. Rodicio, R. Humphry-Baker, E. Mueller, P. Liska, N. Vlachopoulos, M. Graetzel, *J. Am. Chem. Soc.* **1993**, *115*, 6382.
- [20] C. Zhu, Y. Yang, M. Luo, C. Yang, J. Wu, L. Chen, G. Liu, T. Wen, J. Zhu, H. Xia, *Angew. Chem. Int. Ed.* **2015**, DOI: 10.1002/anie.201501349; *Angew. Chem.* **2015**, DOI: 10.1002/ange.201501349.

Received: March 15, 2015

Published online: April 27, 2015