



# A convenient synthesis of pyridine and 2,2'-bipyridine derivatives

Marta Altuna-Urquijo<sup>a</sup>, Alexander Gehre<sup>a</sup>, Stephen P. Stanforth<sup>a</sup>,   and Brian Tarbit<sup>b</sup>

<sup>a</sup>School of Applied Sciences, Northumbria University, Newcastle upon Tyne NE1 8ST, UK

<sup>b</sup>Vertellus Specialities UK Ltd, Seal Sands Road, Seal Sands, Middlesbrough TS2 1UB, UK

## Abstract

$\alpha$ -Chloro- $\alpha$ -acetoxy- $\beta$ -keto-esters **9** were readily prepared from  $\beta$ -keto-esters **6** in good overall yields. These compounds reacted as  $\alpha,\beta$ -diketo-ester equivalents **2** with amidrazones **1** yielding triazines **3**, generally in good yields. Picolinates **10** provided an alternative source of  $\alpha,\beta$ -diketo-ester equivalents **2** when treated with copper(II) acetate. A 'one-pot' reaction of the  $\alpha,\beta$ -diketo-ester equivalents **2** with amidrazones **1** in the presence of 2,5-norbornadiene **5** in boiling ethanol yielded the pyridines **4** and 2,2'-bipyridines **4** ( $R^1=2$ -pyridyl) directly without the need to isolate the corresponding triazines **3**. Triazine **3c** reacted with the azadienophiles **13** and **17** affording the products **16** and **18**, respectively, in good yields.

## Graphical abstract

