

Edith Cowan University
Research Online

ECU Publications 2013

1-1-2013

A blood-based predictor for neocortical A β burden in Alzheimer's disease: results from the AIBL study

Samantha Burnham

Noel Faux

William Wilson

Simon Laws

Edith Cowan University, s.laws@ecu.edu.au

David Ames

See next page for additional authors

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworks2013>

 Part of the [Hematology Commons](#)

10.1038/mp.2013.40

This is an Author's Accepted Manuscript of: Burnham, S., Faux, N., Wilson, W., Laws, S. , Ames, D., Bedo, J., Bush, A., Doecke, J., Ellis, K., Head, R., Jones, G., Kiiveri, H., Martins, R. N., Rembach, A., Rowe, C., Salvado, O., Macaulay, S., Masters, C., & Villemagne, V. (2014). A blood-based predictor for neocortical A β burden in Alzheimer's disease: results from the AIBL study. *Molecular Psychiatry*, 19(4), 519-526. Available [here](#)

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworks2013/527>

Authors

Samantha Burnham, Noel Faux, William Wilson, Simon Laws, David Ames, Justin Bedo, Ashley Bush, James Doecke, Kathryn Ellis, Richard Head, Gareth Jones, Harri Kiiveri, Ralph N. Martins, Alan Rembach, Christopher Rowe, Olivier Salvado, S Lance Macaulay, Colin Masters, and Victor Villemagne

Original Article

A BLOOD BASED PREDICTOR FOR NEOCORTICAL A β BURDEN IN ALZHEIMER'S DISEASE: RESULTS FROM THE AIBL STUDY

Samantha C. Burnham (PhD)^{1*}, Noel G. Faux (PhD)^{2*}, William Wilson (PhD)³, Simon M. Laws (PhD)^{4,5}, David Ames (MD)^{6,7}, Justin Bedo (PhD)⁸, Ashley I. Bush (MD)², James D. Doecke (PhD)⁹, Kathryn A. Ellis (PhD)⁶, Richard Head (PhD)¹⁰, Gareth Jones (MSc)^{11,12}, Harri Kiiveri (PhD)¹, Ralph N. Martins (PhD)^{4,5}, Alan Rembach (PhD)², Christopher Rowe (MD)^{11,12}, Olivier Salvado (PhD)¹³, S. Lance Macaulay¹⁴ (PhD), Colin L. Masters (MD)² and Victor L. Villemagne (MD)¹¹ for the Alzheimer's Disease Neuroimaging Initiative^{15, **} and Australian Imaging, Biomarkers and Lifestyle Study Research Group¹⁶

1CSIRO Preventative Health Flagship: Mathematics, Informatics and Statistics, Floreat, WA, Australia.

2Mental Health Research Institute (MHRI), the University of Melbourne, Parkville, VIC, Australia.

3CSIRO Preventative Health Flagship: Mathematics, Informatics and Statistics, North Ryde, NSW, Australia.

4Centre of Excellence for Alzheimer's Disease Research & Care, School of Medical Sciences, Edith Cowan University, Joondalup, WA, Australia.

5Sir James McCusker Alzheimer's Disease Research Unit (Hollywood Private Hospital), Perth, WA, Australia.

6Academic Unit for Psychiatry of Old Age, Department of Psychiatry, University of Melbourne, Parkville, VIC, Australia.

7National Ageing Research Institute, Parkville, VIC, Australia.

8Victorian Research Laboratory, National ICT of Australia (NICTA), VIC, Australia.

9CSIRO Preventative Health Flagship: Mathematics, Informatics and Statistics, Herston, QLD, Australia.

10CSIRO Preventative Health Flagship: Health and Life Sciences, Urrbrae, SA, Australia.

11Department of Nuclear Medicine and Centre for PET, Austin Health, Heidelberg, VIC, Australia.

12Department of Medicine, Austin Health, The University of Melbourne, Heidelberg, VIC, Australia.

13CSIRO Preventative Health Flagship: Information and Communication Technology, Herston, QLD, Australia.

14CSIRO Preventative Health Flagship: Materials Science and Engineering, Parkville, VIC, Australia.

15adni.loni.ucla.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

16www.aibl.csiro.au/about/aibl-research-team

*These authors contributed equally to the manuscript

**Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.ucla.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: adni.loni.ucla.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Corresponding author: Samantha Burnham

Corresponding author's address: CMIS, CSIRO, Private Bag 5, Wembley, WA 6913, Australia

Corresponding author's phone and fax: +61 (0) 8 9333 6706 and +61 (0) 8 9333 6121

Corresponding author's e-mail address: samantha.burnham@csiro.au

Running head: Neocortical A β Burden Prediction

Support Sources (grants, equipment and drugs):

Core funding for the study was provided by the CSIRO Flagship Collaboration Fund and the Science and Industry Endowment Fund (SIEF) in partnership with Edith Cowan University (ECU), Mental Health Research institute (MHRI), Alzheimer's Australia (AA), National Ageing Research Institute (NARI), Austin Health, CogState Ltd., Hollywood Private Hospital, Sir Charles Gardner Hospital. The study also receives funding from the National Health and Medical Research Council (NHMRC), the Dementia Collaborative Research Centres program (DCRC), The McCusker Alzheimer's Research Foundation and Operational Infrastructure Support from the Government of Victoria. Faux NG is supported by, a National Health and Medical Research Council training fellowship. Laws SM is supported by research fellowships from Edith Cowan University. Bush AI is supported by the NHMRC by a program grant and an Australian Fellowship.

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: Abbott; Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Amorfix Life Sciences Ltd.; AstraZeneca; Bayer HealthCare; BioClinica, Inc.; Biogen Idec Inc.; Bristol-Myers Squibb Company; Eisai Inc.; Elan Pharmaceuticals Inc.; Eli Lilly and Company; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; GE Healthcare; Innogenetics, N.V.; Janssen Alzheimer Immunotherapy

Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Medpace, Inc.; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Servier; Synarc Inc.; and Takeda Pharmaceutical Company. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University of California, San Diego. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of California, Los Angeles. This research was also supported by NIH grants P30 AG010129 and K01 AG030514.

ABSTRACT

Dementia is a global epidemic with Alzheimer's disease (AD) being the leading cause. Early identification of patients at risk of developing AD is now becoming an international priority. Neocortical A β Burden (NAB), as assessed by positron emission tomography (PET) represents one such marker for early identification. These scans are expensive and are not widely available, thus there is a need for cheaper and more widely accessible alternatives. Addressing this need, a blood biomarker based signature having efficacy for the prediction of NAB and that can be easily adapted for population screening, is described. Blood data (176 analytes measured in plasma) and PiB-PET measurements from 273 participants from the Australian Imaging, Biomarkers and Lifestyle (AIBL) study were utilised. Univariate analysis was conducted to assess the difference of plasma measures between high and low NAB groups and cross-validated machine-learning models were generated for predicting NAB. These models were applied to 817 non-imaged AIBL subjects and 82 subjects from the Alzheimer's Disease Neuroimaging Initiative (ADNI) for validation. Five analytes showed significant difference between subjects with high compared to low NAB. A machine-learning model (based on nine markers) achieved sensitivity and specificity of 80% and 82%, respectively for predicting NAB. Validation using the ADNI cohort yielded similar results (sensitivity 79%, specificity 76%). These results show that a panel of blood-based biomarkers is able to accurately predict NAB, supporting the hypothesis for a relationship between a blood-based signature and A β accumulation, therefore providing a platform for developing a population based screen.

INTRODUCTION

The only definitive diagnosis for Alzheimer's disease (AD) is by histological examination at autopsy, with clinical diagnosis based on medical history, clinical examination and neuropsychological testing¹. However, clinical symptoms only present after significant irreversible neurological degeneration has occurred². Thus, there is a need for a simple population based screening test that is able to identify subjects at risk of developing AD at the pre-symptomatic stages³.

The hallmark histological markers of AD are the presence of extracellular β -amyloid ($A\beta$) plaques and intracellular neurofibrillary tangles (NFT) principally comprised of hyperphosphorylated Tau (pTau)⁴. Currently two leading sets of biomarkers are available which show strong correlations with these histological markers of AD. Firstly, imaging markers which bind $A\beta$, (e.g. Pittsburgh Compound B; PiB) coupled with Positron Emission Tomography (PET), that provide semi-quantitative assessments of neocortical $A\beta$ burden (NAB)⁵. The other leading set of biomarkers is derived from cerebral spinal fluid (CSF) and includes measures of $A\beta$ (in particular the $A\beta_{1-42}$ species) that correlate with $A\beta$ in the brain and total Tau and pTau that correlate with NFTs⁶. However, both of these sets of biomarkers have limited use as general population-screening tools for AD. PET imaging is relatively expensive and its availability is limited, whilst the collection of CSF requires a lumbar puncture which is invasive. Widespread population screening requires alternative approaches to estimate NAB derived from more accessible tissues, such as blood.

Given the strong association between NAB and AD pathology and the need of population blood-based screening tests to identify those individuals at risk of developing AD, this study presents a blood biomarker panel whose composite score correlates with NAB, as assessed by PiB-PET. The results are validated using samples taken from an independent study.

METHODS

a) Cohorts:

(i) The Australian Imaging, Biomarkers and Lifestyle (AIBL) Study.

Detailed information on the study design and enrolment procedures has been discussed elsewhere⁷. The AIBL study is a prospective longitudinal study of aging, integrating data from neuroimaging, biomarkers, lifestyle, clinical and neuropsychological analysis. Eligible volunteers, aged over 60 years and fluent in English, were classified into three groups: 1) individuals meeting NINCDS-ADRDA criteria for AD¹, 2) individuals meeting criteria for Mild Cognitive Impairment (MCI)^{8,9} and 3) cognitively healthy individuals (HC). The institutional ethics committees of Austin Health, St Vincent's Health, Hollywood Private Hospital and Edith Cowan University approved the AIBL study, and all volunteers gave written informed consent before participating.

Blood Collection, Processing and Analytes Assays.

Whole blood (80 mL) was collected in the morning (after overnight fasting) by venipuncture (blood processing procedure, is detailed in the Supplementary Methods Section 1 a) of Supplementary Material). EDTA plasma samples were shipped to Myriad Rules-Based Medicine (Myriad RBM; Austin, Texas, www.rulesbasedmedicine.com), on dry ice, for analysis using commercially available multiplexed Luminex human discovery 151MAP panel. All assays were validated according to CLIA standards and no samples were older than 18 months at the time of analysis. The assays used for EDTA plasma A β analysis were both the well-documented double sandwich ELISA technique¹⁰⁻¹² and the commercial luminex based kit (INNO-BIA plasma A β forms, Innogenetics, NV, Belgium), as previously described¹³. EDTA Plasma APOE protein levels were measured using a commercial assay (Pan APOE ELISA, MBL co., Ltd.)¹⁴. Apolipoprotein E (APOE) genotyping using polymerase chain reaction amplification and restriction enzyme digestions¹⁵. Plasma was collected in Sarstedt (01.1608.100 S-Monovette 7.5 mL) lithium heparin tubes for the measurement of seven metal ion levels using induction coupled plasma mass spectrometry (ICP-MS).

(ii) Alzheimer's Disease Neuroimaging Initiative (ADNI):

For validation purposes, data from 82 participants were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database. Details on the study design, enrolment procedures and sample collection are given in Supplementary Methods (Section 1 b) of Supplementary Material).

b) Dataset Quality Control

1090 of the AIBL participants underwent full blood analyses comprised of 57 pathology blood analytes and 169 plasma analytes (151 from the Myriad RBM xMap discovery panel version one, 13 Metals, Apolipoprotein E (APOE) levels, Innogenetics and Mehta based ELISAs for A β 1-40 and A β 1-42). Data points outside 5 standard deviations of the median were treated as missing along with any failed assays (the measurement were below the detectable limit, for the analytes measured on the Myriad RBM multiplex platform) from the blood sample analyses. Analytes with over five percent missing data (see Supplementary Material; Supplementary Table 1, for list), or any subjects with over 50 percent missing data (not applicable in this instance as no AIBL participant who had undergone both imaging and full blood analyses had more than 50% missing data) were removed. This resulted in a working dataset of 176 blood analytes and two ratios (53 Pathology, 111 MyriadRBM, 7 Metals, APOE levels, Innogenetics and Mehta A β 1-40, A β 1-42 and their ratios; listed in Supplementary Table 1) for 1090 AIBL subjects, 273 of whom had undergone PiB-PET imaging. The pattern of missingness in the resulting dataset was random (i.e. the missing data was not biased to a particular clinical diagnostic group) and missing data points were imputed using multiple imputations by chain equation¹⁶ (all data in the working data set was used). Median values of the hundredth iteration of 100 multiple imputations were used. To adopt normality and to ensure uniformity of residuals in the imputation (and the ensuing) analyses all variables were log transformed prior to imputation. The same quality control procedures were applied independently to the validation (ADNI) blood analyte data, resulting in 176 blood analytes and one ratio for 82 subjects, 113 of the analytes and one ratio overlapping with those in the AIBL dataset

c) Image Analysis

273 AIBL subjects and 82 ADNI subjects, for who relevant blood samples had been analysed, underwent PiB-PET imaging. The PiB imaging methodology of the AIBL and ADNI studies is detailed elsewhere^{5,17}. The AIBL and ADNI PiB-PET scans for each participant were spatially normalised to a customized PiB-PET template in the Montreal Neurological Institute reference space using Statistical Parametric Mapping 8 (SPM8; Wellcome Trust Centre for Neuroimaging, London, United-Kingdom). The spatially normalised PiB-PET images were then scaled using the cerebellar grey matter as reference region for the generation of Standardized Uptake Value ratios (SUVR). Regional analysis was obtained using the Automated Anatomical Labeling (AAL)¹⁸ template restricted to the grey matter (GM) segmentation. The GM masking of the AAL ensured that the PiB measurement did not contain any CSF or white matter voxels. Regional measurements were averaged across both hemispheres. Neocortical A β burden (NAB) was expressed as the average SUVR of the mean of frontal, superior parietal, lateral temporal, lateral occipital, and anterior and posterior cingulate regions. A SUVR cut-off of 1.3, determined through a cluster analysis of HC individuals, was used to classify participants as belonging to a high or a low NAB group.

d) Statistical Analysis

Specific details of the statistical software and settings used throughout this section are also given in Supplementary Methods (Section 1 d) of Supplementary Material).

(i) Demographic Evaluation

The demographic make-up of both the AIBL and ADNI imaged cohorts, split by high and low NAB, for Clinical Classification (HC, MCI or AD), age, gender, *APOE* ϵ 4 status, years of education and Clinical Dementia Rate Scale (CDR) sum of boxes are given by Table 1. Demographic differences between the high and low NAB were assessed using a χ^2 test for the categorical variables, analysis of variance (ANOVA) for the continuous variables was applied, due to the non-normality of the CDR sum of boxes the Mann-Whitney U test was applied. Also the demographic make-up of both the AIBL imaged and non-imaged cohorts as well as the ADNI imaged cohort, split by Clinical Classification, are given in Supplementary Table 2.

(ii) Exploratory Analyses

The 176 blood analytes were assessed, using Analysis of Covariance (ANCOVA), to see if their concentrations differed between participants with high versus low NAB. The results were corrected for age, site, gender and *APOE ε4* carrier status, in addition to these main effects, the pairwise interactions between NAB status and age, ApoE ε4 carries status, and gender were also included in the models. To minimize any false positive results the p-values were adjusted for false discovery rate (FDR)¹⁹.

(iii) Variable Selection and Model Generation

Data for the 273 imaged AIBL participants was initially used for variable selection and model generation; data from the ADNI imaged and AIBL non-imaged cohorts was set aside at this stage. Measures of the 176 blood analytes (along with age, gender, *APOE* genotype and years of education) were considered in variable selection and model generation to predict the continuous SUVR values, through cross-validated Random Forest (RF) analysis. The smallest panel of analytes that gave the least squared error in the cross validation predictions of SUVR was chosen for in-depth validation analysis in attempt to assess potential for use as a biomarker panel for the prediction of NAB status.

To ensure that the identified analytes were robust to the analysis protocol employed, the identified panel was assessed by a second analysis protocol which was based on the binary response of NAB status: namely, cross-validated Support Vector Machine (SVM) analysis with a RBF kernel coupled.

In an attempt to specifically quantify the value added by the blood biomarkers, six separate cross-validated RF sub-models were constructed. Three of the sub-models also included a clinical variable (CDR sum of boxes), incorporated to improve model performance and to provide a comparative assessment of the added value of the blood biomarkers. The six sub-models assessed were model 1 (M1) blood-based markers, age, *APOE* genotype and CDR sum of boxes; model 2 (M2) blood-based markers, age and *APOE* genotype; model 3 (M3) blood-based markers and; model 4 (M4) age, *APOE* genotype and CDR sum of boxes; model 5 (M5) age and CDR sum of boxes; model 6 (M6) age and *APOE* genotype. For validation purposes the resultant sub-models were applied to the imaged ADNI cohort to predict high or low NAB status.

To ensure that the signal was specifically associated with NAB and not just with clinical or cognitive status, the best performing of these sub-models was assessed for efficacy in individual clinical diagnosis groups^a. Finally, the best performing of these sub-models was also applied to the non-imaged AIBL cohort. As no actual NAB information was available for this cohort, the percentages of those predicted to have high NAB for each clinical diagnosis group was compared to those of the imaged AIBL cohort and the literature.

Three-fold cross validation with 100 repeats was adopted and sensitivities, specificities and area under the curve (AUC) performance indicators with standard deviations (SD) were calculated. The predictions of SUVR were split into predicted high or low NAB, based on cut offs providing the most equal sensitivities and specificities, for the purpose of reporting performance statistics.

RESULTS

a) Demographics

Table 1 describes the demographic characteristics of the AIBL and ADNI imaged cohorts. There was a large prevalence of MCI and AD subjects, as well as an enrichment of *APOE ε4* carriers, in the high NAB groups when compared with the low NAB groups for both AIBL and ADNI. There was also an enrichment of elderly subjects in the AIBL high NAB group.

b) Exploratory Analysis: Blood-based analyte differences between high and low NAB

Supplementary Table 1 details the age, gender, site and *APOE ε4* carrier status adjusted means of the blood-based analytes for the two NAB groups. After adjusting for false discovery rates, five analytes showed a significant difference between the NAB groups: Immunoglobulin M-1 (IgM-1) and free thyroxine (FT4) were found to be lower in the high

^a Only one imaged AIBL AD subject had low NAB status, therefore accurate performance statistics could not be calculated for the AIBL AD group; for the ADNI cohort there were only 3 HC subjects (all low NAB status) and no AD subjects with low NAB status so performance statistics were not calculated for the individual clinical diagnosis groups in ADNI.

NAB group ($p = 0.019$ and 0.009 respectively, Figure 1, A and B); whilst macrophage inflammatory protein 1 α (MIP1 α), pancreatic polypeptide (PPY) and vascular cell adhesion protein (VCAM-1) were all found to be elevated in the high NAB group ($p = 0.027$, 0.01 and 0.01 respectively, Figure 1 C-E).

c) Variable Selection and Model Generation

(i) Biomarker Identification

The RF variable selection analysis identified a panel of six biomarkers ($A\beta_{1-42}$, chemokine ligand 13 (CXCL-13), IgM-1, interleukin 17 (IL-17), PPY and VCAM-1) as well as Age and *APOE* genotype. Together, these analytes form the model M2. Three of these analytes (IgM-1, PPY and VCAM-1) were also identified as being significantly different between the NAB groups using ANCOVA, Figure 1 B, D and E.

(ii) Performance Statistics

The cross-validated RF model (M2) achieved a sensitivity, specificity and AUC of 79.6% (SD=1.6%), 79.4% (SD=1.4%) and 83.9% (SD=1.0%), respectively; the cross-validated SVM model achieved a similar sensitivity, specificity and AUC of 74.2% (SD=0.8%), 75.7% (SD=1.4%) and 83.7% (SD=1.6%), respectively. Full performance statistics for the six cross-validated RF sub-models are given by Table 2 and Figure 2A. Based on the AUC, it can be seen that addition of the blood-based markers to the models, resulted in increases in performance of 9% above the demographic and CDR sum of boxes alone (M1 *cf.* M4) and nearly 14% above the demographic variables alone (M2 *cf.* M6). The inclusion of CDR sum of boxes improved the models by 4% (M1 *cf.* M2) and 8% (M4 *cf.* M6). The addition of the blood-based markers increased performance by approximately 5% (M2 *cf.* M4 and M3 *cf.* M5) in contrast to the addition of CDR sum of boxes.

M1 achieved a sensitivity, specificity and AUC of 80.0% (SD=1.1%), 74.2% (SD=1.2%) and 78.4% (SD=0.7%), respectively, in the HC group and a sensitivity, specificity and AUC of 80.1% (SD=1.0%), 81.2% (SD=0.9%) and 81.6% (SD=0.5%), respectively, in the MCI group. Thus, relatively consistent performance is displayed across the overall dataset and both the HC and MCI groups.

(iii) Application to the ADNI validation samples

Unfortunately there were no measurements of IL-17 in the validation cohort. Instead, the median IL-17 measurement from the AIBL cohort was substituted for each of the 82 ADNI samples. Then, the six RF sub-models (M1: M6) generated using AIBL samples were applied to the ADNI validation dataset, performance statistics are given by Table 2 and Figure 2B. It can be seen that M1 achieves 84.7% AUC when applied to the ADNI cohort. It should be noted that dropping IL-17 from the AIBL model results in an AUC of 83.9% (SD=1.1%).

(iv) Application to the non-imaged AIBL samples

The RF model M1 was applied to the 817 AIBL participants who had not undergone the imaging protocol, to predict their expected NAB. All AD, 87% of MCI and 35% of HC participants were predicted to have high NAB, compared with 98% AD, 69% MCI and 34% HC deemed to have high NAB by imaging protocol in the AIBL imaged cohort (Figure 3).

DISCUSSION

This study describes how blood-based molecular signatures can be implemented to predict a person's neocortical amyloid burden (NAB). The signature presented here achieves sensitivity, specificity and AUC of 79.6%, 82.4% and 87.6%, respectively, in the AIBL cohort. Validation in a comparable and independent second cohort, ADNI, achieved similar levels of accuracy with sensitivity, specificity and AUC of 78.3%, 76.3% and 84.7%, respectively. The signature is represented by a RF model of a short list of markers, namely: age, *APOE* genotype, $A\beta_{1-42}$, CXCL-13, IgM-1, IL-17, PPY, VCAM-1 and includes a clinical variable (CDR sum of boxes) which slightly improves model performance (increase in AUC of 4%). However, the blood based panel is shown to have efficacy above that of CDR sum of boxes with a 9% additional increase in AUC when the blood panel is added to a model of age, *APOE* genotype and CDR sum of boxes and a 5% comparative increase in AUC if the blood panel is added to age and *APOE* genotype instead of CDR sum of boxes. It should be noted that IL-17 was not included in the validation signature due to assay failure in the ADNI cohort, regardless the performance statistics of the validation signature were comparable with those of the AIBL signature. Further, removal of IL-17 from the AIBL signature (AUC of 83.9%) was still comparable to that of the validation signature. It should also be noted that an SVM model showed similar performance statistics, using the same set of blood-based markers indicating that the set of markers are robust to the choice of algorithm.

Frank and colleagues²⁰ describe an ideal biomarker for AD as: able to detect a fundamental pathological feature, be validated in neuropathologically confirmed cases, have sensitivity and specificity each above 80%, be reliable, reproducible, noninvasive, simple to perform and inexpensive. Here it has been demonstrated that an additional 10% has been added to performance statistics by inclusion of a small number of blood measurements, over a more traditional signature based on demographic and clinical variables. Thus, inclusion of blood based measurements means that a signature for the prediction of NAB in line with the desirable features of a biomarker as described above has been achieved. In translation to a clinical setting it means that only the measurement of a small number of blood analytes along with standard demographic and clinical variables are necessary to obtain an indication of the NAB in a patient. Thus, such a signature, after further validation, would provide a much needed first pass community level screening option which in turn could provide justification for more invasive or costly confirmation tests such as CSF

analysis or PET, which currently do not lend themselves to widespread population screening due to their invasive or costly nature.

The ADNI cohort provided little data with which to validate our signature in the HC class (N=3), however we observed that performance statistics in the imaged AIBL cohort were consistent across clinical diagnosis groups. This indicates that the identified signal is appropriately tracking with NAB status, opposed to solely tracking with clinical or cognitive status. Percentages of individuals predicted to have high NAB were estimated in the non-imaged AIBL cohort for the specific clinical diagnostic groups (Figure 3). In line with the literature²¹⁻²⁴, the model predicted all of AD, 87% of MCI and around a 35% of HC participants to have high NAB. Given that 35% of HC participants were identified to be at risk of having high NAB over 13% of the MCI group, where arguably the HC participants have better clinical and cognitive representations than their MCI counterparts, further indicates that the signal identified is associated with NAB status and not that of clinical or cognitive status.

The inclusion of age and *APOE* genotype markers, commonly corrected for in models associated with AD as they are the two leading risk factors for AD²⁵, is not an unexpected result. Among the other analytes, pancreatic polypeptide has been identified by several other AD proteomic studies²⁶⁻²⁸. Pancreatic polypeptide is produced by the endocrine F cells in the periphery of the pancreatic islets, and levels in plasma dramatically increase in response to food ingestion^{29,30} playing a role in regulating satiety³¹ as well as insulin release and gut emptying³⁰. While PPY levels are positively correlated with age²⁹, we observed a significant increase in the high NAB group, after adjusting for age. The significance of the increase observed in subjects with high NAB and in AD subjects²⁶ is unclear. However, as elevated PPY has been associated with inflammation³², the elevation of PPY in the high NAB group maybe part an immune signature as previously reported³³.

A number of the other biomarkers, CXCL-13, IgM-1, IL-17 and VCAM-1 also have reported involvement in (acute) immune response and/or immune signaling (see OMIM³⁴ records 605149, 147020, 603149, 192225, 109535, 123260). For the classification of AD and HC subjects in other proteomic studies^{26, 28, 33} similar immune based signatures have been reported.

Alzheimer's pathological A β processing is evident in the brain; however, there is ongoing debate if this is also represented in the plasma^{13, 35-39} (Rembach *et al.*)^b. A number of studies have shown that A β species in plasma and blood cells including the plasma ratio of A β ₁₋₄₂/A β ₁₋₄₀ have a mild correlation with NAB, with the direction of the relationship being A β species dependent^{13, 40, 41} (Rembach *et al.*)^a.

The inclusion of biomarkers other than those assessed in this contribution (Supplementary Table 1a) may improve the efficacy of the model presented: in an attempt to explore this, additional blood analytes are being measured within AIBL. As longitudinal monitoring of the AIBL and other cohorts progress, indicators of the efficacy of such models in predicting progression to disease will be able to be examined. Additionally, further validation of the findings presented here for disease specificity in other disease cohorts is essential.

Estimates of NAB using CSF A β measurements have been reported⁴², but these assessments rely on obtaining CSF, a procedure considered invasive by many. Also, there are studies, reporting the efficacy of blood-based signatures in discriminating between AD and aged matched controls^{26-28, 33, 43}. To our knowledge, this contribution is the first study to report efficacy, validated in an independent separate cohort, for a (non-invasive) blood-based predictor of NAB. The clinical implications of fulfilling AD biomarker criteria²⁰ are crucial and especially relevant now that a more sensible paradigm has been proposed as the new diagnostic criteria for AD^{3, 44, 45} MCI⁴⁶, and preclinical AD³ in a model that integrates cognitive, biochemical and imaging biomarkers to provide a better predictive framework. With some further improvement in biomarker selection, the method described here will allow wide and non-invasive population screening, as well as participant selection for therapeutic trials that, after confirmation of high NAB by imaging or CSF, might also be used to follow up these individuals over time or monitor the efficacy of anti-A β therapy.

^bRembach A, Faux N, Watt A, *et al.* Changes in plasma β -amyloid in a longitudinal study of ageing and Alzheimer's disease. *Alzheimer's and Dementia*. In Press.

NAB has been demonstrated by many to be a good predictor for progression to AD^{3, 5, 22, 47-49}. Also biomarker signatures similar to the one presented here can easily be adapted to a low-cost routine test or rolled out as a high throughput population screen. Thus, this work may represent the first step in developing an economical screening tool for early detection of individuals at risk of developing AD, thus allowing earlier disease-specific therapeutic interventions aimed at halting or slowing down this devastating disease.

Acknowledgement Statement (including sponsorship and material support sources):

Core funding for the study was provided by the CSIRO Flagship Collaboration Fund and the Science and Industry Endowment Fund (SIEF) in partnership with Edith Cowan University (ECU), Mental Health Research institute (MHRI), Alzheimer's Australia (AA), National Ageing Research Institute (NARI), Austin Health, CogState Ltd., Hollywood Private Hospital, Sir Charles Gardner Hospital. The study also receives funding from the National Health and Medical Research Council (NHMRC), the Dementia Collaborative Research Centres program (DCRC), The McCusker Alzheimer's Research Foundation and Operational Infrastructure Support from the Government of Victoria. Faux NG is supported by, a National Health and Medical Research Council training fellowship. Laws SM is supported by research fellowships from Edith Cowan University. Bush AI is supported by the NHMRC by a program grant and an Australian Fellowship. The team wishes to thank the participants in AIBL for their commitment and dedication to helping advance research into the early detection and causation of AD and the clinicians who referred patients to the study.

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: Abbott; Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Amorfix Life Sciences Ltd.; AstraZeneca; Bayer HealthCare; BioClinica, Inc.; Biogen Idec Inc.; Bristol-Myers Squibb Company; Eisai Inc.; Elan Pharmaceuticals Inc.; Eli Lilly and Company; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; GE Healthcare; Innogenetics, N.V.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Medpace, Inc.; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Servier; Synarc Inc.; and Takeda Pharmaceutical Company. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University of California, San Diego. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of California, Los Angeles. This research was also supported by NIH grants P30 AG010129 and K01 AG030514.

Conflict of Interest Statement:

We declare that Colin L. Masters and Ashley I. Bush are consultants with Prana Biotechnology. Further, a patent has been filed covering the biomarker algorithm from this work and the author's institutions may benefit from commercialisation of this patent.

REFERENCES

1. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical-Diagnosis of Alzheimers-Disease - Report of the NINCDS-ADRDA Work Group under the Auspices of Department-of-Health-and-Human-Services Task-Force on Alzheimers-Disease. *Neurology* 1984; **34**(7): 939-944.
2. Masters CL, Cappai R, Barnham KJ, Villemagne VL. Molecular mechanisms for Alzheimer's disease: implications for neuroimaging and therapeutics. *Journal of Neurochemistry* 2006; **97**(6): 1700-1725.
3. Sperling RA, Aisen PS, Beckett LA, Bennett DA, Craft S, Fagan AM *et al.* Toward defining the preclinical stages of Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers & Dementia* 2011; **7**(3): 280-292.
4. Price JL, Morris JC. Tangles and plaques in nondemented aging and "preclinical" Alzheimer's disease. *Annals of Neurology* 1999; **45**(3): 358-368.
5. Rowe CC, Ellis KA, Rimajova M, Bourgeat P, Pike KE, Jones G *et al.* Amyloid imaging results from the Australian Imaging, Biomarkers and Lifestyle (AIBL) study of aging. *Neurobiology of Aging* 2010; **31**(8, Sp. Iss. SI): 1275-1283.
6. Blennow K. CSF biomarkers in Alzheimer's disease - use in clinical diagnosis and to monitor treatment effects. *European Neuropsychopharmacology* 2010; **20**: S159-S159.
7. Ellis KA, Bush AI, Darby D, De Fazio D, Foster J, Hudson P *et al.* The Australian Imaging, Biomarkers and Lifestyle (AIBL) study of aging: methodology and baseline characteristics of 1112 individuals recruited for a longitudinal study of Alzheimer's disease. *International Psychogeriatrics* 2009; **21**(4): 672-687.
8. Winblad B, Palmer K, Kivipelto M, Jelic V, Fratiglioni L, Wahlund LO *et al.* Mild cognitive impairment - beyond controversies, towards a consensus: report of the International Working Group on Mild Cognitive Impairment. *Journal of Internal Medicine* 2004; **256**(3): 240-246.
9. Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. Mild cognitive impairment - Clinical characterization and outcome. *Archives of Neurology* 1999; **56**(3): 303-308.
10. Mehta PD, Pirttila T, Mehta SP, Sersen EA, Aisen PS, Wisniewski HM. Plasma and cerebrospinal fluid levels of amyloid beta proteins 1-40 and 1-42 in Alzheimer disease. *Archives of Neurology* 2000; **57**(1): 100-105.
11. Mehta PD, Pirttila T, Patrick BA, Barshatzky M, Mehta SP. Amyloid beta protein 1-40 and 1-42 levels in matched cerebrospinal fluid and plasma from patients with Alzheimer disease. *Neuroscience Letters* 2001; **304**(1-2): 102-106.
12. Lopez OL, Kuller LH, Mehta PD, Becker JT, Gach HM, Sweet RA *et al.* Plasma amyloid levels and the risk of AD in normal subjects in the Cardiovascular Health Study. *Neurology* 2008; **70**(19): 1664-1671.

13. Lui JK, Laws SM, Li QX, Villemagne VL, Ames D, Brown B *et al.* Plasma Amyloid-beta as a Biomarker in Alzheimer's Disease: The AIBL Study of Aging. *Journal of Alzheimers Disease* 2010; **20**(4): 1233-1242.
14. Gupta VB, Laws SM, Villemagne VL, Ames D, Bush AI, Ellis KA *et al.* Plasma apolipoprotein E and Alzheimer disease risk The AIBL study of aging. *Neurology* 2011; **76**(12): 1091-1098.
15. Hixson JE, Vernier DT. Restriction Isotyping of Human Apolipoprotein-E by Gene Amplification and Cleavage with HHA1. *Journal of Lipid Research* 1990; **31**(3): 545-548.
16. Azur MJ, Stuart EA, Frangakis C, Leaf PJ. Multiple imputation by chained equations: what is it and how does it work? *International Journal of Methods in Psychiatric Research* 2011; **20**(1): 40-49.
17. Jagust WJ, Bandy D, Chen KW, Foster NL, Landau SM, Mathis CA *et al.* The Alzheimer's Disease Neuroimaging Initiative positron emission tomography core. *Alzheimers & Dementia* 2010; **6**(3): 221-229.
18. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N *et al.* Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 2002; **15**(1): 273-289.
19. Benjamini Y, Hochberg Y. Controlling The False Discovery Rate - A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-Methodological* 1995; **57**(1): 289-300.
20. Frank RA, Galasko D, Hampel H, Hardy J, de Leon MJ, Mehta PD *et al.* Biological markers for therapeutic trials in Alzheimer's disease - Proceedings of the biological markers working group; NIA initiative on neuroimaging in Alzheimer's disease. *Neurobiology of Aging* 2003; **24**(4): 521-536.
21. Mintun MA, LaRossa GN, Sheline YI, Dence CS, Lee SY, Mach RH *et al.* (11) PIB in a nondemented population - Potential antecedent marker of Alzheimer disease. *Neurology* 2006; **67**(3): 446-452.
22. Pike KE, Savage G, Villemagne VL, Ng S, Moss SA, Maruff P *et al.* beta-amyloid imaging and memory in non-demented individuals: evidence for preclinical Alzheimer's disease. *Brain* 2007; **130**: 2837-2844.
23. Aizenstein HJ, Nebes RD, Saxton JA, Price JC, Mathis CA, Tsopelas ND *et al.* Frequent Amyloid Deposition Without Significant Cognitive Impairment Among the Elderly. *Archives of Neurology* 2008; **65**(11): 1509-1517.
24. Villemagne VL, Pike KE, Darby D, Maruff P, Savage G, Ng S *et al.* A beta deposits in older non-demented individuals with cognitive decline are indicative of preclinical Alzheimer's disease. *Neuropsychologia* 2008; **46**(6): 1688-1697.
25. Blacker D, Haines JL, Rodes L, Terwedow H, Go RCP, Harrell LE *et al.* ApoE-4 and age at onset of Alzheimer's disease: The NIMH genetics initiative. *Neurology* 1997; **48**(1): 139-147.

26. Doecke JD, Laws SM, Faux N, Wilson W, Burnham SC, Lam C-P *et al.* Blood-based protein biomarkers for diagnosis of alzheimer disease. *Archives of Neurology* 2012; 1-8.
27. O'Bryant SE, Xiao G, Barber R, Huebinger R, Wilhelmsen K, Edwards M *et al.* A Blood-Based Screening Tool for Alzheimer's Disease That Spans Serum and Plasma: Findings from TARC and ADNI. *Plos One* 2011; **6**(12).
28. Soares HD, Potter WZ, Pickering E, Kuhn M, Immermann F, Shera D *et al.* Plasma biomarkers associated with the apolipoprotein e genotype and alzheimer disease. *Archives of Neurology* 2012; 1-8.
29. Johns CE, Newton JL, Westley BR, May FEB. Human pancreatic polypeptide has a marked diurnal rhythm that is affected by ageing and is associated with the gastric TFF2 circadian rhythm. *Peptides* 2006; **27**(6): 1341-1348.
30. Schmidt PT, Naslund E, Gryback P, Jacobsson H, Holst JJ, Hilsted L *et al.* A role for pancreatic polypeptide in the regulation of gastric emptying and short-term metabolic control. *Journal of Clinical Endocrinology & Metabolism* 2005; **90**(9): 5241-5246.
31. Druce MR, Small CJ, Bloom SR. Minireview: Gut peptides regulating satiety. *Endocrinology* 2004; **145**(6): 2660-2665.
32. Hallgren R, Lundqvist G. Elevated Levels of Circulating Pancreatic-Polypeptide in Inflammatory and Infectious Disorders. *Regulatory Peptides* 1980; **1**(3): 159-167.
33. Ray S, Britschgi M, Herbert C, Takeda-Uchimura Y, Boxer A, Blennow K *et al.* Classification and prediction of clinical Alzheimer's diagnosis based on plasma signaling proteins. *Nature Medicine* 2007; **13**: 1359-1362.
34. OMIM[®]. Online Mendelian Inheritance in Man. McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, MD), 2012.
35. Cosentino SA, Stern Y, Sokolov E, Scarmeas N, Manly JJ, Tang MX *et al.* Plasma beta-Amyloid and Cognitive Decline. *Archives of Neurology* 2010; **67**(12): 1485-1490.
36. Sundelof J, Giedraitis V, Irizarry MC, Sundstrom J, Ingelsson E, Ronnema E *et al.* Plasma beta amyloid and the risk of Alzheimer disease and dementia in elderly men. *Archives of Neurology* 2008; **65**(2): 256-263.
37. Laske C, Sopova K, Gkotsis C, Eschweiler GW, Straten G, Gawaz M *et al.* Amyloid-beta Peptides in Plasma and Cognitive Decline After 1 Year Follow-Up in Alzheimer's Disease Patients. *Journal of Alzheimers Disease* 2010; **21**(4): 1263-1269.
38. Mayeux R, Honig LS, Tang MX, Manly J, Stern Y, Schupf N *et al.* Plasma A beta 40 and A beta 42 and Alzheimer's disease - Relation to age, mortality, and risk. *Neurology* 2003; **61**(9): 1185-1190.

39. Mayeux R, Tang MX, Jacobs DM, Manly J, Bell K, Merchant C *et al.* Plasma amyloid beta-peptide 1-42 and incipient Alzheimer's disease. *Annals of Neurology* 1999; **46**(3): 412-416.
40. Toledo JB, Vanderstichele H, Figurski M, Aisen PS, Petersen RC, Weiner MW *et al.* Factors affecting A beta plasma levels and their utility as biomarkers in ADNI. *Acta Neuropathologica* 2011; **122**(4): 401-413.
41. Villemagne VL, Perez KA, Pike KE, Kok WM, Rowe CC, White AR *et al.* Blood-Borne Amyloid-beta Dimer Correlates with Clinical Markers of Alzheimer's Disease. *Journal of Neuroscience* 2010; **30**(18): 6315-6322.
42. Weigand SD, Vemuri P, Wiste HJ, Senjem ML, Pankratz VS, Aisen PS *et al.* Transforming cerebrospinal fluid Abeta42 measures into calculated Pittsburgh compound B units of brain Abeta amyloid. *Alzheimers Dement*, vol. 7. 2011 The Alzheimer's Association. Published by Elsevier Inc: United States, 2011, pp 133-141.
43. O'Bryant SE, Xiao G, Barber R, Reisch J, Doody R, Fairchild T *et al.* A Serum Protein-Based Algorithm for the Detection of Alzheimer Disease. *Archives of Neurology* 2010; **67**(9): 1077-1081.
44. Dubois B, Feldman HH, Jacova C, Cummings JL, DeKosky ST, Barberger-Gateau P *et al.* Revising the definition of Alzheimer's disease: a new lexicon. *Lancet Neurology* 2010; **9**(11): 1118-1127.
45. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Jr., Kawas CH *et al.* The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers & Dementia* 2011; **7**(3): 263-269.
46. Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC *et al.* The diagnosis of mild cognitive impairment due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers & Dementia* 2011; **7**(3).
47. Bourgeat P, Chetelat G, Villemagne VL, Fripp J, Raniga P, Pike K *et al.* beta-Amyloid burden in the temporal neocortex is related to hippocampal atrophy in elderly subjects without dementia. *Neurology* 2010; **74**(2): 121-127.
48. Morris JC, Roe CM, Mintun MA. Preclinical Alzheimer's Disease: Relationship with Brain/Cognitive Reserve. *Gerontologist* 2009; **49**: 145-145.
49. Jack CR, Knopman DS, Jagust WJ, Shaw LM, Aisen PS, Weiner MW *et al.* Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurology* 2010; **9**(1): 119-128.

TABLES

Table 1. The Demographic and Clinical make-up of the AIBL and ADNI imaged sub-cohorts. * Interquartile Range (IQR).

		High Neocortical SUVR (>1.3)	Low Neocortical SUVR (<1.3)	p-value
AIBL Imaged Cohort	Number of Participants: [N]	160	113	
	Clinical Diagnosis: [N (%)]	HC: 72 (45) MCI: 40 (25) AD: 48 (30)	HC: 97 (86) MCI: 15 (13) AD: 1 (1)	<0.001
	Age: [mean (SD)]	74.5 (7.7)	69.7 (7.2)	<0.001
	Gender; Males: [N (%)]	79 (49)	53 (47)	0.78
	Years of Education: [N (%)]	<9: 20 (13) 9-12: 63 (39) 13-15: 34 (21) >15: 43 (27)	<9: 8 (7) 9-12: 43 (38) 13-15: 19 (17) >15: 43 (38)	0.05
	CDR Sum of Boxes: [median (IQR*)]	0.5 (0.0 - 3.0)	0.0 (0.0 - 0.0)	<0.001
	<i>APOE</i> ϵ 4 carrier: [N (%)]	106 (66)	32 (28)	<0.001
ADNI Imaged Cohort	Number of Participants: [N]	59	23	
	Clinical Diagnosis: [N (%)]	HC: 0 (0) MCI: 42 (71) AD: 17 (29)	HC: 3 (13) MCI: 18 (78) AD: 2 (9)	<0.01
	Age: [mean (SD)]	78.9 (8.3)	78.4 (7.7)	0.78
	Gender; Males: [N (%)]	20 (34)	7 (30)	0.97
	Years of Education: [N (%)]	<9: 0 (0) 9-12: 10 (17) 13-15: 11 (19) >15: 38 (64)	<9: 0 (0) 9-12: 7 (30) 13-15: 1 (4) >15: 15 (65)	0.15
	CDR Sum of Boxes: [median (IQR*)]	2.0 (1.0 - 3.0)	1.0 (0.5 - 2.0)	0.03
	<i>APOE</i> ϵ 4 carrier: [N (%)]	37 (63)	6 (26)	<0.01

Table 2. The receiver operating characteristics for the efficacy of the cross-validated RF sub-models applied to both the AIBL and the ADNI imaged sub-cohorts

Model	Markers	AIBL Imaged Cohort			ADNI Imaged Cohort		
		Sens% (SD%)	Spec% (SD%)	AUC% (SD%)	Sens %	Spec %	AUC %
M1	Age, <i>APOE</i> Genotype, CXCL-13, IgM-1, IL-17, PPY, VCAM-1, A β ₁₋₄₂ , CDR sum of boxes	79.6 (1.3)	82.4 (1.2)	87.6 (0.7)	78.3	76.3	84.7
M2	Age, <i>APOE</i> Genotype, CXCL-13, IgM-1, IL-17, PPY, VCAM-1, A β ₁₋₄₂	79.6 (1.6)	79.4 (1.4)	83.9 (1.0)	73.9	74.6	81.7
M3	Age, CXCL-13, IgM-1, IL-17, PPY, VCAM-1, A β ₁₋₄₂	73.2 (1.8)	73.3 (1.8)	80.8 (1.7)	68.1	73.6	68.8
M4	Age, <i>APOE</i> Genotype, CDR sum of boxes	71.4 (1.1)	71.5 (1.1)	78.3 (0.8)	72.7	67.9	73.7
M5	Age, CDR sum of boxes	69.0 (2.1)	68.9 (2.2)	76.0 (2.0)	59.1	62.3	64.5
M6	Age, <i>APOE</i> Genotype	66.7 (1.6)	66.7 (1.6)	70.2 (1.3)	77.3	62.3	67.8

FIGURE LEGENDS

Figure 1. Univalent differences of blood analytes between the high and low NAB groups of the imaged AIBL sub-cohort. A: FT4; B: IgM; C: MIP 1 α ; D: PPY; E: VCAM 1. The horizontal line within the box represents the median, the lower and upper boundaries of the box represent the lower and upper quartiles, which define the interquartile range (IQR) and the whiskers are 1.5 times the IQR.

Figure 2. A: ROC curves for the cross-validated RF sub-models applied to the imaged AIBL sub-cohort; B: ROC curves for RF sub-models applied to the imaged ADNI sub-cohort. [Black = M1, Orange = M2, Blue = M3, Green = M4, Pink = M5, Grey = M6].

Figure 3. A: Actual SUVR values for the imaged AIBL sub-cohort split by Clinical Diagnosis; B: Predicted SUVR values for the non-imaged AIBL sub-cohort split by Clinical Diagnosis.





