

Edith Cowan University
Research Online

ECU Publications 2013

1-1-2013

Delivering Genetic Education and Genetic Counseling for Rare Diseases in Rural Brazil

A.X. Acosta

K Abe-Sandes

R Giugliani

Alan H. Bittles

Edith Cowan University, a.bittles@ecu.edu.au

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworks2013>

 Part of the [Genetics Commons](#), [Medical Education Commons](#), and the [Medical Genetics Commons](#)

[10.1007/s10897-013-9570-x](https://doi.org/10.1007/s10897-013-9570-x)

This is an Author's Accepted Manuscript of: Acosta, A., Abe-Sandes, K., Giugliani, R., & Bittles, A. H. (2013). Delivering Genetic Education and Genetic Counseling for Rare Diseases in Rural Brazil. *Journal of Genetic Counseling*, 22(6), 830-834. The final publication is available at Springer via [here](#)

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworks2013/768>

Journal of Genetic Counseling

Delivering genetic education and genetic counseling for rare diseases in rural Brazil --Manuscript Draft--

Manuscript Number:	
Article Type:	Genetic Counseling: A Global Perspective
Keywords:	Genetic education; genetic counseling; rare diseases; founder effect; endogamy; consanguinity; MPS VI
Corresponding Author:	Angelina Xavier Acosta, Ph.D. Federal University of Bahia Salvador, Bahia BRAZIL
First Author:	Angelina Xavier Acosta, Ph.D.
Order of Authors:	Angelina Xavier Acosta, Ph.D. Kiyoko Abé-Sandes Roberto Giugliani Alan H Bittles
Abstract:	<p>Brazil is the largest country in Latin America, with an ethnically diverse, Portuguese-speaking and predominantly Roman Catholic population of some 194 million. Universal health care is provided under the Federal Unified Health System (Sistema Único de Saúde) but, as in many other middle and low income countries, access to medical genetics services is limited in rural and remote regions of the country. Since there is no formally recognized Genetic Counseling profession, genetic counseling is provided by physicians, trained either in medical genetics or a related clinical discipline. A comprehensive medical genetics program has been established in Monte Santo, an inland rural community located in the state of Bahia in Northeast Brazil, with high prevalences of a number of recessive genetic disorders, including non-syndromic deafness, phenylketonuria, congenital hypothyroidism and mucopolysaccharidosis VI (Maroteaux-Lamy syndrome). Genetic education, counseling and treatment are locally provided, with a neonatal screening program for MPSVI currently under trial.</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Delivering genetic education and genetic counseling for rare diseases in rural Brazil

Acosta, A.X.^{1,2,3}, Abé-Sandes, K.^{2,4}, Giugliani, R.^{2,5} & Bittles, A.H.^{3,6}

¹Bahia School of Medicine, Federal University of Bahia, Salvador, BA, Brazil, ²National Institute on Population Medical Genetics – INAGEMP, ³Centre for Comparative Genomics, Murdoch University, Perth, Australia, ⁴Brazil State University of Bahia, Salvador, BA, Brazil, ⁵Department of Genetics, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil, ⁶School of Medical Sciences, Edith Cowan University, Perth, Australia

Running title: Genetic education and genetic counseling in Brazil

Key words: Genetic education, genetic counseling, rare diseases, founder effect, endogamy, consanguinity, MPS VI

All correspondence to:

Dr. A.X. Acosta
Hospital Universitário Professor Edgar Santos
Serviço de Genética Médica
Av Augusto Viana, s/n, 6^a andar, Canela
Salvador-Bahia, Brasil
CEP 40080-160

Tel: +55 71 32838109
Fax: +55 71 32828108
Email: axacosta@hotmail.com

1
2
3
4 **Abstract**
5

6
7 Brazil is the largest country in Latin America, with an ethnically diverse, Portuguese-
8
9 speaking and predominantly Roman Catholic population of some 194 million. Universal health
10
11 care is provided under the Federal Unified Health System (*Sistema Único de Saúde*) but, as in
12
13 many other middle and low income countries, access to medical genetics services is limited in
14
15 rural and remote regions of the country. Since there is no formally recognized Genetic
16
17 Counseling profession, genetic counseling is provided by physicians, trained either in medical
18
19 genetics or a related clinical discipline. A comprehensive medical genetics program has been
20
21 established in Monte Santo, an inland rural community located in the state of Bahia in Northeast
22
23 Brazil, with high prevalences of a number of recessive genetic disorders, including non-
24
25 syndromic deafness, phenylketonuria, congenital hypothyroidism and mucopolysaccharidosis VI
26
27 (Maroteaux-Lamy syndrome). Genetic education, counseling and treatment are locally provided,
28
29
30
31 with a neonatal screening program for MPSVI currently under trial.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **Introduction**
5

6 By global standards Brazil is a huge country with a land area of 8.54 million km² and a
7 population of 194.3 million, which makes it both the largest country in Latin America and the
8 fifth most populous country world-wide. As summarized in Table I, in economic and health
9 terms Brazil is a middle ranked country with an adult literacy rate of 88.6%, a mean annual per
10 capita income of US\$11,000, and 84% of its population living in urban centres. The introduction
11 of birth control programs by the Federal Government in the 1990s resulted in a marked slowing
12 of the annual population growth rate, and the total fertility rate is now 1.9, i.e. below the
13 replacement rate. However, in terms of its age profile Brazil remains a young country, with 25%
14 of the population under 15 years and just 7% over 65 years (PRB, 2012).
15
16
17
18
19
20
21
22
23
24
25
26
27

28 **Health care in Brazil**
29

30 The majority of the population is served by the public Unified Health System (*Sistema*
31 *Único de Saúde* or SUS), which was established in 1988 to ensure equitable access to health for
32 all citizens and is one of the largest public health systems in the world (Brazil, 1990). SUS is
33 organized on a municipal basis, with management provided at state level and financial support
34 from the Federal Ministry of Health. Health expenditure in Brazil in 2007 comprised 8.4% of
35 Gross Domestic Product (WHO, 2010), with 1.84 medical practitioners and 2.26 hospital beds
36 per 1,000 of the population (IBGE, 2009).
37
38
39
40
41
42
43
44
45
46
47

48 The most recent available national statistics indicate neonatal, infant and under-5 year
49 deaths rates of 11/1,000, 20/1,000 and 22/1,000 respectively, while non-communicable diseases
50 now account for 74% of deaths (Table I). The shift to a non-communicable disease profile in
51 Brazil also is reflected in the reported causes of early deaths. In 1980, 38% of infant deaths were
52 ascribed to perinatal causes, with congenital anomalies listed as the fifth most common cause of
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 death and representing just 5% of the total. Given proportional reductions in infectious,
5
6 nutritional, and respiratory diseases, by 2000 congenital malformations had become the second
7
8 most common cause of deaths in children under age 1 year representing 13% of the total, and by
9
10 2006 mortality due to congenital defects had risen to 16.3% of all infant deaths (Brazil, 2010)

11
12
13
14 [Table I here]

15 16 **Ethnicity of the Brazilian population**

17
18 A question on ethnicity was included in the 2000 Census of Brazil, with the options of
19
20 White, Black, *Pardo* (mixed color or race; mulatto or *mestizo*), Asiatic, and Native Brazilian
21
22 Indian. Of those who responded, 54.0% self-declared as White, while 39.9% were *Pardos*, 5.4%
23
24 Black, 0.5% Asiatic, and 0.2% Native Indians. Persons of African ancestry, represented by those
25
26 who self-declared as Black or *Pardo* were mainly resident in the North (69.0%) and Northeast
27
28 (65.8%), whereas the highest percentages of Whites were located in the Southeast (62.4%) and
29
30 South (83.6%) regions of the country (IBGE, 2000).

31 32 33 **Medical genetics services in Brazil**

34
35
36 Most Brazilians do not have access to medical genetic services, as the majority of these
37
38 services are concentrated in tertiary centers located in the most developed Southeast and South
39
40 regions of the country. A number of institutions have sophisticated laboratory infrastructures
41
42 and offer diagnostic testing, including cytogenetic and molecular genetic analyses. Specialists in
43
44 medical genetics tend to aggregate within the major service centers located in the Center-South
45
46 of the country, which in practice means that a majority of Brazilian states have few or no resident
47
48 trained professionals in clinical genetics (Horovitz et al., 2012).

49
50
51 Since access to specialist medical genetics services has been limited, the establishment of
52
53 regional services and the promotion of genetics within primary care were identified as priority
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 issues (Marques-de-Faria et al., 2004; Penchaszadeh, 2004). In response, the Federal
5
6 government formulated a national policy for comprehensive clinical genetics services, which
7
8 included both specialist medical genetics centers and the inclusion of genetics in primary care
9
10 programs (Brazil, 2009). In the latter setting the policy provides for the identification and
11
12 follow-up of families with congenital anomalies and genetically determined diseases, but as yet
13
14 progress in meeting its stated aims has been slow.
15
16
17

18 **Medical genetics training**

19
20
21 By comparison with other clinical specialities, Medical Genetics is a relatively young
22
23 branch of medicine in Brazil with Board Certification in Medical Genetics the responsibility of
24
25 the Brazilian Society of Medical Genetics (Sociedade Brasileira de Genética Médica, SBGM).
26
27 Eleven postgraduate residency programs each of three years' duration have been approved by the
28
29 National Medical Residency Committee (Comissão Nacional de Residência Médica, 2006), all
30
31 co-located with specialist laboratory services in the Center-South of the country (Secretaria de
32
33 Educação Superior, 2009). Professional accreditation in Medical Genetics also can be awarded
34
35 by other medical specialties, e.g. pediatrics, subject to approval by the Brazilian Medical
36
37 Association and the Federal Council of Medicine. During the last 30 years a combined total of
38
39 some 200 physicians have been awarded Board Certification in Medical Genetics through one or
40
41 other of these systems.
42
43
44
45
46
47

48 *Genetic counseling*

49
50 Although health professionals from many backgrounds are involved in providing genetic
51
52 services in Brazil, the profession of Genetic Counselor is not formally recognized and so the
53
54 provision of genetic counseling is almost exclusively restricted to physicians. However, many
55
56 groups, centers and services coordinated by physicians offer informal, non-accredited training
57
58
59
60
61
62
63
64
65

1
2
3
4 programs in genetic counseling to health care professionals drawn from a range of different
5
6 health disciplines, mainly nursing and psychology (Brunoni, 2002). Prior to a medical genetics
7
8 consultation, clients may be interviewed by a non-physician, most commonly health
9
10 professionals with a psychology background, who provide information on the investigations that
11
12 may be undertaken as part of their diagnosis and treatment.
13
14

15 16 **Genetic education and genetic counseling for inherited disorders in a rural setting** 17

18
19 Given the concentration of medical genetics resources in the Center-South of the country,
20
21 specific problems in dealing with inherited disorders in less privileged rural communities
22
23 commonly are encountered. As an example, the inland municipality of Monte Santo is located in
24
25 the state of Bahia, northeast Brazil, some 350 km from Salvador the state capital. According to
26
27 the 2010 Census of Brazil the total population of Monte Santo was 52,338, scattered across some
28
29 200 villages.
30
31

32
33 By comparison with the national average Monte Santo is a poor community, with an
34
35 average annual per capita income of just US\$1,403 and 66% adult literacy, and it is served by a
36
37 single general hospital with 60 beds (IBGE, 2010). Monte Santo was established mainly by
38
39 Portuguese settlers at the end of the 18th century but with substantial Native Indian Amerindian
40
41 and African admixture. Therefore in the 2010 Census respondents self-identified as 37.6%
42
43 White, 57.2% *Pardo*, and 5.2% Black (IBGE, 2010).
44
45
46
47

48 A cluster of cases of mucopolysaccharidosis VI (MPS VI, Maroteaux-Lamy syndrome)
49
50 was provisionally identified in Monte Santo, but there was no information on the prevalence of
51
52 the disorder, which in other parts of the world is regarded as rare with reported incidence rates
53
54 ranging from 1/43,261 in Turkey to 1/1.51 million births in Sweden (Valayannopoulos et al.,
55
56 2010). To investigate the feasibility of establishing genetic services in this generally under-
57
58
59
60
61
62
63
64
65

1
2
3
4 provided community, in 2006 a preliminary scoping visit to Monte Santo was arranged to meet
5
6 with the local civil authorities, medical staff and local residents. Following this visit a
7
8 multidisciplinary project *Genetics in the Sertão* was initiated as a partnership between public and
9
10 private institutions. The aim of the project was to create a coordinated genetic education, genetic
11
12 counseling, treatment and screening program for inherited disorders in the community, with
13
14 oversight provided by a multidisciplinary team of health care professionals, including medical
15
16 geneticists.
17
18
19
20

21 It soon became obvious that the prevalence of MPS VI in Monte Santo was indeed high
22
23 with 13 living cases identified in 11 nuclear families, and genetic analysis showed that all
24
25 affected persons were homozygous for the p.H178L mutation in the arylsulfatase B (ARSB) gene
26
27 (Costa-Motta et al., 2011). The data therefore strongly suggested the expression of a founder
28
29 mutation, maintained and spread through intra-community and intra-familial marriage and
30
31 resulting in a present-day MPS VI prevalence of 1/5,000. On the basis of these results, and with
32
33 the knowledge that other inherited disorders, including non-syndromic deafness, phenylketonuria
34
35 and congenital hypothyroidism, appeared to have an increased prevalence in the Monte Santo
36
37 community, the *Genetics in the Sertão* project was established as an ongoing, community-based
38
39 program.
40
41
42
43
44

45 *Genetic education*

46
47
48 In Brazil, the Community Health Agents Program was established in 1991 as part of the
49
50 Family Health Program. It is seen as an important facet of primary health care strategy, with
51
52 community health agents (CHA) trained to identify, refer, guide and follow-up families in their
53
54 community (Brazil, 2000). As part of the *Genetics in the Sertão* initiative in Monte Santo,
55
56 lectures, workshops and assessments of their training are regularly organized for the CHA, with
57
58
59
60
61
62
63
64
65

1
2
3
4 information on the diseases thought to be most prevalent in the community, reinforcement of the
5
6 importance of early diagnosis, instruction on eliciting relevant information from families, and
7
8 accessing medical advice and appropriate treatment. As community members, the involvement
9
10 of the CHA is fundamental to the success of the *Genetics in the Sertão* program, by continuously
11
12 monitoring the overall health of the population and identifying new disease cases.
13
14

15 16 *Genetic counseling*

17
18
19 General information on the effects of a disorder such as MPS VI is initially provided
20
21 through lectures and workshops conducted at community level, to explain the nature of the
22
23 disease, and the possible implications of family testing. In families with an affected member
24
25 multi-generation pedigrees are constructed, subject to the prior approval of family members and
26
27 on the basis of the information they provide.
28
29

30
31 Because of the highly endogamous nature of the Monte Santo population, individuals
32
33 thought to be carriers of the causative ARSB p.H178L mutation in MPS VI often appear in
34
35 several pedigrees of nuclear families. A blood sample is requested for DNA testing where
36
37 pedigree analysis suggests that an individual may be at 50% or 25% risk of being a carrier for the
38
39 p.H178L mutation, along with a sample from his/her partner and any children over 18 years of
40
41 age. Before such a test is offered its possible implications are again explained, and formal
42
43 permission to sample is separately obtained from each individual.
44
45
46

47 48 *Treatment and neonatal screening*

49
50 In the initial stages of the program, patients with MPS VI had to travel weekly to Salvador, the
51
52 capital of Bahia, to receive enzyme replacement infusions (ERT) with Galsulfase, the only
53
54 specific treatment for MPS VI, which necessitated a round trip of 700 km. However, in April
55
56 2011 an ERT center for MPS VI was inaugurated at the municipal hospital of Monte Santo,
57
58
59
60
61
62
63
64
65

1
2
3
4 where local doctors and nurses have been trained in the provision of treatment, with a medical
5
6 geneticist in regular attendance. As early introduction of therapy seems to provide a better
7
8 prognosis (McGill et al., 2010), a screening program for MPS VI was recently established for
9
10 newborns in Monte Santo.
11
12

13 14 **Conclusions**

15
16 The experience gained in the *Genetics in the Sertão* program has proved invaluable to the
17
18 population Monte Santo, where in the past severe inherited disorders such as MPS VI were
19
20 untreatable and represented a major health burden not only to families with affected members but
21
22 to the community as a whole. In more general terms, *Genetics in the Sertão* has demonstrated
23
24 the feasibility of establishing such a program in an under-privileged rural population, a lesson
25
26 which given community support is applicable to other similar communities across Latin America
27
28 and throughout the developing and middle income world.
29
30
31
32
33

34 35 **Acknowledgements**

36
37 The authors gratefully acknowledge the financial support provided by INAGEMP -
38
39 National Institute of Population Medical Genetics (grant CNPq 573993/2008-4), and the
40
41 Foundation of Research Support of the State of Bahia (FAPESB). Dr. A.X. Acosta was the
42
43 recipient of a National Council of Technological and Scientific Development (CNPq)
44
45 Scholarship, held as Visiting Researcher in the Centre for Comparative Genomics, Murdoch
46
47 University, Australia.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

Brazil (1990). Presidência da República. Lei no. 8.080 de 19 de setembro de 1990. Dispõe sobre as condições para a promoção, proteção e recuperação da saúde, a organização e o funcionamento dos serviços correspondentes e dá outras providências.

Brazil (2000). Ministério da Saúde. A implantação da unidade de saúde da família. Brasília, DF: Departamento de Atenção Básica, Ministério da Saúde. 44p. (cadernos de Atenção Básica, 1).

Brazil (2009). Ministério da Saúde, portaria GM no. 81, 20 de janeiro de 2009—Institui, no âmbito do Sistema Único de Saúde (SUS), a Política Nacional de Atenção Integral em Genética Clínica. Diário Oficial da União 21/01/2009.

Brazil (2010). Ministério da Saúde, Departamento de Informação e Informática do SUS-DATASUS, Sistema de Informações de Saúde. Mortalidade infantil (menores de um ano). Available at: <http://tabnet.datasus.gov.br/cgi/defthtm.exe?sim/cnv/obt10uf.def>

Brazil (2011). DATASUS—Health Information Database. Ministério da Saúde—Departamento de Informática do SUS. Available at: <http://www2.datasus.gov.br/DATASUS/index.php>

Brunoni, D. (2002). Genetic Counseling. *Ciência e Saúde Coletiva*, 7, 101-107.

Comissão Nacional de Residência Médica (2006). Resolução CNRM no. 02, de 15 de maio de 2006. Dispõe sobre requisitos mínimos dos Programas de Residência Médica e dá outras providências. Available at:

http://portal.mec.gov.br/sesu/arquivos/pdf/cnrm/resolcnrm002_2006.pdf

Costa-Motta, F.M.M., Acosta, A.X., Abé-Sandes, K., Bender, F., Schwartz, I.V.D., Giugliani, R., & Leistner-Segal, S. (2011). Genetic studies in a cluster of mucopolysaccharidosis type VI patients in Northeast Brazil. *Molecular Genetics and Metabolism*, 104, 603-607.

1
2
3
4 Horovitz, D.D.G., Ferraz, V.E.F., Dain, S., & Marques-de-Faria, A.P. (2102). Genetic services
5 and testing in Brazil. *Journal of Community Genetics* Published online: 05 May 2012. DOI
6
7 10.1007/s12687-012-0096-y.
8
9
10
11 IBGE (2009). Fundação Instituto Brasileiro de Geografia e Estatística. Estatísticas da saúde:
12 assistência médico-sanitária 2009. Rio de Janeiro; 2010 Available at:
13
14 [www.ibge.gov.br/home/estatistica/ populacao/condicaoodevida/ams/2009/ams2009.pdf](http://www.ibge.gov.br/home/estatistica/populacao/condicaoodevida/ams/2009/ams2009.pdf)
15
16
17
18 IBGE (2000). Instituto Brasileiro de Geografia e Estatística - Brasil: 500 Anos de Povoamento.
19 Rio de Janeiro. Available at: <http://www.ibge.gov.br>.
20
21
22
23 IBGE (2010). Fundação Instituto Brasileiro de Geografia e Estatística; Brasil em Síntese.
24 Available at: [http://www.ibge.gov.br/brasil_em_sintese/ default.htm](http://www.ibge.gov.br/brasil_em_sintese/default.htm)
25
26
27
28 IBGE (2011). Fundação Instituto Brasileiro de Geografia e Estatística— Sinopse do Censo
29 Demográfico 2010. Rio de Janeiro, 2011. Available at:
30
31 <http://www.ibge.gov.br/home/estatistica/populacao/ censo2010/sinopse.pdf>
32
33
34
35 McGill, J.J., Inwood, A.C., Coman, D.J., Lipke, M.L., de Lore, D., Swiedler, S.J., Hopwood, J.J.
36 (2010). Enzyme replacement therapy for mucopolysaccharidosis VI from 8 weeks of age—a
37 sibling control study. *Clinical Genetics*, 77, 492–498.
38
39
40
41
42 Marques-de-Faria, A.P., Ferraz, V.F.E., Acosta, A.X., & Brunoni, D. (2004). Clinical genetics in
43 developing countries: the case of Brazil. *Community Genetics*, 7, 95–105.
44
45
46
47 Penchaszadeh, V.B. (2004). Genetics Services in Latin America. *Community Genetics*, 7, 65-69.
48
49
50 PRB (2012). *World Population Data Sheet*. Population Reference Bureau, Washington DC.
51
52
53 Secretaria de Educação Superior—Comissão Nacional de Residência Médica—Ministério da
54 Educação (2009). Sistema CNRM—Instituição x Programas x Vagas—Programas de Residência
55 em Genética Médica. Available at:
56
57
58
59
60
61
62
63
64
65

1
2
3
4 http://mecsrv04.mec.gov.br/sesu/SIST_CNRM/APPS/inst_especialidades.asp
5

6 UNICEF (2012). *State of the World's Children 2012*. New York, United Nations Children's
7
8 Fund.
9

10 Valayannopoulos, V., Nicely, H., Harmatz, P., & Turbeville S. (2010). Mucopolysaccharidosis
11
12 VI. *Orphanet Journal of Rare Diseases*, 5, 5.
13
14

15 WHO (2010). *World Health Organization. World Health Statistics 2010*. Available at:
16
17 www.who.int/whosis/whostat/EN_WHS10_Full.pdf.
18
19

20 WHO (2011). *Noncommunicable Diseases Country Profiles 2011*. Geneva, World Health
21
22 Organization.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table I

Brazil: Demographic and health summary profile 2012

Population	194.3 million
Urban, %	84
Mean income per person	US\$11,000
Total fertility rate	1.9
Life expectancy at birth (yr):	
Male	70.3
Female	77.1
Neonatal mortality rate	11/1,000
Infant mortality rate	20/1,000
Under 5 mortality rate	22/1,000
Children <5yr under-weight, %	2
Maternal mortality rate	68.7/100,000
Deaths due to non-communicable diseases, %	74

Sources: Brazil (2011); IBGE (2009, 2010); PRB (2012); UNICEF (2012); WHO (2010, 2011).