

## **DIVERSITY IN SOUND PRESSURE LEVELS AND ESTIMATED ACTIVE SPACE OF RESIDENT KILLER WHALE VOCALIZATIONS**

Patrick J. O. Miller<sup>1,2</sup>

<sup>1</sup>Biology Department, Woods Hole Oceanographic Institution, Woods Hole, MA 02543 USA

<sup>2</sup>NERC Sea Mammal Research Unit, University of Saint Andrews, St. Andrews, Fife, KY16  
9QQ, UK.

Email: pm29@st-andrews.ac.uk

Phone: (44) 01334-462-658

Fax: (44) 01334-462-632

Please use second address for correspondence.

Running title: Source levels and active space of killer whale sounds ...

Text word count: 5631

**Abstract** (179 words)

Signal source intensity and detection range, which integrates source intensity with propagation loss, background noise and receiver hearing abilities, are important characteristics of communication signals. Apparent source levels were calculated for 819 pulsed calls and 24 whistles produced by free-ranging resident killer whales by triangulating the angles-of-arrival of sounds on two beamforming arrays towed in series. Levels in the 1-20 kHz band ranged from 131-168 dB re 1 $\mu$ Pa @1m, with differences in the means of different sound classes (whistles:  $140.2 \pm 4.1$  dB; variable calls:  $146.6 \pm 6.6$  dB; stereotyped calls:  $152.6 \pm 5.9$  dB), and among stereotyped call types. Repertoire diversity carried through to estimates of active space, with “long-range” stereotyped calls all containing overlapping, independently-modulated high-frequency components (mean estimated active space of 10-16km in sea state zero) and “short-range” sounds (5-9 km) included all stereotyped calls without a high-frequency component, whistles, and variable calls. Short-range sounds are reported to be more common during social and resting behaviors, while long-range stereotyped calls predominate in dispersed travel and foraging behaviors. These results suggest that variability in sound pressure levels may reflect diverse social and ecological functions of the acoustic repertoire of killer whales.

Keywords: communication, ecology, whistle, pulsed call, repertoire

## Introduction

The amplitude at which a signal is produced, or source level, is of fundamental importance in communication systems, and acoustic signals have the potential to be produced over a wide range of levels (Loftus-Hills & Littlejohn 1971; Gerhardt 1975; Prestwich et al. 1989; Cynx et al. 1998; Bradbury & Vehrencamp 1998; Brumm 2004). For a signaler to benefit from an information transfer, it must produce signals at a sufficient level for an intended receiver to detect and decode the signal. However, signal levels attenuate during transmission from the source to the receiver, which must detect the signal in the presence of competing background noise. The benefits of information transfer, therefore, select for higher amplitude signals because they are more likely to be detected and decoded by an intended receiver (Wiley & Richards 1982). Energetic costs and the increased risk of detection by predators, prey, or social competitors are factors that may select for less intense signals (Prestwich et al. 1989; Ryan et al. 1982; Dabelsteen & Pedersen 1990; McGregor & Dabelsteen 1996).

Apparently in response to these balancing selective pressures, signals appear to be produced at the appropriate intensity for typical distances to target or intended receivers (Wiley & Richards 1982). Certain signal types within a repertoire may be selected to cover a large area while others function over short ranges, reflecting the context and motivation of the signaler (Rowell and Hinde 1962; Dittus 1988; Marten et al. 1977). Blackbirds (*Turdus merula*) switch from loud song to a quieter version of song in close range threat interactions with intruders (Dabelsteen & Pedersen 1990). Several recent studies have also shown that songbirds and non-human primates do not routinely maximize the amplitude of their songs, but adjust the source level in response to background noise (Sinnot et al. 1975; Cynx et al. 1988; Brumm & Todt 2002; Pytte et al. 2003; Brumm et al. 2004) and social competitors (Brumm & Todt 2004).

The range over which a signal can be detected, or 'active space', is not solely a function of its source level, but also the transmission loss the signal suffers between the signaler and the receiver as well as the detection threshold of the receiver in varying levels of background noise. Therefore, variation in source level cannot be assumed to directly result in a matching shift in active space. Techniques for estimating active space have been used for the territorial song display of the red-wing blackbird (*Agelaius phoeniceus*), which is produced at a level sufficient to be heard by neighboring red-wing blackbirds over a range of about two territories away (Brenowitz 1982). The active space of bottlenose dolphin (*Tursiops truncatus*) whistles is estimated to be much larger than that of bird song, partly due to reduced sound absorption in seawater (Janik 2000). Efficient acoustic signal propagation increases the potential for large communication networks to exist in the marine environment (Myrberg 1980; Janik in press).

The goal of this study was to measure the sound pressure levels of pulsed calls and whistles produced by free-ranging resident killer whales, and estimate the maximum range they might be detected by conspecifics in different background noise conditions. Estimating active space links physiological measures of sound production and reception with acoustical ecology and habitat to quantitatively predict the distances over which social communication might take place. Because the active space of a signal relates to its function, analysis of source level and/or active space can be a useful tool to elucidate the function of signals in a repertoire (e.g. Marler 1973; Dittus 1988).

The vocal repertoire of killer whales consist of various burst-pulse calls and whistles (Schevill & Watkins 1966; Watkins 1967; Ford 1989; Ford 1991; Thomsen et al. 2001). The evidence is mixed whether the various signal types in the repertoire or resident killer whales reflect the activities or motivations of signalers, as is documented for many primate sounds (Dittus 1988; Hauser & Marler 1993). No whistle or call type correlates exclusively with any

particular activity, but relative production rates of different call types and whistles vary with broad behavior metrics of the entire group (Bain 1986; Morton et al. 1986; Ford 1989; Thomsen et al. 2002). If the various sound types in the killer whale repertoire primarily serve a contact function of signaling group membership, as proposed by Ford (1989, 1991), one might expect little variation in the estimated active space of different sound types in the repertoire. Conversely, if sound-type production reflects the behavioral context or motivation of the signaler (Bain 1986; Morton 1986; Ford 1989; Thomsen et al. 2002), one may expect to find diversity in the estimated active space of types depending on the context or behavioral state in which they tend to be produced.

To measure the source intensity of a sound, it is necessary to measure the received sound pressure level of a signal and estimate its transmission loss from the source, which (at short ranges) is primarily a function of range. An established approach is to use multiple hydrophones to measure the range from the source to a calibrated receiver (Watkins & Schevill 1974; Watkins et al. 1997; Cato 1998; Mohl et al. 2000; Janik 2000). With this technique, source levels can be measured during normal behavior, although it can be difficult to observe the orientation of underwater signalers to measure to control for signal directionality. Killer whale calls are directional at higher frequencies (Miller 2002; Schevill & Watkins 1966; Bain & Dalheim 1994), so the received level of a sound depends on the orientation of the signaler. The term “apparent source level”, used here, refers to source levels measured when signaler orientation is not known (Madsen et al. 2004). To measure apparent source levels of weak signals, the hydrophones must be positioned near the signaler and have a small inter-hydrophone spacing (e.g. Watkins et al. 1997). Such a small array, if immobile, will have a small coverage area through which animals could pass quite quickly. Larger arrays allow greater spatial coverage, but can only localize relatively intense sounds (e.g. Mohl et al. 2000). In such a case, mean source level represents only a mean of relatively

intense sounds, depending on the array geometry (e.g. Janik 2000). To statistically examine the apparent source levels and estimated active space of sounds over a wide range of levels, I employed a novel towed two-beamformer array design that can be towed near animals, allowing a large 'coverage' area with a small (97m) aperture array.

### **Materials and methods**

Recordings were conducted in Johnstone Strait on September 1 and 7, 1998 while killer whales from several different pods were present. All recordings were taken in low noise conditions: sea state 0 with no other vessels within a kilometer of the animals. The recording system consisted of two evenly spaced 8-element (2.6 m) beamforming arrays that were towed in series with an inter-array spacing of 97m (Fig. 1). The arrays were towed parallel to the direction of movement of the whales at minimum speed (~1.5 m/s) to minimize noise generated by the ship. The signal from each hydrophone was recorded on a channel of two synchronized 16-bit Tascam 8-track digital recorders.

Data from the recorders was digitally transferred as wav files into computer using a precision conversion system that maintains channel synchronization on a sample-by-sample basis across both recorders. Signal analysis was carried out using custom Matlab software (see Miller & Tyack 1998 for details). The analog-signal to wav-file conversion was calibrated using tones measured on a calibrated LeCroy 9314AM oscilloscope. The hydrophone elements were Benthos AQ-2S hydrophones (nominal sensitivity:  $-195\text{dBV}/\mu\text{Pa}$ ) attached to a custom WHOI preamp (43 dB gain flat from 100Hz to 200 kHz).

The sensitivity of one hydrophone in the rear array was calibrated in-situ using a Brüel & Kjær (B&K) 8105 hydrophone and a B&K 2635 preamp set at a sensitivity of  $-160\text{dBV}/\mu\text{Pa}$ . The B&K hydrophone was attached to the array next to one element, and several loud sources were used to measure the relative sensitivity of the array element to the

calibrated hydrophone. The relative sensitivities of the remaining elements were measured using a transient impact sound containing broadband energy from 1-20kHz produced over 100m from the array. The resulting measured sensitivity of  $-152.7$  dBV/ $\mu$ Pa was consistent across hydrophones and closely matched the expected sensitivity based upon separate calibration of the pre-amplifiers and the nominal sensitivity of the hydrophones.

All whale sounds with a received sound pressure level at least 3 dB above background noise on the rear array were extracted from the recordings and a calibrated received spectrum was calculated from 1 to 20 kHz in 1/3-octave bands. One-third octave bands were used because they represent the logarithmic shape of auditory filters in the mammalian ear (Fletcher 1940). Calibrated spectra were calculated from the most intense subsection of each sound, and portions of the sound that were overlapped by extraneous sounds (such as echolocation clicks) or that were attenuated by surface diffraction were not included. The subsection used for the analysis was always greater than 100ms in duration. The +3dB criteria was implemented by assessing the background noise level just prior to the signal onset, and discarding signals where signal plus noise levels were <5dB greater than noise levels alone. The low-frequency cutoff of 1 kHz was chosen because killer whale hearing is much less sensitive below 1 kHz (Hall & Johnson 1972; Symanski et al. 1999), there is little energy in killer whale sounds below 1 kHz (Fig.1; Ford 1987), and it is desirable to eliminate the low-frequency noise below 1 kHz in calculating RL. The high-frequency cutoff of 20 kHz bounded a 1/3-octave band and was chosen to reduce the effects of filter roll-off near the Nyquist frequency of 24 kHz.

Sounds were classified as whistles or calls based upon their harmonic structure (Watkins 1967). Calls were further classified to call type (Fig. 2; Ford 1987; 1991; Miller & Bain 2000). Any calls that did not clearly fall into one of the types described by Ford (1987) were classified as 'variable' calls.

### *Source Level Calculation*

The source level (SL) in dB re 1 $\mu$ Pa @ 1m of each sound was calculated as:

$$SL = RL + 20\log(R) - k$$

where RL is the received level in dB, R is the measured range from the vocalizing whale to the center of the rear array, and k is the near-field anomaly which represents the mean contribution to the field of the multiple bottom and surface reflections. A calibrated source spectrum was calculated for each 1/3-octave band, which was summed to calculate the broadband source level.

This simple transmission loss equation is valid when range is less than water depth and short enough that frequency-dependent absorption can be ignored (Urlick 1981; Marsh & Shulkin 1962). In sea-state zero, k is estimated by Marsh and Shulkin (1962) to be 5.4dB at 2.0kHz and 3.8dB at 8.0kHz (which is the primary energy band in killer whale calls). Since the ranges used here are only the shortest of the possible valid range (out to 400m rather than to 3000m), the proportion of energy in the reflections versus direct path should be corrected to be somewhat less than the typical value given by Marsh and Shulkin. Thus, this study set the value of k to be equal to 3dB.

The -3 dB value reflects the expectation that transmission loss will not be that fully predicted by spherical spreading. This value specifically subtracts non-correlated broadband energy that arrives at the receiving hydrophone after being reflected from the sea surface (Charif et al. 2002) assuming that the path lengths of the direct and surface paths are roughly identical, that reflection from the surface is lossless, and that the reflection will not destructively or constructively interfere with the direct path. Frequency-specific interference was clearly apparent in portions of some of the analyzed signals, and such portions of the sounds were specifically not included in the source level calculation. While the transmission-loss equation used to calculate apparent source level was probably not exact for all sounds



analyzed, it is unlikely to be strongly biased and should be within  $\sim 3$ dB for sounds measured at the ranges reported here.

Range from the rear array to the vocalizing animal was calculated by intersecting the angles-of-arrival of the sound on the two in-series beamforming arrays (Fig. 1). The angle-of-arrival was calculated for each array using a broadband frequency-domain beamforming algorithm that measures the amount of energy coming from each direction at each frequency (Miller & Tyack 1998; Thode et al. 2000b). This technique does not obtain a 3-D localization of a sound because the linear arrays cannot resolve the left-right (and depth) ambiguity. However, the range estimate is valid for any position or depth of the signaler. With an inter-array spacing of 97m, 0 and +3 dB signal-to-noise ratio requirements for the front and rear arrays, respectively, and background noise near 90 dB in the 1-20kHz band, the lower limit of apparent source levels measurable by this system is roughly 127 dB @1m. At the greatest range from the rear array from which sounds were analyzed (410m), sounds with an apparent source level below 142 dB @1m would have insufficient SNR to be analyzed, and may be underrepresented in the sample.

The accuracy of the range estimate was calibrated using a sound source deployed from a stationary boat. Several simultaneous acoustic and visual fixes were taken as the two-array system was driven past the sound source. Visual fixes were taken using a laser-range finder and a hand-held compass as well as by ship's radar. At each of these points the range error was calculated and expressed as the source level error that would result from the range error. Calibration results of 8 separate passes showed that range errors contribute less than  $\pm 3$ dB error to the source level estimate over a 500m x 200m body of water on both sides of the array (Fig. 3). Animal sounds that came from outside the calibrated region were excluded from analysis. Because range errors should not be correlated with errors in transmission loss, the apparent source levels presented here should typically be accurate to within  $\pm 3$ dB.

### *Active space estimation*

The active space of each sound was estimated as the maximum range at which its apparent source level minus transmission loss exceeded the whale's detection threshold in at least one 1/3-octave band under different background noise conditions. The noise scenarios modeled were sea-state 0 and sea state 6 background noise, which have been well characterized and summarized for naturally occurring sea states (Knudsen et al. 1948). In sea state zero, noise levels range from 44 dB re  $1\mu\text{Pa}^2/\text{Hz}$  at 1kHz to 20 dB at 20kHz (see also Erbe 2002). Noise levels in sea state 6 are roughly parallel, in frequency, to those at sea state 0, but are about 26dB higher.

The relatively simple transmission-loss equation used for calculation of source level ignores propagation effects that become prominent over longer distances. The appropriate empirically derived equation for intermediate-range transmission loss in a shallow environment is:

$$\text{TL} = 15\log(\text{R}) + \alpha\text{R} + \alpha_{\text{T}}(\text{R}/\text{H}-1) + 5\log(\text{H}) + 60 - k_{\text{L}}$$

where TL is the transmission loss (dB), R is the range (km),  $\alpha$  is the absorption coefficient of seawater, H is a function of water depth,  $\alpha_{\text{T}}$  is the shallow-water attenuation coefficient and  $k_{\text{L}}$  is a near-field anomaly (Marsh & Shulkin 1962). The absorption coefficient was calculated using an empirically-derived formula (Francois & Garrison 1982). The values for  $\alpha_{\text{T}}$  and  $k_{\text{L}}$  assuming a sand bottom (Erbe 2002) and sea states 0 or 5, as well as the equation to calculate H based on a water depth of 100m are given in Marsh & Shulkin (1962). This equation accounts for spreading loss, frequency-dependent absorption, and surface/bottom interactions.

Transmission loss was subtracted from the source spectrum in each 1/3-octave band at range steps of 100m. A signal was judged to be detectable by another killer whale when the received level in at least one 1/3-octave band exceeded the hearing threshold of killer whales

or 6 dB below the background noise level, whichever was greater. There are few measurements of the hearing abilities of killer whales in noise (Bain and Dalheim 1994) and no explicit measurements of critical ratios, so I used Fletcher's (1940) equal power assumption with a correction for binaural hearing capabilities to estimate a detection threshold in noise (Erbe 2002). Bain and Dalheim (1994) found that masking sources needed to be 4-40dB more intense to mask natural calls when they were placed to the side or behind an animal. Similar results have been found for human ability to detect speech in cluttered environments (Kidd et al. 1998). Sea generated noise arrives from broad azimuth angles, while the conspecific call arrives from one direction primarily. There is also evidence that humans and birds can use energy in different harmonic bands to improve detection thresholds (Hall, 1984; Klump & Langemann, 1995), which is ignored here. Thus, the -6dB signal to ocean-noise criterion adopted here is a best-guess given current information. New information on the ability of a killer whale to detect a call in background noise would be useful to assess the validity of this criterion.

The auditory threshold at the center of each 1/3-octave band was obtained by fitting a third-order polynomial curve to the auditory threshold points reported in the literature (Hall & Johnson 1972; Symanski et al. 1999). This threshold comparison assumes that the energy in a band of noise that is just audible is equal to the threshold energy of the pure tone at the center of the band, as appears to be the case for human speech (Hawkins and Stevens 1950). Because killer whale hearing becomes less sensitive at low frequencies, the detection threshold was always hearing-threshold limited below 9 and 5 kHz in sea-states 0 and 6, respectively (see also Janik 2000).

## Results

I recorded killer whale sounds for a total of five hours on two different days from five different pods (Table 1), though effort was focused on animals from pods A1 and A5. Because the arrays were towed parallel to the direction of movement of the whales and sounds were localized broadside to the array (Fig. 1), animals most often surfaced with a sideways orientation to the receiver. It is therefore reasonable to consider the apparent source levels to primarily represent off-axis levels, with occasional sounds from animals oriented directly toward or away from the receiver.

The same 12 animals from pods A1 and A5 were present on both recording days and the arrays were positioned to obtain sounds primarily from these animals. The animals were traveling in subgroups and vocally active throughout both recording sessions. Although I could not identify the individual producing each sound, calls were often produced by whales in multiple locations in clear vocal exchanges (Miller et al. 2004), so the sample contained calls produced by multiple individuals in the group. The apparent source levels of 819 pulsed calls and 24 whistles were measured. Of the 819 pulsed calls, 60 did not clearly correspond to one of the call types defined by Ford (1987) and were classified as variable calls.

### *Apparent source levels*

The apparent source levels in the 1-20 kHz band of all sounds ranged from 131 to 168 dB @1m (Fig 4). Whistles had the lowest mean ( $\pm$  s.d.) apparent source level (140.2 $\pm$ 4.1 dB re 1 $\mu$ Pa @1m), followed by variable calls (146.6 $\pm$ 6.6 dB), with stereotyped calls being the most intense (152.6 $\pm$ 5.9 dB). These differences by sound type were statistically significant ( $F_{2,840} = 76.1$   $p < 0.0001$ , Tukey post-hoc  $p < 0.0001$ ). Sounds were occasionally observed on only one array when animals were quite close, suggesting that some sounds were produced below the threshold of 127 dB to be localized with the system. All of these faint sounds were either whistles or variable calls, none were stereotyped calls.

Within the stereotyped calls, the eight most common call types produced by pods A1 and A5 accounted for 618 of the 759 stereotyped calls. I compared the apparent source levels and estimated active space of these calls (Fig. 4, see Fig. 2). Significant differences in apparent source level were found across types ( $F_{7,611} = 54.6$ ,  $p < 0.0001$ ), with a mean of 156.5 and 145.4 for the most (N4) and least (N3) intense call types, respectively (Fig. 4). Clustering using Tukey post-hoc probability of 0.01 resulted in four clusters, suggesting diversity in the level of different call types. Though differences were significant, calls of the same type were produced over a rather wide range of source levels, with standard deviations ranging from 2.7 to 5.4 dB for call types N8 and N4, respectively.

#### *Estimated active space*

In sea-state zero noise conditions, estimated active space of all sounds ranged from 1.9 to 25.8 km (Fig. 5). Whistles had the smallest average estimated active space ( $6.4 \pm 2.4$  km), followed by variable calls ( $7.8 \pm 3.7$  km), with stereotyped calls having the largest average estimated active space ( $11.0 \pm 4.7$  km). The mean estimated active space of stereotyped calls was greater than that of variable calls and whistles ( $F_{2,840} = 23.40$ ,  $p < 0.001$ , Tukey post hoc  $p < 0.001$ ), but the contrast between variable calls and whistles was not statistically significant (Tukey post-hoc  $p = 0.46$ ). Significant differences across the eight most common call types were also apparent ( $F_{7,611} = 31.77$ ,  $p < 0.0001$ ), with a mean of  $12.8 \pm 4.7$  km and  $6.1 \pm 1.6$  km for the longest (N2) and shortest range (N3) call types, respectively. Clustering using Tukey post-hoc probability of 0.01 resulted in two clear clusters: a long-range cluster consisting of call types N2, N4, N1, N9, and N5 (combined mean of  $12.4 \pm 4.3$  km), and a short-range cluster consisting of call types N7, N8 and N3 (combined mean of  $6.8 \pm 3.1$  km). Note that *all* the call types in the long-range cluster contain a separately-modulated high frequency component, while *all* those in the short-range cluster do not.

Source level, detection threshold, and water depth variables were perturbed for the long and short-range call groups to determine the sensitivity of the model to the input variables. As expected, the active space estimate was equally sensitive to perturbations in source level and detection threshold. A 6dB increase in source level (or decrease in threshold) resulted in increases in estimated active space of 26% and 37% for long- and short-range call types, respectively, while a 6dB decrease in source level (or increase in threshold) resulted in a corresponding decrease of 24% and 30%. Increasing water depth to 500m resulted in increases of estimated active space of 16% and 9% for long- and short-range call types, respectively, while decreasing water depth to 20m resulted in corresponding decreases of 25% and 19%.

Relative to the sea-state 0 scenario, estimated active space in sea state 6 noise conditions was 75%, 83% and 91% lower for combined stereotyped calls, variable calls and whistles, respectively (Fig. 5, top). Within the eight most common stereotyped calls (Fig. 5, bottom), sea state 6 conditions resulted in reductions of 74% and 81%, respectively for long and short-range call types. The percentage reduction in estimated active space was smallest for the most intense calls, and greatest for the lower-intensity whistles.

## Discussion

The apparent source levels of all measured sounds ranged from 131-168 dB re 1  $\mu$ Pa @1m, with statistically significant differences in the levels of different sound types in the repertoire (Fig. 4). Supported by similar results from the active space model (Fig. 5), the acoustic repertoire of this group of resident killer whales can be partitioned into two primary groups: 'long-range' sounds including call types N4, N2, N5, N9 and N1 (all contain a separately-modulated high-frequency component), and 'short-range' sounds including whistles, variable calls, and call types N7, N8 and N3 (all without a high-frequency

component). Relative to low-noise sea-state 0 conditions, the estimated active space of whistles, variable calls and stereotyped calls was 91%, 83% and 75% less in sea-state 6, respectively, suggesting that repertoire diversity in active space is preserved or possibly enhanced under higher noise conditions.

The range of apparent source levels reported here is similar to those reported for other large delphinids (Watkins et al. 1997; Fish & Turl 1976; Janik 2000). Erbe (2002) used the received level of 105-124 dB as a minimum estimate of killer whale call source levels. Adding 40dB of transmission loss for whales sighted roughly 100-200m away during those recordings, Erbe's (2002) results are consistent with those reported here. The whistles in my sample were more intense than whistles produced by resting spinner dolphins in Hawaii (109-125 dB @ 1m; Watkins & Schevill 1974), but were less intense than those reported for traveling spinner dolphins (mean 150-154 dB; Lammers & Au 2003). Whistles produced by bottlenose dolphins in Moray Firth (134-169 dB) were more intense than the whistles reported here, but were quite consistent with the overall range of sound source levels in this study (Janik 2000). Whistles were roughly 10 dB less intense than burst-pulse sounds in a sample of sounds collected from the melon-headed whale (Watkins et al. 1997).

Despite statistically strong differences across sound types, the range of apparent source levels within each type was large, with >20dB between the most and least intense exemplars in many cases (Fig. 4). Only about 3dB of variability is expected from errors in range measurement or inaccuracies in the transmission loss model. Though signalers may have the ability to modify the source level of a sound depending on their motivation, there may be other features of sound production that add variability to apparent source levels. As killer whale calls are directional at high frequencies (Schevill & Watkins 1966; Bain and Dalheim 1994; Miller 2002), apparent source level should vary as a function of signaler orientation, which we could not control or measure. During the recordings, most whales

surfaced with a sideways orientation to the array, so the average values measured here are certainly below the on-axis source level. Some analyzed calls contained clear high-frequency components, so were likely to be from individuals oriented more directly toward the array (Miller 2002). Additional variation could be due to individual differences in body size, which varies greatly with age and sex of killer whales (Matkin & Leatherwood 1986). Source levels appear to increase with body size across cetacean species (Patterson & Hamilton 1967; Cummings & Thompson 1971; Thode et al. 2000a; Cummings & Holliday 1985; Schevill & Watkins 1972; Cummings et al. 1968; Dawbin & Cato 1992).

The results of the active space estimation suggest that killer whales may be able to detect communication sounds produced by conspecifics over large distances in quiet conditions. Overall estimated active space ranged from 1.9-25.8 km in sea-state 0 conditions. The maximum estimated active space closely matches maximum active space estimates of 20 to 25 km calculated for a bottlenose dolphin whistle using the same transmission loss model in sea state 0 conditions (Janik 2000). While observers have anecdotally noted cases in which killer whales apparently reacted to sounds of other whales greater than 10km distant, carefully controlled observations or experiments to measure the distances over which killer whales react to conspecific sounds are lacking.

The active space model estimates the distance over which communication may occur, and is useful to demonstrate differences within the repertoire, but its accuracy is limited by the lack of information on detection thresholds of calls in noise. This simple model using average conditions ignores variability in the statistics of background noise, propagation conditions, and physiological characteristics of signalers and receivers that undoubtedly lead to a much larger scale of variability in the true active space of each communication signal. Sensitivity analyses suggest that the active space estimates fluctuate by roughly  $\pm 30\%$  with  $\pm 6$  dB changes in SL or auditory sensitivity. Water depth of 100m assumed in the model is



valid over many near-shore areas that killer whales inhabit, but active space estimates increase in deeper water, and decrease in shallower water, due to loss from surface and bottom reflection (Marsh & Shulkin 1962). The depths of the signaler and receiver themselves also influence signal propagation, with lower frequencies being relatively more attenuated when signalers or receivers are near the surface (Mercado & Frazer 1999).

Directionality of signal transmission and hearing sensitivity are also expected to influence active space (Bain and Dalheim 1994). Although I could not directly observe the orientation of signaling animals in this study, most sounds were likely received from an off-axis orientation with some sounds produced by animals facing directly toward or away from the array. Thus, the mean active space estimated in this study is likely to be intermediate to the extremes when an animal is facing directly toward or away from the receiver. While the radiation pattern of killer whale calls has not been measured, the high-frequency component of stereotyped call types are directional, with differences of  $\pm 6$  dB for animals moving toward, versus away from, a receiver (Miller 2002). Assuming the 1-5 kHz band is omni-directional, front-to-back differences of at least  $\pm 30\%$  in active space can be predicted. Actual front-to-back differences are probably greater than this as the 1-5 kHz frequency band may not be omni-directional. The orientation of the receiving animal is also likely to be an important factor in the shape of active space in animals with directional hearing (Bain and Dalheim 1994). Receiver directivity has been measured at high frequencies for dolphins (Au 1993), but little information is available for the low frequencies at which killer whale calls are produced (Bain & Dalheim 1994).

Estimated active space decreased strongly in noisy sea state 6 conditions, with short-range sound relatively more affected than the long-range stereotyped calls. Strong predicted impacts of naturally-occurring noise may imply mechanisms to adaptively modify signals in response to background noise would be selectively advantageous. Mossbridge & Thomas

(1999) found that frequency modulations of Antarctic killer whale calls shifted away from the 2.8-3.8 kHz band seasonally used by calling leopard seals. Killer whales in three different pods increased the duration of their most common call type in the presence of boat noise (Foote et al. 2004). It is important to emphasize that the source levels reported here were produced in sea state 0 conditions with little nearby vessel traffic, except the research vessel itself. Given the documented compensatory mechanisms in call frequency and duration, it seems plausible that killer whale may also have mechanisms for amplitude regulation of their calls in noise (Cynx et al. 1988; Brumm & Todt 2002; Brumm et al. 2004). Measurement of the source level of sounds produced by killer whales in varying noise conditions could test this possibility.

The result that sound pressure levels and active space estimates can be used to partition the acoustic repertoire of resident killer whales into at least two sound classes is strongly corroborated by observations of sound-type production in different behavioral states of killer whale groups. Resident killer whales do not appear to produce any sound type exclusively in association with a particular behavioral category, but do alter the rates of sound-type production in different behavior states (Bain 1986; Morton et al. 1986; Ford 1989; Thomsen et al. 2002). Thomsen et al. (2002) found that resident killer whales produced short-range whistles and variable calls most commonly in socializing contexts, while stereotyped calls predominated in foraging and travel contexts. In his study of behavior and sound production of pods A1, A4, and A5, Ford (1989) found significant increases in the production of the short-range whistles, variable calls, and the N3 call type during beach-rubbing, socializing, and group-resting behaviors. In these behavioral states, animals tend to be closely spaced and interactions between group members appear to be primarily affiliative. The intense, long-range call types N2, N4, N5, and N9 were used more during behaviors where group members are dispersed, including travelling, foraging, when different pods were

meeting, or in large aggregations of animals (Ford 1989, pg 737). In a study of two captive whales from A5 pod, call-type clusters associated with high activity levels (assessed by respiration rate) included call types N4, N2, and N9, while lower-activity clusters contained call types N7, N8, N3 and whistles (Bain 1986). Though there is a clear division in both the estimated active space and behavioral context of production of long-range and short-range sound classes, it is interesting that these animals have a repertoire of sound types within each class. Thus, the functional aspect of active space described here does not fully explain why killer whales produce a repertoire of whistles and burst pulse calls (Ford 1989).

The precise function(s) of calling by killer whales remains unknown. Ford (1989, 1991) proposed that discrete stereotyped call repertoires function primarily to maintain pod cohesion and coordinate activities. Individuals within groups regularly exchange calls, often immediately matching the call type produced by another individual (Miller et al. 2004). Such interactive calling behavior suggests that communication is at least partly directed to other killer whales within the group. Given that group members rarely separate by more than a few kilometers, why should resident killer whales produce calls that can be detected at much greater distances? The simple explanation that signalers should simply always maximize signal level seems inconsistent with the result of this study which demonstrates that killer whales often produce calls with less active space than their maximum capability.

The ability of a killer whale to detect a conspecific at 10's of km may be a consequence of the signal level required for detailed information (group membership, Ford 1991; direction of movement, Miller 2002) to be reliably transferred over shorter typical separation distances to intended receivers within the group (McGregor and Dabelsteen 1996). Alternatively, calling may have inter-group functions such as spatial competition (Miller and Bain 2000) or mate attraction (Yurk et al. 2002) that select for longer-range signals. In the fjord-like inland waters of the coastal NE Pacific, with numerous islands, continuous sections

of water are often considerably shorter than the estimated active space of loud calls. In such a habitat, a direct signal path may often be as important as 'staying within range' for members of stable social groups to maintain acoustic contact with each other. If the intended receivers of killer whale calls are the members of other groups, we might expect calling to occur in 'hot-spots' where acoustic propagation conditions are good or from which there exists a direct path to larger areas of killer whale habitat. A detailed analysis of the geography of call production by these mobile animals may help elucidate intra-group and inter-group functions of calling.

### **Acknowledgements**

Many thanks to Mark Johnson and WHOI's acoustic systems group for assistance with equipment development and data analysis. Field assistance was provided by Erin Gentry, Miranda Buck, Leandra de Sousa, Dennis Schmidt, Nicoletta Biassoni, Morgan Heim, Michael Richlen, Mariya Sweetwyne, and Jodi Udd. Thanks also to David Bain, David Tyre, and Bill & Donna MacKay for help in the field. Darlene Ketten provided invaluable advice on the hearing model. Helpful comments on the manuscript were provided by Henrik Brumm, Peter Tyack, Peter Slater, David Bain, Nicoletta Biassoni, Rebecca Thomas and Vincent Janik. Funding was provided by WHOI's Ocean Ventures Fund and Rinehart Coastal Research Center and a Royal Society fellowship. Observational research was conducted under a permit from the Department of Fisheries and Oceans, Canada, and complies with the "Principles of animal care", publication No. 86-23, revised 1985 of the National Institute of Health.

## References

- Au WWL (1993) *The Sonar of Dolphins*. Springer-Verlag, New York.
- Bain DE (1986) Acoustic behavior of *Orcinus*: sequences, periodicity, behavioral correlates and an automated technique for call classification. In: Kirkevold BC, Lockard JS (eds) *Behavioral Biology of Killer Whales*. Alan R. Liss, Inc., New York, pp 335-371.
- Bain DE, Dalheim ME (1994) Effects of masking noise on detection thresholds of killer whales. In: Loughlin TR (ed) *Marine Mammals and the Exxon Valdez*. Academic Press, San Diego, pp 243-256.
- Bradbury JW, Vehrencamp SL (1998) *Principles of Animal Communication*. Sinauer Press, New York.
- Brenowitz EA (1982) The active space of red-winged blackbird song. *J Comp Physiol A* 147:511-522.
- Brumm H, Todt D (2002) Noise-dependent song amplitude regulation in a territorial songbird. *Anim Behav* 63:891-897.
- Brumm H, Todt D (2004) Male-male vocal interactions and the adjustment of song amplitude in a territorial bird. *Anim Behav* 67:281-286.
- Cato DH (1998) Simple methods of estimating source levels and locations of marine animal sounds. *J Acoust Soc Am* 104:1667-1678.
- Charif RA, Mellinger DK, Dunsmore KJ, Fristrup KM, Clark CW (2002) Estimated source levels of fin whale (*Balaenoptera physalus*) vocalizations: adjustments for surface interference. *Mar Mamm Sci* 18:81-98.
- Cummings WC, Holliday DV (1985) Passive acoustic location of bowhead whales in a population census off Point Barrow, Alaska. *J Acoust Soc Am* 78:1163-1169.

Cummings WC, Thompson PO (1971) Underwater sounds from the blue whale,

*Balaenoptera musculus*. J Acoust Soc Am 50:1193-1198.

Cummings WC, Thompson PO, Cook R (1968) Underwater sounds of migrating gray whales,

*Eschrichtius glaucus* (Cope). J Acoust Soc Am 44:1278-1281.

Cynx J, Lewis R, Tavel B, Tse H (1998) Amplitude regulation of vocalizations in noise by a

songbird, *Taeniopygia guttata*. Anim Behav 56:107-113.

Dabelsteen T, Pedersen SB (1990) Song and information about aggressive responses of

blackbirds, *Turdus merula*: evidence from interactive playback experiments with territory owners. Anim Behav 40:1158-1168.

Dawbin WH, Cato DH (1992) Sounds of a pygmy right whale (*Caperea marginata*). Mar

Mamm Sci 8:213-219.

Deecke VB, Ford JKB, Spong P (1999) Quantifying complex patterns of bioacoustic

variation: use of a neural network to compare killer whale (*Orcinus orca*) dialects. J Acoust Soc Am 105:2499-2507.

Deecke VB, Ford JKB, Spong P (2000) Dialect change in resident killer whales (*Orcinus*

*orca*): implications for vocal learning and cultural transmission. Anim Behav 60:629-638.

Dittus W (1988) An analysis of Toque macaque cohesion calls from an ecological

perspective. In: Todt D, Goedeeking P, Symmes D. (eds) Primate Vocal Communication. Springer-Verlag, New York, pp 31-50.

Erbe C (2002) Underwater noise of whale-watching boats and potential effects on killer

whales (*Orcinus orca*), based on an acoustic impact model. Mar Mamm Sci 18:394-418.

Fish JF, Turl CW (1976) Acoustic Source Levels of Four Species of Small Whales. Naval

Undersea Center Report, NUC-TP 547.

Fletcher H. (1940) Auditory patterns. *Rev Mod Phys* 12:47-65.

Foote AD, Osborne RW, Hoelzel AR (2004) Whale call response to masking boat noise. *Nature* 428:910.

Ford JKB (1987) A catalogue of underwater calls produced by killer whales (*Orcinus orca*) in British Columbia. Canadian Data Report of Fisheries and Aquatic Sciences No. 633.

Ford JKB (1989) Acoustic behavior of resident killer whales (*Orcinus orca*) in British Columbia. *Can J Zool* 67:727-745.

Ford JKB (1991) Vocal traditions among resident killer whales (*Orcinus orca*) in coastal waters of British Columbia. *Can J Zool* 69:1454-1483.

Francois RE, Garrison GR (1982) Sound absorption based upon ocean measurements. Part II: boric acid contribution and equation for total absorption. *J Acoust Soc Am* 72:1879-1890.

Gerhardt HG (1975) Sound pressure levels and radiation patterns of the vocalizations of some north American frogs and toads. *J Comp Physiol A* 102:1-12.

Hall J, Johnson CS (1972) Auditory thresholds of a killer whale *Orcinus orca* Linnaeus. *J Acoust Soc Am* 51:515-517.

Hall JW, Haggard MP, Fernandes, MA (1984) Detection in noise by spectro-temporal pattern analysis. *J Acoust Soc Am* 76 (1):50-56.

Hauser MD, Marler P (1993) Food-associated calls in rhesus macaques (*Macaca mulatta*) I. socioecological factors. *Behav Ecol* 4:194-205.

Hawkins JE, Stevens SS (1950) The masking of pure tones and of speech by white noise. *J Acoust Soc Am* 22:6-13.

- Hoelzel AR, Osborne RW (1986) Killer whale call characteristics: implications for cooperative foraging strategies. In: Kirkevold BC, Lockard JS (eds) Behavioral Biology of Killer Whales. Alan R. Liss, Inc., New York, pp 373-403.
- Janik V (2000) Source levels and the estimated active space of bottlenose dolphin (*Tursiops truncatus*) whistles in the Moray Firth, Scotland. *J Comp Physiol A*. 186:673-680.
- Janik V (in press) Underwater acoustic communication networks in marine mammals. In: McGregor PK (ed) Animal Communication Networks. Cambridge University Press, Cambridge.
- Kidd G Jr, Mason CR, Rohtla TL, Deliwala PS (1998) Release from masking due to spatial separation of sources in the identification of nonspeech auditory patterns. *J Acoust Soc Am* 104 (1):422-431.
- Klump GM, Langemann U (1995) Comodulation masking release in a songbird. *Hear Res* 87:157-164.
- Lammers MO, Au, WWL (2003) Directionality in the whistles of Hawaiian spinner dolphins (*Stenella longirostris*): a signal feature to cue direction of movement? *Mar Mamm Sci* 19:249-264.
- Loftus-Hills JJ, Littlejohn MJ (1971) Mating-call sound intensities of anuran amphibians. *J Acoust Soc Am* 49:1327-1329.
- Knudsen VO, Alford RS, Emling JW (1948) Underwater ambient noise. *J Mar Res* 7:410-429.
- Madsen PT, Kerr I, Payne R (2004) Echolocation clicks of two free-ranging, oceanic delphinids with different food preferences: false killer whales *Pseudorca crassidens* and Risso's dolphins *Grampus griseus*. *J Exp Biol* 207:1811-1823.
- Marler P (1973) A comparison of vocalizations of red-tailed monkeys and blue monkeys, *Cercopithecus ascanius* and *C. mitis*, in Uganda. *Z Tierpsychol* 33:223-247.



- Marsh HW, Schulkin M (1962) Shallow water transmission. *J Acoust Soc Am* 34:863-864.
- Marten K, Quine D, Marler P (1977) Sound transmission and its significance for animal vocalization II. tropical forest habitats. *Behav Ecol Sociobiol* 2:291-302.
- Matkin CO, Leatherwood, S (1986) General biology of the killer whale, *Orcinus orca*: a synopsis of knowledge. In: Kirkevold BC, Lockard JS (eds) *Behavioral Biology of Killer Whales*. Alan R. Liss, Inc., New York, pp 35-95.
- McGregor PK, Dabelsteen T (1996) Communication networks. In: Kroodsma DE, Miller EH (eds) *Ecology and evolution of acoustic communication in birds*. Comstock Publishing Associates, Ithaca, NY, pp 409-425.
- Mercado E III, Frazer LN (1999) Environmental constraints on sound transmission by humpback whales. *J Acoust Soc Am* 106:3004-3016.
- Miller PJO (2002) Mixed-directionality of killer whale stereotyped calls: a direction of movement cue? *Behav Ecol Sociobiol* 52:262-270.
- Miller PJO, Bain DE (2000) Within-pod variation in the sound production of a pod of killer whales (*Orcinus orca*). *Anim Behav* 60:617-628.
- Miller PJO, Shapiro AD, Tyack PL, Solow AR (2004) Call-type matching in vocal exchanges of free-ranging resident killer whales, *Orcinus orca*. *Anim Behav* 67:1099-1107.
- Miller, PJO, Tyack PL (1998) A small towed beamforming array to identify vocalizing resident killer whales (*Orcinus orca*) concurrent with focal behavioral observations. *Deep-Sea Res. II* 45:1389-1405.
- Mohl B, Wahlberg M, Madsen PT (2000) Sperm whale clicks: directionality and source level revisited. *J Acoust Soc Am* 107:638-648.
- Morton AB, Gale JC, Prince RC (1986) Sound and behavioral correlations in captive *Orcinus orca*. In: Kirkevold BC, Lockard JS (eds) *Behavioral Biology of Killer Whales*. Alan R. Liss, Inc., New York, pp 303-333.

- Mossbridge JA, Thomas JA (1999) An “acoustic niche” for Antarctic killer whale and leopard seal sounds. *Mar Mamm Sci* 15:1351-1357.
- Myrberg, A. A. Jr. (1980) Ocean noise and the behavior of marine animals: relationships and implications. In: *Advanced Concepts in Ocean Measurements for Marine Biology*, (Ed. by F. P. Diemer, F. J. Vernberg, D. Z. Mirkes), pp. 461-491. Columbia, SC: University of South Carolina Press.
- Patterson B, Hamilton GR (1967) Repetitive 20 cycle per second biological hydroacoustic signals at Bermuda. In: Tavalga WN (ed) *Marine Bioacoustics*, vol. 1. Pergamon Press, New York, pp 125-145.
- Prestwich KN, Brugger KE, Topping M (1989) Energy and communication in three species of hylid frogs: power input, power output and efficiency. *J Exp Biol* 144:53-80.
- Pytte CL, Rusch KM, Ficken MS (2003) Regulation of vocal amplitude by the blue-throated hummingbird, *Lampornis clemenciae*. *Anim Behav* 66:703-710.
- Rowell TE, Hinde RA (1962) Vocal communication by the rhesus monkey (*Macaca mulatta*). *Proc Zool Soc Lon* 138:279-294.
- Ryan MJ, Tuttle MD, Rand AS (1982) Bat predation and sexual advertisement in a neotropical frog. *Am Nat* 119:136-139.
- Schevill WE, Watkins WA (1966) Sound structure and directionality in *Orcinus* (killer whale). *Zoologica* 51:71-76.
- Schevill WE, Watkins WA (1972) Intense low-frequency sounds from an Antarctic minke whale, *Balaenoptera acutorostata*. *Brevoria* 388:1-8.
- Symanski MD, Bain DE, Kiehl K, Pennington W, Wong S, Henry KR (1999) Killer whale (*Orcinus orca*) hearing: auditory brainstem response and behavioral audiograms. *J Acoust Soc Am* 106:1134-1141.

- Thode AM, D'Spain GL, Kuperman WA. (2000a) Matched-field processing, geoacoustic inversion, and source signature recovery of blue whale vocalizations. *J Acoust Soc Am* 107:1286-1300.
- Thode AM, Norris T, Barlow J. (2000b) Frequency beamforming of dolphin whistles using a sparse three-element towed array. *J Acoust Soc Am* 107:3581-3484.
- Thomsen F, Franck D, Ford JKB (2001) Characteristics of whistles from the acoustic repertoire of resident killer whales (*Orcinus orca*) off Vancouver Island, British Columbia. *J Acoust Soc Am* 109:1240-1246.
- Thomsen F, Franck D, Ford JKB (2002) On the communicative significance of whistles in wild killer whales (*Orcinus orca*). *Naturwissenschaften* 89:404-407.
- Urick RJ (1981) Principles of underwater sound. McGraw-Hill, New York.
- Watkins WA (1967) Harmonic interval: fact or artifact in spectral analysis of pulse trains. In: Tavorla WN (ed) *Marine Bio-acoustics*, vol. 2. Pergamon Press, Oxford, pp 15-42.
- Watkins WA, Daher MA, Samuels A, Gannon DP (1997) Observations of *Peponocephala electra*, the melon-headed whale, in the Southeastern Caribbean. *Car J Sci* 33:34-40.
- Watkins WA, Schevill WE (1974) Listening to Hawaiian spinner porpoises, *Stenella cf. longirostris*, with a three-dimensional hydrophone array. *Mammology* 55:319-328.
- Wiley RH, Richards DG (1982) Adaptations for acoustic communication in birds: sound transmission and signal detection. In: Kroodsma DE, Miller EH (eds) *Acoustic Communication in Birds*, vol. 2. Academic Press, New York, pp 131-181.
- Yurk H, Barrett-Lennard L, Ford JKB, Matkin CO (2002) Cultural transmission within maternal lineages: vocal clans in resident killer whales in southern Alaska. *Anim Behav* 63:1103-1119.

### Figure Legends

**Figure 1.** Schematic of the research vessel and two-array system used to make calibrated recordings. Range from the rear array to a sound source is calculated using measured angles ( $\theta$ , in radians) and the inter-array separation ( $d$ ), which was 97m. The dashed lines and lighter oval illustrate the consistency of the range estimate despite the left-right (and depth) ambiguity of the linear arrays.

**Figure 2.** Spectrograms of the eight most common call types recorded and a whistle. Note that call types N1, N2, N4, N5 and N9 all contain a separately modulated high frequency component, while call types N3, N7 and N8 do not. The effective filter bandwidth was 97 Hz.

**Figure 3.** Calibration results of the range measurement of the 2-array system. The front array is centered on (0,0) and the rear array on (97,0). The  $\Delta$  symbols mark where calibration measurements were taken and the \* symbols mark the position of all killer whale sounds for which an apparent source level was calculated. Calibration measures were taken on both sides of the array system, but are plotted together because the left-right ambiguity is not resolvable using this linear array system. The grayscale background is the interpolated range error on source level (in dB) between measured points. Note that all sounds were within or near the calibrated region, and that the range error did not exceed  $\pm 3$ dB in the region from which apparent source levels were calculated.

**Figure 4.** Left: the apparent source levels of stereotyped calls, variable calls and whistles. Right: source levels of the eight most common stereotyped calls recorded from pods A1 and A5 (see Fig. 2). Note the diversity of average source level among the different sound types.

**Figure 5.** Box plots of estimated active space of whistles, variable calls, and group stereotyped calls (top), and stereotyped calls classified by type (bottom) in sea-state zero (left half of both panels) and sea state six (right half) background noise scenarios. The center of each box is the median value, and the upper and lower box borders are the first quartiles of the data. The whiskers extend to the most extreme datum within the inter-quartile range defined by the box borders, and more extreme data are plotted as symbols. Note the bimodal structure of the estimated active space. The short-range sound class includes whistles, variable calls and call types N3, N7, and N8. The long-range sound class consists of stereotyped call types N1, N2, N4, N5 and N9.

**Table 1.** Group composition and sample size from recording effort. The same 12 animals present on 9/1/98 were also present on 9/8/98. \*The number of animals on 9/8/98 was estimated based upon identified groups as all individuals were not separately identified.

Date:	9/1/98	9/8/98
Pods Present:	A1, A5	A1, A5, B1, C1, I15
# of Animals:	12	46*
Hours Recorded:	2	3
Calls analyzed	334	485
Whistles analyzed	15	9

**Figure 1.**




Figure 2.







Figure 3.



**Figure 4.**




Figure 5.

