

1 **Changes in persistent contaminant concentration and CYP1A1 protein expression in**
2 **biopsy samples from northern bottlenose whales, *Hyperoodon ampullatus*, following the**
3 **onset of nearby oil and gas development**

4
5 Sascha K. Hooker^{a,b}

6 Tracy L. Metcalfe^c

7 Chris D. Metcalfe^c

8 Carolyn M. Angell^d

9 Joanna Y. Wilson^{d,e}

10 Michael J. Moore^d

11 Hal Whitehead^a

12
13 ^aDepartment of Biology, Dalhousie University, Halifax, Nova Scotia, B3H 4J1, Canada.

14 ^bSea Mammal Research Unit, University of St Andrews, FIFE, KY16 8YG, UK

15 ^cEnvironmental and Resource Studies, Trent University, Peterborough, ON, K9J 7B8, Canada

16 ^dBiology Department, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, U.S.A.

17 ^eCurrent address: Department of Biology, McMaster University, Hamilton, Ontario, L8S 4K1,
18 Canada.

19
20
21 Corresponding author: Sascha K. Hooker

22 Email: s.hooker@st-andrews.ac.uk

23 Sea Mammal Research Unit, University of St Andrews, FIFE, KY16 8YG, UK

24 Telephone: 44 (0)1334 467201

25 Fax: 44 (0)1334 462632

1 **Abstract**

2 A small population of endangered northern bottlenose whales (*Hyperoodon ampullatus*) inhabits
3 “The Gully” Marine Protected Area on the Scotian Shelf, eastern Canada. Amid concerns
4 regarding nearby oil and gas development, we took 36 skin and blubber biopsy samples in 1996-
5 97 (prior to major development) and 2002-03 (five years after development began), and 3
6 samples from a population in the Davis Strait, Labrador in 2003. These were analysed for
7 cytochrome P4501A1 (CYP1A1) protein expression (n=36), and for persistent contaminants
8 (n=23). CYP1A1 showed generally low expression in whales from The Gully, but higher levels
9 during 2003, potentially co-incident with recorded oil spills, and higher levels in Davis Strait
10 whales. A range of PCB congeners and organochlorine compounds were detected, with
11 concentrations similar to other North Atlantic odontocetes. Concentrations were higher in
12 whales from The Gully than from the Davis Strait, with significant increases in 4,4’-DDE and
13 *trans*-nonachlor in 2002-03 relative to 1996-97.

14

15 **Capsule**

16 Whale contaminants highlight concerns from oil and gas development near a marine protected
17 area.

18

19 **Keywords:** Bottlenose whale; The Gully; PCB; DDT; cytochrome P4501A.

20

21 **1. Introduction**

22 A small population of approximately 163 northern bottlenose whales (*Hyperoodon ampullatus*)
23 has, as a large part of its habitat, the deeper waters within “The Gully”, a submarine canyon off
24 the coast of eastern Canada (Hooker et al., 2001b; Whitehead and Wimmer, 2005; Wimmer and

1 Whitehead, 2004). Concern about the health of this population of cetaceans stems from the
2 recent designation of The Gully as a Marine Protected Area, and the extraction of oil and gas
3 from the surrounding shelf areas. For about three decades, there has been generally increasing
4 interest in oil and gas exploration and production on the Scotian Shelf. In 1996, the Sable
5 Offshore Energy Project began construction of five production rigs around Sable Island, with
6 one of these lying only 30 km from The Gully (Figure 1). These rigs began production
7 operations in 1998.

8 Northern bottlenose whales are members of the beaked whale family, the Ziphiidae. This family
9 is one of the least known groups of marine mammals, largely due to their offshore habitat, their
10 long dive times, and shyness around vessels. Off the coast of eastern Canada, a population of
11 northern bottlenose whales can be reliably found in The Gully, which appears to be an important
12 foraging location (Hooker et al., 2001b). Unlike most other beaked whales, the northern
13 bottlenose whale is relatively curious, and will approach stationary vessels (Mead, 1989). This
14 has allowed research on the ecology and behaviour of these whales. A photo-identification
15 study has shown that about one third of the population using the waters off the Scotian Shelf
16 reside in The Gully at any time (Gowans et al., 2000b; Whitehead and Wimmer, 2005). This
17 population of whales is thought to be distinct from other populations in the North Atlantic,
18 based on genetic differences (Dalebout et al., 2001; Dalebout et al., 2006) and size differences
19 (Whitehead et al., 1997) observed between these whales and those from the nearest population
20 in the Davis Strait.

21 The Gully has been recognized as an important habitat for several cetacean species (Hooker et
22 al., 1999), but particularly for northern bottlenose whales. Concerns over the small size of the
23 population and its susceptibility to anthropogenic impacts have led to the listing (Committee on
24 the Status of Endangered Wildlife in Canada, 2002), and subsequent designation by the
25 Canadian Government (2006), of this population as “endangered”. After consideration and
26 assessment over much of the last decade (Fenton et al., 2002), The Gully was designated as a
27 Marine Protected Area in May, 2004, in part to protect these whales.

28 There have been few previous reports on the levels and potential impacts of persistent

1 contaminants in northern bottlenose whales. In this study, we compared the concentrations of
2 PCBs and other organochlorine contaminants in skin and blubber biopsy samples collected from
3 whales in The Gully before the start of oil and gas production (i.e. 1996-97) and five years after
4 production began (i.e. 2002-03). We also compared contaminant concentrations in animals from
5 the population in The Gully to concentrations in biopsy samples collected in 2003 from northern
6 bottlenose whales living in the Davis Strait.

7 In addition, in order to evaluate exposure to biologically active contaminants, we analyzed
8 biopsy samples for expression of cytochrome P4501A (CYP1A1) protein. Induction of CYP1A1
9 is a biomarker for exposure to contaminants that interact with the cellular aryl hydrocarbon (Ah)
10 receptor (Stegeman et al., 1992) and induction of CYP1A1 is correlated to toxicity in rodents
11 (Safe, 1986). Typical Ah receptor ligands include the planar halogenated aromatic hydrocarbons
12 such as the non- and mono-ortho substituted PCBs and certain polycyclic aromatic
13 hydrocarbons. CYP1A1 is expressed in multiple cell types of cetacean skin (Angell et al., 2004)
14 and this expression is induced by in vitro exposure to beta-naphthoflavone, a prototypical Ah
15 receptor ligand (Godard et al., 2004).

16

17 **2. Methods**

18 *2.1. Biopsy Collection*

19 Field collections off eastern Canada took place approximately 300 km east of Halifax, Nova
20 Scotia (Figure 1), in The Gully submarine canyon (approximate position: 44°N, 59°W). Biopsy
21 samples were collected from male and female bottlenose whales during August of 1996 and
22 July-August of 1997 (i.e. pre-exploration), and during August of 2002, and August-September
23 of 2003. Field collections in the Davis Strait took place approximately 200 km east of Nain,
24 Labrador (approximate position 57°N, 58°W), where biopsies were collected from male and
25 female whales in August of 2003 (Figure 1).

26 Biopsy sampling was conducted from either a 12-m auxiliary sailing vessel or from a 5-m rigid-

1 hulled inflatable tender. These vessels were operated under power at speeds of 1-4 knots during
2 collection of biopsy samples. Biopsy darts (2.5-cm-long, 0.6-cm-diameter) were fired from a
3 150 lb crossbow (Barnett Wildcat XL) or from a Paxarms air-rifle at ranges of 5 – 15 m, as
4 described by Hooker et al. (2001a). All samples were taken from the flank near the dorsal fin.
5 We attempted to sample adult whales (based on field observations of animal size) and avoided
6 sampling of calves.

7 Each biopsy sample was subdivided for the different types of analysis within an hour of
8 collection to allow matched subsamples from each animal. Subsamples of skin were stored in
9 DMSO for genetic analysis of sex (Gowans et al., 2000a). A biopsy subsample consisting of the
10 skin-blubber interface was stored in neutral buffered formalin for analysis of the expression of
11 CYP1A1. A subsample of blubber from each of the biopsies was stored temporarily in hexane-
12 washed foil in liquid nitrogen, then kept frozen in hexane-washed glassware with teflon-lined
13 lids at –20°C when freezer facilities became available, for analysis of contaminant
14 concentrations.

15 2.2. Biopsy Analysis

16 The sex of the biopsied whales was determined by genetic analysis of the skin using the method
17 described by Gowans et al. (2000a). Expression of CYP1A1 protein was determined using an
18 immunohistochemical method described by Angell et al. (2004). Briefly, the samples (approx.
19 0.5 g) were fixed in 10% neutral buffered formalin, embedded in paraffin and sectioned at 5 µm.
20 Serial sections were deparaffinated, hydrated and stained immunohistochemically using a
21 peroxidase/antiperoxidase detection system (Signet Laboratories, Dedham, MA, USA) with the
22 monoclonal antibody (MAb) 1-12-3 against scup (*Stenotomus chrysops*) CYP1A as the primary
23 antibody and non-specific MAb MOPC or UPC-10 (Sigma-Aldrich, St. Louis, MO, USA) as the
24 control antibody. Mab 1-12-3 recognizes an epitope that occurs in CYP1A1 but not CYP1A2 in
25 mammals (Drahushuk et al., 1998). Amino-9-ethylcarbazole (Signet Laboratories) was used as
26 the chromogenic substrate for visualization of CYP1A1. Sections were counterstained with
27 Mayer's hematoxylin (Sigma-Aldrich) for visualization of nuclei. Stained sections were
28 evaluated under light microscopy for stain occurrence (scale of 0-3) and stain intensity (scale of

1 0-5) in each cell type. CYP1A1 expression was calculated as the product of stain occurrence and
2 intensity to provide a semi-quantitative index (scale 0-15).

3 Blubber from biopsy samples was prepared for analysis of contaminants as described by
4 Metcalfe and Metcalfe (1997). Briefly, blubber samples of < 1g were mixed and ground with
5 sodium sulfate and packed into a glass column for extraction into 50:50 dichloromethane/hexane
6 using a cold-column extraction procedure. The lipids were removed from extracts by gel
7 permeation chromatography (GPC) and the lipid fraction was collected for gravimetric analysis
8 of the lipid content of the sample. The GPC fraction containing contaminants was
9 subfractionated by silica-gel column chromatography to yield Fraction A that contained
10 primarily PCB congeners and some DDE, and Fraction B that contained the rest of the DDE and
11 the majority of the organochlorine analytes.

12 The samples were analyzed by gas chromatography with an electron capture detector (GC-ECD),
13 as described previously (Metcalfe and Metcalfe, 1997). All samples were analyzed for 18 major
14 PCB congeners (denoted as major PCB congeners) with IUPAC numbers 44, 49, 52, 87, 99,
15 101, 105, 110, 118, 138, 149, 153, 156, 170, 180, 194, 195, 209. Samples collected in 2002-03
16 were analyzed for additional minor PCB congeners with IUPAC numbers 17, 18, 28, 31, 33, 47,
17 66, 70, 74, 95, 128, 132, 151, 187, 191, 199, 201, 205, 206. The samples were also analyzed for
18 a range of organochlorine compounds, including 2,4'- and 4,4'-isomers of DDT
19 (dichlorodiphenyltrichloroethane), and its metabolites DDE (dichlorodiphenylchloroethylene)
20 and DDD (dichlorodiphenyldichloroethane), α -HCH (hexachlorocyclohexane), β -HCH, δ -HCH,
21 and γ -HCH, *trans*- and *cis*-chlordane, *trans*- and *cis*-nonachlor, endosulfan, endosulfan II and
22 endosulfan sulfate, heptachlor and heptachlor epoxide, aldrin, dieldrin, endrin, endrin ketone and
23 endrin aldehyde, methoxychlor, mirex, hexachlorobenzene (HCB) and octachlorostyrene. The
24 Limits of Detection (LODs) were between 0.04 and 0.25 ng/mL for individual PCB congeners,
25 between 0.05 and 0.1 ng/mL for DDT and metabolites and between 0.01 and 0.5 ng/mL for all
26 other organochlorine analytes. Procedural blanks and a National Institute for Standards and
27 Technology cod liver oil reference material (SRM 1588) were analyzed for quality
28 control/quality assurance purposes.

1 The total PCB concentration was calculated as the sum of the “major PCB congeners” and the
2 total DDT concentration was calculated as the sum of 4,4'- and 2,4'-isomers of DDT, DDE and
3 DDD. The total HCH concentration was calculated as the sum of the four HCH isomers. Total
4 chlordane was calculated as the sum of *trans*- and *cis*-chlordane and *trans*- and *cis*-nonachlor.
5 Total endosulfan was calculated as the sum of endosulfan and its two metabolites. All analyte
6 concentrations were lipid normalized (i.e. ng/g lipid).

7 2.3. Statistical Analysis

8 Means and standard deviations about the mean were calculated for lipid-normalized
9 concentrations of contaminants, and these data are presented for reference in all tables and
10 figures. Since data were in some cases slightly skewed, we used log-transformed data for all
11 statistical analysis. Thus, in cases where analytes were not detected, we used the detection limit
12 as a default value and surrogate concentrations equivalent to the specific LOD for each congener
13 were substituted prior to log-transformation. Differences in the concentrations of total analytes
14 (total PCBs, total DDTs, etc) in sample groups were tested using a MANOVA with the log-
15 transformed data for the lipid-normalized concentrations of selected contaminants. Differences
16 between CYP1A1 expression in endothelial and smooth muscle cells between years and
17 locations were tested using a MANOVA. Statistical analyses were conducted with SPSS 11.5.

18

19 3. Results

20 Biospy samples were collected from 39 bottlenose whales (18 males, 21 females) during the
21 study. However, since the biopsies did not always contain sufficient material for both types of
22 analyses, CYP1A1 analysis was conducted with 36 samples, and contaminant analysis was
23 conducted with 23 samples (Table 1). Although we attempted to photograph all animals that
24 were sampled, photographs were not always of sufficient quality or animals were not marked
25 well enough for identification. Approximately 50% of the animals that were biopsied were
26 identified from photographs. From photographic identification, it was determined that one
27 animal (a mature male, ID 480) was sampled in both the 1996-97 and 2002-03 time periods

1 (Table 1).

2 Since expression of CYP1A1 in tissues is thought to reflect recent exposure to contaminants that
3 act through the Ah-receptor (Fossi et al., 1994), the CYP1A1 results are presented for each year
4 of collection (Fig. 2). In general, scores for CYP1A1 activity in biopsies from the bottlenose
5 whales were low, with the majority of samples (i.e. 30 of 36 endothelial and 27 of 36 smooth
6 muscle) scoring zero (Table 1). Some limited numbers of animals expressed CYP1A1 in
7 endothelial and vascular smooth muscle cells. Samples collected from The Gully in 2003
8 showed significantly higher expression for CYP1A1 than samples collected in 1996, 1997 or
9 2002 (MANOVA $F = 9.249$, $p < 0.001$). Surprisingly, although only three samples were analysed
10 from whales in the Davis Strait, Labrador, these showed relatively high scores for CYP1A1
11 expression (endothelial 3.5 ± 0.87 and smooth muscle 6.33 ± 2.52 , $n=3$, Fig. 2, Table 1). These
12 scores were significantly greater than those for whales from The Gully (MANOVA, $F = 34.110$,
13 $p < 0.001$). There was no significant difference between expression of CYP1A1 in samples based
14 on sex of whale (MANOVA using location and year as covariates, $F=0.705$).

15 A range of PCB congeners and organochlorine contaminants were detected in the blubber
16 biopsies collected from northern bottlenose whales (Table 2). The classes of compounds present
17 in the highest concentrations were (in order) DDT and its metabolites, PCBs, chlordanes, HCHs,
18 and dieldrin. There was considerable variation in the concentrations of compounds in different
19 individuals. There were significant differences between contaminant concentrations (total
20 PCBs, HCHs, DDTs, chlordanes and endosulfan) for male and female whales (MANOVA using
21 location and sample year as covariates, $F = 7.438$, $p=0.001$). The mean concentrations in
22 females were lower than those in males (Table 2).

23 For most contaminants, the concentrations observed in whales sampled in the Davis Strait were
24 much lower than the concentrations in whales from The Gully (Figure 3, Table 2, MANOVA
25 using sex as a covariate, $F = 6.162$, $p=0.002$).

26 Among whales from The Gully region, there was a significant difference between contaminant
27 concentrations (total PCBs, HCHs, DDTs, chlordanes and endosulfan) in samples collected in

1 1996-97 and those in 2002-03 (MANOVA using sex as a covariate, $F = 11.7$, $p < 0.001$). Of
2 these contaminants, total PCB showed no significant differences, DDTs and chlordanes showed
3 significant increases, while HCHs and endosulfans showed significant decreases (Table 2,
4 Figure 3). Total PCB concentrations in whales from The Gully were 7,273 and 3,791 ng/g lipid
5 for males and females, respectively sampled in 1996-97, and 6,841 and 4,061 ng/g lipid for
6 males and females, respectively sampled in 2002-03. Concentrations of total DDT in whales
7 from The Gully increased by 50% from 14,658 and 4,143 ng/g lipid for males and females,
8 respectively sampled in 1996-97, to 21,571 and 7,601 ng/g lipid for males and females,
9 respectively sampled in 2002-03. Concentrations of total chlordanes in whales from The Gully
10 increased by 100% from 909 and 475 ng/g lipid for males and females, respectively sampled in
11 1996-97, to 1,809 and 1,189 ng/g lipid for males and females, respectively sampled in 2002-03.
12 Closer investigation of the individual components of these classes of compounds indicate that
13 the differences in contaminant concentrations can be primarily attributed to 4,4-DDE and trans-
14 nonachlor (Figures 4 and 5).

15 The same pattern of increase in contaminant concentrations was observed for biopsies from a
16 single male animal that was re-sampled in The Gully area. As shown in Table 2, in this animal,
17 there was little change in total PCBs, and a decrease in total HCH, while total chlordanes
18 increased from 638 to 797 ng/g lipid. However, there was a three-fold increase in total DDT
19 concentration, with a large increase in the concentration of 4,4-DDE. A large increase was also
20 observed in dieldrin concentration, from 500 to 3,432 ng/g lipid (Table 2).

21

22 **4. Discussion**

23 This study uses biopsy sampling to assess contaminant levels in blubber and expression of
24 cytochrome P450 enzymes in the skin of odontocetes. One concern of such studies is that
25 biopsies only sample the superficial blubber layers, and that there may be vertical stratification
26 of lipid types within the blubber mantle (Koopman et al., 1996). However, in blubber removed
27 from baleen whales, lipid-normalized contaminant levels showed no significant differences

1 according to body site or blubber strata (Gauthier et al., 1997). On the other hand, while
2 concentrations of PCBs and organochlorine pesticides in superficial blubber biopsies collected
3 from free-ranging St. Lawrence beluga whales (*Delphinapterus leucus*) were consistent with the
4 concentrations in full depth blubber samples removed from stranded belugas, there were
5 differences between the biopsies and full depth samples in the relative proportions of highly
6 chlorinated PCB congeners, HCB, DDT isomers and mirex (Hobbs et al., 2003). Stratification of
7 dietary fatty acids in the blubber layer of northern bottlenose whales is much less than that
8 observed among other odontocete species, such as harbour porpoises (*Phocoena phocoena*)
9 (Hooker et al., 2001b), so the outer blubber layer may be representative of all blubber strata.
10 Furthermore although outer blubber may be less physiologically dynamic and more structural in
11 function, this may in fact serve as a better measure of integrated long-term persistent
12 contaminant exposure. Therefore, it was assumed that the biopsy method was an appropriate
13 sampling technique for assessing the concentrations of PCBs and organochlorine contaminants
14 in the blubber of bottlenose whales.

15 *4.1 Bottlenose whale contaminant levels*

16 Contaminant concentrations have been previously reported for northern bottlenose whales in two
17 studies. A juvenile male stranded in the North Sea in 1976 contained 37,100 ng/g total PCBs
18 and 14,200 ng/g total DDT in blubber on a wet weight basis (Harms et al., 1978). Oil from a
19 northern bottlenose whale caught in the Canadian fishery in 1962 (which would have been from
20 The Gully region, according to Reeves et al., 1993) contained appreciable amounts of
21 contaminant residues, including total DDT at 11,600 ng/g of oil, and PCBs at 1,000 ng/g of oil
22 (Addison et al., 1972). The DDT in this sample was composed of approximately equal parts of
23 4,4' DDE and 4,4' DDD, but 4,4' DDT was not detected. Some of these differences in
24 contaminant levels between the present study and these earlier reports for bottlenose whales are
25 probably due to recent advances in analytical techniques, such as the congener-specific analysis
26 of PCBs. Unfortunately, this makes any inference regarding historical changes in contaminant
27 levels unreliable.

28 The concentrations of contaminants found in the blubber of northern bottlenose whales are

1 generally consistent with concentrations reported for other large odontocetes in the north
2 Atlantic (Table 3). It is generally accepted that maternal transfer of hydrophobic contaminants in
3 marine mammals reduces the concentrations of these compounds in females relative to males.
4 The large variability seen in the concentrations of contaminants in whales is probably due to
5 differences in sex, age and size of the animals that were biopsied (Jepson et al., 2005; Muir et
6 al., 1996; Ross et al., 2000; Stern et al., 1994; Subramanian et al., 1988; Tuerk et al., 2005;
7 Weisbrod et al., 2000). The DDT/PCB ratio in whales has been suggested to correlate to
8 distance of ecosystem location from the mainland (Aguilar, 1987). Our results demonstrated
9 high DDT/PCB ratios for bottlenose whales, consistent with other beaked whales and pilot
10 whales (*Globicephala melas*), but contrary to Atlantic sperm whales (*Physeter macrocephalus*),
11 arctic narwhal (*Monodon monoceros*) and beluga (Table 3). However, Stern et al. (1994)
12 observed that these ratios in arctic belugas varied with age; especially in females due to
13 differential loss of DDT and PCBs.

14 The concentrations of total DDT and total PCBs in the blubber of bottlenose whales were lower
15 than the concentrations reported in highly contaminated odontocete species (e.g., total PCB
16 concentrations of >250,000 ng/g lipid for killer whales in British Columbia, Canada, Ross et al.,
17 2000; and total PCB concentrations of >60,000 ng/g lipid, and 4,4-DDE concentration >16,000
18 ng/g lipid for belugas in the St. Lawrence estuary, Canada, Muir et al., 1996). Total blubber PCB
19 levels >17,000 ng/g lipid have been associated with an elevated prevalence of infectious
20 diseases in harbor porpoises in the U.K. (Jepson et al., 2005). Schwacke et al. (2002) estimated
21 an excess risk of reproductive failure in bottlenose dolphins (*Tursiops truncatus*) from the coast
22 of the southeast USA with total blubber PCB concentrations of between 4,200 (adult female)
23 and 91,200 ng/g lipid (adult male). Hall et al. (2006) used population models to predict that high
24 concentrations of PCBs in bottlenose dolphins in Florida, USA would depress the growth of the
25 population. However, the concentrations of persistent contaminants, and in particular PCBs, in
26 the bottlenose whales from The Gully region were lower magnitude than these deleterious
27 concentrations.

28 The CYP1A1 analyses showed low levels of expression of this cytochrome P450 enzyme. It is

1 thought that CYP1A1 induction only persists for the duration of exposure (Fossi et al., 1994).
2 Compounds that can induce CYP1A1 include coplanar PCB congeners, halogenated dioxins and
3 dibenzofurans and polynuclear aromatic hydrocarbons (PAHs) (Safe, 2002). Of the
4 contaminants we sampled in the blubber it is only the few mono-*ortho* and non-*ortho* PCBs that
5 would be capable of inducing CYP1A1. The other PCBs and organochlorine contaminants
6 would not induce CYP1A1. Our analysis confirms that contaminants found in the blubber did
7 not correlate with induction of the cytochrome P4501A1 class of enzymes in the skin. This
8 indicates that the planar PCBs detected in the blubber layer were not being actively mobilized
9 and circulated to the skin during or near sampling of these animals. Considering that lower
10 concentrations were found in the animals expressing the highest levels of CYP1A1 (Davis Strait
11 samples) and there was no change in any measured contaminant that also interacts with the Ah
12 receptor in the Gully animals over time, the causative agent inducing CYP1A1 was not
13 measured in this study and remains unknown. No attempt was made to evaluate exposure of the
14 bottlenose whales to PAHs, since these compounds are rapidly metabolized and do not persist in
15 the tissues of vertebrates.

16 *4.2 Changes detected between 1996-97 and 2002-03 for samples from The Gully*

17 There was a significant increase observed in the concentrations of some contaminants (e.g. 4,4-
18 DDE, trans-nonachlor) in whales from The Gully over the five-year time span from 1996-97 to
19 2002-03. There could be many reasons for this observation, including metabolic changes in
20 contaminant proportions, changes in feeding patterns or in the geographic range of the animals.
21 Over time DDT is metabolized into several forms, of which DDE is a derivative that is difficult
22 to degrade or excrete and is thus accumulated (Aguilar, 1984). However, since we have
23 observed increases in total DDTs in addition to some increase in DDE/total(DDT) proportion, it
24 is unlikely that the increased 4,4' DDE observed is simply caused by metabolism. It is also
25 unlikely that these whales changed their feeding patterns or geographic range. A study of the
26 diving behaviour of bottlenose whales in The Gully showed that these whales consistently
27 performed deep dives to the sea-floor, presumably to feed on deep-sea squid (Hooker and Baird,
28 1999; Hooker et al., 2001b). Work on the ranging behavior of individual whales suggests that

1 they are relatively site-faithful (Dalebout et al., 2001; Whitehead et al., 1997; Wimmer and
2 Whitehead, 2004). They do not appear to travel widely and obtain their food from a local
3 source; in fact, some form of territoriality may even be present (Hooker et al., 2002). There have
4 been no observed changes in foraging or ranging behaviour that might drive such difference in
5 contaminant load (personal observations by S.H. and H.W.). The change in contaminant levels
6 over time in these whales is therefore more likely to reflect a temporal change in contaminant
7 levels in the water and/or in prey species. This could be due to alterations in the patterns of
8 oceanic currents, but it cannot be ruled out that the proximity of oil and gas drilling activities
9 may have influenced contaminant patterns. These organochlorine contaminants are not likely to
10 have been released directly from oil rigs or through seismic exploration, but it is possible that oil
11 and gas activities have led to remobilization of persistent contaminants from sediments on the
12 Scotian Shelf.

13 Changes in CYP1A1 expression in Gully samples did not show a consistent pattern of increase
14 between 1996-1997 to 2002-03, in that scores of whales sampled in 2002 showed no difference
15 from those sampled in 1996 and 1997. However, samples in 2003 showed significantly higher
16 CYP1A1 expression than observed during other years of sampling in The Gully (Fig. 2).
17 Coincident with this were several spills of kerosene and streamer fluids during seismic survey
18 work conducted during June-August, 2003 (personal communication from Canada Nova Scotia
19 Offshore Petroleum Board). A total of ten spills of kerosene and streamer fluids were reported,
20 ranging in size from 0.008 – 0.5 cubic metres. Four spills occurred during June, and three in
21 each of July and August. At no time were slicks observed, but fluid was recorded missing when
22 streamers were recovered. Whether such streamer fluids contained compounds which induce
23 CYP1A1 which were not recorded in the blubber contaminants (such as PAHs) is unknown.

24 *4.2 Differences between contaminant levels in The Gully and in the Davis Strait*

25 Blubber contaminant concentrations in northern bottlenose whales from the Davis Strait were
26 significantly lower than concentrations in The Gully whales. This difference could simply
27 reflect differences in the oceanic distribution of contaminants on the Scotian Shelf relative to the
28 Davis Strait in the eastern arctic, consistent with general observations of elevated contaminant

1 exposure at mid-latitudes that is associated with nearby industrialization (Aguilar et al., 2002).
2 However, despite a relatively small sample size, levels of CYP1A1 expression were consistently
3 higher within the three animals sampled from the Davis Strait population in comparison to those
4 sampled from The Gully population.

5 Elevated CYP1A1 expression could theoretically reflect mobilization of blubber fats and the
6 coplanar PCBs therein. If this was occurring, it might suggest that northern bottlenose whales in
7 the Davis Strait were undergoing a period of poor feeding. However, we have no supporting
8 evidence for this hypothesis. Alternatively the higher levels of CYP1A1 exposure in the Davis
9 Strait could be due to exposure to compounds (such as PAHs) which cause CYP1A1 expression,
10 but cannot be detected in tissue. PAHs are rapidly metabolized in marine mammals and are
11 generally present at low to non-detectable concentrations in the tissues of odontocetes (Law and
12 Whinnett, 1992), but studies of beluga whales from the highly contaminated St. Lawrence region
13 have shown evidence of PAH exposure through the presence in tissues of aromatic DNA
14 adducts (Ray et al., 1991). Exposure to PAHs near oil and gas drilling operations is poorly
15 understood and may be a cause for concern. It is possible that there are elevated levels of one or
16 more compounds that interact with the Ah-receptor (e.g., coplanar PCBs, halogenated dioxins
17 and dibenzofurans) that were not quantified in our analysis of blubber contaminants. Elevated
18 levels of persistent contaminants, primarily PCBs have been observed in the arctic, associated
19 with military sites such as the radar station at Saglek Bay, Labrador (approx 250 km from
20 sampling locations) (Kuzyk et al., 2005). CYP1A activity has been found to be relatively high
21 in the tissues of polar bears in the Canadian Arctic (Muir et al., 1999), and these authors
22 suspected that non-*ortho* and mono-*ortho* PCBs were responsible for this activity. There has
23 been recent industrial activity in Voisey's Bay (35 km southwest of Nain, Labrador), where
24 nickel-copper-cobalt mining began in 2004. However, this is over 200km from the region where
25 the northern bottlenose whales were sampled and their deep-water habitat, and so, unless there
26 have been major spills, this is unlikely to have caused a large impact on these whales.

27 *4.3 Conclusions*

28 Blubber contaminants found in biopsies from free-ranging northern bottlenose whales from

1 eastern Canada showed concentrations and patterns consistent with those reported for other
2 North Atlantic cetaceans. These levels were below those suspected of causing health problems
3 in more contaminated odontocetes which is reassuring in the assessment of an “endangered”
4 population of whales in a Marine Protected Area. However, blubber contaminant concentrations
5 were higher for whales sampled in The Gully than for those sampled in the Davis Strait.
6 Furthermore, contaminant levels in The Gully population increased between the 1996-97 and
7 2002-03 time periods for 4,4-DDE and trans-nonachlor, although whether this increase was
8 related to the onset of nearby drilling for gas is unknown.

9 Northern bottlenose whales do not generally appear to show high expression of CYP1A1, nor
10 are they particularly sensitive to CYP1A1 inducers compared to other odontocetes. However,
11 there was a significant difference in CYP1A1 expression observed in 2003 for The Gully
12 population compared to that observed in 1996, 1997 or 2002. Since this was the final year of
13 sampling, it is not known whether this was a single year event, or whether this may have
14 continued beyond 2003. Our results also suggest that animals in the Davis Strait may be
15 exposed to higher concentrations of CYP1A1 inducing contaminants than Gully animals.
16 Repeated sampling of animals in both The Gully and the Davis Strait would be useful to help
17 resolve these questions.

18 Differences between CYP1A1 results and blubber contaminants demonstrate the utility of the
19 comparison between these two techniques. Both have highlighted potential concerns for the
20 Gully marine protected area due to development of the oil and gas industry.

21

22 **Acknowledgements**

23 We are grateful to all the crew members of the *Balaena*, particularly Merel Dalebout, who
24 participated in fieldwork in The Gully and the Davis Strait, and to Phil Clapham and other crew
25 members of the NOAA vessel *Delaware II*, who facilitated fieldwork in The Gully during 2002.
26 Dr. M.L. Dalebout genotyped the biopsy samples to obtain data on the sexes of whales sampled.
27 Dr. S. Gowans and T. Wimmer provided details on whale ID from photographs taken during

1 biopsy sampling. Research was funded by the Natural Sciences and Engineering Research
2 Council (NSERC) of Canada, World Wildlife Fund Canada Endangered Species Recovery Fund,
3 Fisheries and Oceans Canada, the National Geographic Society, the Canadian Federation of
4 Humane Societies and two U.K. Royal Society International Collaborative Awards. S.K.H. was
5 supported by a Canadian Commonwealth Scholarship and Royal Society Dorothy Hodgkin
6 Research Fellowship. C.D.M. was awarded a Discovery grant from the Natural Sciences and
7 Engineering Research Council (NSERC) of Canada. J.Y.W was supported by an NSERC PGS B
8 fellowship and the Woods Hole Oceanographic Institution. A. J. Hall kindly reviewed the
9 manuscript.

10

11

1 **References**

2 Addison, R.F., Zinck, M.E., Ackman, R.G., 1972. Residues of organochlorine pesticides and
3 polychlorinated biphenyls in some commercially produced Canadian marine oils. *Journal*
4 *of the Fisheries Research Board of Canada*, 29, 349-355.

5 Aguilar, A., 1983. Organochlorine pollution in sperm whales, *Physeter macrocephalus*, from
6 temperate waters of the Eastern North Atlantic. *Marine Pollution Bulletin*, 14, 349-352.

7 Aguilar, A., 1984. Relationship of DDE/Sigma-DDT in marine mammals to the chronology of
8 DDT input into the ecosystem. *Canadian Journal of Fisheries and Aquatic Sciences*, 41,
9 840-844.

10 Aguilar, A., 1987. Using organochlorine pollutants to discriminate marine mammal populations:
11 a review and critique of the methods. *Marine Mammal Science*, 3, 242-262.

12 Aguilar, A., Borrell, A., Reijnders, P.J.H., 2002. Geographical and temporal variation in levels
13 of organochlorine contaminants in marine mammals. *Marine Environmental Research*, 53,
14 425-452.

15 Angell, C.M., Wilson, J.Y., Moore, M.J., Stegeman, J.J., 2004. Cytochrome P450 1A1
16 expression in cetacean integument: implications for detecting contaminant exposure and
17 effects. *Marine Mammal Science*, 20, 554-566.

18 Borrell, A., 1993. PCB and DDTs in blubber of cetaceans from the northeastern North Atlantic.
19 *Marine Pollution Bulletin*, 26, 146-151.

20 Dalebout, M.L., Hooker, S.K., Christensen, I., 2001. Genetic diversity and population structure
21 among northern bottlenose whales, *Hyperoodon ampullatus*, in the western North Atlantic
22 Ocean. *Canadian Journal of Zoology*, 79, 478-484.

23 Dalebout, M.L., Ruzzante, D.E., Whitehead, H., Oien, N.I., 2006. Nuclear and mitochondrial
24 markers reveal distinctiveness of a small population of bottlenose whales (*Hyperoodon*
25 *ampullatus*) in the western North Atlantic. *Molecular Ecology*, 15, 3115-3129.

26 Draushuk, A.T., McGarrigle, B.P., Larsen, K.E., Stegeman, J.J., Olson, J.R., 1998. Detection
27 of CYP1A1 protein in human liver and induction by TCDD in precision-cut liver slices
28 incubated in dynamic organ culture. *Carcinogenesis*, 19, 1361-1368.

- 1 Fenton, D.G., Macnab, P.A., Rutherford, R.J., 2002. The Sable Gully marine protected area
2 initiative: history and current efforts. In *Managing protected areas in a changing world*.
3 (eds S. Bondrup-Nielson, N.W.P. Munro, G. Nelson, J.H.M. Willison, T.B. Herman & P.
4 Eagles), pp. 1343-1355. SAMPAA, Wolfville, Canada.
- 5 Fossi, M.C., Leonzio, C., Peakall, D., 1994. The use of nondestructive biomarkers in the hazard
6 assessment of vertebrate populations. In *Nondestructive biomarkers in vertebrates* (eds
7 M.C. Fossi & C. Leonzio), pp. 34-47. Lewis Publishers, Ann Arbor, MI, USA.
- 8 Gauthier, J.M., Metcalfe, C.D., Sears, R., 1997. Validation of the blubber biopsy technique for
9 monitoring of organochlorine contaminants in balaenopterid whales. *Marine*
10 *Environmental Research*, 43, 157-179.
- 11 Godard, C.A.J., Smolowitz, R.M., Wilson, J.Y., Payne, R.S., Stegeman, J.J., 2004. Induction of
12 cetacean cytochrome P4501A1 by b-naphthoflavone exposure of skin biopsy slices.
13 *Toxicological Sciences*, 80, 268-275.
- 14 Gowans, S., Dalebout, M.L., Hooker, S.K., Whitehead, H., 2000a. Reliability of photographic
15 and molecular techniques for sexing northern bottlenose whales (*Hyperoodon ampullatus*).
16 *Canadian Journal of Zoology*, 78, 1224-1229.
- 17 Gowans, S., Whitehead, H., Arch, J.K., Hooker, S.K., 2000b. Population size and residency
18 patterns of northern bottlenose whales (*Hyperoodon ampullatus*) using the Gully, Nova
19 Scotia. *Journal of Cetacean Research and Management*, 2, 201-210.
- 20 Hall, A.J., Hugunin, K., Deaville, R., Law, R.J., Allchin, C.R., Jepson, P.D., 2006. The risk of
21 infection from polychlorinated biphenyl exposure in the harbor porpoise (*Phocoena*
22 *phocoena*): a case-control approach. *Environmental Health Perspectives*, 114, 704-711.
- 23 Harms, U., Drescher, H.E., Huschenbeth, E., 1978. Further data on heavy metals and
24 organochlorines in marine mammals from German coastal waters. *Meeresforschung*, 26,
25 153-161.
- 26 Hobbs, K.E., Muir, D.C.G., Michaud, R., Beland, P., Letcher, R.J., Norstrom, R.J., 2003. PCBs
27 and organochlorine pesticides in blubber biopsies from free-ranging St Lawrence River
28 Estuary beluga whales (*Delphinus leuchas*), 1994-1998. *Environmental Pollution*, 122,
29 291-302.

- 1 Hooker, S.K., Baird, R.W., 1999. Deep-diving behaviour of the northern bottlenose whale,
2 *Hyperoodon ampullatus* (Cetacea: Ziphiidae). Proceedings of the Royal Society of London
3 B, 266, 671-676.
- 4 Hooker, S.K., Baird, R.W., Al-Omari, S., Gowans, S., Whitehead, H., 2001a. Behavioral
5 reactions of northern bottlenose whales (*Hyperoodon ampullatus*) to biopsy darting and
6 tag attachment procedures. Fishery Bulletin, 99, 303-308.
- 7 Hooker, S.K., Iverson, S.J., Ostrom, P., Smith, S.C., 2001b. Diet of northern bottlenose whales
8 inferred from fatty-acid and stable-isotope analyses of biopsy samples. Canadian Journal
9 of Zoology, 79, 1442-1454.
- 10 Hooker, S.K., Whitehead, H., Gowans, S., 1999. Marine protected area design and the spatial
11 and temporal distribution of cetaceans in a submarine canyon. Conservation Biology, 13,
12 592-602.
- 13 Hooker, S.K., Whitehead, H., Gowans, S., Baird, R.W., 2002. Fluctuations in distribution and
14 patterns of individual range use of northern bottlenose whales. Marine Ecology Progress
15 Series, 225, 287-297.
- 16 Jepson, P.D., Bennett, P.M., Deaville, R., Allchin, C.R., Baker, J.R., Law, R.J., 2005.
17 Relationships between polychlorinated biphenyls and health status in harbor porpoises
18 (*Phocoena phocoena*) stranded in the United Kingdom. Environmental Toxicology and
19 Chemistry, 24, 238-248.
- 20 Knap, A.H., Jickells, T.D., 1983. Trace metals and organochlorines in the goosebeaked whale.
21 Marine Pollution Bulletin, 14, 271-274.
- 22 Koopman, H.N., Iverson, S.J., Gaskin, D.E., 1996. Stratification and age-related differences in
23 blubber fatty acids of the male harbour porpoise (*Phocoena phocoena*). Journal of
24 Comparative Physiology B, 165, 628-639.
- 25 Kuzyk, Z.A., Stow, J.P., Burgess, N.M., Solomon, S.M., Reimer, K.J., 2005. PCBs in sediments
26 and the coastal food web near a local contaminant source in Saglek Bay, Labrador. Science
27 of the Total Environment, 351-352, 264-284.
- 28 Law, R.J., Allchin, C.R., Jones, B.R., Jepson, P.D., Baker, J.R., Spurrier, C.J.H., 1997. Metals
29 and organochlorines in tissues of a Blainville's beaked whale (*Mesoplodon densirostris*)

1 and a killer whale (*Orcinus orca*) stranded in the United Kingdom. Marine Pollution
2 Bulletin, 34, 208-212.

3 Law, R.J., Whinnett, J.A., 1992. Polycyclic aromatic-hydrocarbons in muscle-tissue of harbor
4 porpoises (*Phocoena phocoena*) from UK waters. Marine Pollution Bulletin, 24, 550-553.

5 Mead, J.G., 1989. Bottlenose whales *Hyperoodon ampullatus* (Forster, 1770) and *Hyperoodon*
6 *planifrons* Flower, 1882. In Handbook of marine mammals (eds S.H. Ridgway & R.
7 Harrison), Vol. 4: River dolphins and the larger toothed whales, pp. 321-348. Academic
8 Press, London.

9 Metcalfe, T.L., Metcalfe, C.D., 1997. The trophodynamics of PCBs, including mono- and non-
10 ortho congeners, in the food web of North-Central Lake Ontario. Science of the Total
11 Environment, 201, 245-272.

12 Muir, D., Braune, B., DeMarch, B., Norstrom, R., Wagemann, R., Lockhart, L., Hargrave, B.,
13 Bright, D., Addison, R., Payne, J., Reimer, K., 1999. Spatial and temporal trends and
14 effects of contaminants in the Canadian Arctic marine ecosystem: a review. Science of the
15 Total Environment, 230, 83-144.

16 Muir, D.C.G., Ford, C.A., Grift, N.P., Stewart, R.E.A., 1992. Organochlorine contaminants in
17 narwhal (*Monodon monoceros*) from the Canadian Arctic. Environmental Pollution, 75,
18 307-316.

19 Muir, D.C.G., Ford, C.A., Rosenberg, B., Norstrom, R.J., Simon, M., Beland, P., 1996.
20 Persistent organochlorines in beluga whales (*Delphinapterus leucas*) from the St Lawrence
21 River estuary .1. Concentrations and patterns of specific PCBs, chlorinated pesticides and
22 polychlorinated dibenzo-p-dioxins and dibenzofurans. Environmental Pollution, 93, 219-
23 234.

24 Ray, S., Dunn, B.P., Payne, J.F., Fancey, L., Helbig, R., Beland, P., 1991. Aromatic DNA-
25 carcinogen adducts in beluga whales from the Canadian Arctic and Gulf of St-Lawrence.
26 Marine Pollution Bulletin, 22, 392-396.

27 Reeves, R.R., Mitchell, E., Whitehead, H., 1993. Status of the northern bottlenose whale,
28 *Hyperoodon ampullatus*. Canadian Field-Naturalist, 107, 490-508.

- 1 Ross, P.S., Ellis, G.M., Ikonomou, M.G., Barrett-Lennard, L.G., Addison, R.F., 2000. High PCB
2 concentrations in free-ranging Pacific killer whales, *Orcinus orca*: Effects of age, sex and
3 dietary preference. *Marine Pollution Bulletin*, 40, 504-515.
- 4 Safe, S.H., 1986. Comparative toxicology and mechanism of action of polychlorinated dibenzo-
5 P-dioxins and dibenzofurans. *Annual Review of Pharmacology and Toxicology*, 26, 371-
6 399.
- 7 Safe, S.H., 2002. Exposure to POPs and other congeners. In *Biomarkers of environmentally*
8 *associated disease: technologies, concepts, and perspectives* (eds S.H. Wilson & W.A.
9 Suk), pp. 455-470. Lewis Publishers, Ann Arbor, MI, USA.
- 10 Schwacke, L.H., Voit, E.O., Hansen, L.J., Wells, R.S., Mitchum, G.B., Hohn, A.A., Fair, P.A.,
11 2002. Probabilistic risk assessment of reproductive effects of polychlorinated biphenyls on
12 bottlenose dolphins (*Tursiops truncatus*) from the southeast United States coast.
13 *Environmental Toxicology and Chemistry*, 21, 2752-2764.
- 14 Stegeman, J.J., Brouwer, M., Di Giulio, R.T., Forlin, L., Fowler, B.A., Sanders, B.M., Van
15 Veld, P.A., 1992. Enzyme and protein synthesis as indicators of contaminant exposure and
16 effect. In *Biomarkers: biochemical, physiological, and histological markers of*
17 *anthropogenic stress* (eds R.J. Huggett, R.A. Kimerle, P.M.J. Mehrle & H.L. Bergman),
18 pp. 123-138. Lewis Publishers, Ann Arbor, MI, USA.
- 19 Stern, G.A., Muir, D.C.G., Segstro, M.D., Dietz, R., Heide-Jorgensen, M., 1994. PCB's and
20 other organochlorine contaminants in white whales (*Delphinapterus leucas*) from West
21 Greenland: variations with age and sex. *Meddelelser om Gronland, Bioscience*, 39, 245-
22 259.
- 23 Subramanian, A., Tanabe, S., Tatsukawa, R., 1988. Estimating some biological parameters of
24 Baird's beaked whales using PCBs and DDE as tracers. *Marine Pollution Bulletin*, 19,
25 284-287.
- 26 Taruski, A.G., Olney, C.E., Winn, H.E., 1975. Chlorinated hydrocarbons in cetaceans. *Journal*
27 *of the Fisheries Research Board of Canada*, 32, 2205-2209.

1 Tuerk, K.J.S., Kucklick, J.R., McFee, W.E., Pugh, R.S., Becker, P.R., 2005. Factors influencing
2 persistent organic pollutant concentrations in the atlantic white-sided dolphin
3 (*Lagenorhynchus acutus*). Environmental Toxicology and Chemistry, 24, 1079-1087.

4 Weisbrod, A.V., Shea, D., Moore, M.J., Stegeman, J.J., 2000. Bioaccumulation patterns of
5 polychlorinated biphenyls and chlorinated pesticides in Northwest Atlantic pilot whales.
6 Environmental Toxicology and Chemistry, 19, 667-677.

7 Whitehead, H., Gowans, S., Faucher, A., McCarrey, S.W., 1997. Population analysis of northern
8 bottlenose whales in the Gully, Nova Scotia. Marine Mammal Science, 13, 173-185.

9 Whitehead, H., Wimmer, T., 2005. Heterogeneity and the mark-recapture assessment of the
10 Scotian Shelf population of northern bottlenose whales (*Hyperoodon ampullatus*).
11 Canadian Journal of Fisheries and Aquatic Sciences, 62, 2573-2585.

12 Wimmer, T., Whitehead, H., 2004. Movements and distribution of northern bottlenose whales,
13 *Hyperoodon ampullatus*, on the Scotian Shelf and in adjacent waters. Canadian Journal of
14 Zoology, 82, 1782-1794.

15

1 **Figure Legends**

2

3 Figure 1. Map showing location of The Gully and the Davis Strait populations of northern
4 bottlenose whales. Expanded area shows The Gully region in the northwest Atlantic, with the
5 locations of the recently developed oil and gas platforms shown (with names of the platforms).

6

7 Figure 2. Scores for expression of cytochrome P450 1A1 (CYP1A1) in the vascular endothelial
8 layer and smooth muscle layer of biopsy samples from northern bottlenose whales for each year
9 and location sampled. Mean (\pm S.D.) are shown.

10

11 Figure 3. Mean concentrations (\pm S. D.) in ng/g lipid of total PCBs and classes of
12 organochlorine compounds in blubber biopsies taken from bottlenose whales in The Gully in
13 1996-97, versus those in 2002-2003, and those taken from whales in the Davis Strait in 2003.
14 Separate plots are shown for male and female whales.

15

16 Figure 4. Mean concentrations (\pm S. D.) of 2,4- and 4,4-isomers of DDE, DDD and DDT in ng/g
17 lipid from blubber biopsies collected from male and female bottlenose whales in The Gully in
18 1996-97 and in 2002-03, and in the Davis Strait in 2003.

19

20 Figure 5. Mean concentrations (\pm S. D.) of *cis*- and *trans*-isomers of chlordane and nonachlor in
21 ng/g lipid from blubber biopsies collected from male and female bottlenose whales in The Gully
22 in 1996-97 and in 2002-03, and in the Davis Strait in 2003.

23

1 Table 1. Location, dates and times of collection, sex, ID (where determined) and analytical
 2 history for biopsy samples collected from northern bottlenose whales for analysis of
 3 concentrations of persistent contaminants and activity of CYP1A1 enzyme. Scores for
 4 expression of cytochrome P450 1A1 (CYP1A1) in the vascular endothelial layer (Endo) and
 5 smooth muscle layer (Sm muscle) is shown. The CYP1A1 scale ranges from 0 to 15. M =
 6 Male, F = Female.

7

LOCATION	DATE	TIME	SEX	ID	Contaminants	CYP1A1 Endo / Sm muscle
The Gully	27 Aug 96	10:28	M	-	-	0 / 0
	27 Aug 96	11:05	M	143	-	0 / 0
	16 July 97	12:00	F	1289	x	0 / 0
	16 July 97	15:36	F	-	x	0 / 0
	16 July 97	17:20	F	-	x	0 / 0
	16 July 97	19:32	F	54	-	0 / 0
	12 Aug 97	14:48	F	961	-	-
	12 Aug 97	16:07	F	1000	x	0 / 0
	12 Aug 97	18:53	F	1313	-	0 / 0
	12 Aug 97	19:18	F	-	-	0 / 0
	13 Aug 97	14:57	F	1318	-	0 / 0
	13 Aug 97	16:33	F	1315	-	0 / 0
	13 Aug 97	16:44	F	619	-	0 / 0
	14 Aug 97	07:50	F	-	x	0 / 0
	14 Aug 97	10:00	M	-	x	0 / 0
	14 Aug 97	10:07	M	480*	x	0 / 0
	16 Aug 97	08:59	M	1039	x	0 / 0
	16 Aug 97	10:00	M	-	x	0 / 0
	16 Aug 97	12:22	F	1336	-	3 / 0
	16 Aug 97	13:16	M	-	x	0 / 0
	08 Aug 02	14:31	F	677	x	0 / 0
	08 Aug 02	14:42	M	76	x	0 / 0
	08 Aug 02	16:57	M	3	x	0 / 0
	14 Aug 02	09:18	F	-	x	0 / 0
	14 Aug 02	09:27	M	-	-	0 / 0
	14 Aug 02	11:38	F	-	x	0 / 0
	14 Aug 02	12:17	M	-	-	0 / 0
	14 Aug 02	15:08	F	251	x	0 / 0
	14 Aug 02	15:19	F	102	-	-
	29 Aug 03	07:08	M	-	-	0 / 1.5
	29 Aug 03	07:18	M	-	x	2 / 5
	29 Aug 03	12:01	M	-	x	0 / 2.5
	29 Aug 03	12:50	M	824	x	4.5 / 2.5
	29 Aug 03	19:38	M	480*	x	-

	25 Sept 03	14:09	F	-	-	0 / 2
	25 Sept 03	19:04	F	-	-	0 / 1.5
Davis Strait	14 Aug 03	11:53	F	-	x	3 / 6
	16 Aug 03	09:59	M	2209	x	3 / 9
	16 Aug 03	18:16	M	-	x	4.5 / 4

1 * The same male was sampled in both 1996 and 2003.

Table 2. Mean and standard deviations about the mean of concentrations (ng/g lipid) of PCBs and organochlorine compounds in blubber biopsies of northern bottlenose whales from The Gully and the Davis Strait, Labrador. nd = not detected, na = not analyzed.

Analyte	Gully 1996 - 1997		Gully 2002 - 2003		Davis Strait 2003		# 480 1996	# 480 2003		
	males mean n=5	S.D.	females mean n=5	S.D.	males mean n=6	females mean n=4	males mean n=2	female n=1	male n=1	
PCB congeners										
18	na	na	na	10.1	6.5	5.9	4.9	1.5	na	nd
17	na	na	na	0.0	0.0	0.5	1.3	1.8	na	nd
31	na	na	na	15.5	7.8	13.6	nd	-	na	nd
28	na	na	na	0.3	0.1	1.9	9.0	0.5	na	nd
33	na	na	na	-	nd	1.4	nd	-	na	nd
52	534.2	170.7	258.1	278.2	401.7	199.6	161.8	9.7	288.9	286.5
49	145.4	81.9	85.6	72.7	110.4	56.8	42.0	0.1	50.5	10.5
47	127.6	47.6	144.5	na	na	na	na	na	75.4	na
44	109.0	58.9	35.6	43.1	70.0	33.8	40.5	5.5	62.6	38.8
66	16.7	37.2	52.9	na	na	na	na	na	nd	na
74	na	na	na	109.9	93.5	12.3	66.1	25.7	na	58.5
70	na	na	na	17.5	11.0	34.1	18.1	6.0	na	nd
95	na	na	na	272.0	346.5	120.1	160.2	17.6	na	272.6
101	826.2	219.1	403.3	284.0	449.6	286.7	198.1	15.5	539.9	263.4
99	365.5	74.8	197.0	348.0	662.5	647.5	906.8	254.5	257.1	450.0
87	270.2	70.5	122.4	98.6	160.5	99.0	41.5	2.7	169.5	91.8
110	133.0	57.0	121.2	26.4	18.3	108.0	47.3	43.4	94.9	7.1
151	na	na	na	2138.7	1584.1	210.5	2399.2	65.8	na	866.9
149	532.3	98.9	258.5	412.4	636.2	407.5	229.5	42.1	376.0	466.9
118	722.2	161.3	389.9	428.7	651.3	403.7	443.5	152.1	475.1	450.6
153	1209.8	241.6	680.1	842.2	1515.4	768.3	666.5	28.0	861.9	1394.0
132	na	na	na	138.6	175.7	81.2	81.6	10.0	na	149.8
105	134.3	28.6	96.3	85.8	97.6	49.7	129.4	57.4	99.6	78.2
138	1284.4	226.5	671.2	863.1	1387.9	608.3	425.4	133.3	941.6	1424.6
187	na	na	na	187.9	392.1	234.7	166.5	80.4	na	340.1
128	na	na	na	64.3	158.4	191.0	52.4	22.0	na	106.8
156	73.6	22.1	35.3	41.1	55.2	31.4	12.7	3.1	44.7	47.5
180	574.3	169.5	267.9	213.9	443.4	244.7	149.0	33.8	353.5	386.3
191	na	na	na	2.2	1.2	1.3	nd	-	na	nd

mirex	30.0	13.0	19.7	5.6	35.6	15.2	13.9	0.1	19.7	9.9	13.7	29.6	34.3
HCb	236.4	103.7	254.2	223.7	325.2	413.3	46.6	93.2	559.8	16.5	35.8	227.9	269.3
octachlorostyrene	49.0	24.5	13.1	11.7	9.1	14.9	12.1	15.4	nd	-	nd	68.4	nd

1) Total PCBs calculated only for the major PCB congeners in bold, as these were analyzed in samples collected in both 1996-97 and 2002-03.

Table 3. Concentrations of total PCB (Σ PCB) and total DDT (Σ DDT) in the blubber of male and female bottlenose whales from this study in comparison to concentrations reported for the blubber of other odontocetes from the North Atlantic. All totals are shown in ng/g. Either means or ranges are provided. N = Number of animals, m = male, f = female

Species	Location	N	% lipid	Σ PCB ¹	Σ DDT ²	Conc. Units	Source
Northern bottlenose whale, <i>Hyperoodon ampullatus</i>	The Gully 1996-97	5m	40.4	7,270	14,660	lipid	This study
		5f	66.3	3,790	4,140		
	The Gully 2002-03	6m	50.9	6,840	21,570	lipid	This study
Davis Strait 2003	Davis Strait 2003	4f	43.2	4,060	7,600		
		2m	44.6	3,560	3,580	lipid	This study
Cuvier's beaked whale, <i>Ziphius cavirostris</i>	Bermuda	1f	38.3	530	1,080		
		3m	-	9,610	31,190	wet weight	Knap and Jickells, 1983
Blainville's beaked whale, <i>Mesoplodon densirostris</i>	UK	1f	83	4,780	6,150	wet weight	Law et al., 1997
		2m	-	14,000-29,000	38,200-65,100	wet weight	Taruski et al., 1975
Sperm whale, <i>Physeter macrocephalus</i>	Iceland	10m	40.3	10,510	7,800	lipid	Borrell, 1993
		8m	66.4	9,930	5,100	lipid	Aguilar, 1983
Pilot whale, <i>Globicephala melas</i>	Cape Cod, USA	6f	68.9	15,550	7,730		
		21, m&f	-	7,550	18,340	lipid	Weisbrod et al., 2000
Narwhal, <i>Monodon monoceros</i>	Canadian arctic	15m	-	5,180	5,920	wet weight	Muir et al., 1992
		6f	-	2,700	2,540		
Beluga whale, <i>Delphinapterus leucas</i>	Greenland	71m	-	1,800-10,900	1,000-8,100	wet weight	Stern et al., 1994
		67f	-	800-7,200	400-5,400		

1) Σ PCB calculated for various numbers of congeners in each study.

2) Σ DDT calculated from 4,4- and 2,4-isomers for this study and Weisbrod et al. (2000); Σ DDT calculated from only 4,4-isomers for all other studies.

Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5

