

Review Article

Socio-economic determinants of micronutrient intake and status in Europe: a systematic review

Romana Novaković^{1,2,*}, Adriënne Cavelaars², Anouk Geelen², Marina Nikolić¹, Iris Iglesia Altaba³, Blanca Roman Viñas⁴, Joy Ngo⁴, Mana Golsorkhi⁵, Marisol Warthon Medina⁶, Anna Brzozowska⁷, Anna Szczecinska⁷, Diederik de Cock⁸, Greet Vansant⁸, Marianne Renkema², Lluís Serra Majem^{4,9}, Luis Aznar Moreno³, Maria Glibetić¹, Mirjana Gurinović¹, Pieter van't Veer² and Lisette CPGM de Groot²

¹Centre of Research Excellence in Nutrition and Metabolism, Institute for Medical Research, University of Belgrade, Tadeuska Kosuska 1, 11000 Belgrade, Serbia: ²Division of Human Nutrition, Wageningen University and Research Centre, Wageningen, The Netherlands: ³Growth, Exercise, Nutrition and Development (GENUD) Research Group, School of Health Science (EUCS), University of Zaragoza, Zaragoza, Spain: ⁴Community Nutrition Research Centre of the Nutrition Research Foundation, University of Barcelona, Barcelona, Spain: ⁵Department of Sport and Health Sciences, Oxford Brookes University, Oxford, UK: ⁶Centre for Applied Sport and Exercise Sciences, University of Central Lancashire, Preston, UK: ⁷Faculty of Human Nutrition and Consumer Sciences, Warsaw University of Life Sciences (WULS-SGGW), Warsaw, Poland: ⁸Department of Nutrition–Public Health Medicine, IFOCe (Leuven Food Science and Nutrition Research Centre), Catholic University, Leuven, Belgium: ⁹Department of Clinical Sciences, University of Las Palmas de Gran Canaria, Las Palmas de Gran Canaria, Spain

Submitted 22 August 2012: Final revision received 6 March 2013: Accepted 12 March 2013: First published online 11 June 2013

Abstract

Objective: To provide the evidence base for targeted nutrition policies to reduce the risk of micronutrient/diet-related diseases among disadvantaged populations in Europe, by focusing on: folate, vitamin B₁₂, Fe, Zn and iodine for intake and status; and vitamin C, vitamin D, Ca, Se and Cu for intake.

Design: MEDLINE and Embase databases were searched to collect original studies that: (i) were published from 1990 to 2011; (ii) involved >100 subjects; (iii) had assessed dietary intake at the individual level; and/or (iv) included best practice biomarkers reflecting micronutrient status. We estimated relative differences in mean micronutrient intake and/or status between the lowest and highest socio-economic groups to: (i) evaluate variation in intake and status between socio-economic groups; and (ii) report on data availability.

Setting: Europe.

Subjects: Children, adults and elderly.

Results: Data from eighteen publications originating primarily from Western Europe showed that there is a positive association between indicators of socio-economic status and micronutrient intake and/or status. The largest differences were observed for intake of vitamin C in eleven out of twelve studies (5–47%) and for vitamin D in total of four studies (4–31%).

Conclusions: The positive association observed between micronutrient intake and socio-economic status should complement existing evidence on socio-economic inequalities in diet-related diseases among disadvantaged populations in Europe. These findings could provide clues for further research and have implications for public health policy aimed at improving the intake of micronutrients and diet-related diseases.

Keywords
Socio-economic
Micronutrient intake/status
Europe
EURRECA

Many studies have shown that socio-economic status (SES), conventionally measured as education, occupation or income, contributes to inequalities in health^(1,2). Those in lower socio-economic groups have a higher incidence

of premature death, and suffer more from heart diseases and some forms of cancer, than those in more socio-economically advantaged groups⁽²⁾. One important risk factor for these diseases may be the quality of the diet⁽¹⁾.

*Corresponding author: Email romanabgd@yahoo.com

Adequate consumption of micronutrients is one of the key elements of diet quality. Within the EURRECA Network of Excellence (www.eurreca.org), it was shown that the prevalence of micronutrient inadequacies in Europe ranged from 11% to 30% for Cu, folate, Se, iodine, vitamin B₁₂ and vitamin C in adults⁽³⁾. A recent study has shown that diet cost mediates the relationship between socio-economic position (education and income) and diet quality⁽⁴⁾. Since energy-dense and nutrient-poor diets are cheap and more consumed by those with limited economic means and lower educational level^(1,2), it is likely that SES could be associated with variation in micronutrient intake and status. Currently, a knowledge gap exists due to lack of an overview on intake and status of public health-prioritized micronutrients by SES group. Available data that stem from methodologically comparable nutritional surveillances have studied intake and/or status of a few micronutrients within limited life stages, whereas literature reviews have predominantly focused on the relationship between SES and intakes of macronutrients or specific food groups and food patterns (e.g. fat, fruits and vegetables)^(1,5–12). Furthermore, the studies have been performed primarily in adults, while children and the elderly might be more vulnerable to insufficiencies. Therefore the aim of the present study was to provide the evidence base for targeted nutrition policies to reduce the risk of micronutrient/diet-related diseases among disadvantaged populations in Europe, by focusing on intake and status of folate, vitamin B₁₂, Fe, Zn and iodine, and intake of vitamin C, vitamin D, Ca, Se and Cu. We carried out a systematic review on socio-economic differences in micronutrient intake and status in adults, the elderly and children in Europe using education, occupation and income as SES indicators. In addition, we report on availability of the data that addressed our research question.

Methods

Literature search and data extraction

Prioritization of the micronutrients under study was evidence based⁽¹³⁾ and accounted for: (i) the amount of new scientific evidence, particularly from randomized controlled trials; (ii) the public health relevance of micronutrients; and (iii) variations in current micronutrient recommendations for folate, vitamin B₁₂, Fe, Zn and iodine for intake and status, and for vitamin C, vitamin D, Ca, Se and Cu for intake. Socio-economic indicators of interest were educational level, occupational status and income as these are the most conventionally used⁽¹⁴⁾.

A systematic literature search was carried out in MEDLINE and Embase databases to collect studies published from January 1990 up to November 2011 that were performed in apparently healthy populations of all life stages in Europe. Both databases were searched using

database-specific indexing terms for dietary intake and/or nutritional status of at least one of the micronutrients under study, SES (education, occupation or income) and Europe: (micronutrients (listed) OR biomarkers of status (listed)) AND general intake/status terms (diet/or eating/nutritional support/or dietary supplements/or food, fortified/or deficiency diseases/nutritional requirements/or nutritional status/or nutritive value/or nutrition assessment/or diet records) AND socio-economic status terms (socio-economic factors/or poverty/or social class/or educational status/or employment/or unemployment/or income/or occupations/or social conditions) – in title and abstract – AND country (European countries listed). The search strategy can be obtained from the authors upon request.

Studies were considered eligible if they reported on micronutrient intake and/or status across different levels of education, occupation or income. For the final analysis, we used the data where dietary intake was assessed at the individual level and reported as mean intake. Nutritional status was reflected by the level of status markers defined as best practice biomarkers within the context of EURRECA^(15,16).

The search results were screened on the basis of title and abstract by two independent reviewers. For articles that seemed to comply with the study criteria, full texts were obtained and checked for relevance by two independent reviewers. When necessary, the discrepancies in abstract and study selection were discussed, and further processing on inclusion or exclusion was made based on consensus.

Study characteristics (country, life stage and number of subjects, sampling procedure, descriptors of SES) and levels of micronutrient intake/status (mean, standard deviation and/or standard error) and energy consumption were extracted into a database.

Data analysis

We calculated the relative difference (%) in intake and status between the lowest and highest SES groups as a percentage of the value of the highest SES group, for males and females separately unless the paper reported data only for both genders together.

In addition, if studies reported a standard deviation or standard error, the 95% confidence interval of the mean intake/status for SES groups was calculated. The *t* test was performed to evaluate if the difference in intake/status between the lowest and highest SES groups was statistically significant (the level of statistical significance was set at $P < 0.05$).

Means and 95% confidence intervals (when standard deviations/standard errors were available) of intake and status of the lowest and highest SES groups were plotted per micronutrient for males and females separately. Where the results of the *t* test showed a statistically significant difference in intake/status between the lowest

and highest SES groups, study was marked with an asterisk (*; Figs 1–10).

If more than one SES indicator was available in one study, we showed the level of intake/status for one indicator only, i.e. education was given the preference over occupation, and the latter was prioritized over income.

Results

The systematic search in MEDLINE and Embase resulted in 8460 abstracts in total. After screening on the basis of title and abstract, 230 potentially relevant references were identified: 132 on intake and ninety-eight on status. Retrieval of the full papers and their examination disqualified 212 articles mainly because they did not include an internal comparison of intake and/or status between different SES groups, e.g. manual workers *v.* non-manual in the same population group. The remaining eighteen studies included ten studies on intake, three studies on status and one on intake and status in adults, plus four studies on intake in children. Not a single study on status in children was found.

Table 1 gives an overview of data availability per country and study characteristics^(2,17–33).

Most data available were for adults, studying the relationship between vitamin C or Ca intake and education, occupation or income. No single study was found on intake of Cu in adults. Data on biochemical indices of status were very limited and available for adults only: for Fe and Zn only two studies were found per micronutrient, and for iodine status one study was found. The studies included for analysis were performed either in Western Europe or Scandinavia with the exception of one study from Eastern Europe⁽¹⁷⁾. Figures 1–10 show the mean/median intake and status level (with 95% confidence interval of the mean where available) in lowest and highest SES groups in adults/elderly and children. Following each figure, the relative differences between SES groups for intake/status per micronutrient are presented.

Calcium

Number of studies identified: ten studies on intake (Fig. 1).

From six studies on Ca intake in adults, the mean Ca intake in the lowest SES groups was lower than in high-SES groups except for males from Spain⁽²⁰⁾ and males⁽²⁵⁾ and females^(24,25) from the Netherlands. The relative difference ranged from 2% in Swiss females⁽¹⁸⁾ to 14% in UK elderly⁽²⁷⁾. Results on children were more heterogeneous. Higher Ca intake in low SES as compared with high SES was found in Belgian boys⁽²⁸⁾ and in Turkey⁽¹⁷⁾: relative differences were 1% and 19–54%, respectively. Belgian girls and the children from Spain⁽²⁹⁾ and the UK⁽³³⁾ had higher intake in the high-SES groups: the relative difference ranged from 3% to 29%. Statistically significant differences in Ca intake between SES groups

were observed in Irish females⁽²³⁾ and Belgian girls⁽²⁸⁾ ($P < 0.01$).

Vitamin C

Number of studies identified: twelve studies on intake (Fig. 2).

Eight studies on vitamin C intake in adults reported lower mean intake in low-SES groups in comparison to high SES, except in studies in Dutch females^(24,25) and Spanish males⁽²⁰⁾ where relative differences ranged from 3% to 8%. In other studies where higher intake was found in high SES in comparison to low SES, the relative difference was lowest in the Dutch males⁽²⁵⁾ (5%), whereas in the other six studies^(2,20,21,25–27) it ranged from 10% to 48%, the largest difference being found in the UK⁽²⁷⁾ for intake by income.

Four studies on vitamin C intake in children showed lower mean intake in low-SES groups in comparison to high SES, except in Belgian boys⁽²⁸⁾ where the relative difference between SES groups was 12%. In other studies where lower mean intake in low-SES groups in comparison to high SES was reported, relative differences ranged from 6% in Turkish boys⁽¹⁸⁾ up to 33% for Spanish children⁽²⁹⁾.

Statistically significant differences between SES groups were found in studies from Finland⁽²¹⁾, Ireland⁽²³⁾ and Scotland⁽²⁶⁾ for adults and in the study from Spain⁽²⁹⁾ for children ($P < 0.01$).

Iron

Number of studies identified: nine studies on intake, two studies on status (Fig. 3).

Four^(20,23,24,27) out of five studies on Fe intake in adults reported lower intake in low-SES than in high-SES groups (relative differences ranged from 1% to 12%), whereas one study from Spain⁽²¹⁾ reported no difference between SES groups. In children, reported intake was consistently lower in low SES than in high SES (relative differences in the range 2–14%).

Data on Fe status were available only for adults and showed higher values in low SES than in high SES: relative differences were 2% and 7% for Spain⁽²¹⁾ and Norway⁽³²⁾, respectively.

Significant differences in intake/status between SES groups were found in Dutch⁽²⁴⁾ and Irish⁽²³⁾ females and Belgian girls⁽²⁸⁾ ($P < 0.01$).

Folate, vitamin B₁₂, vitamin D, selenium, iodine, zinc, copper

Number of studies identified: folate, vitamin B₁₂ and vitamin D = four studies on intake per each micronutrient (Figs 4–6, respectively); Se = two studies on intake (Fig. 7); iodine = three studies on intake, one study on status (Fig. 8); Zn = four studies on intake, two studies on status (Fig. 9); Cu = one study on intake (Fig. 10).

The two studies available on folate intake in adults^(23,27) and two for children^(29,33) reported lower intake in low SES

Table 1 Characteristics of the study populations, methodology and description of the socio-economic determinants

Study (county, reference and year)	Age range (years)	Characteristics of study population (sampling frame)	Number of participants by gender	Micronutrients	Dietary intake/nutritional status assessment method	Description of the determinant
Intake in adults CH ⁽¹⁸⁾ , 2001	35–74	Community-based random sample of adults in Geneva canton	2929 M; 2767 F	Ca, vitamin D	FFQ	Education: 3 groups (≤ 8 years of education; 9–12 years; ≥ 13 years) Occupation: 3 groups as from British Registrar Classification (high; medium; low)
ES ⁽¹⁹⁾ , 2009	25–60	Randomly selected adults in Andalusia, Spain	1747 M; 1674 F	Fe, Zn	2 \times 24 h recall	Education: 3 groups (no schooling or primary; secondary; university)
ES ⁽²⁰⁾ , 2001	65–95	Institutionalized in urban nursing home and randomly selected free-living subjects	130 M; 218 F	Ca, Fe, vitamin C	FFQ	Education: 2 levels (primary or less; partial secondary to completed university)
FI ⁽²¹⁾ , 1996	25–64	Sub-sample of 3rd FINMONICA risk factor survey (1992)	870 M; 991 F	Vitamin C	3 d FR	Education: 3 groups (0–9 years of education; 10–12 years; > 13 years) Income: household income per consumption unit divided into quartiles
FR ⁽²²⁾ , 2009	35–60	SU.VI.MAX study (1994–96), representative sample of the national population in terms of geographical density and SES	2117 M; 1885 F	Iodine	6 \times 24 h recall	Education: 3 groups (< 13 years of education; 13–14 years; ≥ 14 years)
IE ⁽²³⁾ , 2003	≥ 18	Stratified probability sampling design on the Electoral Register drawn across the Republic's 26 counties	5979 MF	Ca, Fe, vitamin C, vitamin D, vitamin B ₁₂ , Zn, folate, Se	FFQ	Education: 3 groups (non/primary; second level; third level) Occupation: 3 groups (semi-skilled and unskilled labour; non-manual and skilled manual; professional, managerial and technical)
NL ⁽²⁴⁾ , 2003	≥ 19	DNFCS 1997/98, obtained from a panel by a stratified probability sample of the non-institutionalized population	2020 M; 2345 F	Ca, Fe, vitamin C (only females)	2 d FR	Education: 3 groups (SES was based on education and occupation and categorized into (very) low, middle and high)
NL ⁽²⁵⁾ , 2000	≥ 55	The Rotterdam Study (subjects living in a district of Rotterdam, the Netherlands)	2213 M; 3193 F	Ca, vitamin C	FFQ	Education: 4 groups (primary; lower/intermediate general and lower vocational; higher general and intermediate vocational; higher vocational and university)
SCT ⁽²⁶⁾ , 1991	40–59	Data collected from the SHHS from 22 districts in Scotland, recruited through 260 general practitioners	$> 10\,000$ MF	Vitamin C	FFQ	Occupation: divided into 2 groups – manual and non-manual (by husband's occupation for women)
UK ⁽²⁷⁾ , 1999	65–95	Subjects were free-living people drawn from 80 randomly selected postcode sectors, geographically representative of mainland Britain	1000 MF	Ca, Fe, vitamin C, vitamin B ₁₂ , folate	4 d DR	Income: 2 groups (low; high)
UK ⁽²⁾ , 1997	All life stages	National food survey, 1980–1995 (3-stage stratified random sample)	7000 households	Vitamin C	7 d DR conducted by the household member mostly responsible for domestic arrangements	Income: 2 groups (low; high)
Intake in children BE ⁽²⁸⁾ , 2005	13–18	Random sample of adolescents drawn on basis of a multistage cluster sampling technique from private and public schools in Ghent region, Belgium	129 M; 212 F	Ca, vitamin C, Fe	Estimated 7 d FR	Education: 2 groups (general v. vocational school)
ES ⁽²⁹⁾ , 2003	2–5	Infants from two randomly selected day-care centres in Madrid	110 MF	Ca, vitamin C, Fe, vitamin D, vitamin B ₁₂ , folate, iodine, Zn	7 d FR	Education of mothers: 3 groups (no studies or low graduate level; high-school graduate or trade training; not full/full university degree studies)
TR ⁽¹⁷⁾ , 2007	12–13	Schoolchildren attending 3 primary schools, located in 2 districts of Istanbul with higher and lower socio-economic level (multistage sampling method applied to select the schools)	256 M; 248 F	Ca, vitamin C, Fe	3 \times 24 h recall	Income: 2 groups (annual household income of students from lower and higher socio-economic level school districts)
UK ⁽³³⁾ , 2011	10	Children participating in ALSPAC, an ongoing longitudinal cohort study; sample similar to National Census data	3735 M; 3828 F	Ca, vitamin C, Fe, vitamin D, vitamin B ₁₂ , folate, iodine, Zn, Cu, Se	3 d diet diaries (child completed with parental help)	Education of mothers: 3 groups (no schooling or primary; secondary; university)

Table 1 Continued

Study (country, reference and year)	Age range (years)	Characteristics of study population (sampling frame)	Number of participants by gender	Micronutrients	Dietary intake/nutritional status assessment method	Description of the determinant
Status in adults ES ⁽¹⁹⁾ , 2009	25–60	Randomly selected subjects in Andalusia, Spain	170 M; 184 F	Fe, Zn	Hb level; plasma Zn	Education: 3 groups (no schooling or primary; secondary; university)
ES ⁽³⁰⁾ , 2007	18–104	Randomly selected adults in Galicia, Spain	1326 M; 1551 F	Iodine	Urinary iodine	Education: 5 groups (illiterate; uncompleted primary school; primary school; high school; university)
FR ⁽³¹⁾ , 2010	35–60	SU.VI.MAX study (1994–96), representative sample of the national population in terms of geographical density and SES	3127 M; 4425 F	Zn	Plasma Zn	Education: 3 groups (primary; high school; university or equivalent)
NO ⁽³²⁾ , 2006	30–69	Randomly selected non-smoking adults in northern Norway, an iron-mining municipality	690 M	Fe	Serum ferritin	Occupation: miners and non-miners

CH, Switzerland; ES, Spain; FI, Finland; FR, France; IE, Ireland; NL, Netherlands; SCT, Scotland; UK, United Kingdom; BE, Belgium; TR, Turkey; NO, Norway; SU.VI.MAX, Supplementation en Vitamines et Minéraux AntioXydants; SES, socio-economic status; DNFCs, Dutch National Food Consumption Survey; SHHS, Scottish Heart Health Study; ALSPAC, Avon Longitudinal Study of Parents and Children; M, males; F, females; MF, males and females; FR, food record; DR, dietary record.

compared with the high-SES group (relative differences ranged from 7% to 22%). The Irish study in adults and the Spanish study in children, both studying folate intake by educational level, reported significant differences between SES groups ($P < 0.01$).

For vitamin B₁₂ four studies were found. The two studies from the UK for the elderly⁽²⁷⁾ and children⁽³³⁾ showed higher intake in high-SES than in low-SES groups with the relative difference in intake of 6–21%. In contrast, the study in Irish adults⁽²³⁾ and the study in Spanish children⁽²⁹⁾ found higher mean intakes in low-SES groups than in high SES: relative differences were 8–29% and 2%, respectively. The difference in intake by education between SES groups in the Irish study was statistically significantly different ($P < 0.01$).

Results for vitamin D that come from four studies showed consistently lower intakes for the low-SES group. In adults, relative differences were more apparent in Switzerland⁽¹⁸⁾ (13–31%) than in Ireland⁽²³⁾ (4–7%). In children, relative differences in intake between high- and low-SES groups were 4% in the UK⁽³³⁾ and 30% in Spain⁽²⁹⁾, with a statistically significant difference observed in the latter study ($P < 0.05$).

Data on Se, which were reported in one study among Irish adults⁽²³⁾ and one study in UK children⁽³³⁾, showed a slightly lower mean intake in the low-SES group (relative differences 2–5%).


The only study on iodine intake in adults showed no difference in intake in males and a slightly lower intake (relative difference 4%) in the low-SES group in females in France⁽²²⁾. For children, two studies were found and they reported lower intake in the low-SES groups than in high SES: relative differences were 4% and 11% for the UK⁽³³⁾ and Spain⁽²⁹⁾, respectively. Data on iodine status were identified only for Spanish adults⁽³⁰⁾: the median urinary iodine level in the low-SES group was lower than in the high-SES group; the relative difference between the two SES groups was 37%.

Three^(19,29,33) out of four studies on intake of Zn for adults, the elderly and children showed somewhat lower intake in low SES than in the high-SES groups (relative difference ~3%). The Irish study⁽²³⁾ showed inconsistent results on variation in intake between SES groups, depending on gender and SES indicator. Zn status data were available only for adults: the two studies^(19,31) that were found showed 2–5% higher serum Zn level in the low-SES group in comparison to high SES.

The only study on intake of Cu was in children in the UK⁽³³⁾; the intake was higher in the low-SES group than in the high-SES group (relative difference was 6%).

Of the eighteen studies, ten studies used education as an indicator of SES, three applied income and two occupational criteria; three studies used more than one indicator.

Reviewing the results of the different indicators showed that education (especially in females) and income were associated with the largest variations in intake.



Relative differences in intake of Ca for males and females:
 (mean intake in low SES – mean intake in high SES)/mean intake in high SES

		Study (country, year) and SES indicator							
Adults		CH ⁽¹⁸⁾ (2001)	ES ⁽²⁰⁾ (2001)	IE ⁽²³⁾ (2003)		NL ⁽²⁴⁾ (2003)	NL ⁽²⁵⁾ (2000)	UK ⁽²⁷⁾ (1999)	
		Educ	Occup	Educ	Educ	Occup	Educ	Educ	Inc
M		-8.3%	-6.0%	2.6%	-6.2%	-6.5%	-12.0%	1.7%	
F		-8.7%	-2.3%	-2.6%	-11.0%	-2.8%	1.5%	1.9%	
MF									-14.0%
Children		BE ⁽²⁸⁾ (2005)	ES ⁽²⁹⁾ (2003)	TR ⁽¹⁷⁾ (2007)	UK ⁽³³⁾ (2011)				
		Educ	Educ	Inc	Educ				
M		0.7%		19.3%					
F		-28.7%		54.0%					
MF			-3.2%		-5.0%				


Fig. 1 Mean calcium intake (with 95% confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake between the lowest and highest SES group. Countries with studies available: BE, Belgium; CH, Switzerland; ES, Spain; IE, Ireland; NL, Netherlands; TR, Turkey; UK, United Kingdom

Discussion

The present study is the first systematic review addressing differences in levels of micronutrient intake and status between low and high socio-economic groups in European children, adults and elderly using education, occupation and income as indicators of SES. The

results support our hypothesis that social variations are associated with differences in intake and status of certain micronutrients.

The review included all age groups and differences in intake/status were studied separately for each gender where applicable. The subjects were recruited either by stratified random sampling procedures or were representative of the




Relative differences in intake of vitamin C for males and females:
(mean intake in low SES – mean intake in high SES)/mean intake in high SES

Study (country, year) and SES indicator										
Adults	ES ⁽²⁰⁾	FI ⁽²¹⁾	IE ⁽²³⁾	NL ⁽²⁴⁾	NL ⁽²⁵⁾	SCT ⁽²⁶⁾	UK ⁽²⁷⁾	UK ⁽²⁾		
	(2001)	(1996)	(2003)	(2003)	(2000)	(1991)	(1999)	(1997)		
	Educ	Educ	Inc	Educ	Occup	Educ	Educ	Occup	Inc	Inc
	M	8.3%	-27.7%	-16.5%	-28.7%	-18.8%	-5.0%	-10.7%		
	F	-10.1%	-20.3%	-13.0%	-30.3%	-16.4%	2.5%	5.0%	-14.3%	
MF									-47.4%	-23.3%
Children	BE ⁽²⁸⁾	ES ⁽²⁹⁾	TR ⁽¹⁷⁾	UK ⁽³³⁾						
	(2005)	(2003)	(2007)	(2011)						
	Educ	Educ	Inc	Educ						
	M	11.7%		-6.4%						
	F	-13.8%		-11.0%						
MF		-32.6%		-23.9%						

Fig. 2 Mean vitamin C intake (with 95% confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake between the lowest and highest SES group. Countries with studies available: BE, Belgium; ES, Spain; FI, Finland; IE, Ireland; NL, Netherlands; SCT, Scotland; TR, Turkey; UK, United Kingdom

national population, and as such can be considered as exemplary for their countries (Table 1). As one of the outcomes of the systematic review process is observed diversity in study characteristics, there are several aspects

to consider in the interpretation of the results. First, the studies differed in the exact categorization and number of groups used for the three SES indicators: for education studies defined two to four groups, for occupation two or



Relative differences in intake and status of Fe for males and females:
(mean intake/status in low SES – mean intake/status in high SES)/mean intake/status in high SES

		Study (country, year) and SES indicator						
Adults, intake		ES ⁽¹⁹⁾ (2009) Educ	ES ⁽²⁰⁾ (2001) Educ	IE ⁽²³⁾ (2003) Educ	Occup	NL ⁽²⁴⁾ (2003) Educ	UK ⁽²⁷⁾ (1999) Inc	ES ⁽³¹⁾ (2007) Occup
M			3.7 %	–5.3 %	–7.5 %	–4.5 %		
F			–1.0 %	–11.9 %	1.5 %	–9.0 %		
MF		0.0 %					–11.7 %	
Adults, status		ES ⁽¹⁹⁾ (2009) Educ						NO ⁽³²⁾ (2006) Occup
M								7.4 %
MF		2.10 %						
Children, intake		BE ⁽²⁸⁾ (2005) Educ	ES ⁽²⁹⁾ (2003) Educ	TR ⁽¹⁷⁾ (2007) Inc	UK ⁽³³⁾ (2011) Educ			
M		–9.5 %		–2.2 %				
F		–12.5 %		–4.4 %				
MF			–14.1 %		–4.5 %			

Fig. 3 Mean iron intake and status (with 95 % confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake/status between the lowest and highest SES group. Countries with studies available: BE, Belgium; ES, Spain; IE, Ireland; NL, Netherlands; NO, Norway; TR, Turkey; UK, United Kingdom




Fig. 4 Mean folate intake (with 95% confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake between the lowest and highest SES group. Countries with studies available: ES, Spain; IE, Ireland; UK, United Kingdom




Fig. 5 Mean vitamin B₁₂ intake (with 95% confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake between the lowest and highest SES group. Countries with studies available: ES, Spain; IE, Ireland; UK, United Kingdom

three, and for income two to four groups were specified. The size of the absolute and relative differences may be affected by the exact definition of the low- and high-SES groups. Being aware of the inherent differences among studies and after their careful investigation, we made the decision to integrate the results by taking into account both those differences and the same methodological backbone across studies; i.e. a variation between SES groups. The comparisons are made between only two groups, by choosing extreme groups or broad groups⁽¹⁴⁾. Thus, comparing broad SES groups rather than the outer groups

in society may result in an underestimation of existing differences. Another aspect to consider is the impact of the dietary intake method on observed differences in intake. The dietary intake methods used in the identified studies were heterogeneous: FFQ, 24h recall, food record; also, the intake from supplements was included only in three studies^(27,30,33), all of which could contribute to variation in reported mean nutrient intakes. These diverse methods may vary in the extent to which they reflect true usual intake of micronutrients as some of these methods, e.g. non-validated FFQ and diet recorded for less than 3 d, are




Fig. 6 Mean vitamin D intake (with 95% confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake between the lowest and highest SES group. Countries with studies available: CH, Switzerland; ES, Spain; IE, Ireland; UK, United Kingdom




Fig. 7 Mean selenium intake (with 95% confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake between the lowest and highest SES group. Countries with studies available: IE, Ireland; UK, United Kingdom

generally considered inappropriate to estimate a population's usual dietary intake. This is of importance when estimating the prevalence of inadequacy (comparison of absolute mean intake/status levels with reference values). On the other hand, this is less significant for the aim of the

present study, i.e. for observing the variation between SES groups per study as we focused on differences within studies. However, as mentioned in certain publications on socio-economic differences in food consumption, we acknowledge the potential for misreporting of micronutrient




Fig. 8 Mean iodine intake and status (with 95% confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake between the lowest and highest SES group. Countries with studies available: ES, Spain; FR, France; UK, United Kingdom

intake for subjects with higher educational levels as they may be more conscious of nutrition and health issues and possibly exaggerate the consumption of healthy foods, e.g. fruit and vegetables, thus giving rise to elevated mean vitamin and mineral intake levels^(7,34). The issue of reporting errors does not affect the results on micronutrient status; therefore, the information on Fe, Zn and iodine status strengthens the evidence from the observed differences between intake for low and high SES.

Results of the systematic literature search showed that studies on micronutrient intake and/or status associated with SES indicators for children from all European countries and for children, adults and elderly from Central and Eastern Europe (CEE) are very limited. For example, our review included data for Western European populations with the exception of one study on children from Turkey⁽¹⁷⁾. Inclusion of studies from other parts of Europe such as CEE is needed to draw a pan-European conclusion on variations in micronutrient intake/status between groups of different education, occupation and income levels. It is disputable whether similar results would be found for CEE because of the widely divergent socio-economic position within these countries: larger social inequalities, diversity in food costs and availability, etc. A recent publication that

compared micronutrient intake/status in CEE with other European regions showed no striking differences, with the exception of Ca⁽³⁵⁾. More research is needed, however, to fill this gap in available evidence. An explanation for the limited studies we identified throughout Europe is that we focused on the electronic databases MEDLINE and Embase only. This is of particular importance for CEE, as it has been shown that reliable nutritional data from these countries can be found as grey literature: country reports of governmental or academic institutions, theses and dissertations⁽³⁵⁾. Their inclusion was not feasible however, due to the large number of languages to cover. Nevertheless, it is apparent that even though the relationship between socio-economic position and diet has been notably studied in the last few decades, current evidence from key databases is still lacking data on SES-related micronutrient intake and/or status for many European countries and particularly for children and the elderly.

Existing reviews in Europe have not addressed the association between multiple socio-economic indicators and intake and/or status of more than one micronutrient. The largest study⁽³⁶⁾ to date which described intake of five water-soluble vitamins across several European countries reported the association between education and




Fig. 9 Mean zinc intake and status (with 95 % confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake between the lowest and highest SES group. Countries with studies available: ES, Spain; FR, France; IE, Ireland; UK, United Kingdom

intake: it was positive for the intake of riboflavin, vitamin C, vitamin B₆ (in women) and no clear conclusion emerged for vitamin B₁₂. These results are in accordance with the findings for vitamins B₁₂ and C from the present study.

Still, there is a body of evidence on disparities in food habits in association with SES indicators. A review on consumption of fruit and vegetables in fifteen European countries reported lower consumption in low-SES in comparison to high-SES groups⁽⁹⁾; the same was observed in nine European countries for the consumption of cheese, but not for milk⁽¹⁰⁾. These findings support the results on vitamin C and Ca from our study. To estimate if similar results could be found for other vitamins and minerals under study, we examined studies evaluating the variation in intake of nutrient-rich healthy foods in groups with different socio-economic backgrounds. Likewise, whole grains, lean meats, fish, low-fat dairy products and fresh vegetables and fruit were found to be preferably consumed by high-SES groups⁽¹⁾. The most recent study on the relationship between SES and frequency of consumption of seven predefined healthy foods (consumption of fruit, vegetables, wholegrain bread, vegetable-fat spread, vegetable cooking fat, low-fat milk and low-fat cheese) in CEE and Western Europe reported healthier food choices among those with higher education, occupational position

and fewer economic difficulties, as well as heterogeneity in the association between SES and healthy food habits across countries⁽⁶⁾. The present study suggests that the differences are the most apparent in SES groups of different educational background, especially in females. On the other hand, Table 1 shows that for both intake and status for all age groups, it is education that is the most studied indicator: ten studies on education, two studies on education and occupation, one study on education and income, two studies on occupation and three on income. Where studies reported on intake by two indicators^(18,21,23), the relative differences between SES groups observed for education were confirmed for occupation^(18,23) and income⁽²¹⁾, but the variation for the latter two indicators was lower than for intake by education.

The more pronounced effect of education on variation in intake/status in comparison to occupation and income may be due to the fact that the level of education is considered to remain rather stable through the life course⁽³⁷⁾. Furthermore, the education attained in early life is not only a determinant of occupation and income, but also affects the ability to understand and implement dietary guidance messages and risk-reducing dietary behaviours⁽³⁸⁾. Despite the studies in the present review using different procedures in assessing SES, most methods have the same underlying




Fig. 10 Mean copper intake (with 95% confidence interval, where available, represented by vertical bars) of European adults/elderly and children (M, males; F, females; MF, males and females) in groups of lowest (■) and highest (□) socio-economic status (SES) by different SES indicators (educ, education; occup, occupation; inc, income). * denotes a statistically significant difference ($P < 0.05$) in mean intake between the lowest and highest SES group. Countries with studies available: UK, United Kingdom

principle: difference in micronutrient intake/status between SES groups. To the extent that similar principles are indeed addressed, the heterogeneity of results between different studies might be a reflection of these methodological shortcomings; but as long as no clear gold standard is available, the pooling of results in the way we have done is the best possible option. On the other hand, the presence of dissimilarities in SES categorization among studies can have advantages by increasing the generalization of the conclusion⁽³⁹⁾ that there is, in most cases, a positive association observed between all SES indicators under study and micronutrient intake and/or status.

To support policy makers to develop targeted nutritional policies to reduce the risk of diet-related diseases among disadvantaged populations in Europe, a comprehensive overview of nutritional intake and status across different socio-economic strata is needed. However, the current review showed that, contrary to the large body of evidence on socio-economic inequalities in health, very few data on socio-economic differences in micronutrient intake and status are available in the scientific literature. This applies to the whole of Europe and for all age groups, but especially to CEE countries and to children, adolescents and the elderly.

Conclusion

When intake and/or status of priority micronutrients (folate, vitamin B₁₂, Fe, Zn and iodine for intake and

status; vitamin C, vitamin D, Ca, Se and Cu for intake) is compared between low- and high-SES groups, lower values for all micronutrients (except vitamin B₁₂) are found in the low SES category. The largest relative differences between SES groups are observed for intake of Ca, vitamin C, folate and vitamin D (2–29%, 5–47%, 7–22% and 4–31%, respectively), although the data for the latter two were fewer than for the former two nutrients. The results presented in the current review stem from analysis of the original studies as from 1990 up to November 2011. Current evidence suggests that there are inadequate micronutrient intakes in Europe⁽³⁾. Since the relationship between SES and health is dynamic and evolves throughout the life course⁽⁴⁰⁾, it should be regarded that the observed changes in SES inequalities in micronutrient intake and status should be monitored over time. This can be realized via analyses of high-quality nutritional data with EU coverage that apply comparable measures of SES (e.g. from the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study and/or EPIC (European Prospective Investigation into Cancer and Nutrition)) or via the inclusion of comparable socio-economic determinants and measures in currently running and forecasted nutritional surveillance programmes.

Acknowledgements

Sources of funding: The work reported herein has been carried out within the EURRECA Network of Excellence

(<http://www.eurreca.org>), which is financially supported by the Commission of the European Communities, specific Research, Technology and Development (RTD) Programme Quality of Life and Management of Living Resources, within the Sixth Framework Programme, project no. FP6 036196-2. This report does not necessarily reflect the Commission's views or its future policy in this area. This work was in part funded by the Ministry of Education and Science of the Republic of Serbia, project III 41030. *Conflicts of interest:* The authors have no conflicts of interest. *Ethics:* No ethical approval was required for this study. *Authorship responsibilities:* R.N. conceptualised the paper, collected the data and drafted the manuscript. A.C. and A.G. supervised the overall work and assisted with analysis and editing of the manuscript. M.N., I.I.A., B.R.V., J.N., M. Golsorkhi, M.W.M., A.B., A.S., D.d.C., G.V. and M.R. were involved in data collection, reviewing the literature and analysis. L.S.M., L.A.M., M. Glibetić, M. Gurinović, P.v.V. and L.C.P.G.M.d.G. were involved in development of the study hypothesis, editing the working versions of the manuscript and provided advice regarding interpretation of the results.

References

- Darmon N & Drewnowski A (2008) Does social class predict diet quality? *Am J Clin Nutr* **87**, 1107–1117.
- James WP, Nelson M, Ralph A *et al.* (1997) Socioeconomic determinants of health. The contribution of nutrition to inequalities in health. *BMJ* **314**, 1545–1549.
- Roman-Vinas B, Ribas-Barba L, Ngo J *et al.* (2011) Projected prevalence of inadequate nutrient intakes in Europe. *Ann Nutr Metab* **59**, 84–85.
- Aggarwal A, Monsivais P, Cook AJ *et al.* (2011) Does diet cost mediate the relation between socioeconomic position and diet quality? *Eur J Clin Nutr* **65**, 1059–1066.
- Giskes K, Avendano M, Brug J *et al.* (2010) A systematic review of studies on socioeconomic inequalities in dietary intakes associated with weight gain and overweight/obesity conducted among European adults. *Obes Rev* **11**, 413–429.
- Boylan S, Lallukka T, Lahelma E *et al.* (2011) Socioeconomic circumstances and food habits in Eastern, Central and Western European populations. *Public Health Nutr* **14**, 678–687.
- Trichopoulou A, Naska A & Costacou T (2002) Disparities in food habits across Europe. *Proc Nutr Soc* **61**, 553–558.
- Lallukka T, Laaksonen M, Rahkonen O *et al.* (2007) Multiple socio-economic circumstances and healthy food habits. *Eur J Clin Nutr* **61**, 701–710.
- Irala-Estevez JD, Groth M, Johansson L *et al.* (2000) A systematic review of socio-economic differences in food habits in Europe: consumption of fruit and vegetables. *Eur J Clin Nutr* **54**, 706–714.
- Sanchez-Villegas A, Martinez JA, Prattala R *et al.* (2003) A systematic review of socioeconomic differences in food habits in Europe: consumption of cheese and milk. *Eur J Clin Nutr* **57**, 917–929.
- Prattala RS, Groth MV, Oltersdorf US *et al.* (2003) Use of butter and cheese in 10 European countries: a case of contrasting educational differences. *Eur J Public Health* **13**, 124–132.
- World Health Organization (2008) *Closing the Gap in a Generation: Health Equity Through Action on the Social Determinants of Health. Final Report of the Commission on Social Determinants of Health*. Geneva: WHO.
- Cavelaars AE, Doets EL, Dhonukshe-Rutten RA *et al.* (2010) Prioritizing micronutrients for the purpose of reviewing their requirements: a protocol developed by EURRECA. *Eur J Clin Nutr* **64**, Suppl. 2, S19–S30.
- Kunst AE, Bos V & Mackenbach JP; EU Working Group on Socio-economic Inequalities in Health (2001) *Monitoring Socio-economic Inequalities in Health in the European Union: Guidelines and Illustrations. Final report for the Health Monitoring Program of the European Commission*. Rotterdam: Department of Public Health, Erasmus University.
- Food and Agriculture Organization of the United Nations. (1988) *Requirements of Vitamin A, Iron, Folate and Vitamin B₁₂: Report of a Joint FAO/WHO Expert Consultation*. FAO Food and Nutrition Series no. 23. Rome: FAO.
- Ristic-Medic D, Piskackova Z, Hooper L *et al.* (2009) Methods of assessment of iodine status in humans: a systematic review. *Am J Clin Nutr* **89**, issue 6, 2052S–2069S.
- Manios Y, Moschonis G, Kolotourou M *et al.* (2007) Iron deficiency prevalence and dietary patterns by school district in Istanbul. *J Hum Nutr Diet* **20**, 549–557.
- Galobardes B, Morabia A & Bernstein MS (2001) Diet and socioeconomic position: does the use of different indicators matter? *Int J Epidemiol* **30**, 334–340.
- Sanchez C, Lopez-Jurado M, Planells E *et al.* (2009) Assessment of iron and zinc intake and related biochemical parameters in an adult Mediterranean population from southern Spain: influence of lifestyle factors. *J Nutr Biochem* **20**, 125–131.
- Lasheras C, Patterson AM, Casado C *et al.* (2001) Effects of education on the quality of life, diet, and cardiovascular risk factors in an elderly Spanish community population. *Exp Aging Res* **27**, 257–270.
- Roos E, Prattala R, Lahelma E *et al.* (1996) Modern and healthy?: socioeconomic differences in the quality of diet. *Eur J Clin Nutr* **50**, 753–760.
- Valeix P, Faure P, Peneau S *et al.* (2009) Lifestyle factors related to iodine intakes in French adults. *Public Health Nutr* **12**, 2428–2437.
- Friel S, Kelleher CC, Nolan G *et al.* (2003) Social diversity of Irish adults' nutritional intake. *Eur J Clin Nutr* **57**, 865–875.
- Hulshof KF, Brussaard JH, Kruizinga AG *et al.* (2003) Socio-economic status, dietary intake and 10 y trends: the Dutch National Food Consumption Survey. *Eur J Clin Nutr* **57**, 128–137.
- van Rossum CT, van de MH, Witteman JC *et al.* (2000) Education and nutrient intake in Dutch elderly people. The Rotterdam Study. *Eur J Clin Nutr* **54**, 159–165.
- Bolton-Smith C, Smith WC, Woodward M *et al.* (1991) Nutrient intakes of different social-class groups: results from the Scottish Heart Health Study (SHHS). *Br J Nutr* **65**, 321–335.
- Bates CJ, Prentice A, Cole TJ *et al.* (1999) Micronutrients: highlights and research challenges from the 1994–5 National Diet and Nutrition Survey of people aged 65 years and over. *Br J Nutr* **82**, 7–15.
- Matthys C, De HS, Maes L *et al.* (2006) Adolescents' education and their diet recorded by 7-day food records. *Soz Präventivmed* **51**, 80–90.
- Navia B, Ortega RM, Requejo AM *et al.* (2003) Influence of maternal education on food consumption and energy and nutrient intake in a group of pre-school children from Madrid. *Int J Vitam Nutr Res* **73**, 439–445.
- Rego-Iraeta A, Perez-Fdez R, Cadarso-Suarez C *et al.* (2007) Iodine nutrition in the adult population of Galicia (Spain). *Thyroid* **17**, 161–167.

31. Arnaud J, Touvier M, Galan P *et al.* (2010) Determinants of serum zinc concentrations in a population of French middle-age subjects (SU.VI.MAX cohort). *Eur J Clin Nutr* **64**, 1057–1064.
32. Broderstad AR, Smith-Sivertsen T, Dahl IM *et al.* (2006) Serum levels of iron in Sor-Varanger, Northern Norway – an iron mining municipality. *Int J Circumpolar Health* **65**, 432–442.
33. Cribb VL, Jones LR, Rogers IS *et al.* (2011) Is maternal education level associated with diet in 10-year-old children? *Public Health Nutr* **14**, 2037–2048.
34. Macdiarmid J & Blundell J (1998) Assessing dietary intake: who, what and why of under-reporting. *Nutr Res Rev* **11**, 231–253.
35. Novakovic R, Cavelaars AEJM, Bekkering EG *et al.* (2013) Micronutrient intake and status in Central and Eastern Europe compared to other European countries, results from the EURRECA network. *Public Health Nutr* **16**, 824–840.
36. Olsen A, Halkjaer J, van Gils CH *et al.* (2009) Dietary intake of the water-soluble vitamins B₁, B₂, B₆, B₁₂ and C in 10 countries in the European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr* **63**, Suppl. 4, S122–S149.
37. Salonen MK, Kajantie E, Osmond C *et al.* (2009) Role of socioeconomic indicators on development of obesity from a life course perspective. *J Environ Public Health* **2009**, 625168.
38. Kant AK & Graubard BI (2007) Ethnicity is an independent correlate of biomarkers of micronutrient intake and status in American adults. *J Nutr* **137**, 2456–2463.
39. Walker E, Hernandez AV & Kattan MW (2008) Meta-analysis: its strengths and limitations. *Cleve Clin J Med* **75**, 431–439.
40. Ovrum A, Gustavsen G & Rickertsen K (2012) Health inequalities over the adult life course: the role of lifestyle choices. Paper presented at the *International Association of Agricultural Economists (IAAE) Triennial Conference*, Foz do Iguaçu, Brazil, 18–24 August 2012.