

1

2 **The influence of anthropogenic nitrogen loading and meteorological conditions on**
3 **the dynamics and toxicity of *Alexandrium fundyense* blooms in a New York (USA)**
4 **estuary**

5

6 **Theresa K. Hattenrath^a, Donald M. Anderson^b, Christopher J. Gobler^{a*}**

7 ^aStony Brook University, School of Marine and Atmospheric Sciences, Southampton, NY
8 11968, USA

9

10 ^bWoods Hole Oceanographic Institution, Woods Hole, MA, 02543, USA

11

12 *Corresponding author: [Christopher.gobler\(at\)sunysb.edu](mailto:Christopher.gobler@sunysb.edu)
13 PH: 631-632-5043

14

15

16

17

18 **Abstract:** The goal of this two-year study was to explore the role of nutrients and
19 climatic conditions in promoting reoccurring *Alexandrium fundyense* blooms in the
20 Northport-Huntington Bay complex, NY, USA. A bloom in 2007 was short and small (3
21 weeks, 10^3 cells L^{-1} maximal density) compared to 2008 when the *A. fundyense* bloom,
22 which persisted for six weeks, achieved cell densities $>10^6$ cells L^{-1} and water column
23 saxitoxin concentrations $>2.4 \times 10^4$ pmol STX eq. L^{-1} . During the 2008 bloom, both
24 deployed mussels (used as indicator species) and wild soft shell clams became highly
25 toxic (1,400 and 600 μg STX eq./100g shellfish tissue, respectively) resulting in the
26 closure of shellfish beds. The densities of benthic *A. fundyense* cysts at the onset of this
27 bloom were four orders of magnitude lower than levels needed to account for observed
28 cell densities, indicating *in situ* growth of vegetative cells was responsible for elevated
29 bloom densities. Experimental enrichment of bloom water with nitrogenous compounds,
30 particularly ammonium, significantly increased *A. fundyense* densities and particulate
31 saxitoxin concentrations relative to unamended control treatments. The $\delta^{15}\text{N}$ signatures
32 (12 to 23‰) of particulate organic matter (POM) during blooms were similar to those of
33 sewage (10 to 30‰) and both toxin and *A. fundyense* densities were significantly
34 correlated with POM $\delta^{15}\text{N}$ ($p < 0.001$). These findings suggest *A. fundyense* growth was
35 supported by a source of wastewater such as the sewage treatment plant which discharges
36 into Northport Harbor. Warmer than average atmospheric temperatures in the late winter
37 and spring of 2008 and a cooler May contributed to an extended period of water column
38 temperatures optimal for *A. fundyense* growth (12 – 20°C), and thus may have also
39 contributed toward the larger and longer bloom in 2008. Together this evidence suggests

40 sewage-derived N loading and above average spring temperatures can promote intense
41 and toxic *A. fundyense* blooms in estuaries.

42

43 Keywords: *Alexandrium*, anthropogenic nitrogen loading, $\delta^{15}\text{N}$, toxin, climate

44

45 **1. Introduction:**

46 The intensity and impacts of harmful algal blooms (HABs) in coastal ecosystems
47 have increased in recent decades (Anderson, 1994, Glibert et al., 2005; Anderson et al.,
48 2008; Heisler et al., 2008). Blooms of the dinoflagellate *Alexandrium* are common to
49 many coastal regions around the globe and are particularly harmful because they produce
50 saxitoxins, the causative agent of paralytic shellfish poisoning (PSP) (Anderson, 1994;
51 Anderson, 1997; Glibert et al., 2005). Saxitoxins are a family of potent neurotoxins that
52 block sodium channels and cause severe illness or death in humans who consume
53 saxitoxin-contaminated shellfish (Kvitek and Beitler, 1988; Anderson, 1994). The
54 frequency of *Alexandrium* blooms as well as the intensity of these events have been
55 increasing worldwide, and therefore so have PSP outbreaks (Anderson, 1994; Sellner et
56 al., 2003; Glibert et al., 2005). Although it is not certain whether these events can be
57 attributed to an increase in coastal monitoring or to increased anthropogenic nutrient
58 loading to coastal systems (Anderson, 1994; Anderson et al., 2002; Glibert et al., 2005;
59 Heisler et al., 2008), it is clear that these blooms have devastating economic impacts
60 (Anderson et al., 2000; Jin and Hoagland, 2008; Jin et al., 2008). Many consider PSP to
61 be the most widespread of all HAB poisoning syndromes (Hackett et al., 2004; Erdner et
62 al., 2008).

63 *Alexandrium fundyense* blooms are common along the northeast US coast.
64 Paralytic shellfish poisoning in the northeastern US was first documented in Maine in
65 1958 (Anderson, 1997). In 1972, a large *A. fundyense* bloom with cell densities
66 exceeding 10^6 cells L^{-1} spread through the Gulf of Maine and affected coastal regions
67 from Maine to Massachusetts (Mulligan, 1975; Anderson, 1994; Anderson, 1997). Since

68 then these large-scale regional blooms and associated PSP-related shellfish bed closures
69 have been near-annual occurrences in this region (Anderson, 1994; Anderson, 1997;
70 Townsend et al., 2001) to the detriment of the shellfish industry. For example, during the
71 2005 *A. fundyense* bloom in New England, the seafood industry lost more than \$3 million
72 per week in revenue (Jin et al., 2008).

73 In contrast to these widespread coastal outbreaks, *Alexandrium* blooms also occur
74 in isolated embayments along the New England and Long Island coast. These are
75 considered “point source” outbreaks in which localized cyst germination inoculates the
76 overlying waters, with deposition of new cysts at the end of blooms providing the means
77 for the species to bloom again in subsequent years (e.g., Anderson and Morel, 1979;
78 Anderson et al., 1983; Anderson et al., 2008). There is thought to be no significant
79 connection between these small-scale blooms in estuaries and embayments and the large-
80 scale regional blooms described above (Anderson, 1997; Anderson et al., 2008).

81 The presence of *A. fundyense* on Long Island was first documented during the
82 early 1980’s (Anderson et al., 1982; Schrey et al., 1984). At that time, moderate densities
83 of *A. fundyense* ($> 10^2$ cell L⁻¹) were found on the north shore of Long Island in Northport
84 Bay and Mattituck Inlet (Schrey et al., 1984); these blooms, however, were not associated
85 with PSP events (e.g. toxic shellfish or human illness; Anderson et al., 1982; Schrey et
86 al., 1984). At the time, this was attributed to the low toxin content or quota of Long
87 Island isolates, which contain proportionally more of the low potency C toxins than other
88 more potent congeners (Anderson et al., 1994). The assumption was that very dense
89 blooms would be required for there to be dangerous levels of toxicity in shellfish.
90 Although there have been no studies of *A. fundyense* in NY waters since the 1980s, in

91 2006, the detection of elevated saxitoxin in shellfish by the New York State Department
92 of Environmental of Conservation (NYSDEC) prompted the closure of 2,000 acres of
93 shellfish beds in the Northport-Huntington Bay system of Long Island. Blooms recurred
94 in those waters in 2007 and 2008.

95 Factors promoting toxic *A. fundyense* bloom events seem to vary with the
96 ecosystem within which blooms occur (Anderson et al., 2008). Decades of research in
97 the Gulf of Maine have led to the conclusion that the presence and dynamics of *A.*
98 *fundyense* benthic cyst beds and the physical transport of cells controls the dynamics of
99 the widespread regional blooms (Anderson, 1997; Anderson et al., 2005a,c; Stock et al.,
100 2005; Anderson et al., 2008). The low levels of nutrients present during blooms in the
101 open waters of the Gulf of Maine (Townsend et al., 2001; Poulton et al., 2005; Townsend
102 et al., 2005b; Love et al., 2005) and the ability of *A. fundyense* dynamics to be
103 successfully modeled in the absence of a nutrient-dependent growth rate (Stock et al.,
104 2005) suggests nutrients seem to have a smaller, secondary influence on these events
105 (Anderson, 1997; Anderson et al., 2005a,c). In contrast, anthropogenic nutrient loading
106 could have a larger impact on the development of *A. fundyense* blooms in coastal
107 embayments where nutrient concentrations and loads are substantially higher than the
108 Gulf of Maine (Anderson et al., 1983; Penna et al., 2002; Trainer et al., 2003; Poulton et
109 al., 2005; Anderson et al., 2008). Anthropogenic nutrient loading has been associated
110 with an increase in PSP incidences caused by *Alexandrium catenella* in multiple marine
111 ecosystems including shallow, poorly flushed coastal embayments of the northwest US
112 (Trainer et al., 2003). The degree to which *A. fundyense* populations in estuaries are
113 controlled by nutrient loading, cyst beds, or both factors is not well understood.

114 This study documented the dynamics of *A. fundyense* blooms in a coastal region
115 of New York in 2007 and 2008, including a bloom which persisted for six weeks,
116 achieved densities of more than 10^6 cells L^{-1} , and lead to the closure of more than 7,000
117 acres of shellfish beds. The spatial and temporal dynamics of the physical environment,
118 nutrients, toxins, *A. fundyense* cells, and *A. fundyense* cysts are presented in conjunction
119 with experiments examining the impacts of nutrient enrichment on the growth and
120 toxicity of *A. fundyense* populations. The role of nutrient loading and meteorological
121 conditions in the occurrence of *A. fundyense* blooms is subsequently assessed.

122 **2. Materials & Methods:**

123 *2.1. Field sampling and analyses*

124 During 2007 and 2008 sampling was conducted at various locations across the
125 Northport-Huntington Bay complex, located on the north shore of Long Island, NY, USA
126 (Fig. 1, 40.9090°N, 73.4036°W). This system has previously hosted *A. fundyense* cells
127 (Anderson et al., 1982; Schrey et al., 1984; Anderson, 1997) and saxitoxin contaminated
128 shellfish (Karen Chytalo, NYSDEC, personal communication). Within this system,
129 Northport Harbor, located in the southeastern part of the Northport-Huntington Bay
130 complex, was sampled on a weekly basis from April through June at one site in 2007 (site
131 2) and at three locations in 2008 (Fig. 1; sites 2, 7, 8). Other sites, located in Huntington
132 Harbor (site 6) and Centerport Harbor (site 1) were sampled weekly, while 7 other sites
133 (sites 3, 4, and 5 in 2007; sites 3, 4, 10, 11, 16 and LIS in 2008) were sampled during the
134 pinnacle of blooms to document the spatial extent of these events (Fig. 1).

135 At each station, a YSI© probe was used to record surface temperature, salinity
136 and dissolved oxygen. Subsurface water (~0.25m) was filtered for nutrient analysis using

137 precombusted (4 hr @ 450°C) glass fiber filters (GF/F, 0.7 µm pore size) and frozen in
138 acid washed scintillation vials. Filtrate was analyzed colorimetrically for ammonium,
139 nitrate, phosphate, and silicate (Jones, 1984; Parsons et al., 1984) using a
140 spectrophotometric microplate reader. To determine the size distribution of
141 phytoplankton biomass, chlorophyll *a* was fractionated using GF/F (nominal pore size 0.7
142 µm) and polycarbonate filters (2 µm & 20 µm) and measured using standard fluorometric
143 techniques described in Parsons et al. (1984). Whole water samples were preserved in
144 Lugol's iodine. Aliquots were settled in counting chambers and plankton were identified
145 and enumerated using an inverted light microscope (Hasle, 1978). Cells larger than 10
146 µm were identified to at least genus level and grouped as dinoflagellates, diatoms, and
147 ciliates. To assess the $\delta^{15}\text{N}$ signature of plankton communities dominated by *A.*
148 *fundyense*, replicate samples of particulate organic matter (POM) was filtered onto
149 precombusted (4h @ 450°C) GF/F filters, dried for 24 h at 60°C, pelleted, and analyzed
150 for $\delta^{15}\text{N}$ via continuous flow isotope ratio mass spectrometry (IRMS) by David Harris at
151 the UC Davis Stable Isotope Facility.

152 *A. fundyense* cell densities were enumerated using a molecular technique
153 developed by Anderson et al. (2005b). In the field, 2 L of water was pre-sieved through a
154 200 µm mesh to eliminate large zooplankton from the sample and subsequently
155 concentrated onto a 20 µm sieve and backwashed into a centrifuge tube to a volume of 14
156 ml. Samples were preserved in ~2% formaldehyde and refrigerated at 4°C for at least 1
157 hour and no more than 24 hours. After refrigeration, samples were centrifuged at 3000
158 rpm for 11 minutes and the supernatant aspirated without disturbing the cell pellet. The
159 cell pellet was resuspended in 14 ml ice cold methanol and stored at -20°C for up to six

160 months (Anderson et al., 2005b). An aliquot of preserved sample was filtered onto a 5
161 μm polycarbonate track-etched membrane (25mm in diameter). A pre-hybridization
162 buffer was incubated for 5 minutes with each sample and then filtered off of samples. *A.*
163 *fundyense* cells were labeled using oligonucleotide probe NA1 for the North American
164 ribotype *Alexandrium fundyense/catenella/tamarense* with Nu-light™ dye conjugated to
165 the 5' end (5'-/5Cy3/AGT GCA ACA CTC CCA CCA-3'). A hybridization buffer,
166 containing pre-hybridization buffer in addition to probe (a final probe concentration of
167 $4.8 \text{ ng } \mu\text{l}^{-1}$) was added to each sample and allowed to incubate for 1 hour at 50°C .
168 Following incubation, the hybridization buffer was filtered and samples were washed
169 with 0.2X SET for 5 minutes. Filters were then mounted onto a microscope slide and
170 glycerol was added to each filter to prevent fading of the probe. Cells were enumerated
171 using a Nikon epifluorescence microscope with a Cy3™ filter set (Anderson et al.,
172 2005b). As a quality control, measured samples spiked with *A. fundyense* culture (clone
173 GTCA28 or ATNPD7) were hybridized with the oligonucleotide probe and quantified
174 during each analytical run. Oligonucleotide probe quantification of seawater spiked with
175 known densities of *A. fundyense* clone GTCA28 yielded mean recoveries of $87 \pm 16\%$.
176 Light microscope counts of Lugol's stained *A. fundyense* cells yielded large
177 overestimates of population densities compared to oligonucleotide quantification.

178 Toxin concentrations in plankton samples were determined by a competitive
179 enzyme linked immunosorbent assay (ELISA). Several liters of seawater were pre-sieved
180 through a $200 \mu\text{m}$ mesh and subsequently concentrated on a $20 \mu\text{m}$ sieve, backwashed
181 into centrifuge tubes and pelleted. Cell pellets were acidified with 0.1 M HCl and
182 subsequently analyzed for saxitoxin using ELISA kits from R-Biopharm© in 2007 and by

183 Abraxis© in 2008, with toxin concentrations reported in STX equivalents. Each of these
184 kits had varying degrees of cross-reactivities among saxitoxin congeners. Cross-
185 reactivities for the ELISA kits from R-Biopharm© and Abraxis© were as follows: 100%
186 STX, 20% dcSTX, 70% GTX2,3 and 12% NEO, and 100% STX, 29% dcSTX, 23%
187 GTX2,3, 23% GTX5B, 1.3% NEO, and <0.2% GTX1,4, respectively. Analysis of
188 replicated samples by both kits yielded statistically identical results. As a quality control
189 measure, for each analytical run, an *Alexandrium fundyense* culture (GTCA28) known to
190 produce saxitoxins was used as a positive control and *Aureococcus anophagefferens*
191 (CCMP 1984), which does not produce saxitoxins, was used as a negative control. Three
192 times the standard deviation of the negative control was used as the methodological
193 detection limit for each analytical run. Analysis of total saxitoxins in pelleted
194 *Alexandrium fundyense* cultures (clone ATNPD7) via high performance liquid
195 chromatography (HPLC) yielded statistically equivalent levels of total saxitoxin
196 concentrations on a per cell basis to those measured with the both ELISA kits.

197 During November 2007 and 2008 sediment samples were obtained from 17
198 locations across the Northport-Huntington Bay complex (Fig. 1). Surveys were timed to
199 occur following potential fall bloom events and thus quantified cysts represented
200 potential seed populations for the following year (Anderson et al., 2005c). Sediment
201 samples were obtained using a Ponar grab and several subcores from the top 3cm were
202 taken using a modified syringe. All samples were processed according to Anderson et al.
203 (2005c) and stained with primulin (Yamaguchi et al., 1995). Primulin stained cysts were
204 enumerated under an epifluorescent microscope using a 1 ml Sedgewick-Rafter slide.
205 Cyst concentrations were reported in cysts cc⁻¹ of sediment.

206 Meteorological data including wind intensity, wind direction, temperature, and
207 precipitation were obtained from the National Weather Service's monitoring station in
208 Islip, NY, USA which is ~20 km from Northport. For each of these parameters the
209 monthly means for 2007 and 2008 were compared using t-tests. The degree to which all
210 individual water column parameters were correlated to each other was evaluated by
211 means of a Spearman rank order correlation matrix.

212 2.2. Nutrient amendment experiments

213 To assess the impact of nitrogen (N) and phosphorus (P) loading on *A. fundyense*
214 growth and toxin production, a series of nutrient amendment experiments were
215 performed. Triplicate bottles (1.1 L in 2007 and 2.5 L in 2008) were filled with water
216 from Northport Harbor. An unamended control was established along with four
217 treatments in 2007 including 20 μM nitrate, 20 μM ammonium, 10 μM urea (= 20 μM
218 N), and 2 μM phosphate. Due to the response from reduced N in general and ammonium
219 in particular during 2007 experiments, experiments in 2008 included additional
220 treatments: 10 μM ammonium, 40 μM ammonium, 20 μM ammonium combined with 2
221 μM phosphate, and 10 μM glutamine (= 20 μM N). All treatment concentrations were
222 chosen to match those which have previously elicited a growth response in *Alexandrium*
223 cells (Leong et al., 2004) and were similar to peak elevated levels found in Long Island
224 estuaries (Gobler et al., 2004). Bottles were incubated for ~ 48 h at ambient light and
225 temperature after which *A. fundyense* cell enumeration, and toxin quantification were
226 performed via the aforementioned methods. Differences among treatments were
227 elucidated by means of a One-Way ANOVA with multiple comparison tests (i.e. Student-

228 Newman-Keuls) or with an appropriate non-parametric test when normality tests of log
229 transformed data failed.

230 2.3. Toxins in shellfish

231 During both 2007 and 2008, netted bags containing the blue mussel, *Mytilus*
232 *edulis*, from regions not exposed to PSP toxins were hung off piers located adjacent to
233 sampling sites in Northport Harbor and in Huntington Harbor. These mussel bags were
234 deployed in the early spring when temperatures were below those optimal for *A.*
235 *fundyense* growth ($< 10^{\circ}\text{C}$). Mussel bags were collected weekly from each site and
236 mussels were shucked and extracts were prepared using standard techniques (Association
237 of Official Analytical Chemists (AOAC), 1990). Native soft shell clams (*Mya arenaria*)
238 from Northport Harbor were also harvested and extracts were prepared sporadically
239 during the months of April through May. Toxin levels in shellfish were quantified using
240 standard mouse bioassays (AOAC, 1990). Bioassays were performed by NYSDEC staff
241 at the Stony Brook University Health Sciences Center Division of Laboratory Animal
242 Resources by injecting shellfish extracts into mice (strain CD-1).

243 3. Results:

244 3.1. 2007 Northport Harbor *Alexandrium fundyense* bloom

245 During April of 2007 there was a bloom of non-*Alexandrium*, nanophytoplankton
246 (2-20 μm) in Northport Harbor which had chlorophyll levels exceeding $25 \mu\text{g L}^{-1}$ (Fig. 2)
247 and was comprised primarily of diatoms ($95\pm 3\%$ of cells enumerated). During May, as
248 surface temperatures stabilized at $\sim 15^{\circ}\text{C}$, the abundance of nanophytoplankton began to
249 decline and a modest ($>1,000 \text{ cells L}^{-1}$) *A. fundyense* bloom developed (Fig. 2b). *A.*
250 *fundyense* cells were detected in the water column from 8 May to 20 June with cell

251 densities peaking at 2,650 cells L⁻¹ on 23 May (Fig. 2a, Table 1). Elevated toxin levels (>
252 2 pmol STX eq. L⁻¹) in the water column were present through the bloom, with levels
253 peaking at 130 pmol STX eq. L⁻¹ in unison with peak cell densities (Fig. 2a, Table 1).
254 The largest size fraction of chlorophyll (> 20 µm) accounted for 23±0.8% of the total
255 chlorophyll during the bloom peak. Both ammonium and silicate concentrations
256 increased slightly during the bloom compared to before and after the *A. fundyense* bloom
257 as did δ¹⁵N of the total plankton community which reached its annual maximum
258 (9.7±1.2‰) during the peak of the bloom (Fig. 2c, Fig. 3). During the week following
259 peak cell densities, elevated levels of toxins were found in mussels deployed in Northport
260 Harbor (37 µg STX eq./100g shellfish tissue). The *A. fundyense* bloom ended in June as
261 temperatures exceeded 20°C (Fig. 2a,c).

262 During the bloom in Northport Harbor, *A. fundyense* concentrations in Centerport
263 Harbor ranged from 8 to 50 cells L⁻¹ with low pelagic particulate toxin concentrations
264 (1.42-3.73 pmol STX eq. L⁻¹; Table 1). The remaining sites in Northport-Huntington
265 Harbor complex had < 12 cells L⁻¹ and toxin concentrations below 7.1 pmol STX eq. L⁻¹
266 (Table 1). *A. fundyense* cells and toxins were not detected in the water column of the
267 Northport-Huntington Bay system from July through November. During an experiment
268 conducted on 15 May 2007, the addition of ammonium resulted in a 60% increase in *A.*
269 *fundyense* cell densities compared to unamended control treatments (Fig. 4). During a
270 second experiment (30 May), the addition of ammonium resulted in 25% higher
271 particulate toxin concentrations and 70% higher cell densities (Fig. 4).

272 *3.2. Presence of cysts in the Northport-Huntington Bay area: 2007*

273 During a sediment survey conducted on 14 November 2007, *A. fundyense* cysts
274 were present at low levels in the Northport-Huntington Bay complex (0 - 50 cysts cc of
275 sediment⁻¹; Table 1). The highest concentrations of cysts were located in Northport
276 Harbor with concentrations ranging from 18-50 cysts cc⁻¹(sites 2, 7 and 8; Table 1).
277 Maximal cyst concentrations (50 cysts cc⁻¹) were found at site 8 (Table 1) ~0.6km north
278 of the site with maximal cell densities (site 2; Fig. 1). The remainder of the Northport-
279 Huntington Bay system had relatively low cyst concentrations (0-13 cysts cc⁻¹; Table 1).

280 3.3. 2008 Northport Harbor *Alexandrium fundyense* bloom

281 During April and May of 2008, an intense *Alexandrium fundyense* bloom
282 developed and persisted in Northport Harbor for six weeks, during which temperatures
283 ranged from 10-21°C (Fig. 5a,c). During the bloom, the > 20 µm size class accounted for
284 45 ± 1.2% (up to 76% on 16 May) of total chlorophyll *a* (Fig. 5b). The first peak of the
285 bloom occurred on 16 May when 1.2 x 10⁶ *A. fundyense* cells L⁻¹ and 24,662 pmol STX
286 eq. L⁻¹ were present (Table 1). A secondary bloom peak occurred on 23 May (6 x 10⁵ *A.*
287 *fundyense* cells L⁻¹) and a secondary toxin peak occurred three days later on 26 May
288 (7,300 pmol STX eq. L⁻¹; Fig. 5a). Concentrations of nitrate, ammonium, and phosphate
289 were all significantly (p<0.01 for each, t-test) higher before and after the bloom
290 (phosphate 1.5 ± 0.3µM, nitrate 14.1 ± 2.6µM, ammonium 7.0 ±2.0µM) compared to
291 during the bloom peak (6 May through 29 May; phosphate 0.5 ±0.1µM, nitrate 5.0
292 ±1.5µM, ammonium 1.8 ± 1.0µM; Fig. 5c). In contrast, silicate levels gradually rose
293 from 7µM to 32µM from April through June (Fig. 5c). Throughout the bloom period, the
294 δ¹⁵N of particulate organic matter ranged from 12 to 23‰ (Fig. 3). Mussel toxin levels
295 exceeded the regulatory closure limit (80 µg STX eq./100g shellfish tissue) two weeks

296 after the first detection of *A. fundyense* cells and peaked on 27 May (1,400 μg STX
297 eq./100g shellfish tissue) 11 days after peak cell and water column toxin concentrations
298 (Fig. 5a). Native soft shell clams from this area were also highly toxic (600 μg STX
299 eq./100g shellfish tissue). As such, 7,000 acres of shellfish beds in Northport and
300 Huntington Bays were closed to shellfishing for most of May and June 2008. During the
301 demise of the *A. fundyense* bloom, water column temperatures rose above 20°C and 2 –
302 20 μm size fraction chlorophyll *a* levels increased nearly five-fold (Fig. 5b,c).

303 Although other sites in Northport Harbor had the highest levels of *A. fundyense*
304 during the 2008 bloom (sites 7 and 8 cell densities and toxin concentrations = 5.5×10^5
305 cells L^{-1} and 4.5×10^3 pmol STX eq. L^{-1} and 8.8×10^5 cells L^{-1} and 1.9×10^4 pmol STX
306 eq. L^{-1} , respectively; Table 1), elevated cell densities and toxin concentrations were also
307 present throughout the Northport-Huntington Bay system (Table 1). Centerport Harbor
308 (site 1), had peak cell densities of 7,170 cells L^{-1} and toxin concentrations of 183 pmol
309 STX eq. L^{-1} (Table 1). *A. fundyense* cell densities in Huntington Harbor (site 6) peaked
310 at 24,900 cells L^{-1} with corresponding toxin concentrations of 312 pmol STX eq. L^{-1}
311 (Table 1). After the occurrence of peak cell densities in Huntington Harbor, high levels
312 of toxin were quantified in deployed mussels (161 μg STX eq./100g shellfish tissue).
313 Peak cell densities occurred across Northport-Huntington Bay between 16-26 May with
314 $>10^4$ cells L^{-1} found throughout the system and over 8×10^3 cells L^{-1} in Long Island
315 Sound (Table 1).

316 3.4. Nutrient amendment experiments: 2008

317 The response of *A. fundyense* populations to nutrient amendments changed
318 through the course of the bloom. During experiments conducted at the beginning (30

319 April) and the demise of the *A. fundyense* bloom (2 June), there were no significant
320 changes in toxin concentrations in response to nutrient amendments (Fig. 6). However
321 during these same experiments, the addition of ammonium (10 μM on 30 April; 20 μM
322 on 2 June) significantly increased *A. fundyense* densities compared to the control ($p < 0.01$,
323 Student-Newman-Keuls; Fig. 6). On 6 May, the addition of ammonium (40 μM) yielded
324 a significant ($p < 0.001$, Student-Newman-Keuls) increase in both *A. fundyense* densities
325 and toxin concentrations by 4-fold and 8-fold, respectively, compared to controls. At the
326 same time, the addition of smaller concentrations of ammonium (10 and 20 μM) yielded
327 smaller, but significant ($p < 0.01$, Student-Newman-Keuls), increases in toxin (5-fold and
328 2-fold higher compared to controls, respectively) relative to the unamended control but
329 did not significantly alter cell densities. During the experiment conducted on 12 May the
330 enrichment of each nitrogenous compound produced significantly higher toxin
331 concentrations (3 – 10 fold increase compared to controls; $p < 0.001$, Student-Newman-
332 Keuls; Fig. 6). During the same experiment, *A. fundyense* densities were also
333 significantly ($p < 0.05$, Student-Newman-Keuls) enhanced by the additions of glutamine,
334 nitrate and ammonium (10 and 40 μM); other N compounds (urea, ammonium (20 μM),
335 and ammonium + phosphate) increased *A. fundyense* densities (60-80%), but not
336 significantly (Fig. 6). During late May (19 May, 26 May) the addition of N (all
337 nitrogenous compounds on 26 May, and only nitrate and urea on 19 May) yielded
338 modest, but non-significant increases (10 – 60%) in *A. fundyense* densities compared to
339 controls. During the 19 May experiment, the addition of ammonium (20 μM) and urea
340 resulted in modest (50% and 33%) increases in toxin, while toxin levels were

341 significantly ($p < 0.05$, Student-Newman-Keuls) enhanced by the addition of nitrate and
342 ammonium compared to the control during the 26 May experiment (Fig. 6).

343 Toxin concentrations normalized per cell were significantly increased by nutrient
344 enrichment in four of the six experiments conducted in 2008 ($p < 0.05$, Student Newman-
345 Keuls; Fig. 7). The exceptions were the first (30 April) experiment during which cell-
346 normalized toxin levels were unchanged and the final (2 June) experiment during which
347 the addition of N and P significantly decreased levels ($p < 0.05$, Student Newman-Keuls;
348 Fig. 7). Experiments conducted on both 12 May and 26 May resulted in the most
349 significant increases in toxin per cell for all N and P additions (2 – 4 times higher;
350 $p < 0.05$, Student-Newman-Keuls) with the exception of urea on 26 May (Fig. 7). In
351 contrast, only ammonium enrichment significantly increased cell-normalized toxin levels
352 during the 6 May and 19 May experiments ($p < 0.05$; Fig. 7).

353 3.5. Presence of cysts in the Northport-Huntington Bay area: 2008

354 The cyst survey conducted on 11 November 2008 indicated that *A. fundyense*
355 cysts were present at nearly every site in the Northport-Huntington Bay complex and
356 abundances were nearly an order of magnitude higher than those present in November
357 2007 (Table 1). Cyst concentrations were the highest in Northport Harbor with
358 concentrations ranging from 220 to 745 cysts cc^{-1} . As was the case in 2007, site 8 had
359 the highest cyst concentrations (745 cysts cc^{-1} , Table 1). Sites located just outside of
360 Northport Harbor also had elevated cyst concentrations compared to 2007 (20 - 115 cysts
361 cc^{-1} , Table 1). The western part of the Northport-Huntington Bay complex generally had
362 lower cyst concentrations (0-15 cysts cc^{-1} , Fig.1, Table 1) compared to the eastern part of
363 the bay.

364 3.6. Meteorological conditions: Winter and spring 2007 v 2008

365 Atmospheric temperatures were significantly ($p < 0.001$, t-test) warmer in February
366 2008 ($1.3 \pm 0.8^\circ\text{C}$) than February 2007 ($-2.5 \pm 0.7^\circ\text{C}$) as well as 1°C warmer than the long
367 term monthly mean (0.3°C) (Fig. 8). Furthermore, March 2008 ($4.7 \pm 0.5^\circ\text{C}$) was 1.1°C
368 warmer than March 2007 ($3.6 \pm 1.0^\circ\text{C}$) and slightly warmer than the long term monthly
369 mean (4.2°C). April 2008 ($10.9 \pm 0.7^\circ\text{C}$) was significantly ($p = 0.05$, t-test) warmer than
370 April 2007 ($8.7 \pm 0.9^\circ\text{C}$) as well as 1.5°C warmer than the long term monthly mean (9.4°C)
371 (Fig. 8). In May 2008, temperatures ($14.1 \pm 0.6^\circ\text{C}$) were cooler than both May 2007
372 ($16.0 \pm 0.8^\circ\text{C}$) and the long term monthly mean (15°C) (Fig. 8). During April of 2008,
373 winds blew persistently from the SE ($160 \pm 18.7^\circ$), whereas April 2007 winds came from
374 the SW ($238 \pm 19.8^\circ$; $p = 0.006$, t-test). There were no significant differences in
375 precipitation or wind intensity between 2007 and 2008 compared to long-term averages
376 for the months of January through June.

377 **4. Discussion:**

378 4.1. 2007 & 2008 *Alexandrium fundyense* bloom toxicity and intensity

379 This study documented the dynamics of two contrasting blooms, one of which
380 achieved cell densities greater than 10^6 cells L^{-1} and resulted in the closure of 7000 acres
381 of shellfish beds in Northport, NY. The *A. fundyense* bloom in 2008 was dramatically
382 more intense and toxic than the bloom in 2007, with toxin and cell concentrations (means
383 = $5,816$ pmol STX eq. L^{-1} ; $353,184$ cells L^{-1}) in May 2008 being two and three orders of
384 magnitude higher ($p < 0.001$, t-test) than those in May 2007; toxin levels were
385 significantly correlated ($r^2 = 0.942$, $p < 0.001$) with *A. fundyense* abundances during both
386 years. The sustained high densities of *A. fundyense* during the peak of the 2008 bloom

387 ($>10^5$ cells L^{-1}) were higher than those typically found in coastal embayments or open
388 waters of the Gulf of Maine where blooms are annual occurrences and cell densities are
389 usually below 10^4 cells L^{-1} (Townsend et al., 2001; Love et al., 2005; Poulton et al., 2005;
390 Townsend et al., 2005a, b). Similar concentrations ($>10^5$ cells L^{-1}) of *A. fundyense* have
391 also been observed in the Nauset Marsh System on Cape Cod (D.M. Anderson,
392 unpublished data). While absolute toxin levels in Northport Harbor (up to 24,662 pmol
393 STX eq. L^{-1}) were also higher than those reported in Maine (400 pmol STX eq. L^{-1} ;
394 Poulton et al., 2005), the toxin contents or quotas in Northport Harbor (6.2 – 58.8 fmol
395 STX eq. cell $^{-1}$; Fig. 9) were substantially lower than those of *Alexandrium* populations
396 from the Gulf of Maine (36 – 325 fmol cell $^{-1}$; Poulton et al., 2005), a finding consistent
397 with the known north-south gradient in cell toxicity along the western Atlantic coast
398 (Maranda et al., 1985; Anderson et al., 1990a; Anderson et al., 1994; Bricelj and
399 Shumway, 1998), and with the dominance of low-potency saxitoxin congeners in
400 populations from Long Island and Connecticut waters (Anderson et al., 1994). Despite
401 the lower toxicity cells in NY, the large bloom in 2008 caused blue mussels (*Mytilus*
402 *edulis*), and native soft shell clams (*Mya arenaria*) in Northport Bay to become highly
403 toxic (1,400 and 600 μ g STX eq./100g shellfish tissue, respectively) causing the closure
404 of >7,000 acres of shellfish beds for nearly two months (Karen Chytalo, NYSDEC,
405 Marine Division).

406 4.2. The relative importance of nitrogen, cysts, and meteorological conditions promoting 407 New York *Alexandrium fundyense* blooms:

408 The dynamics of *A. fundyense* blooms in Northport Harbor and the differences
409 between the magnitude of the 2007 and 2008 blooms might be controlled by multiple

410 factors including cyst beds, meteorological conditions, and nutrient loading. Benthic cyst
411 concentrations in November 2008 were an order of magnitude greater than those present
412 in November 2007 ($p < 0.001$, t-test) and the spatial extent of cysts also expanded in 2008
413 likely due, in part, to the larger bloom that year compared to 2007. In the Gulf of Maine,
414 cyst seed bed distribution and cyst densities in combination with physical circulation
415 patterns are used to model blooms since cysts provide the inocula for future events
416 (Anderson, 1997; Anderson et al., 2003, 2005a,c; Stock et al., 2005; McGillicuddy et al.,
417 2005). The cyst densities found in Northport Harbor during 2007 were more than an
418 order of magnitude lower than those found in the Gulf of Maine and the Bay of Fundy
419 (Anderson et al., 2005c), suggesting that cysts may be less important to bloom dynamics
420 in this system. This hypothesis is affirmed by comparing the density of cysts in
421 November 2007 to the abundance of cells in May 2008. The highest cyst densities in
422 2007 (50 cysts cc^{-1}) would yield a vegetative population of only 125 cells L^{-1} if all cysts
423 in the top cm of sediment emerged successfully and simultaneously into the 4 m water
424 column. Since this cell abundance is four orders of magnitude smaller than vegetative
425 cell densities observed in 2008 (10^6 cells L^{-1}), *in situ* growth of vegetative cells likely
426 played an important role in the development of the 2008 bloom (Anderson 1998).

427 Meteorological conditions likely affected bloom dynamics in Northport Harbor.
428 Vegetative *A. fundyense* cells are known to grow maximally from 12 to 20 °C (Yentsch et
429 al., 1975; Anderson et al., 1983) and during 2007 and 2008, *A. fundyense* blooms
430 developed when Northport Harbor temperatures were between 10 and 20°C, with
431 temperatures close to 15°C yielding the highest cell densities. The spring of 2008 was
432 warmer than 2007 as during 2007, temperatures persisted between 15°C and 20°C for

433 only three weeks whereas in 2008, temperatures stabilized near 15°C for almost six weeks
434 (mid-April – June), giving the 2008 population more time to bloom. In contrast to early
435 spring, May 2008 temperatures were cooler than both May 2007 and the long term May
436 mean, which likely aided in keeping water temperatures in the optimal range for *A.*
437 *fundyense* growth allowing the large *A. fundyense* bloom to develop. In addition to
438 influencing pelagic cell dynamics, warmer temperatures during early spring 2008 likely
439 stimulated the germination of *A. fundyense* cysts (Anderson and Morel, 1979; Anderson,
440 1998) earlier than in 2007. Wind patterns may have also influenced the 2008 *A.*
441 *fundyense* bloom. During April of 2008, winds blew from the SE, whereas April 2007
442 winds came from the SW. While the SW winds in 2007 might have kept water within
443 Northport Harbor, winds in April 2008 may have spread cells throughout the Northport-
444 Huntington Bay complex and thus may have contributed to the more widespread bloom
445 in that year. Atmospheric conditions such as wind direction have often been found to
446 control the spread and persistence of *Alexandrium* blooms (Anderson and Morel, 1979;
447 Garcon et al., 1986; Anderson, 1997; Townsend et al., 2005a,b).

448 N played a central role in supporting *A. fundyense* blooms in Northport Harbor.
449 During the 2008 bloom, there were significant ($p < 0.01$, t-test) declines in phosphate,
450 nitrate and ammonium concentrations during the *A. fundyense* bloom (6 May through 29
451 May) compared to before and after the bloom, suggesting that there was a larger nutrient
452 demand due to the higher biomass and more prolonged bloom in 2008. Furthermore,
453 nitrate concentrations were significantly ($p < 0.01$, t-test) lower in 2008 ($5.12 \pm 1.58 \mu\text{M}$)
454 compared to 2007 ($12.4 \pm 1.86 \mu\text{M}$) and ammonium concentrations were also lower in
455 2008 ($0.58 \pm 0.17 \mu\text{M}$; 6 May to 26 May) compared to 2007 ($1.34 \pm 0.51 \mu\text{M}$; 8 May to 5

456 June). These observations suggest N was more likely to be limiting to the *A. fundyense*
457 bloom in 2008 compared to 2007. High biomass *A. taylori* blooms in the Mediterranean
458 which are influenced by anthropogenic N loading have caused a drawdown of nutrients
459 similar to that observed in Northport in 2008 (Penna et al., 2002).

460 Nutrient amendment experiments performed during 2007 and 2008 demonstrated
461 that N loading can affect *A. fundyense* densities and toxicity, and affirms that N was
462 important in supporting the large 2008 bloom. Overall, the addition of N (glutamine,
463 nitrate, ammonium and/or urea) resulted in increased *A. fundyense* densities and/or toxin
464 concentrations compared to control treatments during every 2008 experiment. These
465 increases were frequently significant in 2008 (83% of experiments), when ambient
466 inorganic N concentrations were lower, suggesting this bloom was N stressed. On
467 average, the additions of ammonium and glutamine, specifically, resulted in the highest
468 *A. fundyense* densities and toxin concentrations when compared to the addition of other N
469 species when pooling together all experiments conducted in both 2007 and 2008.
470 However, the addition of ammonium most frequently yielded statistically significant
471 increases in *A. fundyense* densities and toxin concentrations compared to control
472 treatments (66% and 50% of experiments in 2008), suggesting that ammonium may
473 promote the formation of toxic *A. fundyense* blooms. The strong response to glutamine
474 also suggests that dissolved organic N and amino acids such as glutamine may play an
475 important role in supporting *Alexandrium* blooms as they are known to do for other
476 HABs (Mulholland et al., 2002; Gobler et al., 2004).

477 The effects of nutrients on the 2008 *A. fundyense* bloom was also evident from
478 cell normalized toxin concentrations found in the field and during experiments. Variation

479 in toxin content per cell of natural bloom populations and isolates from the Gulf of Maine
480 has been previously attributed to nutrient limitation, with N limited cells generally
481 displaying lower levels of toxin (Anderson et al., 1990a,b; Poulton et al., 2005). During
482 the 2008 *A. fundyense* bloom, cell toxicity was high (34.5 - 58.8 fmol STX eq. cell⁻¹) at
483 the beginning and end of the bloom (April and June) but was significantly lower during
484 the peak of the bloom (15.2±5.1 fmol STX eq. cell⁻¹; 6 May - 29 May; p<0.001, Student-
485 Newman-Keuls; Fig 9). Since values of 51.9 ± 29.5 fmol STX eq. cell⁻¹ were measured in
486 nutrient replete cultures of *A. fundyense* strains (n=3) isolated from Northport, this field
487 pattern supports the hypothesis that *A. fundyense* populations were nutrient replete at the
488 end and beginning of the bloom, but nutrient stressed during May. Nutrient amendment
489 experiments displayed similar variations in toxin concentrations normalized per cell, with
490 significant increases in toxin per cell during experimental N loading in general and
491 ammonium loading in particular. The ability of ammonium to consistently increase
492 cellular toxin content has also been observed in *A. tamarensis* cultures (Leong et al.,
493 2004), supporting the hypothesis that ammonium promotes toxic *A. fundyense* blooms.
494 The decreases in toxin per cell during the bloom peak could indicate N-stress causing a
495 cellular partitioning of resources (Leong et al., 2004), with more N put toward growth
496 and less toward toxin production during the peak of the bloom since saxitoxin is a N-rich
497 molecule, containing 7 N atoms (with the decarbamoyl derivatives having 6 N atoms;
498 Samsur et al., 2006).

499 N played an important role in the development and toxicity of *A. fundyense*
500 blooms in Northport, and the Scudder Beach Sewage Treatment Plant, which discharges
501 0.4 million gallons of effluent daily into Northport Harbor, may have been an important

502 N source which supported these blooms (discharge pipe at 40.8965°N, 73.3567°W, Fig.1;
503 Paul Harding, NYSDEC, personal communication). During periods when chlorophyll *a*
504 levels and presumably nutrient demands were low, DIN concentrations in Northport
505 Harbor frequently exceeded 25µM, suggesting there is a strong source of N in this region.
506 The active uptake of sewage-derived N was evident in the isotopic signatures of
507 particulate organic nitrogen (PON) from Northport Harbor as $\delta^{15}\text{N}$ values ranged from 12
508 to 23‰ during large *A. fundyense* blooms. This range overlaps with wastewater derived
509 N (10 to 30 ‰; Kendall, 1998; Bianchi, 2007), and is significantly higher than levels
510 measured in particulate organic matter (POM) of the adjacent waters of Long Island
511 Sound (7 to 9 ‰). Furthermore, toxin and *A. fundyense* densities were significantly
512 correlated to $\delta^{15}\text{N}$ of POM ($r^2=0.63$ and 0.68 , respectively; $p<0.001$) indicating POM was
513 the most enriched in ^{15}N during bloom events. These findings, combined with the ability
514 of N enrichment to significantly increase the abundance and toxicity of *A. fundyense*
515 supports the hypothesis that N from the Scudder Beach wastewater treatment plant or
516 some other sources of highly enriched wastewater supported the proliferation of these
517 blooms. Similarly, anthropogenic nutrient loading has been associated with an increase
518 in PSP incidences caused by *A. catenella* in multiple marine ecosystems including
519 shallow, poorly flushed coastal embayments of the northwest US (Trainer et al., 2003).

520 Nutrient loading has been cited as a factor responsible for promoting multiple
521 HABs around the world (Anderson et al., 2002; Penna et al., 2002; Trainer et al., 2003;
522 Poulton et al., 2005; Glibert et al., 2006; Anderson et al., 2008; Heisler et al., 2008).
523 However, the degree to which *A. fundyense* blooms are related to anthropogenic nutrient
524 loading to coastal systems has been unclear (Anderson, 1994; Anderson et al., 2002,

525 2008; Glibert et al., 2005). This study demonstrated that N enrichment was capable of
526 significantly increasing *A. fundyense* cell densities, particulate toxin levels, and the levels
527 of toxin per cell. Moreover, the isotopic N signature of POM during blooms was
528 consistent with those found in wastewater. This data set combined with the proximity of
529 a sewage treatment plant to the occurrence of this bloom indicates that estuarine *A.*
530 *fundyense* blooms can be promoted by anthropogenic N loading. It is possible that
531 anthropogenic nutrient loading plays a similar role in the development of *A. fundyense*
532 blooms in coastal embayments around the world, although this phenomenon has not been
533 well studied.

534

535 *Acknowledgements-* We gratefully acknowledge F. Koch, A. Marcoval, J. Goleski, A.
536 Burson, M. Harke, T. Davis, S. Angles, C. Wall, Y.Z. Tang, C. Lehmann and R.
537 Hattenrath for their assistance in the field and with sample processing. We would also
538 like to thank B. Keafer, K. Norton and D. Kulis for assistance with the oligonucleotide
539 method, cyst sampling methodologies as well as HPLC analysis of saxitoxin samples.
540 This work was supported by a grant from EPA's Long Island Sound Study, New York
541 Sea Grant, and the New York State Department of Environmental Conservation (to CJG)
542 and from the NOAA Sea Grant Program (Grant No. NA06OAR4170021 (R/B-177)) to
543 DMA.

544

545 **References**

- 546 Anderson, D.M., 1994. Red tides. *Sci. Am.* 271(2),52-58.
547
- 548 Anderson, D.M., 1997. Bloom dynamics of toxic *Alexandrium* species in the northeastern
549 US. *Limnol. Oceanogr.* 42,1009-1022.
550
- 551 Anderson, D.M., 1998. Physiology and bloom dynamics of toxic *Alexandrium* species,
552 with emphasis on life cycle transitions. In: Anderson, D.M., Cembella, A.D.,
553 Hallegraeff, G.M. (Eds), *The Physiological Ecology of Harmful Algal Blooms*.
554 Springer-Verlag, Heidelberg, pp. 29-48.
555
- 556 Anderson, D.M., Burkholder, J.M., Cochlan, W.P., Glibert, P.M., Gobler, C.J., Heil,
557 C.A., Kudela, R.M., Parsons, M.L., Rensel, J.E.J., Townsend, D.W., Trainer,
558 V.L., Vargo, G.A., 2008. Harmful algal blooms and eutrophication: Examining
559 linkages from selected coastal regions of the United States. *Harmful Algae* 8,39-
560 53.
561
- 562 Anderson, D.M., Chisholm, S.W., Watras, C.J., 1983. Importance of life cycle events in
563 the population dynamics of *Gonyaulax tamarensis*. *Mar. Biol.* 76,179-189.
564
- 565 Anderson, D.M., Fukuyo, Y., Matsuoka, K., 2003. Cyst methodologies. In: Hallegraeff,
566 G.M., Anderson, D.M., Cembella, A.D. (Eds) *Manual on Harmful Marine*
567 *Microalgae*, Monographs on Oceanographic Methodology, 11, UNESCO, pp.
568 165-190.
569
- 570 Anderson, D.M., Glibert, P.M., Burkholder, J.M., 2002. Harmful algal blooms and
571 eutrophication: nutrient sources, composition, and consequences. *Estuaries*
572 25,704-726.
573
- 574 Anderson, D.M., Hoagland, P., Kaoru, Y., White, A.W., 2000. Estimated Annual
575 Economic Impacts from Harmful Algal Blooms (HABs) in the United States.
576 Woods Hole Oceanographic Inst Tech Rept, WHOI 2000-11. (99 pp)
577
- 578 Anderson, D.M., Keafer, B.A., Geyer, W.R., Signell, R.P., Loder, T.C., 2005a. Toxic
579 *Alexandrium* blooms in the western Gulf of Maine: The plume advection
580 hypothesis revisited. *Limnol. Oceanogr.* 50(1), 328-345.
581
- 582 Anderson, D.M., Kulis, D.M., Doucette, G.J., Gallagher, J.C., Balech, E., 1994.
583 Biogeography of toxic dinoflagellates in the genus *Alexandrium* from the
584 northeastern United States and Canada. *Mar. Biol.* 120, 467-478.
585
- 586 Anderson, D.M., Kulis, D.M., Keafer, B.A., Gribble, K.E., Marin, R., Scholin, C.A.,
587 2005b. Identification and enumeration of *Alexandrium* spp. from the Gulf of
588 Maine using molecular probes. *Deep-Sea Res. II* 52,2467-2490.
589

- 590 Anderson, D.M., Kulis, D.M., Orphanos, J.A., Ceurvels, A.R., 1982. Distribution of the
591 toxic dinoflagellate *Gonyaulax tamarensis* in the southern New England region.
592 Est. Coast. Shelf Sci. 14, 447-458.
593
- 594 Anderson, D.M., Kulis, D.M., Sullivan, J.J., Hall, S., 1990a. Toxin composition
595 variations in one isolate of the dinoflagellate *Alexandrium fundyense*. Toxicon
596 28,885-893.
597
- 598 Anderson, D.M., Kulis, D.M., Sullivan, J.J., Hall, S., Lee, C., 1990b. Dynamics and
599 physiology of saxitoxin production by the dinoflagellates *Alexandrium* spp. Mar.
600 Biol. 104, 511-524.
601
- 602 Anderson, D.M., Morel, F.M.M., 1979. The seeding of two red tide blooms by the
603 germination of benthic *Gonyaulax tamarensis* hypnocyts. Est. Coast. Mar. Sci.
604 8,279-293.
605
- 606 Anderson, D.M., Stock, C.A., Keafer, B.A., Bronzino Nelson, A., Thompson, B.,
607 McGillicuddy, D.J., Keller, M., Matrai, P.A., Martin, J., 2005c. *Alexandrium*
608 *fundyense* cyst dynamics in the Gulf of Maine. Deep-Sea Res. II 52, 2522-2542.
609
- 610 Association of Official Analytical Chemists. In: Horowitz, W. (Ed) Official Methods of
611 Analysis. Washington, DC, pp. 881-882.
612
- 613 Bianchi, T.S., 2007. Biogeochemistry of Estuaries. Oxford Press.
614
- 615 Bricelj, V.M., Shumway, S.E., 1998. Paralytic shellfish toxins in bivalve molluscs:
616 Occurrence, transfer kinetics, and biotransformation. Rev. Fish Sci. 6(4), 315 –
617 383.
618
- 619 Erdner, D.L., Dyble, J., Parsons, M.L., Stevens, R.C., Hubbard K.A., Wraebel, M.L.,
620 Moore, S.K., Lefebvre K.A., Anderson, D.M., Bienfang, P., Bidigare, R.R.,
621 Parker, M.S., Moeller, P., Brand, L.E., Trainer, V.L., 2008. Centers for Oceans
622 and Human Health: a unified approach to the challenge of harmful algal blooms.
623 Environmental Health 7(Suppl 2):S2
624
- 625 Garcon, V.C., Stolzenbach, K.D., Anderson, D.M., 1986. Tidal flushing of an estuarine
626 embayment subject to recurrent dinoflagellate blooms. Estuaries 9,179-187.
627
- 628 Glibert, P.M., Anderson, D.M., Gentien, P., Graneli, E., Sellner, K., 2005. The global,
629 complex phenomena of harmful algal blooms. Oceanography 18(2),132-141.
630
- 631 Glibert, P.M., Harrison, J., Heil, C., Seitzinger, S., 2006. Escalating worldwide use of
632 urea- a global change contributing to coastal eutrophication. Biogeochemistry
633 77,441-463.
634

- 635 Gobler, C.J., Deonaraine, S.N., Leigh-Bell, J., Downes Gastrich, M., Anderson, O.R.,
636 Wilhelm, S.W., 2004. Ecology of phytoplankton communities dominated by
637 *Aureococcus anophagefferens*: The role of viruses, nutrients, and
638 microzooplankton grazing. *Harmful Algae* 3,471-483.
639
- 640 Gobler, C.J., Lonsdale, D.J., Boyer, G.L., 2005. A synthesis and review of causes and
641 impact of harmful brown tide blooms caused by the alga, *Aureococcus*
642 *anophagefferens*. *Estuaries* 28,726-749.
643
- 644 Hackett, J.D., Anderson, D.M., Erdner, D.L., Bhattacharya, D., 2004. Dinoflagellates: a
645 remarkable evolutionary experiment. *Am. J. Bot.* 91(10): 1523-1534.
646
- 647 Hasle, G.R., 1978. The inverted microscope method. *Monogr. Oceanogr. Meth.* 6, 88-96.
648
- 649 Heisler, J., Glibert, P.M., Burkholder, J.M., Anderson, D.M., Cochlan, W., Dennison,
650 W.C., Dortch, Q., Gobler, C.J., Heil, C.A., Humphries, E., Lewitus, A., Magnien,
651 R., Marshall, H.G., Sellner, K., Stockwell, D.A., Stoecker, D.K., Suddleson, M.,
652 2008. Eutrophication and harmful algal blooms: A scientific consensus. *Harmful*
653 *Algae* 8, 3-13.
654
- 655 Jin, D., Hoagland, P., 2008. The value of harmful algal bloom predictions to the
656 nearshore commercial shellfish fishery in the Gulf of Maine. *Harmful Algae* 7,
657 772-781.
658
- 659 Jin, D., Thunberg, E., Hoagland, P., 2008. Economic impact of the 2005 red tide event on
660 commercial shellfish fisheries in New England. *Ocean Coast. Manage.* 51,420-
661 429.
662
- 663 Jones, M.N., 1984. Nitrate reduction by shaking with cadmium: alternative to cadmium
664 columns. *Water Res.* 18, 643-646.
665
- 666 Kendall, C., 1998. Tracing Nitrogen Sources and Cycling in Catchments. In: Kendall, C.,
667 McDonnell, J.J., (Eds) *Isotope Tracers in Catchment Hydrology*. Elsevier Science
668 BV, Amsterdam, pp. 519-576.
669
- 670 Kvitek, R.G., Beitler, M.K., 1988. A case for sequestering of paralytic shellfish toxins as
671 a chemical defense. *J. Shellfish Res.* 7(4), 629-636.
672
- 673 Leong, S.C.Y., Murata, A., Nagashima, Y., Taguchi, S., 2004. Variability in toxicity of
674 the dinoflagellate *Alexandrium tamarense* in response to different nitrogen
675 sources and concentrations. *Toxicon* 43, 407-415.
676
- 677 Love, R.C., Loder, T.C., Keafer, B.A., 2005. Nutrient conditions during *Alexandrium*
678 *fundyense* blooms in the western Gulf of Maine, USA. *Deep-Sea Res. II* 52,2450-
679 2466.
680

- 681 Maranda, L., Anderson, D.M., Shimizu, Y., 1985. Comparison of toxicity between
682 populations of *Gonyaulax tamarensis* of eastern North American waters. Estuar.
683 Coast. Shelf Sci. 21,401-410.
684
- 685 McGillicuddy, D.J., Anderson, D.M., Lynch, D.R., Townsend, D.W., 2005. Mechanisms
686 regulating large-scale seasonal fluctuations in *Alexandrium fundyense* populations
687 in the Gulf of Maine: Results from a physical-biological model. Deep-Sea Res. II
688 52, 2698-2714.
689
- 690 Mulholland, M.R., Gobler, C.J., Lee, C., 2002. Peptide hydrolysis, amino acid oxidation,
691 and nitrogen uptake in communities seasonally dominated by *Aureococcus*
692 *anophagefferens*. Limnol. Oceanogr. 47(4), 1094-1108.
693
- 694 Mulligan, H.F., 1975. Oceanographic factors associated with New England red-tide
695 blooms. In: LoCicero, V.R. (Ed), Toxic Dinoflagellate Blooms. Proceedings of
696 the International Conference (1st), Massachusetts Science and Technology
697 Foundation. pp. 23-40.
698
- 699 Parsons, T.R., Maita, Y., Lalli, C.M., 1984. A manual of chemical and biological
700 methods for seawater analysis. Pergamon Press, Oxford.
701
- 702 Penna, A., Giacobbe, M.G., Penna, N., Andreoni, F., Magnani, M., 2002. Seasonal
703 blooms of the HAB dinoflagellate *Alexandrium taylori* Balech in a new
704 Mediterranean area (Vulcano, Aeolian Islands) Mar. Ecol. 23,320-328.
705
- 706 Poulton, N.J., Keafer, B.A., Anderson, D.M., 2005. Toxin variability in natural
707 populations of *Alexandrium fundyense* in Casco Bay, Maine – evidence of
708 nitrogen limitation. Deep Sea Res. II 52,2501-2521.
709
- 710 Samsur, M., Yamaguchi, Y., Sagara, T., Takatani, T., Arakawa, O., Noguchi, T., 2006.
711 Accumulation and depuration profiles of PSP toxins in the short-necked clam
712 *Tapes japonica* fed with the toxic dinoflagellate *Alexandrium catenella*. Toxicon
713 48,323-330.
714
- 715 Schrey, S.E., Carpenter, E.J., Anderson, D.M., 1984. The abundance and distribution of
716 the toxic dinoflagellate, *Gonyaulax tamarensis*, in Long Island Estuaries.
717 Estuaries 7, 472-477.
718
- 719 Sellner, K.G., Doucette, G.J., Kirkpatrick, G.J., 2003. Harmful algal blooms: causes,
720 impacts and detection. J. Ind. Microbiol. Biotechnol. 30,383-406.
721
- 722 Stock, C.A., McGillicuddy Jr., D.J., Solow, A.R., Anderson, D.M., 2005. Evaluating
723 hypotheses for the initiation and development of *Alexandrium fundyense* blooms
724 in the western Gulf of Maine using a coupled physical-biological model. Deep-
725 Sea Res II 52,2715-2744.
726

- 727 Townsend, D.W., Pettigrew, N.R., Thomas, A.C., 2001. Offshore blooms of the red tide
728 dinoflagellate *Alexandrium* sp., in the Gulf of Maine. Cont. Shelf Res. 21,347-
729 369.
730
- 731 Townsend, D.W., Bennett, S.L., Thomas, M.A., 2005a. Diel vertical distributions of the
732 red tide dinoflagellate *Alexandrium fundyense* in the Gulf of Maine. Deep-Sea
733 Res II 52,2593-2602.
734
- 735 Townsend, D.W., Pettigrew, N.R., Thomas, A.C., 2005b. On the nature of *Alexandrium*
736 *fundyense* blooms in the Gulf of Maine. Deep-Sea Res II 52, 2603-2630.
737
- 738 Trainer, V.L., Eberhart, B.T.L., Wekell, J.C., Adams, N.G., Hanson, L., Cox, F., Dowell,
739 J., 2003. Paralytic shellfish toxins in Puget Sound, Washington State. J. Shellfish
740 Res. 22, 213-223.
741
- 742 Yamaguchi, M., Itakura, S., Imai, I., Ishida, Y., 1995. A rapid and precise technique for
743 enumeration of resting cysts of *Alexandrium* spp. (Dinophyceae) in natural
744 sediments. Phycologia 34, 207–214.
745
- 746 Yentsch, C.M., Cole, E.J., Salvaggio, M.G., 1975. Some of the growth characteristics of
747 *Gonyaulax tamarensis* isolated from the Gulf of Maine. In: LoCicero, V.R. (Ed)
748 Proceedings of the International Conference (1st), Massachusetts Science and
749 Technology Foundation, pp. 163-180.
750

751

List of Figures

- 752 **Figure 1.** Site locations in Northport-Huntington Bay complex; located on the north
 753 shore of Long Island, NY, USA. Cyst sampling locations include sites 1-17
 754 whereas pelagic samples were obtained from sites 1-8, 10, 11, 16 and LIS, SD=
 755 Sewage discharge pipe from Scudder Beach Sewage Treatment Plant.
- 756 **Figure 2.** Dynamics of: A) Pelagic toxin (pmol STX eq. L⁻¹) and *Alexandrium*
 757 *fundyense* densities (cells L⁻¹ x 10²), off-scale values are indicated by the black
 758 arrow, B) size fractioned chlorophyll *a* (µg L⁻¹), and C) inorganic nutrient
 759 concentrations (µM) and temperature (°C) in Northport Harbor during spring
 760 2007. Points are means while error bars represent SD.
- 761 **Figure 3.** δ¹⁵N (‰) values of particulate organic nitrogen from Northport Harbor during
 762 spring 2007 and 2008. The ranges of levels measured in particulate organic
 763 matter in Long Island Sound are depicted by the grey bar. Nitrogen from
 764 wastewater typically ranges from 10-30‰ (Kendall 1998, Bianchi 2007). Points
 765 are means while error bars represent SD.
- 766 **Figure 4.** *Alexandrium fundyense* densities (cells L⁻¹ x 10²) and toxin concentrations
 767 (pmol STX eq. L⁻¹) at the end of nutrient amendment experiments conducted
 768 during May of 2007. Bars are means while error bars represent SD of triplicate
 769 measurements.
- 770 **Figure 5.** Dynamics of: A) Pelagic toxin (pmol STX eq. L⁻¹ x 10³), *Alexandrium*
 771 *fundyense* densities (cells L⁻¹ x 10⁶) and toxin concentrations (µg STX eq. 100g⁻¹
 772 x 10³) in deployed blue mussels (*Mytilus edulis*) as determined by mouse
 773 bioassay, off-scale values are indicated by the black arrow, B) size fractioned
 774 chlorophyll *a* (µg L⁻¹), and C) inorganic nutrient concentrations (µM) and
 775 temperature (°C) in Northport Harbor (site 2) during spring 2008. Points are
 776 means while error bars represent SD.
- 777 **Figure 6.** *Alexandrium fundyense* densities (cells L⁻¹) and toxin concentrations (pmol
 778 STX eq. L⁻¹) following experimental nutrient amendments during April - June
 779 2008. Bars are means while error bars represent SD of triplicate & duplicate
 780 (saxitoxin concentrations) measurements. C= control, P= phosphate, N= nitrate,
 781 U= urea, A= ammonium (10, 20 and 40 indicate different concentrations added in
 782 µM), G= glutamine, and A+P= ammonium + phosphate.
- 783 **Figure 7.** Toxin per cell (fmol STX eq. cell⁻¹) following experimental nutrient
 784 amendments during April - June 2008. Bars are means while error bars represent
 785 SD of duplicate measurements. C= control, P= phosphate, N= nitrate, U= urea,
 786 A= ammonium (10, 20 and 40 indicate different concentrations added in µM), G=
 787 glutamine, and A+P= ammonium + phosphate.
- 788 **Figure 8.** Atmospheric temperatures (°C) observed during the winter and spring of 2007
 789 and 2008 compared to long term monthly means from Islip, NY, USA. Bars are
 790 monthly means while error bars represent SE.
- 791 **Figure 9.** *Alexandrium fundyense* densities (cells L⁻¹ x 10⁶) and toxin concentrations per
 792 cell (fmol STX eq. cell⁻¹) for Northport Harbor (site 2) in 2008. Points are means
 793 while error bars represent SD (error bars for toxin concentrations per cell
 794 represent propagated SD). The area highlighted in grey represents the range of

795 total toxin concentrations per cell (fmol STX eq. cell⁻¹) measured in nutrient
796 replete cultures of *Alexandrium fundyense* isolated from Northport Bay.

797

List of Tables

798

799 **Table 1.** Peak *Alexandrium fundyense* densities (cells L⁻¹) and pelagic toxin
800 concentrations (pmol STX eq. L⁻¹) in Northport- Huntington Bay, NY for 2007
801 (May 15th-30th) and 2008 (May 16th-26th), and mean cyst concentrations (cysts cc⁻¹)
802 in Northport-Huntington Bay, NY sediments during November of 2007 and
803 2008. Values in parentheses are standard deviations.

804

805

806

807


808

809

810


811

812



813


814 **Figure 1.** Site locations in Northport- Huntington Bay complex; located on the north
 815 shore of Long Island, NY, USA. Cyst sampling locations include sites 1-17 whereas
 816 pelagic samples were obtained from sites 1-8, 10, 11, 16 and LIS, SD= Sewage discharge
 817 pipe from Scudder Beach Sewage Treatment Plant.
 818



819

820 **Figure 2.** Dynamics of: A) Pelagic toxin (pmol STX eq. L⁻¹) and *Alexandrium*
 821 *fundyense* densities (cells L⁻¹ x 10²), off-scale values are indicated by the black arrow, B)
 822 size fractionated chlorophyll *a* (μg L⁻¹), and C) inorganic nutrient concentrations (μM)
 823 and temperature (°C) in Northport Harbor during spring 2007. Points are means while error
 824 bars represent SD.

825



826

827 **Figure 3.** $\delta^{15}\text{N}$ (‰) values of particulate organic nitrogen from Northport Harbor during
 828 spring 2007 and 2008. The ranges of levels measured in particulate organic matter in
 829 Long Island Sound are depicted by the grey bar. Nitrogen from wastewater typically
 830 ranges from 10-30‰ (Kendall 1998, Bianchi 2007). Points are means while error bars
 831 represent SD.


832

833

834


835

836




837

838 **Figure 4.** *Alexandrium fundyense* densities (cells L⁻¹ x 10²) and toxin concentrations
 839 (pmol STX eq. L⁻¹) at the end of nutrient amendment experiments conducted during May
 840 of 2007. Bars are means while error bars represent SD of triplicate measurements.




841
 842 **Figure 5.** Dynamics of: A) Pelagic toxin (pmol STX eq. L⁻¹ x 10³), *Alexandrium*
 843 *fundyense* densities (cells L⁻¹ x 10⁶) and toxin concentrations (μg STX eq. 100g⁻¹ x 10³) in
 844 deployed blue mussels (*Mytilus edulis*) as determined by mouse bioassay, off-scale
 845 values are indicated by the black arrow, B) size fractioned chlorophyll *a* (μg L⁻¹), and C)
 846 inorganic nutrient concentrations (μM) and temperature (°C) in Northport Harbor (site 2)
 847 during spring 2008. Points are means while error bars represent SD.



848


849 **Figure 6.** *Alexandrium fundyense* densities (cells L⁻¹) and toxin concentrations (pmol
 850 STX eq. L⁻¹) following experimental nutrient amendments during April - June 2008.
 851 Bars are means while error bars represent SD of triplicate & duplicate (toxin
 852 concentrations) measurements. C= control, P= phosphate, N= nitrate, U= urea, A=
 853 ammonium (10, 20 and 40 indicate different concentrations added in μM), G= glutamine,
 854 and A+P= ammonium + phosphate.



855
856
857
858
859
860
861

Figure 7. Toxin per cell (fmol STX eq. cell⁻¹) following experimental nutrient amendments during April - June 2008. Bars are means while error bars represent SD of duplicate measurements. C= control, P= phosphate, N= nitrate, U= urea, A= ammonium (10, 20 and 40 indicate different concentrations added in μ M), G= glutamine, and A+P= ammonium + phosphate.

862



863


864

865 **Figure 8.** Atmospheric temperatures (°C) observed during the winter and spring of 2007

866 and 2008 compared to long term monthly means in Islip, NY, USA. Bars are monthly

867 means while error bars represent SE.

868



869
870

871 **Figure 9.** *Alexandrium fundyense* densities (cells L⁻¹ x 10⁶) and toxin concentrations per
872 cell (fmol STX eq. cell⁻¹) for Northport Harbor (site 2) in 2008. Points are means while
873 error bars represent SD (error bars for toxin concentrations per cell represent propagated
874 SD). The area highlighted in grey represents the range of total toxin concentrations per
875 cell (fmol STX eq. cell⁻¹) measured in nutrient replete cultures of *Alexandrium fundyense*
876 isolated from Northport Bay.

877
878
879
880
881
882
883
884
885
886
887
888
889
890
891

892 **Table 1.** Maximal *Alexandrium fundyense* densities (cells L⁻¹) and pelagic toxin
 893 concentrations (pmol STX eq. L⁻¹) in Northport- Huntington Bay, NY for 2007 (May
 894 15th-30th) and 2008 (May 16th-26th), and mean cyst concentrations (cysts cc⁻¹) in
 895 Northport-Huntington Bay, NY sediments during November of 2007 and 2008. Values
 896 are means with standard deviations in parentheses.
 897
 898

Northport-Huntington Bay						
Site	<i>A. fundyense</i> (cells L ⁻¹)		water column toxin (pmol STX eq. L ⁻¹)		<i>A. fundyense</i> cysts (cc ⁻¹)	
	2007	2008	2007	2008	2007	2008
1	50 (9)	7,166 (983)	3.73 (0.68)	183 (60.8)	3 (3)	25 (7)
2	2650 (81)	1,199,567 (435,248)	130 (3.90)	24,662 (564)	18 (12)	345 (35)
3	9 (4)	4,429 (578)	3.04 (0.14)	98.6 (0.57)	13 (10)	20 (14)
4	0 (0)	13,580 (2,623)	2.62 (0.06)	399 (31.8)	0	10 (14)
5	11 (8)	-	3.01 (0.31)	-	0	0
6	12 (0)	24,850 (1,072)	7.14 (0.66)	312 (22.7)	5 (7)	0
7	-	554,167 (41,908)	-	4,483 (11.3)	26 (4)	220 (28)
8	-	887,600 (352,422)	-	19,521 (3152)	50 (21)	745 (176)
9	-	-	-	-	20 (21)	285 (35)
10	-	31,675 (16,581)	-	379 (36.1)	3 (3)	115 (35)
11	-	14,733 (0)	-	449 (63.9)	8 (3)	75 (7)
12	-	-	-	-	1 (1)	35 (21)
13	-	-	-	-	3 (3)	25 (7)
14	-	-	-	-	1 (1)	35 (21)
15	-	-	-	-	0	30 (42)
16	-	28,178 (10,019)	-	335 (36.0)	0	15 (7)
17	-	-	-	-	0	10 (0)
LIS	-	8,244 (82)	-	422 (26.9)	-	-

899

900

901

902

903

904