

In press, *Proc. Nat'l. Acad. Sci. U.S.A.*

Rediscovery of the Nucleolinus, a Dynamic RNA-Rich Organelle Associated with the Nucleolus, Spindle, and Centrosomes

Mary Anne Alliegro,^a Jonathan J. Henry,^b and Mark C. Alliegro^a

^a Josephine Bay Paul Center, The Marine Biological Laboratory, Woods Hole,
Massachusetts 02543, and ^b Department of Cell & Developmental Biology, University
of Illinois, Urbana, Illinois 61801

Corresponding Author: Mark C. Alliegro
Josephine Bay Paul Center
Marine Biological Laboratory
7 MBL Street
Woods Hole, MA 02543
508-289-7419
malliegro@mbi.edu

Manuscript information: Research Report
Major category: Biological Sciences
Minor categories: Cell Biology
27 pages with 6 figures
2 Supplemental Figures
2 Supplemental videos

Abstract

The nucleolus is an RNA-rich compartment, closely apposed to or embedded within the nucleolus. Discovered over 150 years ago, fewer than two dozen articles have been published on the nucleolus, probably because complex histochemical stains are required for its visualization in the great majority of cells. The nucleolus has been reported in invertebrate oocytes, mammalian and amphibian epithelial cells, neurons, and several transformed cell lines. A prominent nucleolus, clearly visible with transmitted light microscopes at 10x magnification, is present in each oocyte of the surf clam, *Spisula solidissima*. We observed a consistent relationship between the nucleolus and the developing meiotic apparatus following *Spisula* oocyte activation. Through sonication and sucrose gradient fractionation of purified oocyte nuclei, we isolated nucleoli, extracted their RNA, and prepared an *in situ* riboprobe (NLI-1) that is associated specifically with the nucleolus, confirming its unique composition. Other *in situ* observations revealed a NLI-1 and nucleolar association with the developing spindle and centrosomes. Laser microsurgery that targeted the nucleolus resulted in failed meiotic cell division in parthenogenetically activated oocytes and failed mitosis in fertilized oocytes. Although the nucleolus may be a forgotten organelle, its demonstrated role in spindle formation suggests it deserves renewed attention.

Introduction

The nucleolus was described by Montgomery in eggs of the nudibranch mollusc, *Montagua*, in his 1898 monograph, “Comparative Cytological Studies, with Especial Regard to the Morphology of the Nucleolus” (1). As the title indicates, the study focuses on the appearance and behavior of the nucleolus. In it, Montgomery presents his own thoughts on the significance of the nucleolus as well as the observations of other investigators, including Louis Agassiz, over the previous 40 years (2-4). These range from the attachment of no morphological significance to the structure, to suggestions that the nucleolus is a microorganism enclosed in the nucleus and a direct morphological progenitor of the centrosome. Twenty-two years later, in his “Observations on an Intra-Nucleolar Body”, Carleton experimented with Cajal’s formol-silver nitrate technique to visualize the nucleolus in cat and frog intestinal epithelial cells (5). In some cases, he reported, the nucleolus was the only silver-impregnated structure within the entire nucleus. As of 90 years ago, then, the nucleolus was clearly distinguished from the surrounding or adjacent nucleolus, but only via histochemistry. Carleton, based on morphological observations of fixed, sectioned tissues, suggested that the nucleolus divided into two during prophase. Allen (6,7) subsequently proposed a “spindle forming role” for the nucleolus based on his observations of live *Spisula* oocytes, but no photomicrographs or other data were presented to directly support this hypothesis.

The most comprehensive analysis of the nucleolus appears in a series of reports by Love and colleagues from the late 1950s to the early 1970s (8-14). Love and Liles

developed a method for the differentiation of nucleoprotein complexes by deamination and staining with a combination of toluidine blue and ammonium molybdate (8). Using a variety of normal and tumorigenic mammalian cells, histochemical studies showed that the nucleolus stained metachromatically against the uniformly green background of the nucleolus. The authors observed that the metachromatic compartment increased significantly during prophase and then disappeared through anaphase and telophase. Love and Wildy later reported that the first detectable abnormality in Herpes virus-infected HeLa cells was an enlargement of nucleolus followed by their extrusion from the nucleolus into the nucleoplasm (9). In Ehrlich ascites and other cells, changes in nucleolus were correlated with perturbations in cell division (9). As did Allen (6,7), the authors speculated on a functional relationship between the nucleolus and cell division. Love (10) reported that changes in nucleolar morphology occur in aneuploid and virally-transformed cells, and Mironescu *et al.* (15) observed similar results in rat liver cells after administration of liver carcinogens. A few other reports discussing the nucleolus can be found in the literature, but for the most part the organelle is mentioned only in passing (16-18).

The nucleolus as described by Carleton, Love, Allen, and others has thus been identified in many cell types and species. In mammals alone, the list includes corneal and gut epithelium, sympathetic neurons, amnion, liver, kidney, pancreas, and more. Yet, our lack of knowledge regarding nucleolar composition [other than to say it contains RNA and no detectable DNA (13)] and function make it impossible to know what role(s) the organelle plays in cell physiology. An understanding of nucleolar dynamics,

bioactivity, and relationship to other nuclear bodies would likely have fundamental impact on cell biology. With this report we introduce the first specific molecular marker for the nucleolus, test the long-standing hypothesis that it functions in cell division, and aim to reinvigorate the study of this little known and enigmatic organelle.

Results

The nucleolus is spatially related to the forming spindle axis and pole

Mature *Spisula* oocytes are arrested in prophase I of meiosis. They contain a large tetraploid germinal vesicle (nucleus) and a prominent nucleolus (Fig. 1). Closely apposed to the nucleolus is a prominent nucleolus. There are no centrosomes in unactivated oocytes; neither centrosomes, spindle, nor microtubules are detectable by light, immunofluorescence, or electron microscopy until several minutes *after* activation (19). When meiosis is resumed after fertilization or parthenogenetic activation, the nuclear envelope breaks down and the nucleolus become morphologically indistinct. The nucleolus persists as a discrete structure for several minutes after dissipation of the nucleolus, and then, in agreement with observations on somatic cells (8), eventually becomes undetectable by phase contrast or differential interference contrast (DIC) microscopy before meiosis I spindle assembly (Figs. 2,3). During this process, centrosomes are thought to form *de novo* in the oocyte. To effect unequal cleavage for polar body formation, the spindle is positioned eccentrically in the oocyte cytoplasm. One spindle pole and centrosome come to lie in the oocyte interior, and the second spindle pole and centrosome migrate to the cell periphery, just beneath the plasma membrane. The point at which the polar bodies are ejected marks the first cleavage plane in *Spisula* and the animal pole of the embryo for many molluscs.

We tracked the nucleolus after oocyte activation by time lapse DIC microscopy and consistently observed that its last visible position in the cytoplasm coincided with the interior spindle pole (Fig. 2 and Supplemental Video 1). Because its movement after activation usually consisted of only localized oscillations, spindle position and the eventual site of polar body extrusion could be approximated by the position of the nucleolus before oocyte activation. However, in some instances the nucleolus was transported over striking distances in a brief time. Fig. 3 and Supplemental Video 2 show an example in which the nucleolus was translocated over 12 μ m in less than one minute. Again, it was the last visible position of the nucleolus and not its starting point that correlated with spindle location. Although results such as this are only correlative, they nevertheless provide a new, living view of nucleolar dynamics. Taken together with descriptions made by earlier investigators (6,7), they support the hypothesis that the nucleolus is spatially associated with the developing spindle and spindle pole.

The nucleolus is a unique cellular compartment containing centrosome- and spindle-associated RNAs

It has been reported that specific RNAs (cnRNAs) are associated with centrosomes (20-22). In *Spisula* oocytes, most of these are not detectable prior to oocyte activation, but several were found to be present in unactivated oocytes as a distinct hybridization patch associated with the nucleolus (21). As the nuclear envelope breaks down, the newly formed centrosomes are found to be embedded within, or closely apposed to, the

nucleolar patch. We now report on NLi-1 RNA, the first specific molecule of any kind to be isolated from and localized to the nucleolus *in situ*.

Nucleolar RNA was extracted from isolated nucleoli obtained in a two step process described in detail in the *Methods* section. Importantly, the first step is isolation of intact nuclei, which eliminates a large proportion of cytoplasmic contamination. At this stage of the preparation there is virtually no debris, and the only structures visible by DIC microscopy are the nuclear envelope, within which lie chromosomes and intact nucleoli (Supplemental Fig. 1A and B). The nucleolus dissipates early in the process, during initial cell lysis. After mechanical disruption of isolated nuclei, the highly dense nucleoli are isolated through several sucrose gradient steps and washes. The final preparation contains intact nucleoli and a small amount of debris which, by their appearance and comparison with intermediate steps in the preparation, are likely to be damaged nucleoli (Supplemental Fig. 1C).

NLi-1 is a >1kb contig assembled from isolated, cloned nucleolar RNA (Supplemental Fig. 2; Genbank accession HM004235). The contig comprised over 50% of all non-ribosomal clones sequenced. NLi-1 contains a potential 555nt open reading frame encoding a highly basic polypeptide of 184 amino acids with a predicted molecular mass of 20kDa and a pI of 10.5. It is unclear at present whether this gene is translated into protein in oocytes. No significant conserved domains or orthologues were found in nucleic acid or protein databases. The apparent lack of homology is not a function of *Spisula* divergence or underrepresentation of mollusk entries in sequence databases,

because BLAST analysis of other *Spisula* sequences readily reveals orthologues across a broad spectrum of phylogenetic groups (20). It is also important to highlight the absence of similarity to rRNA. There are a number of database entries for *Spisula* rRNAs as well as rRNAs derived from other mollusks. The lack of identity in our analysis, at even low stringency, indicates that NLi-1 is not rRNA.

In the unactivated oocyte, NLi-1 is exclusively localized to the nucleolus (Fig. 4A). There is an absence of label from all other cytoplasmic and nuclear structures including the nucleolus and chromosomes. This is in firm agreement with histochemical studies suggesting the nucleolus is a distinct cellular compartment (5,8). Following oocyte activation, as the nucleolus begins to dissipate and centrosomes are formed, NLi-1 is seen as a discrete, though slightly more diffuse patch in the cytoplasm (Fig. 4B). Centrosomes are often seen to be resting upon (or sometimes encased within) the NLi-1 hybridization patch, or in its immediate proximity. Slightly later, at 8-10 minutes postactivation, NLi-1 often appears as a narrow streak coinciding with the developing spindle axis. By 10 minutes postactivation, the developing spindle is situated within the expanded NLi-1 zone, giving the impression of an NLi-1 spindle matrix (Fig. 4C). The identification of a specific nucleolar RNA intimately associated with the developing spindle serves as a molecular correlate of the time-lapse images obtained from living oocytes, and summons the idea posed by early investigators that the nucleolus functions in spindle formation and/or cell division. We next performed a series of laser microsurgery experiments to directly test this long-standing hypothesis.

The nucleolus is required for normal cell division

The effect of laser microsurgery on the nucleolus was examined by DIC microscopy to determine if this disrupts meiosis. We use the term "targeted" and not "ablated" because, although clear cell division defects (described below) and a visible welt were produced, gross nucleolar structure was not destroyed in these experiments. Cells were labeled with bis-benzamide and observed by fluorescence microscopy to track the fate of chromatin. Oocytes in which the nucleolus was targeted, approximately one nucleolus-diameter ($\sim 5\mu\text{m}$) away from the experimental target site, were used as controls. In addition, non-lasered oocytes were used as controls to visualize normal meiotic progression, and also as event timers (i.e., cells were fixed at the time non-lasered cells exhibited polar bodies). Details of these experiments are provided in the *Methods* section. The results show a significant disparity between experimental and control groups (Fig. 5A-C). Only 19% of oocytes in which the nucleolus was targeted successfully completed meiosis. Interestingly, a single small membrane bleb the size of a normal polar body was sometimes observed in these oocytes, but this "polar body" was devoid of chromatin. We do not know if these blebs were the result of aborted cytokineses, or if they were an experimental artifact. As will be noted below, the spindle in these oocytes was often grossly disorganized, so we could not determine with confidence any spatial relationship or polarity in relation to these membrane blebs. In nucleolus-targeted (control) cells, a chromatin-containing polar body of normal appearance was evident in 91% of cases.

In subsequent experiments, oocytes were fixed at different times after laser microsurgery and KCl-activation to determine the morphological basis of the cell division defect. Samples were stained with anti- γ tubulin antibody to assess the presence of centrosomes, anti- α tubulin to examine spindle morphology, and bis-benzamide to visualize chromosome congression and segregation. The effects of microsurgery were multiple and manifested primarily at later time points (Fig. 5D-K). Approximately two-thirds of cases involved gross malformation of the meiotic figure. Chromatin often appeared as a disorganized cluster associated with a tangle of microtubules. In other cases the meiotic spindle appeared somewhat normal in shape, but was diminished in size and microtubule content as judged by anti-tubulin immunofluorescence. In a few cases progression through meiosis appeared to be arrested in metaphase at times when control cells had advanced from late anaphase through telophase and cytokinesis. It is possible that this range of results is due to slight variability in laser targeting, but we consider it more likely they are due to real, pleiotropic effects of nucleolar perturbation. At earlier time points the differences between control and experimental groups were subtle and difficult to substantiate, however supernumerary and sometimes smaller, less intensely stained centrosomes were observed.

Unperturbed, parthenogenetically activated oocytes complete meiosis but do not enter mitosis. By fertilizing oocytes rather than activating with KCl, we could address the questions of whether meiotic spindle organization could be restored by sperm-derived elements, and whether laser microsurgery on the nucleolus had effects upon mitosis as well as meiosis. The results were clear in addressing both questions. First, we found that

oocytes laser targeted on the nucleolus and then fertilized were able to proceed through meiosis I and II (Fig. 6). This occurred in 8/8 cases. However after completing meiosis, 0/8 of these zygotes formed mitotic centrosomes or spindles. In controls (nucleolar-targeted), 8/9 oocytes contained a normal complement of centrosomes and formed a mitotic spindle. In total, the results reported here indicate that the nucleolus is required for progression through meiosis in activated oocytes. When fertilized with sperm, however, elements of the nucleolus are not required for meiosis, but are required for entry into the subsequent rounds of mitosis.

Discussion

Based on morphological observations of nucleolar dynamics, investigators spanning the years between the 1870s and 1970s proposed a role for this structure in cell division. Some reported centrosome-like behavior of the nucleolus, including self-divisional properties, dysregulation related to aneuploidy, and proximity to the spindle pole. The more recent discovery that some RNAs later found in centrosomes localize to the nucleolus *before* centrosomes assemble suggested that it may even serve as a cytological precursor of the division center (21), a view first expounded over a century ago (23). While it seems clear that the nucleolus provides a component or components critical for the normal function of the centrosomes and spindle apparatus, its role is likely more complicated than the straightforward progenitor-progeny relationship proposed by Lavdowsky (4). Otherwise, laser targeting of the nucleolus should have resulted in the absence of mitotic, as well as meiotic centrosomes.

Also attesting to the complexity of nucleolar function is the observation that laser microsurgery had no effect on meiosis after fertilization, but still affected the subsequent mitosis (Fig. 6). This is intriguing, and may have bearing on the differences between *de novo* and replicative modes of centrosome formation, at least in the zygote. Several models may explain this result, but one that is consistent with our observations and others on the early development of *Spisula* is that some components required for biogenesis of meiotic centrosomes are present in the cytoplasm of unactivated oocytes, and others are located in the nucleolus. Thus, when we lasered the nucleolus and activated oocytes

with KCl, γ -tubulin-containing foci (centrosomes) formed, but apparently did not function to form a spindle. Fertilization, however, allowed sperm-derived elements to either replace maternally-derived centrosomes wholesale, or substitute for nucleolar elements that normally complement the cytoplasmic components. Since laser-treated, fertilized oocytes complete meiosis but cannot subsequently undergo mitosis, either (a) sperm possess enough of these putative complementing elements for only one round of cell division, (b) sperm-derived complementing elements still require nucleolar input to function through multiple cell cycles, or (c) they are suppressed by maternal factors normally present until the end of meiosis. This latter possibility is supported by the fact that, following meiosis, the original (meiotic) centrosomes in *Spisula* zygotes are suppressed and the job of organizing subsequent cell divisions is transferred to the replication-competent sperm-derived MTOC (24).

Whether the nucleolus should be considered an organelle in its own right must be evaluated using a number of criteria, including structural and functional differentiation from the nucleolus and its ubiquity in cells and across species. It may be closely apposed to, or in some cases embedded within the nucleolus, but our observations lead us to agree with Montgomery (1), Carleton (5), and others (8-15) that it is certainly a unique cellular compartment. Supporting this are behavioral, molecular, and functional data: (a) In meiosis as well as during cell fractionation, the nucleolus persists as a distinct morphological entity after complete dissolution of the nucleolus transpires; (b) the nucleolus contains molecules (minimally, a subset of cnRNAs (21) and NLi-1 RNA) that are completely excluded from the nucleolus; and (c) based on our laser irradiation

experiments in which the nucleolus or nucleolus was targeted, the nucleolus plays a role in cell division that is experimentally separable from nucleolar function (damaging the nucleolus had profound effects on the cell division apparatus, damaging the nucleolus did not). This is an important point in light of studies indicating the nucleolus, beyond its well known metabolic functions, plays a direct role in centrosome biogenesis and cell cycle regulation (25,26). It is possible that cell division defects induced in the work of Gaulden and Perry (25) and Ugrinova et al. (26) are actually attributable to the nucleolus. The inability to detect its presence in these and most other cell models has certainly left it unaccounted for in the interpretation of past results. Regardless of any role the nucleolus may play in cell cycle regulation, the functional relationship between the nucleolus and cell division, speculated upon for more than a century, is presently confirmed. To know the precise physiological role of the nucleolus will require a more thorough knowledge of the nature and fate of its macromolecular components, an area we are only beginning to explore.

Methods and Materials

Gravid *Spisula solidissima* were obtained from the Marine Resources Center at the Marine Biological Laboratory, Woods Hole, MA. Gametes were collected by dissection. Oocytes were rinsed several times in 0.45 μ m filtered sea water and activated with 0.14 volumes of 0.5M KCl. All brightfield and immunofluorescence imaging was performed with a Nikon 90i microscope equipped with DIC, epifluorescence, and a Photometrics CoolSNAP *fx* digital camera. Anti α - and γ -tubulin immunofluorescence (for microtubules and centrosomes, respectively) were performed as described in an earlier report (20).

Isolation and cloning of NLi-1 RNA

Nucleolini are isolated in a two-phase process. The first phase is isolation of oocyte nuclei based on methods described by Maul and Avdalovic (27). Starting with 8 -10mL of packed, unactivated oocytes distributed between four 50mL conical centrifuge tubes, the cells are resuspended by addition of 20mL glycerol phosphate buffer (1M glycerol, 20mM NaPO₄, pH 8.0), incubated at room temperature for three minutes, and sedimented at 1,750xg for 15 seconds. The oocytes are resuspended to 40mL in ice cold lysis buffer (0.5M hexylene glycol, 0.75mM MgCl₂, 1mM HEPES, pH 7.5) and incubated at room temperature with occasional mixing by inversion. They are monitored until complete lysis of cells, leaving only intact nuclei. Unlike the nucleolinus, the nucleolus is unstable in lysis buffer and rapidly dissolves. Nuclei are stabilized by addition of 8mL ice cold

40% sucrose with gentle mixing, and the tubes are placed on ice. The lysate is underlain with 1.5mL 10% sucrose buffer (equal parts lysis buffer and 20% sucrose) and centrifuged for eight minutes at 780xg, 4°C, in a swinging bucket rotor. The resultant soft, white pellet is collected by pipetting into a microcentrifuge tube, frozen in liquid nitrogen and stored at -80°C for later use.

Nucleolini are then harvested from isolated nuclei. Eight frozen nuclear pellets are thawed on ice, resuspended with 1mL 10% sucrose buffer each, pooled, divided into two 15mL conical centrifuge tubes, and disrupted by sonication on ice with three bursts of 5 seconds each at 11W with a microprobe sonicator. The samples are underlain with 2mL 40% sucrose buffer (40% sucrose in lysis buffer) and centrifuged for 10 minutes at 500xg. The resulting pellet is washed twice by resuspension in ice cold PBS and sedimentation in a microcentrifuge for one minute at maximum speed. RNA was extracted using Qiagen RNeasy reagents, reverse transcribed using universal primers described previously,²⁰ and blunt-cloned into Invitrogen PCR-script plasmid. Over 3,500 clones have been sequenced to date, approximately 10% of which are non-ribosomal. Characterization of isolated nucleolini and complete analysis of all potential nucleolar RNAs and proteins is ongoing.

In situ hybridization

Methods for *in situ* hybridization and immunofluorescence were similar to those described in an earlier report (20) with an important modification. In the present study,

oocytes were fixed for one hour at room temperature in 4% formaldehyde containing 0.6% Brij 58, 5mM EGTA, and 1mM MgCl₂, buffered with 100mM PIPES, pH 6.8 (28). They were then rinsed once briefly in PBS with 0.1% Tween-20 (PBST), equilibrated in PBST for 30 minutes, dehydrated in a series of ethanols, and stored at -20°C in 70% ethanol. Previously, the nucleolus and nucleolus were not visible by DIC at the end of the hybridization protocol. The new fixation method preserved nucleolar and nucleolar structure so that RNA label could be correlated with DIC-visible structures. Samples were hybridized with digoxigenin-labeled nucleolus-1 RNA probe for 3 days at 60°C. Alkaline phosphatase reaction product was visualized after development at 4°C with gentle rocking.

Laser microsurgery

For laser microsurgery, 8-10 oocytes were transferred into a 3µL drop of 0.45µm-filtered sea water (buffered with 10mM TRIS, pH 8.0) on a 1.0mm thick glass slide with 0.2mm spacers and a 1.2mm thick cover. A 150µs pulsed 1480nm laser set at 300mW was directed into a 5µm spot through the 40x objective using a Hamilton Thorne (Beverly, MA) XYClone laser system mounted on a Nikon 90i microscope. A visible lightbeam sight guide of the same diameter was used to effect precise targeting. These conditions were selected in preliminary experiments using a graded series of pulse parameters, and determined mild enough so that no overt physical damage to the oocyte was observed (as confirmed by controls). Controls were irradiated using these same parameters, but at one nucleolus-diameter away from the experimental target, striking

within the nucleolus. Following irradiation, oocytes were quickly transferred to sea water containing 70mM KCl to activate meiosis (or sperm to fertilize). A similar number of untreated oocytes were transferred to a separate KCl-containing well to serve as non-lasered controls and event timers. Activated oocytes were fixed at various times and labeled with bis-benzamide to visualize chromatin segregation and/or prepared for immunohistochemistry as described above. Because these cells are nearly spherical, it was necessary to rotate oocytes during observation to avoid mistaking an end-on view of a normal spindle for a malformed cluster of chromatin and microtubules. We reiterate here for clarity that at the time of laser irradiation (unactivated oocyte stage), no centrosomes or spindle are present. These elements arise only *after* fertilization or parthenogenetic activation.

Acknowledgements

The authors thank Dr. Robert Palazzo for reading and commenting on the manuscript, Melissa Clarkson for suggesting a fixation protocol to preserve nucleolini through the *in situ* hybridization protocol, and Zachary Swartz for his contribution to a method for their isolation. The laser system used in this study was generously lent by Hamilton Thorne Biosciences with assistance from Lily Reed. This work was supported by grants from the NIH (GM088503) and NSF (MCB0843092) to MCA. J.H. acknowledges support from the National Science Foundation (IOB 05-16799) and the Directors of the MBL Embryology Course for providing laboratory space and facilities.

References

1. Montgomery TH (1898) Comparative cytological studies, with especial regard to the morphology of the nucleolus. *J Morph* 15:265.
2. Agassiz L (1857) *Contributions to the Natural History of the United States of America*. First Monograph. Part IIL Embryology of the Turtle. Boston.
3. Haeckel E (1874) *Anthropogenie oder Entwicklungsgeschichte des Menschen*. Leipzig.
4. Lavdowsky M (1894) Von der Entstehung der chromatischen und achromatischen Substanzen in den tierischen und pflanzlichen *Zellen Merkel und Bonnefs Anat* Hefte.
5. Carleton HM (1920) Observations on an intra-nucleolar body in columnar epithelium cells of the intestine. *Q J Microsc Sci* s2-64:329-341.
6. Allen RD (1951) The role of the nucleolus in spindle formation. *Biol Bull* 101:214.
7. Allen RD (1953) Fertilization and artificial activation in the egg of the surf calm, *Spisula solidissima*. *Biol Bull* 105:213-239.
8. Love R, Liles RH (1959) Differentiation of nucleoproteins by inactivation of protein-bound amino groups and staining with toluidine blue and ammonium molybdate. *J Histochem Cytochem* 7:164-181.
9. Love R, Wildy,P (1963) Cytochemical studies of the nucleoproteins of HeLa cells infected with herpes virus. *J Cell Biol* 17:237-254.
10. Love R (1966) Anisonucleolinosi in mammalian cell cultures. *Natl Cancer Inst Monographs* 23:167-180.

11. Love R, Bharadwaj TP (1959) Two types of ribonucleoprotein in the nucleolus of mammalian cells. *Nature* 183:1453-1454.
12. Love R (1965) Differences in the internal structure of nucleoli in diploid and non-diploid transformed or neoplastic cells *in vitro*. *Exptl Cell Res* 40:188-192.
13. Love R, Walsh RJ (1968) Nucleolar morphology in normal diploid, neoplastic, and aneuploid cells *in vitro*. *Cancer Res* 30:990-997.
14. Love R, Soriano RZ (1971) Correlation of nucleolini with fine structural constituents of cultured normal and neoplastic cells. *Cancer Res* 31:1030-1037.
15. Mironescue S, Encut I, Mironescu K, Liciu F (1968) Nucleolar behavior in regenerating liver of rats receiving intra-abdominal injections of azo dyes and thioacetamide. *J Natl Cancer Inst* 40:917-933.
16. Gates RR (1942) Some observations regarding the nucleolus and cytoplasm in living marine eggs. *Biol Bull* 82:47-51.
17. Kopac MJ, Mateyko GM (1959) Malignant nucleoli: cytological studies and perspectives. *Ann New York Acad Sci* 73:237-282.
18. Dan K, Ito S (1984) Studies on unequal cleavage in molluscs: I. Nuclear behavior and anchorage of a spindle pole to cortex as revealed by isolation technique. *Dev Growth Differ* 26:249-262.
19. Palazzo RE, Vaisberg E, Cole RW, Rieder CL (1992) Centriole duplication in lysates of *Spisula solidissima* oocytes. *Science* 256:219-221.
20. Alliegro MC, Alliegro MA, Palazzo RE (2006) Centrosome-associated RNA in surf clam oocytes. *Proc Natl Acad Sci USA* 103:9034-9038.

21. Alliegro MC, Alliegro MA (2008) Centrosomal RNA correlates with intron-poor nuclear genes in *Spisula* oocytes. *Proc Natl Acad Sci USA* 105:6993-6997.
22. Le'cuyer E *et al.* (2007) Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. *Cell* 131:174–187.
23. Quoted from Montgomery (1): “Lavdovsky ('94) concludes that the nucleolus is the centrosome in the process of formation, but he failed to observe the steps by which the body develops into a centrosome.”
24. Wu X, Palazzo RE (1999) Differential regulation of maternal vs. paternal centrosomes. *Proc Natl Acad Sci USA* 96:1397-1402.
25. Gaulden ME, Perry RP (1958) Influence of the nucleolus on mitosis as revealed by ultraviolet microbeam irradiation. *Proc Natl Acad Sci USA* 44:553-559.
26. Ugrinova I *et al.* (2007) Inactivation of nucleolin leads to nucleolar disruption, cell cycle arrest and defects in centrosome duplication. *BMC Mol Biol* 8:66.
27. Maul GG , Avdalovic N (1980) Nuclear envelope proteins from *Spisula solidissima* germinal vesicles. *Exp Cell Res* 130:229-240.
28. Lee KG (2002) Rapid visualization of microtubules in blood cells and other cell types in marine model organisms. *Biol Bull* 203:204-206.

Figure Legends

Figure 1. DIC image of the nucleolus within an unactivated *Spisula* oocyte. A large tetraploid nucleus (germinal vesicle; GV) is present in the unactivated oocyte, within which lie a prominent, spherical nucleolus (arrow) and nucleolus (arrowhead). The nuclear envelope begins to disintegrate within the first two minutes of activation and is indistinguishable by 10-12 minutes post-activation. The nucleolus becomes morphologically indistinct within approximately five minutes post-activation, and the nucleolus persists for several minutes beyond that time (see Figs. 2 and 3, for examples). Size bar = 15 μ m.

Figure 2. Spatial relationship of the nucleolus to the developing spindle and pole in a living oocyte. A, at approximately two minutes post-activation, the nucleus (germinal vesicle; GV), already appearing ragged around the edges, the nucleolus, and nucleolus are all clearly visible. B, by seven minutes the nucleolus is no longer visible, but the nucleolus still persists as a well-defined structure. C, D, the disintegration of the nucleolus (arrowhead) is traced through to its last visible position as cytoplasm continues to infiltrate the central area of the oocyte. During this time the nucleolus oscillates slightly in all three planes, but remains in the same vicinity despite significant cytoplasmic rotation (see Supplemental Video 1). E, the emerging polar body (arrow) together with the site mark for the nucleolus (arrowhead) delineate the axis and poles of the first meiotic spindle. Elapsed time in minutes and seconds is indicated in each frame.

Figure 3. Transport of the nucleolus through the cytoplasm. In this case the nucleolus (black arrowhead) was transported from its original position (marked by the black dot in B-E) and, along with the emerging polar body (white arrowhead in E), delineate the long axis of the spindle. During the 50 second time span shown in panels B-D, the nucleolus moved over 12 μ m ($> 0.24\mu\text{m}/\text{second}$) through the cytoplasm. This series of time frames is taken from Supplemental Video 2.

Figure 4. *In situ* localization of NLi-1 RNA. In panel A (unactivated oocyte), the nucleolus labels intensely with the NLi-1 RNA probe (purple). The borders of the nucleolus are difficult to distinguish after the three-day hybridization regime, and are therefore highlighted with black arrows in this figure. Chromosomes are visualized with bis-benzamide (smaller, pale-blue structures, two of which are labeled with white arrows). Observations on non-hybridized cells labeled with bis-benzamide and hybridized cells not labeled with bis-benzamide confirm that there is no overlap between these two compartments. That is, nucleoli contain no detectable DNA [as reported earlier by Love and Walsh (13)] and chromosomes contain no detectable NLi-1 signal. As the nucleolus begins to “dissolve” at 6 minutes post-activation (B), a strong but slightly more diffuse hybridization patch remains in the oocyte. γ -tubulin immunofluorescence in the same cell (green) reveals the newly-formed centrosomes (arrowheads) closely apposed to the nucleolus. By 12 minutes post-activation (C) the entire cell division apparatus is associated with an NLi-1 “matrix”. Spindle microtubules are present at this time, but cannot be visualized in these oocytes due to lability of α -tubulin antigenicity following

the hybridization regimen (including 3 days at 60°C). A-C are overlays of black and white DIC images with color brightfield images (alkaline phosphatase staining) and immunofluorescence. "Sense" probes were used for negative controls in all *in situ* hybridizations, and were completely devoid of label.

Figure 5. Laser ablation of the nucleolus results in failed meiotic cell division. Panel A is an example of successful meiosis I in a control (nucleolar-targeted) oocyte fixed at 31 minutes post-activation and labeled with bis-benzamide. Polar body chromatin is clearly observed on the cell periphery (arrow), segregated from the remaining chromatin of the 2° oocyte. Panel B shows an oocyte targeted at the nucleolus and fixed 31 minutes post-activation. In this case chromatin is scattered in the oocyte cytoplasm, but in other examples (see panel I, below) remained clustered. Panel C shows the percentage of cells determined to successfully complete meiosis I (using images such as these) following laser ablation of the nucleolus (Nli) compared to those receiving no laser treatment (Non-), or control cells in which the nucleolus was targeted. D-K show oocytes that have been double-labeled with bis-benzamide (blue) and anti- α -tubulin (green) to reveal spindle configuration in control (D-G) and nucleolus-targeted (H-K) oocytes. Arrows in D (DIC) and E (bis-benzamide) indicate the newly-formed first polar body. F (anti-tubulin) and G (overlay of E and F) show a normal metaphase II configuration. In the nucleolar-targeted cell shown in H-K, clustered chromatin is associated with a disorganized knot of microtubules. No polar body is evident.

Figure 6. Mitosis failure in fertilized zygotes following laser microsurgery on the

nucleolinus. Oocytes in which nucleolini were laser-targeted and then fertilized (vs. KCl-activated, as in the previous set of experiments) were able to complete meiosis; sperm components were apparently able to substitute for damaged maternal-nucleolar components. However, pronuclear fusion and mitotic spindle formation that normally follow polar body ejection were inhibited. A-E is a control zygote, F-J is a zygote targeted on the nucleolinus. A, DIC image showing polar bodies (arrowheads) and a full cleavage furrow (arrows), indicating completion of meiosis and first mitosis. Cleavage does not occur in the nucleolar-targeted zygote (F). B and G, bis-benzamide staining reveals polar bodies (arrowheads) in both control and experimental, although the latter are not apparent in the DIC image in F. Daughter cell nuclei (arrows in B) can be seen in control zygotes, small, unfused male and female pronuclei appear to persist in experimental (arrows in G). Anti- α -tubulin staining (green) reveals the developing asters and spindle of second cleavage in controls (C), but not in experimental. Likewise, centrosomes are readily apparent in controls labeled with anti- γ -tubulin (D, arrowheads), but are absent in nucleolar-targeted zygotes (I). E and J, composite fluorescence images of panels B-D and G-I, respectively.




Figure 1
Alliegro et al.




Figure 2
Alliegro *et al.*




Figure 3
Alliegro *et al.*




Figure 4
Alliegro *et al.*




Figure 5
Alliegro *et al.*





Figure 6
Alliegro *et al.*



Supplemental Figure 1. Isolation of nucleolini from unactivated *Spisula* oocytes. The first step is isolation of intact nuclei (A,B). The nucleolus is labile and no longer visible at this step in the isolation. Only nuclear envelopes, condensed prophase chromosomes (arrows), and nucleolini (arrowheads) are detectable by DIC. Fine granular material within nuclei can sometimes be seen. Nucleolini are remarkably dense and easily pellet through a 50% sucrose cushion at low *g*-force after disruption of GVs by sonication, leaving all other debris behind in the supernatant. C, isolated nucleolini.

1 AACTNAGGGGGAAAAATGTTGAAAGCATGTAGGCAGCTCGAGGGCTTAGATGATCTCTTG
N X G G K M L K A C R Q L E G L D D L L
61 TTTGCCGGGTCTGTAGGCGGTGAGTGAAATCAGTGGTTTTTAACTGCGTGTAGCTCATG
F A G S V G G E W K S V V F N C V * L M
121 TTAACGTCTTATGATAGTCATGCAATCGGCGTAGTACTCCACGGAGGTCAGGCTCCCTTA
L T S Y D S H A I G V V L H G G Q A P L
181 CCGAGAGGTACGTACTTTTCGCACAGGGCATCAGGGTCCGGCCTCCGAAACACCTTGGCA
P R G T Y F S H R A S G S G L R N T L A
241 GTGGCAACCAAGTCGCCGAGAGGGCAATCGACGCCCGCCGAACCGTACACATGGTTAGAT
V A T K S P R G Q S T P A E P Y T W L D
301 GGCGGGTGTCTGCCGTGGTTGCGTAGATCCATAGGATATACTCTATGTAACAGTAGTGTC
G G C L P W L R R S I G Y T L C N S S V
361 GTGCCTATGGACTCGCTTCCGCACTCGGTGCAATATTCGCCGCGTAAGCGGCTCAGGTCC
V P M D S L P H S V Q Y S P R K R L R S
421 CGAGAAGCATTGTCCGCCGCTCCTTCCGCTAGTAACACAGCCGGACAGGTCCTGCCGATT
R E A L S A A P S A S N T A G Q V L R I
481 CGACACCGCTTTAAGCGGATGTTGCAAGGCGAACGAACGAACGNTCACTCGATCGATATT
R H R F K R M F E G E R T N X H S I D I
541 ACACACCACCCATGATATCATCTATGGCGTGTGTGCTGTAAACAACCATTAATTAATNAG
T H H P *
601 TGGGCTCCCATCGAGGAAGTAGTCCACTAAGACACGTCGGCCCCCTGCTCCCCGGCCTAGT

661 AGCCTCAGTCGTAGAGAAGTGTGTCGAAANGTTTGGGTGGTAGCAGCGGGAAGGCAGTC
721 AGCAATGCCGTAAGCCGACTGGTTTTRTGTCTAGTAGTGGTGGAAACGCCGCGAAAAGCA
781 TGGCCAGGAGGTCTGTTCCACCAGCAGCGTAGTAGCGTCCACCACAGCGCGAAAAACTGT
841 TGGCCGTTTCTAGTAGGCGTCCGTTGGCTAGCTAGCATGCCCATGTAGACGAGTAGCGTACC
901 GACCGTGACGCGAACATACATCGGGTAGGCCGGCTGGCTGCGGGCAAACCAAAAATCTTA
961 CGGAGGGAGAGCGCGAGTCCAGTACCTCCAGGCAAGTGGCGG

Supplemental Figure 2. NL-1 RNA partial sequence, including translation of largest potential open reading frame.