

# **CHEMISTRY**

---

## **A EUROPEAN JOURNAL**

---

### **Supporting Information**

© Copyright Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, 2010

### **Using a Threading-Followed-by-Swelling Approach to Synthesize [2]Rotaxanes**


**Jia-Ling Ko,<sup>[a]</sup> Shau-Hua Ueng,<sup>[a]</sup> Ching-Wei Chiu,<sup>[a]</sup> Chien-Chen Lai,<sup>[b]</sup> Yi-Hung Liu,<sup>[a]</sup>  
Shie-Ming Peng,<sup>[a]</sup> and Sheng-Hsien Chiu<sup>\*[a]</sup>**

chem\_201000074\_sm\_miscellaneous\_information.pdf

	Page
<sup>1</sup> H and <sup>13</sup> C NMR spectra of all new compounds	S2–S57
<sup>1</sup> H NMR spectra of an equimolar mixture of macrocycle <b>12</b> and thread <b>1</b> -H·PF <sub>6</sub>	S58
The <sup>1</sup> H NMR spectra of the swelling synthesis of rotaxane <b>14</b> -H·PF <sub>6</sub>	S59
The <sup>1</sup> H NMR spectra of the swelling synthesis of rotaxane <b>28</b> -H·PF <sub>6</sub>	S60
The <sup>1</sup> H NMR spectra of the swelling synthesis of rotaxane <b>32</b> -H·PF <sub>6</sub>	S61
The <sup>1</sup> H NMR spectra of rotaxane <b>29</b> -H·PF <sub>6</sub> heated at 353 K in CD <sub>3</sub> SOCD <sub>3</sub>	S62
The <sup>1</sup> H NMR spectra of rotaxane <b>31</b> -H·PF <sub>6</sub> heated at 353 K in CD <sub>3</sub> SOCD <sub>3</sub>	S63
The <sup>1</sup> H NMR spectra of rotaxane <b>33</b> -H·PF <sub>6</sub> heated at 353 K in CD <sub>3</sub> SOCD <sub>3</sub>	S64
The <sup>1</sup> H NMR spectra of rotaxane <b>15</b> -H·PF <sub>6</sub> heated at 353 K in CD <sub>3</sub> SOCD <sub>3</sub>	S65



<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul



**STANDARD 1H OBSERVE**

Pulse Sequence: s2pul



<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul



199.062

172.744

77.319  
77.000  
76.688


62.345

36.334  
27.022  
22.857  
18.813  
14.077  
13.226




STANDARD 1H OBSERVE

Pulse Sequence: s2pul





<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE

Pulse Sequence: s2pu1




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




7





STANDARD 1H OBSERVE

Pulse Sequence: s2pul





<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




**STANDARD 1H OBSERVE**

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




## STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE




Pulse Sequence: s2pul



STANDARD 1H OBSERVE

Pulse Sequence: s2pul






13C (ppm)

100


150

50




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE

Pulse Sequence: s2pu1






<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




400MHz H<sup>1</sup> CD<sub>3</sub>NO<sub>2</sub>




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




400MHz H<sup>1</sup> CD<sub>3</sub>NO<sub>2</sub>




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul





400MHz H<sup>1</sup> CD<sub>3</sub>Cl<sub>3</sub>




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




400MHz H<sup>1</sup> CD<sub>3</sub>Cl<sub>3</sub>




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




**STANDARD 1H OBSERVE**

Pulse Sequence: s2pu1




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




**STANDARD 1H OBSERVE**

Pulse Sequence: s2pul






<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul



STANDARD 1H OBSERVE

Pulse Sequence: s2pu1



<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul

200.428



131.880

126.400

119.468


77.319  
77.000  
76.688

30.222  
28.315  
26.817  
24.940  
21.648  
17.164  
15.529




STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul



STANDARD 1H OBSERVE

Pulse Sequence: s2pul



<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul

202.457



137.840  
129.927  
121.030

114.307

77.319  
77.000  
76.688




43.806

29.143  
27.296  
22.515  
21.740  
17.300  
14.883




STANDARD 1H OBSERVE

Pulse Sequence: s2pul






<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pu1




STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE

Pulse Sequence: s2pul




<sup>13</sup>C OBSERVE

Pulse Sequence: s2pul




STANDARD 1H OBSERVE


Pulse Sequence: s2pul




**<sup>1</sup>H NMR spectra (400 MHz, CD<sub>3</sub>CN, 298 K) of a) macrocycle 12, b) an equimolar mixture of 12 and 1-H·PF<sub>6</sub> (10mM), and c) 1-H·PF<sub>6</sub>.**




**The swelling synthesis of [2]rotaxane from 12 and 1-H·PF<sub>6</sub> (10 mM) in CD<sub>3</sub>NO<sub>2</sub> at 318 K**




**The swelling synthesis of [2]rotaxane from DB24C8 and 20-H·PF<sub>6</sub> (70 mM) in CDCl<sub>3</sub> at 318 K**




**The swelling synthesis of [2]rotaxane from 13 and 20-H·PF<sub>6</sub> (70 mM) in CD<sub>3</sub>NO<sub>2</sub> at 318 K**




**<sup>1</sup>H NMR spectra (400 MHz, CD<sub>3</sub>SOCD<sub>3</sub>, 298 K) of a) DB24C8, b) [2]rotaxane [29-H][PF<sub>6</sub>] and c) heating solution b) at 343 K for 2 h.**



**<sup>1</sup>H NMR spectra (400 MHz, CD<sub>3</sub>SOCD<sub>3</sub>, 298 K) of a) macrocycle 12, b) [2]rotaxane [31-H][PF<sub>6</sub>] and c) heating solution b) at 343 K for 2 h.**



**<sup>1</sup>H NMR spectra (400 MHz, CD<sub>3</sub>SOCD<sub>3</sub>, 298 K) of a) macrocycle 13, b) [2]rotaxane [33-H][PF<sub>6</sub>] and c) heating solution b) at 343 K for 2 h.**



**<sup>1</sup>H NMR spectra (400 MHz, CD<sub>3</sub>SOCD<sub>3</sub>, 298 K) of a) macrocycle 12, b) [2]rotaxane [15-H][PF<sub>6</sub>] and c) heating solution b) at 343 K for 2 h.**

