

Role of calcineurin in *Porphyromonas gingivalis*-induced myocardial cell hypertrophy and apoptosis

Shin-Da Lee¹, Wei-Wen Kuo², Ding-Yu Lin³, Ter-Hsin Chen⁴, Wu-Hsien Kuo⁵, Hsi-Hsien Hsu⁶, Jian-Zhi Chen⁷, Jer-Yuh Liu³, Yu-Lan Yeh⁸ & Chih-Yang Huang^{3,*}

¹School of Physical Therapy, Chung-Shan Medical University, Taichung, Taiwan; ²Department of Biological Science and Technology, China Medical College, Taichung, Taiwan; ³Institute of Biochemistry, Chung-Shan Medical University, No. 110, Section 1, Chien Kuo N. Rd., Taichung 40203, Taiwan, R.O.C.; ⁴Graduate Institute of Veterinary Public Health, National Chung-Hsing University, Taichung, Taiwan; ⁵Division of Gastroenterology, Department of Internal Medicine, Armed-Force, Taichung General Hospital, Taichung, Taiwan; ⁶Division of colorectal surgery, Mackay Memorial Hospital, Taipei, Taiwan; ⁷Division of Cardiology, Show-Chwan Memorial Hospital, Changhua, Taiwan; ⁸Department of Pathology, Changhua Christian Hospital, Changhua, Taiwan

Received 17 July 2005; accepted 15 November 2005
© 2005 National Science Council, Taipei

Key words: periodontitis, *Porphyromonas gingivalis*, myocardial cell, apoptotic, cell death, calcineurin signal pathways

Summary

Background and objective: Periodontal pathogen *Porphyromonas gingivalis* (*P. gingivalis*) increased cardiomyocyte hypertrophy and apoptosis whereas *Actinobacillus actinomycetemcomitans* and *Prevotella intermedia* had no effects. The purpose of this study is to clarify the role of calcineurin signaling pathway in *P. gingivalis*-induced H9c2 myocardial cell hypertrophy and apoptosis. **Methods:** DNA fragmentation, nuclear condensation, cellular morphology, calcineurin protein, Bcl2-associated death promoter (Bad) and nuclear factor of activated T cell (NFAT)-3 protein products in cultured H9c2 myocardial cell were measured by agarose gel electrophoresis, DAPI, immunofluorescence, and Western blotting following *P. gingivalis* and/or pre-administration of CsA (calcineurin inhibitors cyclosporin A). **Results:** *P. gingivalis* not only increased calcineurin protein, NFAT-3 protein products and cellular hypertrophy, but also increased DNA fragmentation, nuclear condensation and Bad protein products in H9c2 cells. The increased cellular sizes, DNA fragmentation, nuclear condensation, and Bad of H9c2 cells treated with *P. gingivalis* were all significantly reduced after pre-administration of CsA. **Conclusion:** Our findings suggest that the activity of calcineurin signal pathway may be initiated by *P. gingivalis* and further lead to cell hypertrophy and death in culture H9c2 myocardial cells.

Introduction

The accumulation of epidemiologic, *in vitro*, animal and clinical evidence suggests that periodontal infection may be a contributing risk factor for

heart diseases [1]. Periodontitis pathogens may disseminate through the blood to infect the vascular endothelium and contribute to the occurrence of atherosclerosis and risk of myocardial ischemia and infarction [2]. *Porphyromonas* (*P.*) *gingivalis*, a major pathogen of chronic adult periodontitis, exhibits several potential virulence properties which could play a role in the development of cardiovascular disease as mediators of low

*To whom correspondence should be addressed. Phone: +886-4-24730022 ext. 11682; Fax: +886-4-23248195; E-mail: chuang1@csmu.edu.tw

density lipoprotein oxidation, foam cell formation, and rupture of atherosclerotic plaque [3]. The balance between cell death and cell survival is a tightly controlled process, especially in terminally differentiated cells, such as the cardiomyocyte [4]. Therefore, it becomes a very important issue whether *P. gingivalis* will induce cardiomyocyte apoptosis and hypertrophy. In our previous studies, *P. gingivalis* increased cellular sizes, apoptosis, and the activated apoptotic caspase-3, caspase-8, and caspase-9 proteins of cardiomyocyte H9c2 cells whereas *Actinobacillus actinomycescomitans* and *Prevotella intermedia* had no effects. [5]. Besides, *P. gingivalis* was demonstrated to adhere and to invade aortic and heart endothelial cells [6]. However, the role of calcineurin myocardial hypertrophy and apoptosis induced by *P. gingivalis* is still unclear.

Cardiac hypertrophy may initially be a beneficial adaptation that normalizes wall stress, but the prolonged cardiac hypertrophy is a leading cause of sudden death and heart failure [7]. Apoptosis had been reported to contribute to the loss of myocardial cells in cardiomyopathy, and is recognized as a predictor of adverse outcomes in subjects with cardiac diseases or heart failure [8]. Cardiac myocyte apoptosis may occur in a variety of diseases, such as hypertension, myocardial infarction, endocrine disorders, toxicants, and bacterial endocarditis [9, 10]. Apoptosis is characterized by DNA-fragmentation, nuclear condensation, and shifting the balance between anti-apoptotic and pro-apoptotic proteins towards the latter. [11]. Members of the Bcl2 family are important regulators of programmed cell death and individual members that can suppress (e.g. Bcl2) or promote (e.g. Bad) apoptosis. The Bcl2-associated death promoter (Bad), a pro-apoptotic member of the Bcl-2 family, modulate death signaling and lead to release of cytochrome *c* from the mitochondria into the cytoplasm, which results in DNA fragmentation and cell death by activation of caspase 9 and 3 [12].

Calcineurin is a calmodulin-dependent, serine/threonine-specific phosphatase function within a signal transduction pathway that regulates gene expression and biological responses in many important cell types [13]. Calcineurin was considered as an essential mediator of cardiac hypertrophy and apoptosis, and in most studies, calcineurin inhibitors attenuated cardiac hypertrophy *in vivo* and also attenuated apoptosis of cardiac myocytes

[14]. Calcineurin plays a critical role in the apoptosis of cardiac myocytes through dephosphorylating Bad, thus enhancing Bad heterodimerization with Bcl-xL or other Bcl-2 family and promoting apoptosis. [14, 15]. Calcineurin transmits signals to the nucleus through the dephosphorylation and translocation of nuclear factor of activated T cell (NFAT) transcription factors. [13, 16]. Calcineurin and its direct downstream transcriptional effector NFAT-3 potentially play a regulatory role for maladaptive hypertrophy and heart failure [16]. In human heart failure, the calcineurin signaling pathway was activated not only by an increased activity of calcineurin but also by the shift from phosphorylated NFAT-3 to the nucleus dephosphorylated NFAT-3 [17]. Once activated, calcineurin will dephosphorylate NFAT-3 and Bad, allowing translocation of these proteins to the nucleus and mitochondria, respectively, both of which may potentially lead to myocardial pathologic hypertrophy and apoptosis [15]. However, it is unclear whether calcineurin signaling pathways are involved in *P. gingivalis*-associated myocardial cell hypertrophy and apoptosis.

The purpose of the current study is, first to determine the effect of periodontal pathogen *P. gingivalis* on calcineurin, Bad, and NFAT-3 protein levels, second to clarify whether *P. gingivalis*-induced H9c2 myocardial cell hypertrophy is mediated via calcineurin signaling pathway, and third to evaluate whether calcineurin/Bad signaling pathways is related to *P. gingivalis*-induced H9c2 myocardial cell apoptosis. We hypothesized that calcineurin activity may be related to *Porphyromonas gingivalis*-induced myocardial cell hypertrophy and apoptosis.

Methods

Porphyromonas gingivalis preparation

Porphyromonas (P.) gingivalis obtained from American Type Culture Collection (ATCC 33277) was maintained on brain heart infusion (BHI) enriched with Vitamin K3 (menadione, 0.5 mg/l) and hemin (5 mg/l) in an atmosphere of 5% CO₂, 10% H₂, and 85% N₂ at 37 °C. Turbidimetric method was used to evaluate bacterial cultures with 2 McFarland standard (6×10⁸ colony forming units (CFUs)), the upper aqueous

layer was collected following centrifugation at 10,000 g, filtered through 0.22 μm filter, and stored at -80°C . The upper aqueous layers of *P. gingivalis* bacterial mediums were diluted to 10-fold and 100-fold.

Cell culture

H9c2 cardiac cells were obtained from American Type Culture Collection (ATCC CRL-1446) and were cultured in Dulbecco's modified essential medium supplemented with 10% fetal bovine serum, 2 mM glutamine, 100 units/ml penicillin, 100 $\mu\text{g}/\text{ml}$ streptomycin, and 1 mM pyruvate in humidified air (CO_2 5%) at 37°C . H9c2 cells were cultured in serum-free medium for 12 h, and then treated with control (serum-free medium with minimal essential medium) or CsA (0.5 $\mu\text{M}/\text{ml}$; calcineurin inhibitors cyclosporin A) for 1 h and post-treated with *P. gingivalis* bacterial 10-fold, or 100-fold diluted bacterial mediums. The incubation was continued for another 0.5, 1, 2 or 24 h, and then the cells were harvested and extracted for the analysis.

Actin-immunofluorescence (rhodamine-phalloidin)

H9c2 cells were fixed in fresh 3.7% paraformaldehyde and permeabilized with 2% Triton X-100 for 10 min, and blocked with PBS buffer with 1% bovine serum albumin (PBS) for 15 min. Actin filaments were stained by rhodamine-phalloidin. Cells were examined and photographed using a Zeiss Axioskop and a confocal microscope. The cell sizes were magnified 100 times and analyzed by digital image analysis, using Image-Pro Plus software.

DNA fragmentation

H9c2 cells were lysed in 50 μl of lysis buffer (50 mM Tris-HCl (pH 7.4), 20 mM EDTA, 1% IGEPAL-630) followed by incubation with 1% SDS and 5 $\mu\text{g}/\mu\text{l}$ RNase (Roche Molecular Biochemicals, Mannheim, Germany) for 2 h at 56°C and 2.5 $\text{g}/\mu\text{l}$ proteinase K (Roche) for 2 h at 37°C , and only fragmented DNA was extracted. DNA was ethanol-precipitated and finally resuspended in distilled water. The fragmented DNA was electrophoretically fractionated on 1.5% agarose gel and stained with ethidium bromide.

Nuclear condensation stained with DAPI

H9c2 cells grown on 6 mm plate were washed twice with PBS and fixed for 30 min in 4% buffered paraformaldehyde. The cells were then stained with 1 $\mu\text{g}/\text{ml}$ 4',6-diamidino-2-phenylindole dihydrochloride (DAPI, Roche) for 30 min. The stained cells were examined with fluorescence microscopy. The nuclei condensation was expressed by percentage (%) of cell death (nuclei condensation) and expressed by relative percentage (%) of *P. gingivalis* induced cell death (nuclei condensation).

Western blot analysis

Cultured H9c2 cells were scraped and washed once with phosphate-buffered saline. Cell suspension was then spun down, and cell pellets were lysed for 30 min in lysis buffer (50 mM Tris (pH 7.5), 0.5 M NaCl, 1.0 mM EDTA (pH 7.5), 10% glycerol, 1 mM BME, 1% IGEPAL-630 and proteinase inhibitor cocktail tablet (Roche)) and spun down 12,000 rpm for 10 min. Then, the supernatants were applied to new eppendorf tube for Western blot analysis. Proteins from H9c2 cell line was then separated in 12% gradient SDS-PAGE and transferred to nitrocellulose membranes. Nonspecific protein binding was blocked in blocking buffer (5% milk, 20 mM Tris-HCl, pH 7.6, 150 mM NaCl, and 0.1% Tween 20) and blotted with specific antibodies of Bcl2-associated death promoter (Bad), and NFAT-3 (Santa Cruz Biotechnology, Santa Cruz, CA, USA), and α -tubulin (Neo Markers, Fremont, CA, USA) as indicated for each experiment in the blocking buffer at 4°C overnight.

Experimental protocols

Protocol 1

To quickly assess whether calcineurin signaling pathways were regulated by the periodontitis pathogen *P. gingivalis* in myocardial cell, calcineurin total protein and its direct downstream transcriptional effector, NFAT-3, and Bad proteins were measured by Western blotting in H9c2 cells cultured in serum-free medium (control) for 4 h or serum-free medium plus 10-fold diluted ($10\times P. g$) for 0.5 h (more details in Figure 1).




Figure 1. (a) The calcineurin, Bad, NFAT-3 protein products were measured by Western Blotting in H9c2 cells cultured in serum-free medium (control) for 4 h and then treated with or without 10-fold diluted *Porphyromonas gingivalis* (10× P. g) for 0.5 h. An internal control of α -tubulin was measured. (b) Bars represent the relative densitometry analysis of calcineurin/ α -tubulin, bad/ α - tubulin and NFAT-3/ α -tubulin to the control levels and indicate mean values \pm SD. ($n=6$) * $p < 0.05$, ** $p < 0.01$, significant different between *P. gingivalis* and control.

Protocol 2

To assess whether calcineurin signaling pathways mediate *P. gingivalis*-induced myocardial cell hypertrophy, the cellular sizes were measured by actin-immunofluorescence in H9c2 myocardial cells treated with either serum-free medium (control), or 10-fold diluted *P. gingivalis* for 24 h with or without calcineurin inhibitors. Beside *P. gingivalis* induced calcineurin and NFAT-3 in H9c2 cells were observed for 24 h (more details in Figure 2).

Protocol 3

To evaluate whether calcineurin is related to *P. gingivalis*-induced Bcl2-associated death promoter (Bad), the protein products of calcineurin and Bad were measured by Western blotting analysis in myocardial H9c2 cells with pre-administration of serum-free medium (control) or calcineurin inhibitors (CsA) for 1 h and with post-administration of bacterial medium containing 10-fold diluted *P. gingivalis* medium for 0.5, 1, and 2 h (more details in Figure 3).

Protocol 4

To assess whether calcineurin signaling pathways mediate *P. gingivalis* induced myocardial cell apoptosis, DNA fragmentations were measured twice by agarose gel electrophoresis in H9c2 myocardial cells with pre-administration of serum-free medium (control) or calcineurin inhibitors (CsA) for 1 h and with post-administration of bacterial medium containing 10-fold diluted *P. gingivalis* medium for 24 h (more details in Figure 4).

Protocol 5

To assess whether calcineurin signaling pathways mediate *P. gingivalis* induced myocardial cell apoptosis, nuclei condensation was measured by DAPI (4,6-diamidino-2-phenylindole) staining in myocardial H9c2 cells with pre-administration of serum-free medium (control) or calcineurin inhibitors (CsA) for 1 h and with post-administration of bacterial medium containing 10-fold or 100-fold diluted *P. gingivalis* medium for 24 h (more details in Figure 5).

Statistics

The nuclei condensation and cellular size of myocardial H9c2 cell were compared among groups treated with either control, 10-fold diluted *P. gingivalis*, and pre-treated calcineurin inhibitors using one-way analysis of variance (ANOVA) with pre-planned contrast comparison with control group and 10-fold diluted *P. gingivalis* group. In all cases, a difference at $p < 0.05$ was considered statistically significant. All data presented in the bar figures are means \pm SD ($n=6$).

Results

Total calcineurin, activated Bad and NFAT-3 proteins were induced by *P. gingivalis*

Western blot analysis revealed that total calcineurin, activated Bad and NFAT-3 proteins were gradually increased in H9c2 cells after administration of 10-fold diluted *P. gingivalis* (10× P. g) for 0.5 h. compared with cultured in serum-free medium (control) (Figure 1).




Figure 2. (a) Calcineurin and NFAT-3 protein products induced 10-fold diluted *Porphyromonas gingivalis* (10 \times P. g) for 24 h. (b) Bars represent the relative densitometry analysis of calcineurin/ α -tubulin and NFAT-3/ α -tubulin to the control levels and indicate mean values \pm SD. ($n=6$) * p <0.05, significant different between *P. gingivalis* and control. (c) Cellular morphology was measured by actin-immunofluorescence in H9c2 myocardial cells treated with either control (Con, left panel), 10-fold diluted *Porphyromonas gingivalis* medium only (10 \times P. g, middle panel), or pre-treated calcineurin inhibitors cyclosporin A (CsA, 10 μ M/ml) for 1 h, and then post-treated 10-fold diluted *P. gingivalis* medium for 24 h (10 \times P. g + CsA, right panel) (d) Bars represent the relative cellular sizes to control and indicate mean values \pm SD. ($n=6$) ## p <0.01, significant different of cellular sizes between *P. gingivalis* and control, ** p <0.01, significant different of cellular areas between *P. gingivalis* and pre-treated signaling pathway inhibitors plus *P. gingivalis*.

Calcineurin mediate *P. gingivalis*-induced myocardial cell hypertrophy

P. gingivalis-induced total calcineurin and NFAT-3 were not only shown in early 0.5 h but was also observed for 1, 2, 4, 8, 12, and 24 h (Figure 2a,2b). Actin-immunofluorescence revealed that the increased cellular sizes by 85% were ob-

served in myocardial H9c2 cells treated with 10-fold diluted *P. gingivalis* medium for 24 h compared with serum-free control (Figure 2c). The *P. gingivalis* (10-fold diluted)-induced myocardial cell hypertrophy were significantly reduced by pre-administration of CsA (calcineurin inhibitor, 49% relative to cellular size of 10 \times P. g) (Figure 2d).





Figure 3. (a) The represented protein products of mitochondrial Bcl2-associated death promoter (Bad) were measured by Western Blotting in H9c2 cells cultured in serum-free medium (control) for 4 h or treated with 10-fold diluted *P. gingivalis* ($10\times P. g$) for 0.5 h, 1 h, and 2 h with or without pre-administration of calcineurin inhibitors cyclosporin A (CsA, 0.5 μ M/ml) before 1 h. An internal control of α -tubulin was measured. (b) Bars represent the relative densitometry analysis of Bad/ α -tubulin to the control level and indicate mean values \pm SD. ($n=6$) # $p < 0.05$, ## $p < 0.01$, significant different between *P. gingivalis* and control, * $p < 0.05$, ** $p < 0.01$, significant different between *P. gingivalis* and pre-treated CsA plus *P. gingivalis*.



Marker	+				
Serum free	+				
10xP.g		+	+	+	+
CsA				+	+

Figure 4. Represented DNA fragmentations were measured by agarose gel electrophoresis in H9c2 myocardial cells. H9c2 cells were treated with either serum free for 4 h (control), pre-treated with or without calcineurin inhibitors cyclosporin A (CsA, 0.5 μ M/ml) for 1 h, and then treated with bacterial medium containing 10-fold diluted *Porphyromonas gingivalis* ($10\times P. g$) for 24 h.

Calcineurin mediate *P. gingivalis* induced Bad measured by Western blotting

Western blot analysis revealed that mitochondrial Bcl2-associated death promoter (Bad) protein products were increased after administration of 10-fold diluted *P. gingivalis* ($10\times P. g$) for 0.5, 1, and 2 h (Figure 3). The increased activated Bad protein induced by 10-fold diluted *P. gingivalis* in H9c2 cell were significantly reduced after pre-administration of CsA (calcineurin inhibitor) (Figure 3).

Calcineurin mediate *P. gingivalis* -induced myocardial cell apoptosis measured by DNA fragmentation

DNA ladder formation revealed that H9c2 myocardial cell underwent DNA fragmentation and apoptosis on exposure to either bacterial medium containing 10-fold diluted *P. gingivalis* ($10\times P. g$) for 24 h compared to the serum free control (Figure 4). The 10-fold diluted *P. gingivalis* induced DNA fragmentation was attenuated by CsA (calcineurin inhibitor).




Figure 5. (a) Represented nuclei condensation expressed by % of cell death (nuclei condensation) was measured by DAPI (4,6-diamidino-2-phenylindole) (1 $\mu\text{g/ml}$) staining in H9c2 myocardial cells treated with either control (Con, upper left panel), 10-fold diluted *Porphyromonas gingivalis* medium (10X P. g, upper middle panel), pre-treated calcineurin inhibitors cyclosporin A (CsA; 0.5 $\mu\text{M/ml}$) plus post-treated 10-fold diluted bacterial medium (CsA + 10 \times P. g, upper right panel), 100-fold diluted *P. gingivalis* medium (100 \times P. g, lower left panel), pre-treated CsA plus post-treated 100-fold diluted *P. gingivalis* bacterial medium (CsA + 100 \times P. g, lower right panel) (b) Graphed by % of cell death. # $p < 0.05$ # $p < 0.01$, significant different of % cell death (nuclei condensation) between *P. gingivalis* and control, ** $p < 0.01$, significant different of % cell death (nuclei condensation) between *P. gingivalis* and pre-treated calcineurin inhibitors plus *P. gingivalis*.

Calcineurin mediate P. gingivalis-induced myocardial cell apoptosis measured by condensed nuclei with DAPI staining

The condensation of nuclei stained with DAPI, indicating cell apoptosis, was significantly increased after administration of 10-fold diluted *P. gingivalis* for 24 h (10 \times P. g, $p < 0.01$) and 100-fold diluted *P. gingivalis* for 24 h (100 \times P. g, $p < 0.05$) compared with control (Con) (Figure 5). The 100-fold diluted *P. gingivalis*-induced condensed nuclei in H9c2 cell show no differences between with or without CsA (calcineurin inhibitor), whereas the increased

condensed nuclei induced by 10-fold diluted *P. gingivalis* were significantly ($p < 0.01$) reduced after pre-administration of CsA.

Discussion

Major findings

Our main findings can be summarized as follows: (1) Total calcineurin, activated Bad and NFAT-3 protein in H9c2 cells were upregulated by *P. gingivalis*. (2) *P. gingivalis* increased H9c2 cellular sizes, DNA

fragmentation, and nuclear condensation in H9c2 myocardial cells, which imply *P. gingivalis* induced myocardial cellular hypertrophy and apoptosis. (3) The increases in cellular sizes, DNA fragmentation, nuclear condensation, and pro-apoptotic Bad protein of H9c2 cells treated with *P. gingivalis* were all reduced after pre-administration of CsA. In sum, *P. gingivalis*-induced myocardial hypertrophy and apoptosis appear to be mediated by calcineurin/NFAT or calcineurin/Bad signaling pathway. These findings provide a potential link between *P. gingivalis* and calcineurin activity. After integrating previous theories and our current findings into one, we hypothesize that that myocardial cell hypertrophy and myocardial cell apoptosis will be induced by *P. gingivalis* via calcineurin/NFAT-3 pathway and calcineurin/Bad pathway, respectively (shown in Figure 6).

Experimental limitation

P. gingivalis, a gram-negative anaerobe, is involved in the pathogenesis of periodontal disease and plays important roles in eliciting cellular responses in various ways [18]. *P. gingivalis* possesses a variety of virulence factors including gingipains, fimbrillin peptides, capsule polysaccharides, lipopolysaccharides, hemagglutinating and hemolysing activities, toxic products of metabolism, outer membrane vesicles, and other enzymes [19]. Because of the widespread virulence factors, *P. gingivalis* can produce myocardial cell apoptosis via various virulence factors or metabolic byproducts. Therefore, we have to add a note of caution that any effect noted in the present investigation

cannot be isolated to any specific virulence factor or any metabolic byproduct. Nevertheless, the goal of the present study was to clarify the role of calcineurin signaling pathway in *P. gingivalis*-induced myocardial cell apoptosis. Clearly, additional experiments using a reductionist approach, such as investigating gingipains alone or lipopolysaccharides alone, will be required in order to identify which specific virulence factors or metabolic byproducts directly cause myocardial cell apoptosis.

P. gingivalis and cardiovascular system

Some studies indicated that *P. gingivalis* did affect the cardiovascular system, such as the coronary arteries [3, 20]. *P. gingivalis* was suggested to involve in atherosclerotic lesion formation [20] and was found in the aorta of some cardiovascular patients who had undergone open-heart surgery [21], the latter of which suggests that periodontal pathogen *P. gingivalis* did invade the cardiovascular system and might involve cardiovascular diseases [21]. Most cross-sectional, prospective epidemiological studies, animal model studies, and cellular studies have demonstrated that periodontal disease (*P. gingivalis*) may be associated with endothelial cell, atherosclerosis formation, and coronary heart disease [1–3, 6, 20, 21]. Myocardial ischemia and infarction are often preceded by acute thromboembolic events induced by periodontal disease, such as *P. gingivalis* [2]. In our previous studies, *P. gingivalis* increased apoptosis and the activated apoptotic caspase-3, caspase-8, and caspase-9 proteins of cardiomyocyte H9c2 cells as well as *P. gingivalis*-induced apoptotic activity of H9c2 cells were involved in p38, ERK, PI3K, and calcineurin pathway, but might play a protective role via JNK pathway [5]. In the current study, we observed the two major calcineurin-mediated pathways in *P. gingivalis*-induced myocardial cell hypertrophy and apoptosis.

P. gingivalis and calcineurin

Calcineurin, a Ca^{2+} -calmodulin-dependent protein phosphatase is one of the links between Ca^{2+} signals and regulation of gene transcription in cardiac muscle. Calcineurin has been implicated in cardiovascular disease and is activated by numerous stimuli, such as hypertension, ischemic/




Figure 6. After integrating previous theories and our current findings into one, we hypothesized that myocardial cell hypertrophy and myocardial cell apoptosis may be induced by *porphyromonas gingivalis* via calcineurin/NFAT-3 pathway and calcineurin/Bad pathway, respectively.

re-perfusion, hypoxia, pressure-overload stress, increase in cardiac pacing frequency [14, 22–24] and periodontitis pathogens *P. gingivalis* found in the current study. The current findings indicated that *P. gingivalis*-induced H9c2 cell hypertrophy and apoptosis was mediated by calcineurin signaling pathways. Herein, our findings imply that calcineurin signaling pathway in myocardial cell can be activated by one more cellular stimulus, *P. gingivalis*, i.e. periodontal infection. To our knowledge, no other study looking into the direct or indirect interaction between *P. gingivalis* and calcineurin activity in all types of cells or systems. This finding might provide a potential link between *P. gingivalis* and calcineurin signaling pathway. Therefore, more studies in the interaction between *P. gingivalis* and calcineurin signaling pathway in cardiovascular systems are needed.

Role of calcineurin in P. gingivalis-induced myocardial cell hypertrophy

The heart is a dynamic organ capable of adapting its size and architecture in response to various stressors or bacterial infection [25, 26]. Our findings showed that myocardial cell sizes, total calcineurin, and activated NFAT-3 were all increased in H9c2 cells by 10-fold diluted *P. gingivalis* challenge and the increased myocardial cell sizes-induced by *P. gingivalis* were reduced by calcineurin inhibitor, suggesting that myocardial cell hypertrophy will induced by *P. gingivalis* via calcineurin/NFAT-3 pathway. The current findings provide a potential link between *P. gingivalis* and calcineurin/NFAT-3 or calcineurin/Bad pathway in myocardial cell hypertrophy or apoptosis. Calcineurin/NFAT was shown to induce up-regulation of the Fas death apoptotic pathway in neuronal cells [27]. Therefore, there is a limitation in our study to clarify the relative role of NFAT or Bad in hypertrophy or apoptosis. Further studies are required to confirm the role of *P. gingivalis* on cardiac hypertrophy or heart failure via calcineurin/NFAT-3 pathway.

Calcineurin role in P. gingivalis-induced myocardial cell apoptosis

Apoptosis is characterized by DNA-fragmentation, nuclear condensation, and shifting the balance between anti-apoptotic and pro-apoptotic

proteins towards the latter [14, 15]. The mitochondrial Bcl2 family plays an important role in cardiac apoptosis. Shifting the balance between anti-apoptotic and pro-apoptotic Bcl2 family proteins towards the latter (e.g. increased Bad proteins) will cause releasing cytochrome *c* into cytosol and activate caspase 9, which will activate caspase 3 that is responsible for cleavage action and DNA fragmentation. Our findings showed that periodontal pathogen *P. gingivalis* increased DNA fragmentation, nuclear condensation, and activated Bad protein in H9c2 cells. All increases were attenuated by calcineurin inhibitor, which suggest that *P. gingivalis* induce H9c2 myocardial cell apoptosis via calcineurin signaling pathway and *P. gingivalis*-induced myocardial cell apoptosis may be involve in Bad related apoptotic pathways.

Calcineurin plays a critical role in the pro-apoptosis of cardiac myocytes through dephosphorylating pro-apoptotic protein Bad to promote apoptosis. Our findings showed that the increased Bad proteins induced by 0.5–2 h *P. gingivalis* challenges were attenuated by calcineurin inhibitors, which can be explained by the supposition that calcineurin inhibitors may inhibit dephosphorylation of Bad and decrease activated Bad protein level. Therefore, we can generally conclude that calcineurin activity may be initiated by *P. gingivalis* and may further activate Bad related apoptotic pathways to lead cell death in culture H9c2 myocardial cells. However, 100-fold diluted *P. gingivalis* induced myocardial cell nuclear condensation were not inhibited by CsA in the current study. Therefore, we have to make a cautious note that more diluted *P. gingivalis*-induced myocardial cell apoptosis may not be mediated via calcineurin signaling pathway.

Further hypotheses for clinical application

Our findings can provide a future hypothesis that myocardial cell hypertrophy and apoptosis may potentially gradually occur in patients with periodontitis infected by *P. gingivalis*, which might be mediated by calcineurin signaling pathways. In order to prevent potential cardiac diseases, such as pathologic cardiac hypertrophy or heart failure, in periodontitis patients infected by *P. gingivalis*, the infected patients have been suggested that they should ideally maintain excellent oral hygiene

and suppress *P. gingivalis* infection as much as possible, and our findings further proposed that calcineurin signaling pathways inhibitors may be a good candidate for developing a future therapeutic approach to prevent *P. gingivalis*-induced cardiac hypertrophy and/or heart failure. Of course, further clinical studies are needed to prove or disprove our hypotheses for clinical application.

References

- Genco R., Offenbacher S. and Beck J., Periodontal disease and cardiovascular disease: epidemiology and possible mechanisms. *J. Am. Dent. Assoc.* 133(Suppl): 14S–22S, 2002.
- Herzberg M.C. and Weyer M.W., Dental plaque, platelets, and cardiovascular diseases. *Ann. Periodontol.* 3: 151–160, 1998.
- Kuramitsu H.K., Qi M., Kang I.C. and Chen W., Role for periodontal bacteria in cardiovascular diseases. *Ann. Periodontol.* 6: 41–47, 2001.
- Fortuno M.A., Lopez N., Gonzalez A. and Diez J., Involvement of cardiomyocyte survival-apoptosis balance in hypertensive cardiac remodeling. *Expert Rev. Cardiovasc. Ther.* 1: 293–307, 2003.
- Lee S.D., Wu C.C., Kuo W.W., Lin J.A., Huang C.M., Lu M.C., Chen L.M., Hsu H.H., Wang C.K. and Chang S.H., Porphyromonas gingivalis-related cardiac cell apoptosis was majorly co-activated by p38 and ERK pathways. *J. Periodontol. Res.* 2006 (in press).
- Deshpande R.G., Khan M.B. and Genco C.A., Invasion of aortic and heart endothelial cells by Porphyromonas gingivalis. *Infect. Immun.* 66: 5337–5343, 1998.
- Takano H., Zou Y., Akazawa H., Toko H., Mizukami M., Hasegawa H., Asakawa M., Nagai T. and Komuro I., Inhibitory molecules in signal transduction pathways of cardiac hypertrophy. *Hypertens. Res.* 25: 491–498, 2002.
- Narula J., Pandey P., Arbustini E., Haider N., Narula N., Kolodgie F.D., Dal Bello B., Semigran M.J., Bielsa-Masdeu A., Dec G.W., Israels S., Ballester M., Virmani R., Saxena S. and Kharbanda S., Apoptosis in heart failure: release of cytochrome c from mitochondria and activation of caspase-3 in human cardiomyopathy. *Proc. Natl. Acad. Sci. USA* 96: 8144–8149, 1999.
- Gill C., Mestral R. and Samali A., Losing heart: the role of apoptosis in heart disease—a novel therapeutic target? *Faseb J.* 16: 135–146, 2002.
- Dhalla N.S., Ziegelhoffer A., Singal P.K., Panagia V. and Dhillon K.S., Subcellular changes during cardiac hypertrophy and heart failure due to bacterial endocarditis. *Basic Res. Cardiol.* 75: 81–91, 1980.
- Saraste A., Morphologic criteria and detection of apoptosis. *Herz* 24: 189–195, 1999.
- Bishopric N.H., Andrecka P., Slepak T. and Webster K.A., Molecular mechanisms of apoptosis in the cardiac myocyte. *Curr. Opin. Pharmacol.* 1: 141–150, 2001.
- Schulz R.A. and Yutzey K.E., Calcineurin signaling and NFAT activation in cardiovascular and skeletal muscle development. *Dev. Biol.* 266: 1–16, 2004.
- Saito S., Hiroi Y., Zou Y., Aikawa R., Toko H., Shibasaki F., Yazaki Y., Nagai R. and Komuro I., beta-Adrenergic pathway induces apoptosis through calcineurin activation in cardiac myocytes. *J. Biol. Chem.* 275: 34528–34533, 2000.
- Wang H.G., Pathan N., Ethell I.M., Krajewski S., Yamaguchi Y., Shibasaki F., McKeon F., Bobo T., Franke T.F. and Reed J.C., Ca²⁺-induced apoptosis through calcineurin dephosphorylation of BAD. *Science* 284: 339–343, 1999.
- Wilkins B.J., Dai Y.S., Bueno O.F., Parsons S.A., Xu J., Plank D.M., Jones F., Kimball T.R. and Molkenkin J.D., Calcineurin/NFAT coupling participates in pathological, but not physiological, cardiac hypertrophy. *Circ. Res.* 94: 110–118, 2004.
- Diedrichs H., Chi M., Boelck B., Mehlhorn U. and Schwinger R.H., Increased regulatory activity of the calcineurin/NFAT pathway in human heart failure. *Eur. J. Heart Fail.* 6: 3–9, 2004.
- Curtis M.A., Kuramitsu H.K., Lantz M., Macrina F.L., Nakayama K., Potempa J., Reynolds E.C. and Aduse-Opoku J., Molecular genetics and nomenclature of proteases of Porphyromonas gingivalis. *J. Periodontol. Res.* 34: 464–472, 1999.
- Kantarci A. and Van Dyke T.E., Neutrophil-mediated host response to Porphyromonas gingivalis. *J. Int. Acad. Periodontol.* 4: 119–125, 2002.
- Kuramitsu H.K., Kang I.C. and Qi M., Interactions of Porphyromonas gingivalis with host cells: implications for cardiovascular diseases. *J. Periodontol.* 74: 85–89, 2003.
- Stelzel M., Conrads G., Pankuweit S., Maisch B., Vogt S., Moosdorf R. and Flores-de-Jacoby L., Detection of Porphyromonas gingivalis DNA in aortic tissue by PCR. *J. Periodontol.* 73: 868–870, 2002.
- Nagata K., Somura F., Obata K., Odashima M., Izawa H., Ichihara S., Nagasaka T., Iwase M., Yamada Y., Nakashima N. and Yokota M., AT1 receptor blockade reduces cardiac calcineurin activity in hypertensive rats. *Hypertension* 40: 168–174, 2002.
- Lakshmikuttyamma A., Selvakumar P., Kakkar R., Kanthan R., Wang R. and Sharma R.K., Activation of calcineurin expression in ischemia-reperfused rat heart and in human ischemic myocardium. *J. Cell. Biochem.* 90: 987–997, 2003.
- Tavi P., Pikkariainen S., Ronkainen J., Niemela P., Ilves M., Weckstrom M., Vuolteenaho O., Bruton J., Westerblad H. and Ruskoaho H., Pacing-induced calcineurin activation controls cardiac Ca²⁺ signalling and gene expression. *J. Physiol.* 554: 309–320, 2004.
- Tomlinson C.W. and Dhalla N.S., Alteration in calcium metabolism in cardiac hypertrophy and failure caused by bacterial infection. *Recent Adv. Stud. Cardiac Struct. Metab.* 12: 191–198, 1976.
- Ravingerova T., Barancik M. and Strniskova M., Mitogen-activated protein kinases: a new therapeutic target in cardiac pathology. *Mol. Cell Biochem.* 247: 127–138, 2003.
- Jayanthi S., Deng X., Ladenheim B., McCoy M.T., Cluster A., Cai N.S. and Cadet J.L., Calcineurin/NFAT-induced up-regulation of the Fas ligand/Fas death pathway is involved in methamphetamine-induced neuronal apoptosis. *Proc. Natl. Acad. Sci. USA* 102: 868–873, 2005.