

ADVANCED DEVELOPMENTS IN RADAR

4-7 December 1995, Washington, D.C. Continuing Engineering Education Program. George Washington University, Academic Center, Room T-308, 801 22nd Street NW, Washington, DC 20052. Tel: (800) 424-9773 (US); (800) 535-4567 (Canada); Fax: (202) 872-0645; e-mail: ceepinfo@seas.gwu.edu.

MODERN TARGET SIGNATURE MEASUREMENT, EXPLOITATION AND COHERENT SIGNAL PROCESSING

8-10 January 1996, Washington, DC.
27-29 March 1996, Washington, DC. Johns Hopkins University, Whiting School of Engineering, Organizational Effectiveness Institute, 14515 Barkwood Drive, Rockville, MD 20853. Tel: (800) 683-7267, or (301) 871-9608; Fax: (301) 871-4942; E-mail: info.oci@apl.jhu.edu.

COMPUTATIONAL TECHNIQUES FOR EM APPLICATIONS

1-4 April 1996, Cambridge, England (Lecturers: J. B. Davies, R. D. Graglia, R. Mittra, D. R. Wilton). J.B. Davies, Dept. of Electrical Engineering, University College London, Torrington Place, London WC1E 7JE, England. Tel: (+44/0) 44-171-388-7849; e-mail: b.davies@eleceng.ucl.ac.uk; or Dr. Raj Mittra, Dr. Raj Mittra, Tel: (217) 333-1200; Fax: (217) 333-8986; e-mail: rmittra@decwa.ecc.uiuc.edu.

AP-S VIDEO COURSES

QUASI-OPTICAL SYSTEM DESIGN FOR MILLIMETER WAVELENGTHS.

by Paul F. Goldsmith, one 1-hour tape plus booklet, Order Number HVO 0211-3, \$59.95 plus \$5.00 shipping & handling.

PATTERN SYNTHESIS FOR ANTENNA ARRAYS, by Robert S. Elliott, one 1.5-hour tape plus notes, Order Number EVO 4036, \$25.00.

ADAPTIVE PROCESSING ANTENNA SYSTEMS, by William F. Gabriel, two tapes (3 hours) plus notes, Order Number EVO 4051, \$40.00.

LATERAL ELECTROMAGNETIC WAVES, by Ronald W.P. King, two tapes (4 hours), Order Number EVO 4010, \$30.00.

LAWRENCE LIVERMORE NATIONAL LABORATORY MOVIE PROJECT, by R.W Cole and E.K. Miller (eds.), two tapes (2.5 hours) plus notes, Order Number EVO 4044, \$50.00.

ANTENNA FAR-FIELD PATTERNS FROM NEAR-FIELD MEASUREMENTS, by Yahya Rahmat-Samii, one 1.25-hour tape, Order Number EVO 4028, \$15.00.

A PRIMER ON THE REFLECTOR ANTENNA ART, by Willard V. T. Rusch, two tapes (3 hours), Order Number EVO 4085, \$30.00.

These VHS tapes can be ordered by order number, author, and title from the IEEE Service Center, Attn: Carolyn Yankowski, Course/Video-tape Coordinator, Educational Dept., 445 Hoes Lane, PO Box 1331, Piscataway, NJ 08855-1331. Call (908) 562-5493 to verify current shipping charges and prices before ordering. Checks payable to the IEEE should be enclosed with your order. Add sales tax for shipments to CA, DC, NJ, and NY.

Database Of "In Vivo" Measurements for Quantitative Microwave Imaging and Reconstruction Algorithms Available

[Editor's note: The availability of the following experimental data, appropriate for use in testing quantitative imaging and reconstruction algorithms, was announced at the joint AP-S/URSI-AB Special Session on "Image Reconstruction from Real Data," organized at the Newport Beach Symposium by Bob McGahan, Ralph Kleinman, and Mike Fiddy. The data set used as a common test for the algorithms discussed—the so-called "Ipswich Data"—may still be available from Bob McGahan (see his address, etc. in the Chapter News column in this issue). Such data sets, particularly when based on carefully made and documented measurements, are very valuable for testing both direct- and inverse-scattering prediction and reconstruction algorithms. The *Magazine* invites those workers who have such data sets, and are willing to share them, to submit an announcement and description, such as that given below, for publication in the *Magazine*. WRS]

Workers in Spain and France have developed equipment and a series of reconstruction algorithms for microwave imaging purposes. Special attention has been devoted in this work to non-invasive control of deep hyperthermia treatments. These workers are currently engaged in a program aiming to integrate a 434 MHz circular scanner for the coaxial TEM applications of the Academic Hospital of Utrecht (NL). Both groups have developed experimental prototypes, namely a planar-microwave camera [1], operating in France, and a circular scanner, in Spain [2]. They have also developed the corresponding tomographic-reconstruction algorithms for qualitative spectral and quantitative spatial iterative purposes [3-5]. As a result, a series of simulated and measured data are available. It seems interesting to extend these data to other workers for assessing the overall robustness of quantitative reconstruction algorithms. At present, this database contains near-field data from Spain. If you want to use it, please contact us at the e-mail addresses below. Details on how to access the database, along with a description of the data, follow.

How to access the data

The data files can be retrieved from an FTP server by the following scheme (the README.DOC file contains text similar to the above, describing the data):

\$ ftp voltor.upc.es

220 voltor FTP server (Version 16.2 Mort Apr 29 20:45:42 GMT 1991) ready.

Connected to VOLTOR.UPC.ES.

Name (VOLTOR.UPC.ES:nadine): anonymous

331 Guest login ok, send ident as password.

Password:

230 Guest login ok, access restrictions apply.

FTP> cd pub/BIOIMA

250 CWD command successful.

FTP> ascii

200 Type set to A.
 FTP> get README.DOC
 200 PORT command successful.
 150 Opening ASCII mode data connection for README (5046 bytes).
 226 Transfer complete.
 local: README.DOC remote: README.DOC
 5046 bytes received in 00:00:15.35 seconds

The data

You should obtain the following files:

README.DOC
 FANTCENT.ASC
 BRAGREG.ASC

Each of the files with the .ASC extension are scattered fields measured in the cylindrical system. The data in FANTCENT.ASC were obtained from a phantom consisting of two Plexiglas cylinders, filled with different concentrations of ethyl alcohol, as explained below. Dimensions are given in millimeters. Figures 1 and 2 show the phantom for which the measurements were taken, and a reconstruction obtained from it. Cylinder A was filled with a 96% solution of ethyl alcohol, with $\epsilon^* = 10 - j8.3$. Cylinder B was filled with a 4% ethyl alcohol solution, with $\epsilon^* = 73 - j11$. The data in BRAGREG.ASC were taken from a human forearm.

File format

All data files contain a matrix of 64×64 complex data values. The files are written in ASCII text, so you can inspect the data with a text editor. There is no header: just data. Each complex element is written in a line as floating-point text, in the format

[real part]	[imaginary part]	pair of antenna numbers
0.0	0.0	(0,0)
0.0	0.0	(0,1)
....
2.301653e-1	-6.499484e-1	(0,17)
....
0.0	0.0	(64,64)

Note that the number of the pair is not included in the file.

Illumination

The field amplitude corresponds to an incident cylindrical wave, produced by an axial current of $I = 1 A$:

$$E_{inc} = -(1/4)\omega\mu_0IH_0(k_0r)$$

where $I = 1$; r is the distance between the emitter and the point where the field is evaluated; k_0 is the wavenumber for the immersion medium; ω is the pulsation; H_0 is the second-order Hankel function; and μ_0 is the vacuum permeability.

The circular array is composed of 64 detectors. Hence, a maximum of 64 views can be used for the reconstruction. A view is obtained by using one antenna as an emitter, and 33 as receivers located on the opposite side, on a semicircle, as shown in Figure 3. The radius of the array is 0.125 m; the frequency is 2.33 GHz; and





Figure 1. The phantom used for the measurements.



T=25°C ± 0.5°C - f=(2.330±0.001) GHz
 spatial res. : Δ=3.5mm - 33 receivers - 64 views

Figure 2. A reconstruction obtained from the phantom data.




Figure 3. The arrangement of the transmitting and receiving antenna locations around the phantom. Note that when each antenna is used as a transmitter, 33 of the antennas are used as receivers.

the complex permittivity of water at 25° C is taken to be $77.3 - j8.66$.

References

1. J. Ch. Bolomey, Ch. Pichot, "Microwave tomography: from theory to practical imaging systems," *International Journal of Imaging Systems Technology*, 2, 1990, pp. 144-156.

