

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

130,000

International authors and editors

155M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Metrics Based on Information Entropy to Evaluate Landscape Complexities

*Sérgio Henrique Vannucchi Leme de Mattos,
Luiz Eduardo Vicente, Andrea Koga Vicente,
Cláudio Bielenki Junior, Maristella Cruz de Moraes,
Gabriele Luiza Cordeiro and José Roberto Castilho Piqueira*

Abstract

Information entropy concept is the base for many measures used to evaluate the complexity of complex environmental systems. Its application has great potential to evaluate landscape organization and dynamics, especially if we consider that there is a direct relation between their patterns and processes: the spatial arrangement (structure) of units within a mosaic reflects on system functions. Consequently, changes on structure reflects on functions and *vice versa*. Here, we exemplify how three measures based on information entropy – LMC and SDL complexity measures and H_e/H_{\max} variability measure – could be applied to evaluating the degree of complexity of a landscape and its components by associating their heterogeneity with the diversity of information acquired from the remote sensors' images. For this, we developed two scripts for a Geographical Information System (QGIS): (1) *CompPlex H_eROI*, that compares the complexity of a landscape patch with others and also with their transition areas; and (2) *CompPlex Janus*, which analyzes how complexity varies in the landscape over space and time, generating landscape complexity maps. We also use LMC and SDL complexity measures and H_e/H_{\max} variability measure to evaluate complexity time series of environmental variables, as rain and temperature, which allow to evaluate how their variations along time and space affects landscape dynamics. Therefore, application of such metrics in multi-temporal studies of landscape dynamics provides indicators of landscape resilience and the degree of conservation or degradation of its different fragments due to anthropic impacts related to land uses.

Keywords: complexity, Information entropy, landscape metrics

1. Introduction

From the perspective of the Complexity Paradigm [1], the landscape can be interpreted as a complex environmental system that is established from the interdependence relationships of the physical-natural system (that is, by the elements and processes present in nature) and the socioeconomic system (that is, the

elements and processes linked to human societies in their cultural, economic and social aspects). When these two systems interact, they are considered as subsystems of a system with a higher level of ecological organization: the landscape. Landscape can be considered the 2nd level of ecological organization, characterized by a set of interrelated ecosystems and formed, as the other levels, by the interactions between society and nature (**Figure 1**).

As is typical of complex adaptive system, the landscape presents non-linear negative and positive feedback processes generated from self-organization of its elements in interaction networks. The structure and dynamics of the landscape are affected and affect the other three levels (holons) constituents of this holarchic organization through bottom-up and top-down processes (**Figure 1**). The evidence of these interactions can be seen in landscape's patterns, as in this type of system, there is a direct relationship between its patterns and processes [4]: the spatial arrangement (structure) of units within a mosaic influence system functions. Consequently, changes in structure reflects on functions and vice versa, therefore affecting landscape resilience and integrity.

Thus, the complexity of a landscape and of the units that comprise the mosaic can be associated with the heterogeneities of its spatial, temporal, and structural patterns, with greater complexities being represented by patterns located in regions of intermediate heterogeneity in a gradient that goes from totally ordered patterns up to those completely disordered [5, 6]. To capture this typical signature of complex environmental systems not only qualitatively, it is necessary to use indicators capable of representing it in a quantitative way. This can be done through measures based on information entropy, which can be applied to assess the structure and dynamics of landscapes, as done by Mattos et al. [7] and Piqueira et al. [8] in




Figure 1. Holarchic organization levels of complex environmental systems (inspired in: [2, 3]).

relation to ecological interactions between their populations or by Piqueira & Mattos [9] for abiotic factors present in them.

Here, the application of measures based on information entropy for two purposes is demonstrated. The first is to assess the complexity of climatic time series. The other is both in comparing the complexity of a landscape patch with others and also with their transition areas, as well as allowing to verify as complexity varies in the landscape over space and time.

2. Use of measures based on information entropy to evaluate landscapes complexity

Remote sensor images can be used to identify landscape patterns in two different ways. The first one is related to the degree of roughness caused by the variability of tone or color, which provides image geometry and texture from targets [10–12]. The other is by examining the image spectral features, analyzing the values (e.g., surface reflectance, radiance, digital numbers) for each wavelength interval (i.e., band) vs. matrix pixels obtained for a specific target, which provides the spectral signature of a specific target [11, 13].

Several methods have been developed for both texture and spectral analysis to recognize patterns in remote sensing data [14–16]. Generally, many measures (also called metrics) derived from these methods have the same purposes: identifying similar patterns that occur in different places and distinguishing different patterns within a landscape. Although theoretically simple, in practice, this objective is not always easily achieved. Here, we discuss the use of three metrics derived from information entropy to measure the complexity of landscape patterns and show their applications to some case studies.

2.1 Information entropy applied to landscape patterns recognition and the evaluation of their level of complexity

Several metrics applied to textural and spectral analysis try to capture landscape patterns by using approaches deriving from theories and methods associated to the complexity paradigm, such as General System Theory, Cybernetic, Theory of Dissipative Structures, Hierarchy Theory, Percolation Theory, Self-Organized Criticality, Catastrophe Theory and Fractal Geometry ([17–20]).

Information entropy and other measures derived from it are also extensively used to quantify landscape heterogeneity and, consequently, to evaluate its organization level and complexity [18, 21]. According to Shiner et al. [22], there are three broad categories in which complexity measures based on information entropy may be classified: the first is composed of measures that consider complexity as a direct function of disorder (as is the case of very popular Shannon diversity index). So, measures of this category attribute lower values of complexity to ordered states and higher values to disordered states [22, 23]. Another category inverts this interpretation by associating higher complexity to most ordered states [22].

However, both measure categories are considered inadequate since there is no real complexity in situations that present zero or maximum entropy [23]. This fact is particularly applicable in Landscape Ecology, since, as mentioned by Parrot [24], more spatially complex landscapes are those in which the spatial pattern is situated in regions of intermediary heterogeneity, between order and disorder patterns. Thus, the maximum complexity would be located between these two extreme situations, which could be mathematically expressed as a convex function of disorder, as are the measures belonging to the third category defined by Shiner et al. [22].

This is the case of the LMC and SDL measures, proposed initially by López-Ruiz et al. [25] and Shiner et al. [22], respectively. In both definitions, the two complementary parameters – disorder and order – are combined to obtain a complexity measure.

2.1.1 H_e/H_{max} , LMC and SDL complexity measures and their application to remote sensing images

Shannon's Information Theory [26] could be applied to reflectance data in a remote sensing image band. These data are represented by their discretization in single digital numbers (DN), each DN representing a pixel value related to the intensity of the radiation in a particular wavelength at the sensor [11]. As the occurrence of certain DN values becomes more likely than other values, the entropy of the image decreases.

The variability measure H_e/H_{max} is related to Shannon entropy, and it belongs to the first category mentioned by Shiner et al. [22]; therefore, considering that complexity increases as a function of increasing the system disorder. This measure is useful to verify if a landscape and its patches are near the ordered/homogeneous or the disordered/heterogeneous patterns. To use this measure, it is necessary first to define system extension (N), given by the system's total number of possible states. In the case of remote sensing images, N corresponds to the number of different DN values present in the region of interest (ROI). As the maximum entropy value of a ROI could only be reached when the occurrence of the DNs values (*i.e.*, states) is equiprobable, the maximum entropy (H_{max}) is calculated considering all DNs values with the same probability as follows (Eq.(1)):

$$H_{max} = \log_2 N. \quad (1)$$

Dividing the number of pixels that have a determined DN value by the total DN values present in the ROI, we have the probability p of the i^{th} DN value of occurrence of this value within the ROI. The Boltzmann-Gibbs-Shannon entropy (H_e) for ROI is then calculated as (Eq.(2)):

$$H_e = - \sum_{DN \in N} P_{(DN)} \log_2 P_{(DN)}. \quad (2)$$

Finally, the variability measure (V) is obtained by dividing the information entropy calculated (H_e) by the maximum entropy (H_{max}), as follows (Eq.(3)):

$$V = \frac{H_e}{H_{max}}. \quad (3)$$

It can be deduced that complexity values for this measure range between 0 and 1, with complexity values associated with disorder (thermodynamic equilibrium).

Differently from the variability measure H_e/H_{max} , SDL and LMC belong to the third category of complexity measures defined by Shiner et al. [22], considering that the highest complexity is situated between order/homogeneous and disorder/heterogeneous patterns, that is, regions of intermediary heterogeneity associated with a high degree of self-organization. A convex function of information entropy may mathematically represent this assumption.

SDL measure is composed of two terms: disorder and order, *i.e.* (Eq.(4)):

$$SDL = (H_e/H_{max})[1 - (H_e/H_{max})]. \quad (4)$$

For the LMC measure, the order term is substituted by another term, called disequilibrium (D), which measures the distance between the system probability and the uniform distribution [27] (Eq.(5)):

$$D = \sum_{i=1}^N \left(P_{(DN)} - \frac{1}{N} \right)^2. \quad (5)$$

Consequently, LMC is given by (Eq.(6))

$$LMC = (H_e/H_{max})[1 - (H_e/H_{max})] \quad (6)$$

or by (Eq.(7)):

$$LMC = (H_e/H_{max})D \quad (7)$$

Zero is the minimum value for both measures, while 0.25 and 0.15 are the maximum values for SDL and LMC, respectively. These maximum values occur when the DN distribution is uniform [27].

To apply these measures based on information entropy to the remote sensing images, we developed two scripts in Python language to be executed as plugins in QGIS, an open-source Geographic Information System. The first one is CompPlex H_e ROI, which calculates H_e/H_{max} , LMC, and SDL complexity measures of a ROI and compares them with others patches and their transition areas. The other plugin is CompPlex Janus, composed of a sliding window that runs through the image, calculating those three measures for the set of pixels inside it. CompPlex Janus then generates complexity maps, allowing verification as complexity varies in the landscape over space and time.

Here we present examples illustrating the application of CompPlex H_e ROI to evaluate the complexity of several ROIs (Example 1) and the use of CompPlex Janus to evaluate the spatial distribution of landscape patterns complexity (Example 2), highlighting in both cases the efficiency of the measures based on the information entropy presented.

2.1.1.1 Example 1: CompPlex H_e ROI applied to evaluate patterns of different land uses

In this example, we show how CompPlex H_e ROI had been applied to evaluate, by using metrics based on information entropy, the complexity of spatial patterns with different land uses present in two river neighbor basins located at municipality of São Carlos (São Paulo state, Brazil – **Figure 2**), especially as indicators of the resilience of its green areas, to help establishing a free space system for this region. Land uses inside these river basins had been identified using images from CBERS 4 remote sensor (**Figure 3**), and six categories of use were selected to be evaluated by CompPlex H_e ROI. Results obtained for ROIs of these categories are shown in **Table 1**, where they are compared for each measure and each band used.

Colors associated with values presented in **Table 1** help to identify tendencies of each land use complexity pattern. In general, ROIs of exposed soil, urban areas, and pasture have high values for H_e , H_{max} , and H_e/H_{max} measures and low values for SDL and LMC measures, indicating that these land uses have more disordered patterns. In turn, agricultural use varies from low to relatively high values for H_e , H_{max} , and H_e/H_{max} measures, but has, in most cases, low values for SDL and LMC measures. Finally, vegetation areas have low values for the first three measures (H_e , H_{max} and H_e/H_{max}) and high values for SDL and LMC measures that use convex




Figure 2.
Localization of study area in São Carlos's municipality (São Paulo state, Brazil).




Figure 3.
Land use and land cover on study area (municipality of São Carlos, São Paulo state, Brazil).

function of disorder to associate more complexity with patterns situated in a zone between ordered and disordered patterns. Therefore, these results are coherent and consistent with the Landscape Ecology assumption that more complexity is found in intermediary heterogeneity patterns [5, 6], as is the case of vegetation areas present in the two neighbor river basins studied. High values for SDL and LMC obtained by these areas could be related to their high levels of self-organization and resilience.

Land use	ROI	He			Hmax			He/Hmax			SDL			LMC		
		Band 5	Band 6	Band 7	Band 5	Band 6	Band 7	Band 5	Band 6	Band 7	Band 5	Band 6	Band 7	Band 5	Band 6	Band 7
Exposed soil	ROI 1	4,4811	3,9741	3,5066	5,0444	4,6439	4,2479	0,8883	0,8358	0,8255	0,0992	0,1234	0,1441	0,0256	0,0366	0,0485
	ROI 2	4,1565	3,5415	3,0735	5,2095	4,5850	4,2479	0,7979	0,7724	0,7235	0,1613	0,1758	0,2000	0,0361	0,0527	0,0696
	ROI 3	4,2545	3,6888	3,2378	5,0444	4,3923	4,0000	0,8434	0,8398	0,8094	0,1321	0,1345	0,1542	0,0358	0,0504	0,0666
	ROI 4	4,0900	3,4612	2,9686	4,3923	3,9069	3,4594	0,9323	0,8859	0,8581	0,0631	0,1011	0,1218	0,0211	0,0413	0,0466
	ROI 5	3,3000	2,6311	2,5105	4,7549	4,1699	3,7004	0,6928	0,6310	0,6784	0,2128	0,2328	0,2182	0,0719	0,1054	0,0980
Urban area	ROI 1	4,6000	4,1853	3,8554	5,0875	4,7549	4,3923	0,9061	0,8802	0,8778	0,0851	0,1054	0,1073	0,0159	0,0228	0,0274
	ROI 2	5,0000	4,4723	4,0663	5,6724	5,3576	5,1293	0,8802	0,8348	0,7928	0,1054	0,1379	0,1643	0,0145	0,0234	0,0322
	ROI 3	4,7000	4,4006	4,1164	5,7004	5,7814	5,4263	0,8279	0,7612	0,7586	0,1425	0,1818	0,1831	0,0234	0,0315	0,0374
	ROI 4	5,1000	4,8281	4,5210	6,1293	6,1699	6,0444	0,8344	0,7825	0,7480	0,1382	0,1702	0,1885	0,0173	0,0253	0,0334
	ROI 5	6,0000	5,9472	5,6884	6,4919	6,5236	6,3399	0,9306	0,9116	0,8969	0,0646	0,0805	0,0924	0,0072	0,0099	0,0134
	ROI 6	4,3000	3,8912	3,4527	4,7004	4,5236	4,1699	0,9061	0,8602	0,8280	0,0851	0,1203	0,1424	0,0182	0,0319	0,0453
	ROI 7	4,8000	4,3885	4,0597	5,5546	6,4757	6,3750	0,7389	0,6777	0,6368	0,1929	0,2184	0,2313	0,0240	0,0355	0,0437
Forest vegetation	ROI 1	3,3000	2,8502	2,5804	4,4594	4,0000	3,7004	0,7327	0,7125	0,6973	0,1958	0,2048	0,2111	0,0869	0,1109	0,1092
	ROI 2	3,7000	3,2672	2,7327	4,6439	4,3219	3,8074	0,7944	0,7560	0,7177	0,1633	0,1845	0,2026	0,0570	0,0694	0,0904
	ROI 3	3,4000	2,8545	2,5770	4,9542	4,9069	4,5850	0,6779	0,5817	0,5621	0,2184	0,2433	0,2461	0,0684	0,0906	0,1017
	ROI 4	3,3000	2,8845	2,4381	4,7549	4,2479	3,8074	0,6880	0,6790	0,6404	0,2146	0,2179	0,2303	0,0815	0,0920	0,1112
	ROI 5	3,7000	3,3071	2,8197	4,8074	4,3923	4,1699	0,7684	0,7529	0,6762	0,1779	0,1860	0,2190	0,0532	0,0633	0,0895
	ROI 6	3,5000	3,2881	2,8041	4,5850	4,3923	3,9069	0,7631	0,7486	0,7177	0,1808	0,1882	0,2026	0,0571	0,0616	0,0835
	ROI 7	3,2000	2,8528	2,4169	4,8580	4,5236	4,2479	0,6535	0,6307	0,5690	0,2264	0,2329	0,2452	0,0793	0,0943	0,1117
	ROI 8	3,5000	3,0592	2,6443	4,9542	4,6439	4,0875	0,7122	0,6588	0,6469	0,2050	0,2248	0,2284	0,0659	0,0794	0,0950
	ROI 9	2,8000	2,5271	2,2851	4,3923	4,0875	3,5850	0,6276	0,6183	0,6374	0,2337	0,2360	0,2311	0,1100	0,1139	0,1247
	ROI 10	2,6000	2,2068	1,9570	3,8074	3,3219	2,8074	0,6828	0,6643	0,6971	0,2166	0,2230	0,2112	0,1168	0,1316	0,1251
	ROI 11	3,7000	3,1969	2,8469	4,9069	4,5850	4,3219	0,7505	0,6973	0,6587	0,1873	0,2111	0,2248	0,0760	0,0979	0,1146
	ROI 12	2,8000	2,3646	2,2609	4,1699	3,5850	3,4594	0,6766	0,6596	0,6335	0,2188	0,2245	0,2264	0,1153	0,1384	0,1272
	ROI 13	3,0000	2,6754	2,3351	4,1699	3,9069	3,7004	0,7188	0,6848	0,6310	0,2021	0,2159	0,2328	0,0757	0,1000	0,1121
	ROI 14	2,9000	2,6716	2,2181	4,2479	3,9069	3,4594	0,6725	0,6838	0,6412	0,2202	0,2162	0,2301	0,0963	0,1000	0,1232
	ROI 15	2,4000	2,0496	1,9656	4,0875	3,5850	3,0000	0,5805	0,5717	0,6552	0,2435	0,2449	0,2259	0,1433	0,1516	0,1380
	ROI 16	2,7000	2,6331	2,0771	3,9069	3,7004	3,3219	0,6934	0,7116	0,6253	0,2126	0,2052	0,2343	0,0902	0,0926	0,1244
	ROI 17	3,6190	3,2078	2,9394	4,9542	4,5236	4,0875	0,7305	0,7091	0,7191	0,1969	0,2063	0,2020	0,0837	0,0955	0,0985
	ROI 18	2,6816	2,1261	2,0039	4,2479	3,7004	3,4594	0,6313	0,5745	0,5793	0,2328	0,2444	0,2437	0,1124	0,1364	0,1429
	ROI 19	2,2629	1,8992	1,8473	3,4594	3,1699	2,8074	0,6541	0,5991	0,6580	0,2262	0,2402	0,2250	0,1148	0,1474	0,1311
	ROI 20	3,5891	3,1214	2,7833	4,7004	4,0875	3,4594	0,7636	0,7637	0,8045	0,1805	0,1805	0,1573	0,0778	0,0890	0,0853
	ROI 21	3,6965	3,2385	2,6494	4,2479	3,7004	3,1699	0,8702	0,8752	0,8358	0,1130	0,1093	0,1372	0,0403	0,0448	0,0693
	ROI 22	2,5271	1,9190	1,8315	3,8074	3,1699	3,1699	0,6637	0,6054	0,5778	0,2232	0,2389	0,2440	0,1287	0,1422	0,1488
ROI 23	3,1657	2,8982	2,2734	4,1699	3,7004	3,1699	0,7592	0,7832	0,7172	0,1828	0,1698	0,2028	0,0923	0,0805	0,1129	
Agriculture	ROI 1	3,4584	2,9410	2,3150	4,0875	3,7004	3,1699	0,8461	0,7948	0,7303	0,1302	0,1631	0,1970	0,0453	0,0621	0,1011
	ROI 2	3,6998	3,3780	2,6080	4,0875	3,8074	3,1699	0,9052	0,8872	0,8227	0,0858	0,1001	0,1458	0,0294	0,0366	0,0699
	ROI 3	4,0709	3,6021	2,9829	5,0000	4,7004	4,2479	0,8142	0,7663	0,7022	0,1513	0,1791	0,2091	0,0375	0,0451	0,0687
	ROI 4	4,7789	4,0867	3,7128	5,6724	5,5546	5,4594	0,8425	0,7357	0,6801	0,1327	0,1944	0,2176	0,0171	0,0338	0,0416
	ROI 5	3,2443	2,6515	2,1540	4,3923	3,9069	3,3219	0,7386	0,6787	0,6484	0,1931	0,2181	0,2280	0,0621	0,0995	0,1225
	ROI 6	4,3082	3,7854	3,2445	5,3219	5,0000	4,5236	0,8095	0,7571	0,7172	0,1542	0,1839	0,2028	0,0298	0,0427	0,0597
Mining	ROI 1	5,8503	5,2216	4,5579	6,3038	6,0000	5,4919	0,9281	0,8703	0,8299	0,0668	0,1129	0,1411	0,0070	0,0142	0,0250
Pasture	ROI 1	4,3831	3,6957	3,2239	5,3576	5,0000	4,6439	0,8181	0,7391	0,6942	0,1488	0,1928	0,2123	0,0253	0,0442	0,0577
	ROI 2	3,9380	3,5371	2,9939	4,6439	4,2479	3,9069	0,8480	0,8327	0,7663	0,1289	0,1393	0,1791	0,0297	0,0396	0,0395
	ROI 3	4,0871	3,6439	3,0653	4,6439	4,0875	3,5850	0,8801	0,8915	0,8550	0,1055	0,0967	0,1239	0,0237	0,0253	0,0408
	ROI 4	4,2984	3,6619	3,1821	5,0000	4,5850	4,4594	0,8597	0,7987	0,7136	0,1206	0,1608	0,2044	0,0249	0,0418	0,0615
	ROI 5	4,5809	4,1065	3,6705	5,6724	5,4919	5,1293	0,8076	0,7477	0,7156	0,1554	0,1886	0,2035	0,0269	0,0397	0,0523
	ROI 6	4,0211	3,4431	3,0533	5,0000	4,3923	3,8074	0,8042	0,7839	0,8019	0,1575	0,1694	0,1588	0,0389	0,0532	0,0585
	ROI 7	4,3997	4,0811	3,5683	4,6439	4,3219	3,9069	0,9474	0,9443	0,9133	0,0498	0,0526	0,0791	0,0127	0,0139	0,0232
	ROI 8	4,4676	3,9592	3,4447	5,3576	5,0000	4,7004	0,8339	0,7918	0,7328	0,1385	0,1648	0,1958	0,0219	0,0337	0,0497
	ROI 9	4,1668	3,5807	3,0151	4,5850	4,0000	3,5850	0,9088	0,8952	0,8410	0,0829	0,0938	0,1337	0,0219	0,0329	0,0533
	ROI 10	4,2233	3,7916	3,1975	4,5236	4,0875	3,5850	0,9336	0,9276	0,8919	0,0620	0,0671	0,0964	0,0135	0,0206	0,0323
	ROI 11	3,8417	3,4896	2,9152	4,3219	3,9069	3,3219	0,8889	0,8932	0,8775	0,0988	0,0954	0,1075	0,0284	0,0332	0,0463

Legends:

- for H_e , H_{max} and H_e/H_{max} : ■ = high value ■ = low value
- for SDL and LMC: ■ = low value ■ = high value

Table 1.

Results obtained by CompPlex H_e ROI for six land use and land cover categories in the study area (municipality of São Carlos, São Paulo state, Brazil).

2.1.1.2 Example 2: CompPlex Janus and landscape complexity maps

To exemplify how CompPlex Janus works to generate landscape complexity maps, here is presented a case study of sensor images from the Assis Ecological Station and its boundaries (located at Assis, São Paulo State, Brazil – **Figure 4**). Several tests varying the sliding window size, sensor band, and number of color classes had been performed to compare results obtained by H_e , H_e/H_{max} , SDL, and LMC measures. Some of these results are shown in **Figures 5** and **6**.

Comparing the four examples of maps of H_e , we observe that for sliding window of 3x3 pixels sides (**Figure 5A** and **B**), this measure highlights borders among different land uses and, especially for the image of band 3 (**Figure 5A**), the apparent homogeneity of natural vegetation is broken. For this same measure, but for sliding window of 9x9 pixels sides results are shown in **Figure 5C** and **D**. On the other side, these edges are blurred for the 9x9 pixel window. However, areas with higher values for this measurement (visualized by more intense red tones) are found around natural vegetation areas, possibly indicating areas of greater risk to their integrity and resilience.




Figure 4.
Localization of Assis Ecological Station and surroundings (municipality of Assis, State of São Paulo, Brazil).




Figure 5.
Some complexity maps for H_e and H_e/H_{max} entropy measures generated by CompPlex Janus to study area.

For H_e/H_{max} measure, we can perceive a significant difference between maps generated by a window of 3x3 pixels sides (Figure 5E and F) and those of 9x9 pixels (Figure 5G and H). Due to the reduced amount of pixels in the smaller window, there is less diversity of information, and the interval between minimum and maximum values is high. As occurred with the H_e measurement, a larger window (9x9 pixels) generated a more extensive range for the minimum and maximum values, highlighting possible areas that represent greater risks to natural vegetation.

In Figure 6A-H, we show, respectively, some results obtained by Complex Janus to SDL and LMC measure. For a window of 3x3 pixels sides (Figure 6A, B, E and F), these measures allow identifying punctual areas with higher values within natural vegetation regions. This effect is best observed on images generated by windows of




Figure 6.
 Some complexity maps for SDL and LMC entropy measures generated by CompPlex Janus to study area.

9x9 pixels (Figure 6C, D, G, and H), where areas of natural vegetation have a more variable value gradient and areas with other land uses are ‘homogenized’ with low values for both measures. In general, SDL and LMC measures assign higher values to natural vegetation, consistent with the assumption of Landscape Ecology that more complex patterns are associated with intermediary spatial heterogeneity.

3. Measures based on information entropy applied to analyze climatic time series

Information entropy measures are also useful to verify complexity (in the sense of variability) of time series, as shown by Piqueira and Mattos [9]. To exemplify how these measures can be utilized for this purpose, here we show an application of H_e/H_{max} , SDL, and LMC measures for a time series corresponding to the maximum daily temperatures that occurred in each January from 1980 to 2017 in the municipality of São Carlos (São Paulo State, Brazil).

To calculate the measures, daily maximum temperature data for each January of the entire time series were used to define its quartiles. Then, for the January data for each year, we check the number of days that belonged to each quartile, which allowed us to calculate the probability p for each interval. The system extension (N) corresponded to the number of quartiles that presented at least one data.

With these values, it was possible to calculate the measures H_e/H_{max} , SDL, and LMC for data from January of each year, according to the equations previously presented. The results obtained are shown in Table 2. To compare the performance of measures and to try to identify any patterns from results for each measure we group each of them in decreasing order of value and organize in four classes. Table 3 shows pairwise comparisons between the measures to verify whether or not a given year occupies the same class and the same position for both measures. Through these comparisons, it is evident that H_e/H_{max} and SDL have the same behavior, while LMC measure behavior differs from them, revealing the differences in the relations between order and disorder terms present in these measures’ equations.

Month/Year	He/Hmax	SDL	LMC	El Niño
jan/80	0,996121	0,003864	0,002764	Moderate
jan/81	0,891832	0,096468	0,058234	-
jan/82	0,842241	0,132871	0,104075	-
jan/83	0,955452	0,042563	0,028501	Strong
jan/84	0,883083	0,103247	0,083392	-
jan/85	0,899002	0,090798	0,056753	-
jan/86	0,929433	0,065587	0,041346	-
jan/87	0,943894	0,052958	0,036096	Moderate
jan/88	0,913992	0,078611	0,057778	Strong
jan/89	0,807586	0,155391	0,098112	-
jan/90	0,898222	0,091419	0,053043	-
jan/91	0,888074	0,099399	0,072774	-
jan/92	0,946830	0,050343	0,038179	Strong
jan/93	0,987089	0,012744	0,008988	Weak
jan/94	0,887049	0,100193	0,059767	-
jan/95	0,958728	0,039569	0,028682	-
jan/96	0,975916	0,023504	0,017010	-
jan/97	0,750346	0,187327	0,128636	-
jan/98	0,815862	0,150231	0,111003	Strong
jan/99	0,959517	0,038844	0,024712	-
jan/00	0,982448	0,017244	0,010990	-
jan/01	0,770428	0,176869	0,128872	-
jan/02	0,995932	0,004051	0,002850	-
jan/03	0,958728	0,039569	0,028682	Moderate
jan/04	0,882989	0,103319	0,063169	-
jan/05	0,944311	0,052588	0,030216	-
jan/06	0,906762	0,084544	0,059208	-
jan/07	0,933882	0,061746	0,037657	Moderate
jan/08	0,920452	0,073220	0,048609	-
jan/09	0,869816	0,113236	0,087570	-
jan/10	0,933882	0,061746	0,037657	Moderate
jan/11	0,987089	0,012744	0,008988	-
jan/12	0,872595	0,111173	0,076878	-
jan/13	0,964841	0,033923	0,024765	-
jan/14	0,987089	0,012744	0,008988	-
jan/15	0,934765	0,060979	0,119966	-
jan/16	0,989680	0,010213	0,006951	Strong
jan/17	0,998848	0,001150	0,000780	-

For H_e/H_{max} : = low value = high value
 For SDL and LMC: = low value = high value

Table 2. Results obtained by applying entropy measures for daily maximum temperature data of municipality of São Carlos (São Paulo state, Brazil) for each January from 1980 to 2017.

Group	Comparaison		Comparaison		Comparaison	
	He/Hmax*	SDL**	He/Hmax*	LMC**	SDL**	LMC**
1	jan/97	jan/97	jan/97	jan/01	jan/97	jan/01
	jan/01	jan/01	jan/01	jan/97	jan/01	jan/97
	jan/89	jan/89	jan/89	jan/15	jan/89	jan/15
	jan/98	jan/98	jan/98	jan/98	jan/98	jan/98
	jan/82	jan/82	jan/82	jan/82	jan/82	jan/82
	jan/09	jan/09	jan/09	jan/89	jan/09	jan/89
	jan/12	jan/12	jan/12	jan/09	jan/12	jan/09
	jan/04	jan/04	jan/04	jan/84	jan/04	jan/84
	jan/84	jan/84	jan/84	jan/12	jan/84	jan/12
	jan/94	jan/94	jan/94	jan/91	jan/94	jan/91
2	jan/91	jan/91	jan/91	jan/04	jan/91	jan/04
	jan/81	jan/81	jan/81	jan/94	jan/81	jan/94
	jan/90	jan/90	jan/90	jan/06	jan/90	jan/06
	jan/85	jan/85	jan/85	jan/81	jan/85	jan/81
	jan/06	jan/06	jan/06	jan/88	jan/06	jan/88
	jan/88	jan/88	jan/88	jan/85	jan/88	jan/85
	jan/08	jan/08	jan/08	jan/90	jan/08	jan/90
	jan/86	jan/86	jan/86	jan/08	jan/86	jan/08
	jan/07	jan/07	jan/07	jan/86	jan/07	jan/86
	jan/10	jan/10	jan/10	jan/92	jan/10	jan/92
3	jan/15	jan/15	jan/15	jan/07	jan/15	jan/07
	jan/87	jan/87	jan/87	jan/10	jan/87	jan/10
	jan/05	jan/05	jan/05	jan/87	jan/05	jan/87
	jan/92	jan/92	jan/92	jan/05	jan/92	jan/05
	jan/83	jan/83	jan/83	jan/95	jan/83	jan/95
	jan/95	jan/95	jan/95	jan/03	jan/95	jan/03
	jan/03	jan/03	jan/03	jan/83	jan/03	jan/83
	jan/99	jan/99	jan/99	jan/13	jan/99	jan/13
	jan/13	jan/13	jan/13	jan/99	jan/13	jan/99
4	jan/96	jan/96	jan/96	jan/96	jan/96	jan/96
	jan/00	jan/00	jan/00	jan/00	jan/00	jan/00
	jan/93	jan/93	jan/93	jan/93	jan/93	jan/93
	jan/11	jan/11	jan/11	jan/11	jan/11	jan/11
	jan/14	jan/14	jan/14	jan/14	jan/14	jan/14
	jan/16	jan/16	jan/16	jan/16	jan/16	jan/16
	jan/02	jan/02	jan/02	jan/02	jan/02	jan/02
	jan/80	jan/80	jan/80	jan/80	jan/80	jan/80
jan/17	jan/17	jan/17	jan/17	jan/17	jan/17	

- belongs to same group and occupy the same position
- belongs to same group but occupy different position
- doesn't belong to the same group

* From minor value to major value
 ** From major value to minor value

Table 3. Results obtained by applying entropy measures for daily maximum temperature data of municipality of São Carlos (São Paulo state, Brazil) for each January from 1980 to 2017.

4. Conclusions

In addition to exploring results obtained from case studies here presented, we intended to show why and how measures based on information entropy can contribute to understanding complexity of landscapes patterns and processes. As shown in the first example, H_e/H_{\max} , SDL, and LMC are complexity measures that represent useful tools for evaluating landscape patterns. H_e/H_{\max} allows identifying ordered and disordered targets, while SDL and LMC are related to intermediary heterogeneity patterns presented by landscape patches. Comparing the landscape metrics used here with the spectral decomposition methods proposed in Mustard and Sunshine [13], they prove to be quite efficient in comparing the complexity of the patterns of different patches as well as their variation over the entire landscape. Based on this example, the application of such metrics is proposed for multi-temporal studies of landscape dynamics, for evaluating resilience and the degree of degradation of different fragments, for estimating the degree of the anthropic impact due to alterations on land usage, among other applications.

In the second example, we highlight the use of these measures to evaluate complexity in climatic time series. Our future studies involve the application of these measures as alternatives for classical statistical analysis, using them to assess the influences of both natural processes, such as El Niño and La Niña, and those resulting from anthropic processes, such as the increase in temperature and frequency of extreme weather events, such as severe droughts and heavier rains.

Author details

Sérgio Henrique Vannucchi Leme de Mattos^{1*}, Luiz Eduardo Vicente²,
Andrea Koga Vicente², Cláudio Bielenki Junior¹, Maristella Cruz de Moraes¹,
Gabriele Luiza Cordeiro¹ and José Roberto Castilho Piqueira³


¹ Environmental Complex Systems Laboratory, Department of Hydrobiology, Biological and Health Sciences Center, Federal University of São Carlos (UFSCar), São Carlos, SP, Brazil

² ABC Platform, Embrapa Environment, Jaguariúna, SP, Brazil

³ Department of Telecommunications and Control Engineering, Polytechnic School of the University of Sao Paulo (USP), São Paulo, SP, Brazil

*Address all correspondence to: sergiomattos@ufscar.br

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] MORIN, E. **From the concept of system to the paradigm of complexity.** Journal of Social and Evolutionary Systems. Volume 15, Issue 4, 1992, Pages 371-385.
- [2] CHRISTOFOLETTI, A. **Modelagem de Sistemas Ambientais.** Editora Blucher, 1999. 256 Pages.
- [3] KOESTLER, A. **The Ghost in the Machine.** Macmillan Length. 384 Pages.
- [4] TURNER, M.G. & GARDNER, R.H. **Quantitative methods in landscape ecology: an introduction.** In: *Quantitative methods in landscape ecology.* New York: Springer Verlag, 1990. p.3-14.
- [5] ANAND, M. et al. **Ecological Systems as Complex Systems: Challenges for an Emerging Science.** Diversity 2010, 2, 395-410; doi:10.3390/d2030395.
- [6] PARROTT, L; **Measuring ecological complexity.** Ecological Indicators, November 2010, 10(6):1069-1076. doi: 10.1016/j.ecolind.2010.03.014.
- [7] MATTOS, Sérgio Henrique Vannucchi Leme de; PIQUEIRA, José Roberto Castilho ; VASCONCELLOS NETO, João ; ORSATTI, Fernando Moya. **Measuring q-bits in three-trophic level systems.** Ecological Modelling, v. 200, n.1-2, p. 183-187, 2007.
- [8] PIQUEIRA, J.R.C., MATTOS, S.H.V. L; VASCONCELOS NETO, J. **Measuring complexity in three-trophic level systems.** Ecological Modelling, v. 220, p. 266-271, 2009.
- [9] PIQUEIRA, J. R. C.; MATTOS, S.H.V. L. **LMC and SDL complexity measures: a tool to explore time series.** Complexity, v. 2019, p. 1-8, 2019.
- [10] AVERY, T.E., BERLIN, G.L. **Fundamentals of remote sensing and airphoto interpretation.** New York: Macmillan, 1992. 472p.
- [11] JONES, H.G., VAUGHAN, R.A. **Remote Sensing of Vegetation: Principles, Techniques, and Applications.** UP Oxford, 2010. 384 Pages.
- [12] MUSICK, H.B. & GROVER, H.D. **Image textural measures as indices of landscape pattern.** In: Turner, M.G. & Gardner, R.H. Quantitative methods in landscape ecology. New York: Springer Verlag, 1990. p.77-103.
- [13] MUSTARD, J.F.; SUNSHINE, J.M. **Spectral analysis for Earth Science: investigations using remote sensing data.** In: RENCZ, A.N. Remote sensing for the Earth Sciences. New York: John Wiley & Sons, 1999.
- [14] ASNER, G.P. et al. **Remote sensing of selective logging in Amazonia- Assessing limitations based on detailed field observations, Landsat ETM+, and textural analysis.** Remote Sensing of Environment, v.80, 2002, p. 483-496.
- [15] GARRIGUES, S. *et al.* **Influence of landscape spatial heterogeneity on the non-linear estimation of leaf area index from moderate spatial resolution remote sensing data.** Remote Sensing of Environment, v.105, n.4.2006, p.286-298.
- [16] MIRANDA, F.P. *et al.* **Analysis of JERS-1 (Fuyo-1) SAR data for vegetation discrimination in northwestern Brazil using the semivariogram textural classifier (STC).** *International Journal of Remote Sensing*, v.17, 1996.
- [17] BAK, P. **How nature works: the science of self-organized criticality.** New York: Springer-Verlag, 1997. 212p.

[18] FARINA, A. **Principles and methods in landscape ecology.** Londres: Chapman & Hall, 1998. 235p.

[19] LI, B.L. **Fractal geometry applications in description and analysis of patch patterns and patch dynamics.** Ecological Modelling, v.132, 2000. p.33-50.

[20] NAVEH, Z. & LIEBERMAN, A. **Landscape ecology.** New York: Springer Verlag, 1994. 360p.

[21] FEAGIN, R.A. **Heterogeneity versus homogeneity: a conceptual and mathematical theory in terms of scale-invariant and scale-covariant distributions.** Ecological Complexity, v.2, 2005, p.339–356.

[22] SHINER, J.S. et al. **Simple measure of complexity.** Physical Review, 1999, v.59, n.2, Pages 1459-1464.

[23] KANEKO, K., TSUDA, I. **Complex Systems: chaos and beyond.** Berlin: Springer Verlag, 2001. 257p.

[24] PARROT, L. **Quantifying the complexity of simulated spatiotemporal population dynamics.** Ecological Complexity, v.2, 2005, p.175-184.

[25] LÓPEZ-RUIZ, R. *et al.* A statistical measure of complexity. **Physical Letter A**, v.209, 1995, p.321-326.

[26] SHANNON, C. E. **A mathematical theory of communication.** The Bell System Technical Journal, Vol. 27, pp. 379–423, 623–656, July, October, 1948.

[27] PIQUEIRA, J.R.C. **A comparison of LMC and SDL complexity measures on binomial distributions.** Physica. A (Print), v. 444, p. 271-275, 2016.