

# Northumbria Research Link

Citation: Geirdal, Amy K. Østertun, Price, Daicia, Schoultz, Mariyana, Thygesen, Hilde, Ruffolo, Mary, Leung, Janni and Bonsaksen, Tore (2021) The Significance of Demographic Variables on Psychosocial Health from the Early Stage and Nine Months after the COVID-19 Pandemic Outbreak. A Cross-National Study. *International Journal of Environmental Research and Public Health*, 18 (8). p. 4345. ISSN 1660-4601

Published by: MDPI

URL: <https://doi.org/10.3390/ijerph18084345> <<https://doi.org/10.3390/ijerph18084345>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/45979/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria  
University**  
NEWCASTLE





**UniversityLibrary**



Article

# The Significance of Demographic Variables on Psychosocial Health from the Early Stage and Nine Months after the COVID-19 Pandemic Outbreak. A Cross-National Study

Amy K. Østertun Geirdal <sup>1,\*</sup>, Daicia Price <sup>2</sup>, Mariyana Schoultz <sup>3</sup>, Hilde Thygesen <sup>4,5</sup>, Mary Ruffolo <sup>2</sup>, Janni Leung <sup>6</sup>  and Tore Bonsaksen <sup>5,7</sup> 

<sup>1</sup> Department of Social Work, Faculty of Social Sciences, Oslo Metropolitan University, 0167 Oslo, Norway

<sup>2</sup> School of Social Work, University of Michigan, Ann Arbor, MI 48109, USA; daiciars@umich.edu (D.P.); mruffolo@umich.edu (M.R.)

<sup>3</sup> Department of Health and Life Sciences, Northumbria University, Newcastle upon Tyne NE1 8ST, UK; Mariyana.schoultz@northumbria.ac.uk

<sup>4</sup> Department of Occupational Therapy, Prosthetics and Orthotics, Faculty of Health Science, Oslo Metropolitan University, 0167 Oslo, Norway; hilde.thygesen@oslomet.no

<sup>5</sup> Faculty of Health Studies, VID Specialized University, 4306 Sandnes, Norway; tore.bonsaksen@inn.no

<sup>6</sup> Faculty of Health and Behavioural Science, University of Queensland, Brisbane, QLD 4072, Australia; j.leung1@uq.edu.au

<sup>7</sup> Department of Health and Nursing Science, Faculty of Social and Health Sciences, Inland Norway University of Applied Sciences, 2418 Elverum, Norway

\* Correspondence: amyoge@oslomet.no



**Citation:** Geirdal, A.K.Ø.; Price, D.; Schoultz, M.; Thygesen, H.; Ruffolo, M.; Leung, J.; Bonsaksen, T. The Significance of Demographic Variables on Psychosocial Health from the Early Stage and Nine Months after the COVID-19 Pandemic Outbreak. A Cross-National Study. *Int. J. Environ. Res. Public Health* **2021**, *18*, 4345. <https://doi.org/10.3390/ijerph18084345>

Academic Editor: Paul B. Tchounwou

Received: 31 March 2021

Accepted: 17 April 2021

Published: 20 April 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** This cross-national study explored stability and change in mental health, quality of life, well-being and loneliness during the early stage and nine months after the implementation of COVID-19 pandemic social distancing measures and periodic lockdowns as adjusted by demographic variables. In the USA, the UK, Australia and Norway, 7284 individuals responded to the invitation to take part in two cross-sectional web-based surveys (April and November 2020), including questions about sociodemographic variables and psychosocial outcomes. Independent *t*-tests and generalized linear models (GLM) and estimated marginal means were used to analyze differences between subgroups and countries, multiple linear regression analyses were conducted on the psychosocial outcome measures by demographic variables and time in each country and mean responses presented by time after adjusting for all demographic variables in the model. Age, gender, civil status, education, employment, place of work and living area were all significant factors for psychosocial health across the countries. Differences in mental health, quality of life, well-being and loneliness were found between the countries in both April and November 2020, while time did not contribute to reducing the toll in any of the four countries over the nine-month period.

**Keywords:** COVID-19; coronavirus; psychosocial health; mental health; quality of life; well-being; loneliness; cross-national study

## 1. Introduction

During the COVID-19 pandemic, the main policy for public behavior has been periodic lockdowns and social distancing [1] including recommendations of an appropriate distance between individuals outside one's own household. To prevent viral spread, people were asked to stay at home as much as possible and only do "necessary" travels. Businesses defined as non-essential, for example, gyms and furniture stores were temporarily shut down at different points throughout the pandemic. Schools and universities have been either closed for in-person classroom learning or moved to distance learning, remote teaching and online instruction formats [2]. Many people have transitioned to working remotely [3], and those working at their workplaces, e.g., in healthcare, have had to take great precautions and follow strict infection control guidelines [4].

A review article based on 19 cross-sectional studies that focused on the association between the COVID-19 pandemic and mental health revealed that being female was associated with a higher degree of mental distress than among men [5] and that being older was associated with experiences of better mental health. The finding is in line with earlier studies, e.g., Carstensen et al. [6] and Neubauer et al. [7] showed that older people in general report fewer negative emotions than younger people, and stressful events are perceived as less unpleasant. According to Löwe et al. [8], older people tend to score lower on depression and anxiety measures.

According to Xiong et al. [5], the COVID-19 pandemic represents a threat to mental health worldwide, independent of a country's status as low-, middle-, or high-income. This is in line with a previous study [9] that found that in a number of western countries, COVID-19 bears a significant impact on psychosocial health including a higher level of emotional distress, poorer quality of life and well-being and increased feeling of loneliness. Other studies, as summarized in a review study [10], measured well-being and psychological distress in the general public and confirmed worse psychological well-being and higher scores of psychological distress including increased levels of depression and anxiety and high levels of post-traumatic stress disorder (PTSD) [10,11].

In the initial phase after the COVID-19 outbreak, Wang et al. [12] performed a longitudinal study looking at the impact of COVID-19 on mental health in the general population (January/February (T1) and February/March (T2) 2020) and found stable levels of psychological distress between T1 and T2. British longitudinal studies based on the same population before and after the outbreak of COVID-19 concluded that prolonged deterioration in mental health occurred after the COVID-19 outbreak [13,14].

However, most studies with a focus on mental health, quality of life and well-being due to the COVID-19 outbreak are cross-sectional studies that do not address how time may impact psychosocial health. Based on this and the lack of knowledge related to the importance of time on psychosocial health, we performed two consecutive cross-sectional studies that allowed for a comparison between timepoints using appropriate statistical tools.

### *Study Aim*

The aim of this study was to explore stability and change in mental health, quality of life, well-being and loneliness from the early stage and nine months after the COVID-19 pandemic outbreak as adjusted by demographic variables in two cross-national samples.

## **2. Material and Methods**

### *2.1. Design and Procedures*

The study used a cross-sectional survey design; the survey was distributed through social media in each of the involved countries in April (T1) and November (T2) 2020. The same survey was used at both timepoints with minor adjustments for the second administration of the survey [3,9]. The landing sites for the survey were established at the researchers' universities: OsloMet—Oslo Metropolitan University, Norway; University of Michigan, USA; Northumbria University, UK; and the University of Queensland, Australia. The initiator of the project was A.K.Ø.G. from OsloMet. Due to ethical considerations and permissions in each of the countries, each country had its own project lead. The survey was simultaneously codeveloped by the researchers in two languages: Norwegian and English. Language and cultural differences were considered during the survey development process, meaning that the countries' phrasing of each item conveyed the same meaning when considering culturally embedded meanings of words and phrases.

### *2.2. Inclusion and Exclusion*

To be included in the study, participants had to be 18 years or older, understand Norwegian or English, live in Norway, the USA, the UK or Australia and be able to

access the electronic survey. There were no exclusion criteria. Altogether, 7284 individuals participated by responding to the online form.

### 2.3. Measures

#### 2.3.1. Sociodemographic Characteristics and Time of Survey

Sociodemographic variables included country of residence (NO, USA, UK, AUS); gender (male (1), female (2)); age (18–24 years (1), 25–34 years (2), 35–44 years (3), 45–54 years (4), 55–64 years (5), 65–74 years (6), over 75 years (7)); highest completed level of education (Bachelor's degree or below (1), Master's/Doctoral degree (2)); civil status (living with a spouse or partner, no (0), yes (1)); having children, yes (1), no (0)); employment (full-time (1) or part-time employment (2)), place of employment/workplace (healthcare (1), trade industry (2), construction industry (3), education (4), public safety (5), other (6)) and living area (rural or farming area (1), town/suburb (2), city (3)) (Table 1).

**Table 1.** Overview of the sociodemographic characteristics.

Sociodemographic Characteristics	Answering Options
Country of residence	NO USA UK AUS
Gender	Women Men Other
Age	<24 years 25–34 years 35–44 years 45–54 years 55–64 years 65–74 years >75 years
Civil status	Married/cohabiting Not married/cohabiting
Having children	Yes No
Education	≤Bachelor ≥Master
Employment	Full-time Part time
Place of work	Healthcare Trade industry Education Public safety, police, fire services, etc. Other
Living area	Rural or farming area Town/suburb City

Time of survey included April 2020 (1) and November 2020 (2).

#### 2.3.2. Mental Health, Quality of Life, Well-Being and Loneliness

General Health Questionnaire 12 (GHQ-12) is widely used as a self-report measure of mental health [15–17]. A large number of studies in the general adult, clinical, work and student populations have provided support for its validity across samples and contexts [16,18–24] and translated from English to several other languages, including Norwegian [25,26]. Six items of the GHQ-12 are phrased positively (e.g., 'able to enjoy day-to-day

activities') and six—negatively (e.g., 'felt constantly under strain'). The person indicates the degree to which the item content was experienced during the two preceding weeks using four response categories ('less than usual' (0), 'as usual' (1), 'more than usual' (2) or 'much more than usual' (3)); score range, 0–36. Positively formulated items are recoded prior to analysis. Higher scores indicate poorer mental health (more psychological distress). Case-level scores (the person indicating 'more than usual' or 'much more than usual' on at least four of the 12 items) indicate a level of emotional distress where treatment may be needed [27]. Cronbach's alpha ranged between 0.88–0.92 in the current samples across the countries, and the Pearson correlation coefficient between the questions and the total value showed significance  $<0.05$  indicating the questionnaire's validity in the total sample and across the four national samples.

Cantril's self-anchoring ladder (CL) is a self-administered overall quality of life (QoL) questionnaire with one question, 'How is your life?', asking the person to rate his or her present experience of life on a scale anchored by their own identified values [28] and used when comparing satisfaction with life between groups and populations [29–33]. The response alternatives are between 0 and 10 with 0 = worst possible QoL and 10 = best possible QoL [28]. The cut-off score was chosen at 6 and above for good overall quality of life [28]. The CL has been reported to have good validity and stability and reasonable reliability [34–37].

Psychosocial well-being (PSW) assesses an individual's psychological experience of well-being and consists of ten items. The measure includes five positive and five negative statements with item scores ranging between 1 (best well-being) and 5 (worst well-being) [38]. Cronbach's alpha was excellent, ranging from 0.89–0.90 across the country samples. Validity tests in the current questionnaire indicated good validity and it is good evidence to propose a one-factor solution to the PSW scale as used in this study [9] and consistent with previous validation studies [38].

The Loneliness Scale [39] consists of six statements, all of which are rated from 0 ('totally disagree') to 4 ('totally agree'). It measures two different aspects of loneliness, 'emotional loneliness' and 'social loneliness'. Previous factor analyses found the six statements to load on two different factors, and therefore they should be treated as constituting two different scales reflecting the two different aspects of loneliness [39,40]. For both scales, the score range is 0–12 with higher scores indicating more loneliness. Using a one-factor solution to measure overall loneliness may also be appropriate (score range, 0–24), depending on the level of conceptual nuance required. In this study, only overall loneliness was included in the analysis. Higher sum scores indicated higher overall loneliness. Cronbach's alpha was good, 0.80 for overall loneliness.

#### 2.4. Statistical Analysis

Mental health, overall quality of life, psychosocial well-being and overall loneliness were calculated for each included demographic subgroup (gender, age, civil status, having children, education level, employment, workplace and place of living) within each country at each timepoint.

Depending on the number of group categories, group differences were examined using the one-way analysis of variance (ANOVA) and the independent *t*-test. The Pearson correlation coefficient *r* was used to assess the strength between categorical data.

In order to examine whether changes in mental health, overall quality of life, psychosocial well-being and loneliness were different between the four countries, the time  $\times$  country variable was computed in SPSS and entered as a predictor (Step 2) in the regression analysis. The sociodemographic variables gender, age, civil status, having children, education and employment were included in Step 1. In cases of significant interaction effects (time  $\times$  country), the subsequent regression analyses were performed separately for each country.

Generalized linear models (GLM) were used to conduct multiple linear regression analyses on psychosocial measures (GHQ, CL, PSW, Loneliness) by demographic variables

and time (November vs. April) in each country. Estimated marginal means were used to present mean responses by time after adjusting for all demographic variables in the model. Unstandardized beta (B) and standard error (ES) were reported for measures of the effect of associations between the independent and dependent variables. Statistical significance was set at  $<0.05$ . Missing values were handled by case-wise deletion.

### 2.5. Ethics

The data collected in this study were anonymous. The researchers adhered to all relevant regulations in their respective countries concerning ethics and data protection. The study was approved by OsloMet (20/03676) and the regional committees for medical and health research ethics (REK; ref. No. 132066) in Norway, reviewed by the University of Michigan's Institutional Review Board for Health Sciences and Behavioral Sciences (IRB HSBS) and designated as exempt (HUM00180296) in the USA, by Northumbria University Health Research Ethics Committee (HSR1920-080) in the UK, as well as in Australia (HSR1920-080 2020000956).

## 3. Results

### 3.1. Participants

The total sample from Survey 1 (T1) was 3810 [3,9], including 771 (20%) individuals from Norway, 1393 (37%) from the USA, 1373 (36%) from the UK and 210 (7%) from Australia [3,9].

The participants included in Survey 2 (T2) were 3474 individuals from Norway ( $n = 547$ , 16%), the USA ( $n = 2130$ , 61%), the UK ( $n = 640$ , 18%) and Australia ( $n = 157$ , 4%); altogether, 7284 individuals from both surveys. Overall, there were more women than men in the total sample (77% versus 20%, respectively) and the individuals who responded were spread across age groups. Sixty percent lived with a spouse or partner, 60% had children and 64% had a Bachelor's degree or lower levels of education. Sixty-eight percent had either a full-time or a part-time job, and of these individuals, 18% worked in healthcare, 4%—in trade industry, 14%—in education and 7%—in other critical functions in the society (public safety, police, fire services, etc.); 24% reported other areas of work. Eleven percent lived in rural areas, 51%—in a town or a suburb, while 38% lived in a city.

### 3.2. Mental Health, Overall Quality of Life, Well-Being and Loneliness at T1 and T2 Combined

Table 2 displays the levels of psychosocial health according to subgroups. Women reported a significantly higher level of emotional distress ( $p < 0.001$ ) and poorer well-being ( $p < 0.001$ ) than men. The lower the age of the individual, the higher the level of emotional distress ( $p < 0.001$ ), poorer quality of life ( $p < 0.001$ ) and well-being ( $p < 0.001$ ) and higher the feeling of loneliness ( $p < 0.001$ ). Participants living together with a spouse or partner reported less emotional distress ( $p < 0.001$ ), better quality of life ( $p < 0.001$ ) and well-being ( $p < 0.001$ ) and less loneliness ( $p < 0.001$ ) than their counterparts. The same overall pattern was found among those who had children ( $p < 0.001$  for all outcomes, respectively), had a Master's degree or higher education ( $p < 0.01$  for CL,  $p < 0.001$  for GHQ, PSW and Loneliness, respectively) and were in full-time employment ( $p < 0.001$  for GHQ and PSW,  $p < 0.01$  for Loneliness). People working in trade industry exhibited the highest level of emotional distress ( $p < 0.001$ ) and felt most lonely ( $p < 0.01$ ), while those living in rural or farming areas reported less emotional distress ( $p < 0.001$ ), better quality of life ( $p < 0.001$ ) and a higher level of well-being ( $p < 0.001$ ) than those living in towns/suburbs or cities.

**Table 2.** Mental health, overall well-being, quality of life and loneliness in sample subgroups nine months after the COVID-19 outbreak in the total sample from April and November 2020.

	N = 7284	GHQ		CL		PSW		Loneliness	
		Mean (SD)	p	Mean (SD)	p	Mean (SD)		Mean (SD)	
Gender									
Women	5580	16.8 (6.6)	<0.001	6.2 (2.2)	0.5	2.8 (0.8)	<0.001	10.5 (4.9)	<0.001
Men	1489	14.8 (7.1)		6.3 (2.5)		2.6 (0.9)		10.2 (5.3)	
Other	215	18.45 (8.2)		6.1 (2.2)		3.1 (0.9)		12.2 (5.3)	
Age									
<24 years	655	19.3 (7)	<0.001	5.8 (2.4)	<0.001	3.1 (0.8)	<0.001	11.8 (4.7)	<0.001
25–34 years	1478	18.1 (6.5)		5.9 (2.2)		2.9 (0.8)		10.7 (4.7)	
35–44 years	1327	16.9 (6.8)		6.1 (2.2)		2.8 (0.8)		10.3 (5.)	
45–54 years	1299	15.9 (6.6)		6.3 (2.3)		2.7 (0.8)		9.9 (5)	
55–64 years	1112	15.1 (6.6)		6.5 (2.3)		2.6 (0.8)		9.6 (4.8)	
65–74 years	830	13.4 (6.1)		6.8 (2.4)		2.4 (0.8)		9.1 (4.6)	
>75 years	182	12.8 (5.2)		6.6 (2.4)		2.3 (0.8)		8.6 (4.5)	
Civil status									
Married/cohabiting	4379	15.8 (6.5)	<0.001	6.4 (2.2)	<0.001	2.6 (0.8)	<0.001	9.7 (4.8)	<0.001
Not married/cohabiting	2638	17.3 (7.2)		5.9 (2.4)		3 (0.9)		11.6 (5.1)	
Having children									
Yes	4344	15.4 (6.5)	<0.001	6.5 (2.8)	<0.001	2.6 (0.8)	<0.001	9.8 (4.9)	<0.001
No	2940	17.6 (6.9)		5.9 (2.3)		3 (0.8)		10.7 (4.8)	
Education									
≤Bachelor	4644	16.8 (7.1)	<0.001	6.1 (2.4)	<0.01	2.8 (0.9)	<0.001	10.7 (4.9)	<0.001
≥Master	2570	15.7 (6.2)		6.4 (2.2)		2.6 (0.8)		9.3 (4.7)	
Employment									
Full-time	3580	16.2 (6.5)	0.001	6.2 (2.2)	0.9	2.7 (0.8)	<0.001	10.3 (4.9)	<0.01
Part time	1411	17.3 (6.7)		6.1 (2.3)		2.9 (0.8)		10.8 (5.1)	
Place of work									
Healthcare	882	16 (6)	<0.001	6.3 (2.3)	0.4	2.7 (0.8)	<0.05	9.5 (4.6)	<0.01
Trade industry	176	17.2 (7.4)		5.8 (2.6)		2.9 (0.9)		10.7 (5.2)	
Education	697	16.9 (6.2)		6.3 (2.1)		2.8 (0.7)		10.2 (4.9)	
Public safety, police, fire services, etc.	370	14.7 (6.2)		6.3 (2.2)		2.7 (0.8)		10.5 (5.2)	
Other	1379	16.7 (6.9)		6.2 (2.2)		2.8 (0.8)		10.4 (4.9)	
Living area									
Rural or farming area	794	15.4 (6.7)	<0.001	6.4 (2.3)	<0.001	2.6 (0.9)	<0.001	10 (5.2)	0.9
Town/suburb	3672	16.4 (6.8)		6.3 (2.3)		2.7 (0.8)		10.3 (4.8)	
City	2744	16.9 (6.9)		6.0 (2.3)		2.8 (0.8)		10 (4.9)	

Demographic variables



### 3.3. Mental Health, Overall Quality of Life, Well-Being and Loneliness in and between Countries and Time

The levels of emotional distress when measured as mean and standard deviation were higher in the Norwegian sample ( $p < 0.001$ ) and at the same level in the three other countries in November 2020 compared to April 2020 (Table 3). However, fewer individuals had scores equal to the case level in November compared to April in all the four countries ( $p < 0.001$ ). Alternatively, a higher number of individuals reported poorer quality of life on the case level for quality of life in November compared to April ( $p < 0.001$ , all countries), and in line with this, the mean scores of quality of life (CL) were significantly reduced in all countries ( $p < 0.001$ ) except the UK. Psychosocial well-being was reported significantly poorer in the UK and Norway in November compared to April ( $p < 0.001$ ), while the Australian and USA samples reported significantly better well-being ( $p < 0.05$  and  $p < 0.001$ , respectively) in the same timeframe (Table 3). All the countries' samples except the Australian one reported a significantly higher level of loneliness between the timepoints. Significant differences between the countries were found for all outcomes in both April and November.

### 3.4. Regression Analysis

When assessing whether changes in the outcomes across the two timepoints differed between the countries, regression analysis showed that the time by country interaction effects were significant for all outcomes (GHQ  $p < 0.001$ , CL  $p < 0.001$ , PSW  $p < 0.001$ , Loneliness  $p < 0.001$ ) when controlled for the included demographic variables as described above (data not shown).

### 3.5. Multiple Linear Regression

When the generalized linear models (GLM) were used to conduct multiple linear regression analyses on the psychosocial measures (GHQ, CL, PSW, Loneliness) by demographic variables and time (November vs. April) in each country, both demographic variables and time of survey explained a significant effect of the values of GHQ, CL, PSW and Loneliness (Table 4).

The analysis showed that age and civil status were the independent factors that showed the strongest unique effects across all the countries.

When controlling for all demographic variables, time had a significant contribution on the higher level of emotional distress and poorer well-being in the Norwegian sample ( $p < 0.01$  and  $p < 0.01$ , respectively) and of poorer well-being in the UK samples ( $p < 0.01$ ) (Table 4).

The estimated marginal mean responses by time after adjusting for all demographic variables in the model are shown in Figure 1.

**Table 3.** Comparison of mental health, overall quality of life, psychosocial well-being and loneliness between the national samples and between the countries in April and November 2020.

		Norway			USA			UK			Australia			Total	
		April N = 771	November N = 547	<i>p</i>	April N = 1393	November N = 2130	<i>p</i>	April N = 1373	November N = 640	<i>p</i>	April N = 273	November N = 157	<i>p</i>	N = 3810 Differences between countries April 2020	N = 3474 Differences between countries November 2020
GHQ	Mean (SD)	13.7 (6.5)	14.9 (7)	<0.001	16.3 (6)	16.4 (6.6)	0.10	18.5 (7.6)	18.6 (7)	0.5	16.1 (6.8)	15.2 (6.9)	0.16	<0.001	<0.001
	Case (N/%)	375/49	242/44	0.06	950/70	1033/56	<0.001	1015/74	373/68	<0.01	172/63	55/42	<0.001	<0.001	<0.001
CL	Mean (SD)	6.7 (1.9)	6.1 (2.4)	<0.001	6.6 (2.1)	6.3 (2.8)	<0.05	6.1 (2.3)	6 (2.2)	0.12	6.8 (2.2)	5.8 (2.4)	<0.001	<0.05	<0.05
	Case (N/%)	204/27	210/38	<0.001	414/30	1186/56	<0.001	498/36	324/51	<0.001	64/24	92/59	<0.001	<0.001	<0.001
PSW	Mean (SD)	2.2 (0.7)	2.6 (0.9)	<0.001	2.9 (0.4)	2.7 (0.9)	<0.001	2.8 (0.3)	3 (0.8)	<0.001	2.9 (0.3)	2.7 (0.9)	0.05	<0.001	<0.001
Loneli- ness	Overall loneliness (Mean/SD)	7.8 (4.6)	8.4 (5)	<0.01	10.2 (4.5)	10.8 (4.9)	<0.001	11.0 (4.8)	11.5 (4.5)	<0.05	9.4 (4.7)	9.7 (5.1)	0.5	<0.001	<0.001

Table 4. Multiple linear regression analysis in each country.

Psychosocial Variables by Demographic Variables and Time	Norway		USA		UK		AUS	
	B (SE)	p	B (SE)	p	B (SE)	p	B (SE)	p
<b>GHQ</b>								
Gender (women)	1.33 (0.67)	0.047	1.24 (0.4)	0.002	1.93 (0.65)	0.003	1.54 (1.24)	0.213
Age	−0.99 (0.24)	<0.001	−0.69 (0.14)	<0.001	−0.38 (0.21)	0.065	−0.26 (0.4)	0.514
Civil status (married/cohabiting)	−1.48 (0.59)	0.012	0.34 (0.38)	0.373	−2.37 (0.51)	<0.001	−1.52 (1.11)	0.17
Children (yes)	−1.09 (0.68)	0.113	−0.62 (0.39)	0.115	0.72 (0.77)	0.352	−2.01 (1.27)	0.113
Education (Master's or above)	0.89 (0.57)	0.12	−0.46 (0.36)	0.193	−0.3 (0.66)	0.653	−2.39 (1.06)	0.024
Employment (part-time)	1.16 (0.63)	0.066	0.28 (0.43)	0.514	0.81 (0.54)	0.133	2.29 (0.97)	0.018
Place of work (healthcare as ref.)								
Other	1.08 (0.71)	0.13	0.58 (0.46)	0.204	1.02 (0.6)	0.086	1.71 (1.11)	0.126
Public safety, police, fire services, etc.	0.35 (1.09)	0.75	−0.16 (0.62)	0.802	0.57 (0.77)	0.463	0.5 (1.5)	0.738
Education	0.54 (0.88)	0.536	0.98 (0.53)	0.066	0.86 (0.83)	0.301	4.13 (1.89)	0.029
Trade industry	−1 (1.13)	0.376	−0.55 (0.72)	0.447	2.77 (1.1)	0.012	1.56 (1.95)	0.426
Living area (rural as ref.)	0.11 (0.44)	0.8	0.25 (0.27)	0.367	0.3 (0.36)	0.399	0.27 (1.06)	0.798
Time (November vs. April)	1.83 (0.68)	0.007	0.02 (0.42)	0.962	1.15 (0.68)	0.09	−2.06 (1.14)	0.072
<b>CL</b>								
Gender (women)	−0.1 (0.26)	0.714	−0.17 (0.16)	0.267	−0.14 (0.21)	0.511	1.05 (0.44)	0.016
Age	0.29 (0.09)	0.003	0.11 (0.06)	0.039	0.03 (0.07)	0.674	0.01 (0.14)	0.933
Civil status (married/cohabiting)	0.66 (0.23)	0.005	−0.15 (0.15)	0.331	0.59 (0.17)	0.001	0.48 (0.39)	0.227
Children (yes)	0.34 (0.27)	0.213	0.2 (0.16)	0.193	0.15 (0.25)	0.55	0.01 (0.45)	0.994
Education (Master's or above)	0.11 (0.22)	0.628	−0.09 (0.14)	0.53	−0.05 (0.22)	0.821	−0.03 (0.38)	0.94
Employment (part-time)	0.06 (0.25)	0.806	0.36 (0.17)	0.035	0.14 (0.18)	0.43	0 (0.34)	0.993
Place of work (healthcare as ref.)								
Other	−0.25 (0.28)	0.381	−0.19 (0.19)	0.305	−0.16 (0.2)	0.426	0.39 (0.39)	0.322
Public safety, police, fire services, etc.	−0.24 (0.43)	0.58	−0.46 (0.24)	0.059	0.02 (0.26)	0.943	0.47 (0.53)	0.377
Education	−0.06 (0.34)	0.861	−0.09 (0.21)	0.684	−0.19 (0.27)	0.479	−0.52 (0.67)	0.434
Trade industry	0.58 (0.44)	0.184	−0.48 (0.29)	0.102	−0.48 (0.36)	0.18	0.74 (0.7)	0.295
Living area (rural as ref.)	0 (0.18)	0.996	0.05 (0.11)	0.626	−0.09 (0.12)	0.456	0.16 (0.37)	0.672
Time (November vs. April)	0.09 (0.29)	0.756	0.12 (0.17)	0.463	0.12 (0.22)	0.599	−0.61 (0.41)	0.134

Table 4. Cont.

Psychosocial Variables by Demographic Variables and Time	Norway		USA		UK		AUS	
	B (SE)	p	B (SE)	p	B (SE)	p	B (SE)	p
<b>PSW</b>								
Gender (women)	0.15 (0.08)	0.065	0.12 (0.05)	0.012	0.13 (0.07)	0.082	0.18 (0.15)	0.214
Age	−0.1 (0.03)	0.001	−0.11 (0.02)	<0.001	−0.06 (0.02)	0.007	−0.07 (0.05)	0.113
Civil status (married/cohabiting)	−0.34 (0.07)	<0.001	−0.14 (0.05)	0.003	−0.46 (0.06)	<0.001	−0.26 (0.13)	0.043
Children (yes)	−0.17 (0.08)	0.038	−0.1 (0.05)	0.034	0.03 (0.09)	0.761	−0.24 (0.15)	0.101
Education (Master's or above)	−0.04 (0.07)	0.572	−0.09 (0.04)	0.055	−0.04 (0.07)	0.621	−0.28 (0.12)	0.027
Employment (part-time)	0.14 (0.08)	0.076	0.06 (0.05)	0.239	0.08 (0.06)	0.183	0.28 (0.11)	0.014
Place of work (healthcare as ref.)								
Other	0.1 (0.09)	0.26	0.07 (0.06)	0.193	0.09 (0.07)	0.192	0.05 (0.13)	0.724
Public safety, police, fire services, etc.	0.11 (0.13)	0.395	0.08 (0.08)	0.28	0.01 (0.09)	0.897	−0.12 (0.18)	0.481
Education	0.02 (0.11)	0.857	0.14 (0.07)	0.031	0.04 (0.09)	0.704	0.46 (0.22)	0.04
Trade industry	−0.01 (0.14)	0.918	0.06 (0.09)	0.505	0.21 (0.12)	0.096	0.21 (0.23)	0.356
Living area (rural as ref.)	−0.02 (0.05)	0.691	0.01 (0.03)	0.78	−0.01 (0.04)	0.804	−0.05 (0.12)	0.715
Time (November vs. April)	0.23 (0.08)	0.006	−0.01 (0.05)	0.889	0.19 (0.08)	0.011	−0.31 (0.13)	0.02
<b>Loneliness</b>								
Gender (women)	0.47 (0.49)	0.337	−0.13 (0.31)	0.664	−0.43 (0.41)	0.297	−0.77 (0.87)	0.376
Age	−0.36 (0.18)	0.045	−0.33 (0.11)	0.003	−0.2 (0.13)	0.131	−0.26 (0.28)	0.35
Civil status (married/cohabiting)	−2.23 (0.43)	< 0.001	−0.86 (0.3)	0.004	−2.41 (0.32)	< 0.001	−0.76 (0.78)	0.327
Children (yes)	−0.84 (0.5)	0.097	−0.26 (0.31)	0.396	−0.56 (0.5)	0.264	−0.04 (0.89)	0.962
Education (Master's or above)	−0.08 (0.42)	0.847	−0.79 (0.28)	0.004	0.04 (0.42)	0.915	−1.66 (0.74)	0.026
Employment (part-time)	0.54 (0.46)	0.242	0.2 (0.34)	0.552	0.75 (0.35)	0.031	0.68 (0.68)	0.314
Place of work (healthcare as ref.)								
Other	0.56 (0.52)	0.284	0.43 (0.36)	0.231	0.43 (0.38)	0.257	0.35 (0.78)	0.657
Public safety, police, fire services, etc.	0.93 (0.8)	0.245	0.68 (0.49)	0.165	0.47 (0.49)	0.339	0.01 (1.05)	0.997
Education	−0.28 (0.65)	0.666	0.54 (0.41)	0.193	0.36 (0.53)	0.501	3.12 (1.32)	0.018
Trade industry	−0.57 (0.84)	0.497	0.03 (0.57)	0.96	1.35 (0.71)	0.057	3.94 (1.37)	0.004
Living area (rural as ref.)	−0.16 (0.32)	0.612	−0.1 (0.21)	0.627	−0.20 (0.23)	0.381	0.36 (0.74)	0.627
Time (November vs. April)	0.71 (0.5)	0.157	0.34 (0.33)	0.301	0.64 (0.44)	0.145	−1.07 (0.8)	0.18






Figure 1. Cont.



**Figure 1.** Adjusted means. See Section 2.3.2. Mental health, quality of life, well-being and loneliness in the measurement units used.

#### 4. Discussion

The most important findings in this paper were that the levels of emotional distress, quality of life, well-being and loneliness either got poorer or were stable but at a poor level over time. The Norwegian sample reported a higher level of emotional distress, decreased quality of life, poorer well-being and higher level of loneliness over the time on the group level. All outcome variables were mainly stable in the three other countries in the same time period. Nonetheless, the Norwegian sample had significantly better scores in all outcome measures than the comparing countries.

In ordinary times, a high level of emotional distress (GHQ) may indicate reduced functional impairment equal to a mental disorder. However, in this particular study, it is clear that these findings are associated with the ongoing COVID-19 pandemic. Even if the level of caseness is reduced compared to the reports in April 2020, it is still much higher than the 20% within a year in all countries due to mental distress before the COVID-19 outbreak [41–45]. It is of interest though that the caseness is reduced throughout the samples. Even if it is to be assumed that social distancing still has an important impact on mental health, it might be that people have become more used to the situation and are adapting to using different coping strategies.

According to the OECD [46], the average level of quality of life for western countries, including those in this study, was 7.4 in 2018 as measured with CL. However, life is dynamic and changing, and it seems that the situation has gone from better to worse not only when compared to the reported levels in general populations in 2018, but also between the results in April and November 2020 in our study. While 27% (Norway), 30% (USA), 36% (UK) and 24% (Australia) reported caseness for poorer quality of life in April, the findings for the same countries were 38%, 56%, 51% and 59%, respectively, in November. A cut-off score  $\leq 5$  is defined as suffering due to the experienced overall quality of life [28], meaning the experienced overall quality of life is moderate or inconsistent. This means that people are struggling in their present situation and may have negative thoughts and views about the future. The pandemic's social distancing protocols had been in place for several months by the Survey 2 measurement, and the reduced quality of life reported reflects individuals struggling with uncertainty about the future. The reduced quality of life may be associated with the social distancing protocols that reduced social relationships and meetings with colleagues and friends outside one's household, promoted engagement in remote work and implemented limitations on travelling. On the other hand, participants might have gotten accustomed to the changed situation that in some ways may have reduced the initial heavy burden of mental distress experienced early in the pandemic. In line with this, the difference between the countries when it comes to reported well-being might be understood as an adaptation to the ongoing pandemic. At the same time, it is to be expected that most people will over time return to what was known as a normal level of both mental health and experienced quality of life.

The fact that the Norwegian sample reported a higher level of emotional distress in November than in April does not remove the significant differences between the countries in November. While the Norwegian sample still reported significantly less emotional distress than the other countries, it might have been attributed to cultural differences and the level of supports provided in the welfare systems. Differences in perceived trust in how political systems, government and health authorities handle the pandemic and meet challenges due to the COVID-19 pandemic may also contribute to these results similar to what is described in previous research [47].

Prior research, e.g., by Rosenfiled et al. [48], underscored that the gender status does not necessarily have a significant impact on overall mental health, but that it may be related to different kinds of mental health issues. Women, for example, report more internalizing diseases like depression and anxiety, while men are more likely to report more externalizing disorders like antisocial behavior and drug abuse [48,49]. These prior findings are consistent with the current study since the GHQ measures of psychological distress include anxiety and depression. In addition, our findings are in line with the

review study based on 19 cross-sectional studies measuring mental health related to gender in the time of COVID-19 [5]. The review study underscores that identifying as female is associated with more vulnerability for mental distress than identifying as male.

Age and civil status in particular were variables associated with significant contribution to emotional distress, quality of life, well-being and loneliness across the countries. Higher age seems to be a buffer against higher emotional distress, poorer quality of life, well-being and loneliness. The impact on mental health seems to take a higher toll on younger people and this finding is in line with research that points out that younger people have more negative emotions and experience stressful events as more unpleasant than older people [6,7], and older people in general score lower on depression and anxiety [8]. An explanation for the results in the current study might be that young people normally look forward, and in the present situation they may feel their daily lives and the future are set on hold or crumbling away due to loss of social interactions. According to Myklestad et al. [50], support from friends is generally associated with lower levels of distress among adolescents. Older people, on the other hand, need, perhaps, to achieve less, they can look back to good memories and joy and know through life experience that most stressful times will end, even if it takes time. Older people have a variety of tools in the toolbox to manage a particular situation of stress when compared to the younger population. Another possible explanation might be that in the countries studied, older people are the first in line for vaccination and on a faster track to resuming a normal life. However, pre-pandemic studies suggested that social isolation and older age were associated with loneliness [51], which is seen as a key public health issue [52,53]. On the other hand, previous research also showed that the number of contacts is not important for psychological distress among elderly people, but social support is associated with decreased emotional distress [54].

Living with a spouse or a partner was associated with less emotional distress, better quality of life and well-being and less loneliness and could simply be explained by the support of having a close person to confide in, and that living with a spouse or a partner is a buffer against psychosocial stress. This study did not examine mental health with regard to living with other people or a pet. It is to be assumed that for some of those not living with a spouse or a partner living with someone or having a pet would be of big importance.

Respondents with a Bachelor's degree or below exhibited higher levels of emotional distress, poorer quality of life and well-being and more loneliness than those with a Master's degree or higher education. However, when controlled for other demographic variables, education had a significant contribution to mental health, well-being and loneliness in the Australian sample, and to well-being and loneliness in the USA sample. While education did not contribute to any significant outcomes in the Norwegian sample, it could be explained by an overall higher educational level in the Norwegian population than, for example, in the sample from the USA. An alternative speculation may be that health inequalities by education level are less pronounced in Norway. On the general level, education is assumed to buffer the effect of a disability, in this case associated with a global situation. A reason for this might be the more educated are more equipped, including cognitive skills, to cope with the consequences of a particular cause of stress [55].

The individuals who were employed full-time exhibited less emotional distress, better well-being and less loneliness than those who were employed part-time. In the multiple hierarchical regression, employment had a significant unique contribution to reduced emotional distress in all the countries over time, which is in line with former cross-sectional studies [3] and former studies underscoring that employment, in general, is associated with improved mental health and quality of life [56]. Having a job often means that people have a social network and experience social inclusion through their work. In line with the findings, it might also be understood that higher levels of employment may result in better psychosocial health over time.

Healthcare workers exhibited a high level of emotional distress, reduced well-being and loneliness, which is consistent with former studies [4], but in this study, even higher scores on these measures were reported by those employed in the trade industry. While



healthcare workers are constantly exposed to the increased risk of physical contact with infections and have had to take greater precautions and follow strict infection control protocols, trade workers may not have had adequate infection control equipment and supports apart from hand washing, antibacterial cleaning practices in the work setting, masks and keeping distance. Another reason for the higher scores in the trade industry may be associated with the uncertainty about the potential loss of jobs due to closedowns. Teachers, which is another group with high scores, have very limited possibilities for keeping distance in the workplace and may have increased concerns of infection, sickness and quarantine.

Interestingly, living in a rural or farming area had a positive impact on emotional distress, quality of life and well-being. This does not necessarily mean that there were no infections in these areas, but it may suggest that the toll of the pandemic was smaller in rural areas than in suburbs or cities

Time contributed significantly to emotional distress (GHQ). This may be understood as the Norwegian sample was the only country with significantly different scores between April and November 2020. In the countries with higher but stable scores, time had no significant impact. On the other hand, it is to be concluded that, overall, time did not contribute to lowered levels and reduce the toll on mental health, quality of life, well-being or loneliness in any of the four countries.

The pandemic is still causing social distance, quarantine, and isolation protocols in many places around the world. More research is needed on how COVID-19-related risk variables and concerns are associated with time so that appropriate support systems could be put in place to improve public psychosocial health. In light of this, health and social workers should be aware of the importance of using professional skills to promote better psychosocial health in meeting individuals with needs, such as young people and those living alone. For example, and as underscored by Dorado et al. [57], social work interventions are thought to have a key role when it comes to improving psychosocial factors such as well-being among individuals suffering from COVID-19-produced consequences. Further research could have a more specific focus on how health workers and social workers engage people in such a need. Understanding how risk factors present during the COVID-19 pandemic are associated with an individual's mental health, quality of life, well-being and loneliness warrants future research as well.

#### *Study Strength and Limitations*

The strength of this study was the cross-national comparison and the relatively high number of respondents. However, some limitations should be taken into consideration. First, the data were collected using cross-sectional online surveys, which means assumptions about causations cannot be made, only associations. Secondly, the respondents received invitations to participate through social media, and the responses thereby do not include individuals that do not use social media. This may limit the ability to generalize the results to the general population in the four countries included in the study. The sample included respondents from all age groups, but fewer from the oldest age group, as well as more women than men. This may represent a skewed population. To avoid this, in future studies, more sophisticated sampling methods could be used. Third, our findings imply that social distancing policies are significant for psychological distress, quality of life, well-being and loneliness. The degree of disease outbreak and social distancing policies differed between the countries and the states in, for example, the USA and may have included additional factors on individual levels, as well as on the regional and country level. For example, the restriction levels in the four countries at T1 and T2 varied between moderate and high levels, including closures of schools and universities, variations of "stay at home" orders and closedowns, restrictions regarding gatherings of people from different households, closed borders, directions to stay at home other than for essential reasons and no unnecessary travel. Both in April and November, the number of infected people and deaths varied between the countries with the USA and the UK having higher rates than

Norway and Australia, and both surveys, T1 and T2, took place before the vaccination rollouts. These issues could have influenced the respondents' answers.

## 5. Conclusions

Overall, significant differences in psychosocial health across Norway, the USA, the UK and Australia emerged in this study. Individuals in the younger age groups struggled more than the older groups with mental health, quality of life, well-being and loneliness. Being female, not married or cohabiting, lower level of education, unemployed, working in healthcare or the trade industry and living in urban areas are all factors significantly associated with poorer psychosocial health across the countries. However, when adjusting for months into the pandemic (time) and other demographic variables, age and living together with a spouse or a partner contributed most to the psychosocial outcomes in all the countries. Differences in mental health, quality of life, well-being and loneliness were found between the countries in both April and November 2020. The toll of the COVID-19 pandemic on mental health did not reduce in any of the four countries between April and November 2020.

**Author Contributions:** Conceptualization, A.K.Ø.G., D.P., M.S., H.T., M.R., J.L. and T.B.; methodology, A.K.Ø.G., D.P., M.S., H.T., M.R., J.L. and T.B.; validation, A.K.Ø.G., D.P., M.S., H.T., M.R., J.L. and T.B.; formal analysis, A.K.Ø.G., T.B. and J.L.; investigation, A.K.Ø.G., D.P., M.S., H.T., M.R., J.L. and T.B.; data curation, A.K.Ø.G.; writing—original draft preparation, A.K.Ø.G.; writing—review and editing, A.K.Ø.G., D.P., M.S., H.T., M.R., J.L. and T.B.; visualization, A.K.Ø.G.; project administration, A.K.Ø.G. All authors have read and agreed to the published version of the manuscript.

**Funding:** The research did not receive external funding.

**Institutional Review Board Statement:** The study was conducted according to guidelines of the Declaration of Helsinki and approved by the Regional Committee for Medical and Health Research Ethics in Norway (#132066, April 2020).

**Informed Consent Statement:** Informed consent was obtained from all individuals involved in the study.

**Data Availability Statement:** The data presented in this study are available upon request from the corresponding author on completion of the research. The data are not published/available due to ongoing publications stemming from the project and different data protection regulations in the four countries involved in the study.

**Acknowledgments:** The authors thank all the respondents for their participation.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. World Health Organization. *Coronavirus Disease (COVID-19) Advice for the Public*; WHO: Geneva, Switzerland, 2021.
2. Williamson, B.; Eynon, R.; Potter, J. *Pandemic Politics, Pedagogies and Practices: Digital Technologies and Distance Education during the Coronavirus Emergency*; Taylor & Francis: Abingdon, UK, 2020.
3. Ruffolo, M.; Price, D.; Schoultz, M.; Bonsaksen, T.; Thygesen, H.; Geirdal, A.Ø. Employment Uncertainty and Mental Health during the COVID-19 Pandemic Initial Social Distancing Implementation: A Cross-National Study. *Glob. Soc. Welf.* **2021**, 1–10. [[CrossRef](#)]
4. The Lancet. COVID-19: Protecting health-care workers. *Lancet* **2020**, 395, 922. [[CrossRef](#)]
5. Xiong, J.; Lipsitz, O.; Nasri, F.; Lui, L.M.W.; Gill, H.; Phan, L.; Chen-Li, D.; Iacobucci, M.; Ho, R.; Majeed, A.; et al. Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *J. Affect. Disord.* **2020**, 277, 55–64. [[CrossRef](#)] [[PubMed](#)]
6. Carstensen, L.L.; Pasupathi, M.; Mayr, U.; Nesselroade, J.R. Emotional experience in everyday life across the adult life span. *J. Personal. Soc. Psychol.* **2000**, 79, 644. [[CrossRef](#)]
7. Neubauer, A.B.; Smyth, J.M.; Sliwinski, M.J. Age differences in proactive coping with minor hassles in daily life. *J. Gerontol. Ser. B* **2019**, 74, 7–16. [[CrossRef](#)] [[PubMed](#)]
8. Löwe, B.; Wahl, I.; Rose, M.; Spitzer, C.; Glaesmer, H.; Wingenfeld, K.; Schneider, A.; Brähler, E. A 4-item measure of depression and anxiety: Validation and standardization of the Patient Health Questionnaire-4 (PHQ-4) in the general population. *J. Affect. Disord.* **2010**, 122, 86–95. [[CrossRef](#)] [[PubMed](#)]

9. Geirdal, A.Ø.; Ruffolo, M.; Leung, J.; Thygesen, H.; Price, D.; Bonsaksen, T.; Schoultz, M. Mental health, quality of life, wellbeing, loneliness and use of social media in a time of social distancing during the COVID-19 outbreak: A cross country comparative study. *J. Ment. Health* **2021**. [CrossRef]
10. Vindegaard, N.; Benros, M.E. COVID-19 pandemic and mental health consequences: Systematic review of the current evidence. *Brain Behav. Immun.* **2020**, *89*, 531–542. [CrossRef]
11. Bonsaksen, T.; Heir, T.; Schou-Bredal, I.; Ekeberg, Ø.; Skogstad, L.; Grimholt, T.K. Post-Traumatic Stress Disorder and Associated Factors during the Early Stage of the COVID-19 Pandemic in Norway. *Int. J. Environ. Res. Public Health* **2020**, *17*, 9210. [CrossRef]
12. Wang, C.; Pan, R.; Wan, X.; Tan, Y.; Xu, L.; McIntyre, R.S.; Choo, F.N.; Tran, B.; Ho, R.; Sharma, V.K.; et al. A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav. Immun.* **2020**, *87*, 40–48. [CrossRef]
13. Daly, M.; Sutin, A.R.; Robinson, E. Longitudinal changes in mental health and the COVID-19 pandemic: Evidence from the UK Household Longitudinal Study. *Psychol. Med.* **2020**, 1–10. [CrossRef] [PubMed]
14. Niedzwiedz, C.L.; Green, M.J.; Benzeval, M.; Campbell, D.; Craig, P.; Demou, E.; Leyland, A.; Pearce, A.; Thomson, R.; Whitley, E.; et al. Mental health and health behaviours before and during the initial phase of the COVID-19 lockdown: Longitudinal analyses of the UK Household Longitudinal Study. *J. Epidemiol. Community Health* **2021**, *75*, 224–231.
15. Goldberg, D.P.; Gater, R.; Sartorius, N.; Ustun, T.B.; Piccinelli, M.; Gureje, O.; Rutter, C. The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychol. Med.* **1997**, *27*, 191–197. [CrossRef] [PubMed]
16. Goodwin, L.; Ben-Zion, I.; Fear, N.T.; Hotopf, M.; Stansfeld, S.A.; Wessely, S. Are reports of psychological stress higher in occupational studies? A systematic review across occupational and population based studies. *PLoS ONE* **2013**, *8*, e78693. [CrossRef] [PubMed]
17. Hankins, M. The reliability of the twelve-item general health questionnaire (GHQ-12) under realistic assumptions. *BMC Public Health* **2008**, *8*, 355. [CrossRef] [PubMed]
18. Aalto, A.-M.; Elovainio, M.; Kivimäki, M.; Uutela, A.; Pirkola, S. The Beck Depression Inventory and General Health Questionnaire as measures of depression in the general population: A validation study using the Composite International Diagnostic Interview as the gold standard. *Psychiatry Res.* **2012**, *197*, 163–171. [CrossRef] [PubMed]
19. Adlaf, E.M.; Gliksman, L.; Demers, A.; Ma, B.N.-T. The Prevalence of Elevated Psychological Distress Among Canadian Undergraduates: Findings from the 1998 Canadian Campus Survey. *J. Am. Coll. Health* **2001**, *50*, 67–72. [CrossRef]
20. Donath, S. The validity of the 12-item General Health Questionnaire in Australia: A comparison between three scoring methods. *Aust. N. Z. J. Psychiatry* **2001**, *35*, 231–235. [CrossRef]
21. Firth, J. Levels and sources of stress in medical students. *Br. Med. J.* **1986**, *292*, 1177–1180. [CrossRef]
22. Gorter, R.; Freeman, R.; Hammen, S.; Murtooma, H.; Blinkhorn, A.; Humphris, G. Psychological stress and health in undergraduate dental students: Fifth year outcomes compared with first year baseline results from five European dental schools. *Eur. J. Dent. Educ.* **2008**, *12*, 61–68. [CrossRef]
23. Malt, U.F. The validity of the General Health Questionnaire in a sample of accidentally injured adults. *Acta Psychiatr. Scand.* **1989**, *80*, 103–112. [CrossRef]
24. Nerdrum, P.; Rustøen, T.; Rønnestad, M.H. Student Psychological Distress: A psychometric study of 1750 Norwegian 1st-year undergraduate students. *Scand. J. Educ. Res.* **2006**, *50*, 95–109. [CrossRef]
25. Hystad, S.W.; Johnsen, B.H. The Dimensionality of the 12-Item General Health Questionnaire (GHQ-12): Comparisons of Factor Structures and Invariance Across Samples and Time. *Front. Psychol.* **2020**, *11*. [CrossRef]
26. Malt, U.F.; Mogstad, T.E.; Refnir, I.B. Goldberg's General Health Questionnaire. *Tidsskr. Nor. Laegeforen.* **1989**, *109*, 1391–1394. [PubMed]
27. Goldberg, D.; Oldehinkel, T.; Ormel, J. Why GHQ threshold varies from one place to another. *Psychol. Med.* **1998**, *28*, 915–921. [CrossRef] [PubMed]
28. Cantril, H. *The Pattern of Human Concerns*; Rutgers University Press: New Brunswick, NJ, USA, 1965.
29. Aasprang, A.; Andersen, J.R.; Våge, V.; Kolotkin, R.L.; Natvig, G.K. Psychosocial functioning before and after surgical treatment for morbid obesity: Reliability and validation of the Norwegian version of obesity-related problem scale. *PeerJ* **2015**, *3*, e1275. [CrossRef]
30. GallupWorldPoll. Country Averages of Self-Reported Life Satisfaction (Question: Cantril Ladder). 2017. Available online: <http://worldhappiness.report/> (accessed on 11 June 2020).
31. Mazur, J.; Szkulciecka-Dębek, M.; Dzielska, A.; Drozd, M.; Małkowska-Szkućnik, A. What does the Cantril Ladder measure in adolescence? *Arch. Med. Sci. AMS* **2018**, *14*, 182. [CrossRef] [PubMed]
32. Ortiz-Ospina, E.; Roser, M. Happiness and Life Satisfaction. 2017. Available online: <https://ourworldindata.org/happiness-and-life-satisfaction> (accessed on 11 June 2020).
33. Steptoe, A.; Deaton, A.; Stone, A.A. Subjective wellbeing, health, and ageing. *Lancet* **2015**, *385*, 640–648. [CrossRef]
34. Atkinson, T. The stability and validity of quality of life measures. *Soc. Sci. Med.* **1982**, *10*, 113–132. [CrossRef]
35. Geirdal, A.Ø.; Dheyauldeen, S.; Bachmann-Harildstad, G.; Heimdal, K. Quality of life in patients with hereditary hemorrhagic telangiectasia in Norway: A population based study. *Am. J. Med. Genet. Part A* **2012**, *158*, 1269–1278. [CrossRef] [PubMed]
36. Jenkins, L.S.; Brodsky, M.; Schron, E.; Chung, M.; Rocco, T., Jr.; Lader, E.; Constantine, M.; Sheppard, R.; Holmes, D.; Mateski, D.; et al. Quality of life in atrial fibrillation: The atrial fibrillation follow-up investigation of rhythm management (AFFIRM) study. *Am. Heart J.* **2005**, *149*, 112–120. [PubMed]

37. Levin, K.A.; Currie, C. Reliability and validity of an adapted version of the Cantril Ladder for use with adolescent samples. *Soc. Indic. Res.* **2014**, *119*, 1047–1063. [CrossRef]
38. Kaasa, S.; Mastekaasa, A.; Naess, S. Quality of Life of Lung Cancer Patients in a Randomized Clinical Trial Evaluated by a Psychosocial Well-Being Questionnaire. *Acta Oncol.* **1988**, *27*, 335. [CrossRef] [PubMed]
39. Gierveld, J.D.J.; Tilburg, T.V. A 6-item scale for overall, emotional, and social loneliness: Confirmatory tests on survey data. *Res. Aging* **2006**, *28*, 582–598. [CrossRef]
40. Bonsaksen, T.; Opseth, T.M.; Misund, A.R.; Geirdal, A.Ø.; Orsolya Reka, F. The De Jong Gierveld Loneliness Scale Used with Norwegian Clubhouse Members: Psychometric Properties and Associated Factors. *Int. J. Psychosoc. Rehabil.* **2019**, *22*, 88–100.
41. Folkehelseinstituttet (FHI). Psykiske Lidelser hos Voksne (Mental Illness among Adults). 2018. Available online: <https://www.fhi.no/nettpub/hin/psykisk-helse/psykiske-lidelser-voksne/> (accessed on 11 June 2020).
42. Mental Health First Aid (MHFA). Mental Health Statistics. 2020. Available online: <https://mhfaengland.org/mhfa-centre/research-and-evaluation/mental-health-statistics/> (accessed on 13 June 2020).
43. National Alliance on Mental Illness (NAMI). Mental Health by the Numbers. 2020. Available online: <https://nami.org/mhstats> (accessed on 13 June 2020).
44. Parliament of Australia. Mental Health in Australia: A Quick Guide. 2019. Available online: [https://www.aph.gov.au/About\\_Parliament/Parliamentary\\_Departments/Parliamentary\\_Library/pubs/rp/rp1819/Quick\\_Guides/MentalHealth](https://www.aph.gov.au/About_Parliament/Parliamentary_Departments/Parliamentary_Library/pubs/rp/rp1819/Quick_Guides/MentalHealth) (accessed on 13 June 2020).
45. Office for national Statistics. Coronavirus and the Social Impacts on Great Britain: 16 April 2020. Available online: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthandwellbeing/bulletins/coronavirusandthesocialimpactsongreatbritain/16april2020> (accessed on 13 June 2020).
46. OECD. *How's Life 2020. Measuring Well-Being*; OECD Publishing: Paris, France, 2020.
47. Liao, J.Q.; Fielding, R. Uncertain news: Trust and preventive practices in respiratory infectious diseases. *Eur. Psychol.* **2014**, *19*, 4. [CrossRef]
48. Rosenfield, S.; Lennon, M.C.; White, H.R. The self and mental health: Self-salience and the emergence of internalizing and externalizing problems. *J. Health Soc. Behav.* **2005**, *46*, 323–340. [CrossRef]
49. Rosenfield, S.; Mouzon, D. Gender and Mental Health. In *Handbook of the Sociology of Mental Health*; Springer: Berlin/Heidelberg, Germany, 2013; pp. 277–296.
50. Myklestad, I.; Røysamb, E.; Tambs, K. Risk and protective factors for psychological distress among adolescents: A family study in the Nord-Trøndelag Health Study. *Soc. Psychiatry Psychiatr. Epidemiol.* **2012**, *47*, 771–782. [CrossRef]
51. Berg-Weger, M.; Morley, J.E. Loneliness in Old Age: An Unaddressed Health Problem. *J. Nutr. Health Aging* **2020**, *24*, 243–245. [CrossRef]
52. Hebblethwaite, S.; Young, L.; Martin Rubio, T. Pandemic Precarity: Aging and Social Engagement. *Leis. Sci.* **2020**, 1–7. [CrossRef]
53. Holt-Lunstad, J. The potential public health relevance of social isolation and loneliness: Prevalence, epidemiology, and risk factors. *Public Policy Aging Rep.* **2017**, *27*, 127–130. [CrossRef]
54. Cairney, J.; Krause, N. The social distribution of psychological distress and depression in older adults. *J. Aging Health* **2005**, *17*, 807–835. [CrossRef] [PubMed]
55. Drapeau, A.; Marchand, A.; Beaulieu-Prévost, D. Epidemiology of Psychological Distress. In *Mental Illnesses—Understanding, Prediction and Control*; L'Abate, L., Ed.; InTech Europe: Rijeka, Croatia, 2012.
56. Evans, J.; Repper, J. Employment, social inclusion and mental health. *J. Psychiatr. Ment. Health Nurs.* **2000**, *7*, 15–24. [CrossRef] [PubMed]
57. Barbé, A.D.; Viejo, J.M.P.; Rodríguez-Brioso, M.D.M.; Gallardo-Peralta, L.P. Emotional well-being and resilience during the COVID-19 pandemic: Guidelines for social work practice. *Int. Soc. Work* **2021**. [CrossRef]