
6 Conclusions

The results from this work illustrate the complex wave generation and propagation processes that occur across the NW Mediterranean and the complexity of the regional models, and lead the way to further improvements on regional wave modelling.

As described in Chapter 2, coastal wave storms in the region were confirmed to be highly affected by the spatial and temporal variability of the local wind fields. Isolated H_s peaks were registered at some of the instruments along the transect but not at all of them, and important differences in PWD were also identified, all related to the patterns of the wind field. Bimodal events were also frequent. Even though integrated parameters follow the theoretical growth in terms of H_s and T_p increasing with fetch, they might also be affected by the high spatial and temporal variability of the wind and wave fields, and be a source of error when modelling wind-waves in the area.

In Chapter 3, the local features and the high spatial and temporal variability of the wind fields were signalled to be responsible for the large inaccuracies of the operational wave predictions (e.g. up to -50% under-prediction of H_s at the storm peak). Wave modelling was consequently expected to improve by increasing the temporal and spatial resolution of atmospheric and wave models in the region. The study indicated that bimodal sea conditions could play an important role in this region, although its consequences were unfortunately not fully understood yet.

In Chapter 4, it is shown that during a particular storm event, under fetch-limited growth conditions, wind and wave models with higher spatial and temporal resolutions improved H_s predictions: the storm peaks were properly resolved and their magnitude more closely approximated. However, H_s was still under-predicted by up to 1m (30%), and the reason could not be attributed to the modelled wind fields, which were over-predicted (35%). T_p was also under-predicted by about 25%. The results pointed to an inaccurate parameterization of the transfer of energy from the wind to the waves.

Finally, in Chapter 5, the wind fields were further studied. The wind input was modified by (1) using wind observations, (2) forcing the wave model with the maximum WS along the fetch, (3) enhancing the WS by a factor of 1.1, and (3) using growth and sink expressions for which C_D increases with rougher seas (JAN formulation). The results were diverse: from the consistent under-prediction obtained when using in-situ observations (up to -78% for H_s and -38% for T_p), to the better results obtained when using higher WS (as good as +1% for H_s and -15% for T_p). But they all pointed out the need to better understand the energy input and growth processes in such fetch-limited wave growth and highly variable wind conditions. For modelling purposes, the mechanisms for energy input should be further studied in order to improve the air-sea coupling. Because of the seldom presence of swell and the high variability of the area, it was suggested that

wave growth could be deviated from the theoretical growth curves and new parameterizations might be needed.

The results from this work were presented in:

Alomar, M., Bolaños, R., Sairouni, A., Ocampo-Torres, F., and Sanchez-Arcilla, A.: High resolution wave modelling in fetch-limited conditions, in: WISE Meeting, Ensenada, Mexico, 27th-30th April 2009, 2009.

Alomar, M., Bolaños, R., Sairouni, A., Ocampo-Torres, F., and Sanchez-Arcilla, A.: Modelado de oleaje en alta resolución en condiciones de fetch limitado. in: Jornadas Españolas de Costas y Puertos, Santander, Spain, 27-28 May 2009, 2009.

Alomar, M., Bolaños, R., Sanchez-Arcilla, A., Sairouni, A., and Ocampo-Torres, F.: Uncertainties in wave modeling for fetch-limited growth conditions, in: 33rd IAHR Congress: Water engineering for a sustainable environment, Vancouver, Canada, 9-14th August 2009, 2009.

Alomar, M., Sanchez-Arcilla, A., and González-Marco, D.: Coastal wave storms. Observations and simulations in the NW Mediterranean, in: 10th Plinius Conference: Mediterranean Storms, Nicosia, Cyprus, 22-24 September 2008, 2008.

7 Future work

This work is part of a PhD thesis which aims to better understand wave generation and dissipation processes in the study region in order to improve wave forecasts in this coastal zone. Eventually, this knowledge will be used within the RIMA project to define a risk index for the different coastal activities such as port and marine operations, fish farming, recreational and fishing activities, etc.

The work presented here concluded with the need to understand the source of under-prediction associated to KOM's formulation in SWAN, and the urgency to study the source of inaccuracy in JAN's formulation. Consequently, a better understanding of the model and an in-depth study of the observations is needed.

Indeed, it is essential to go back to the observations and to study in detail the generation and growth processes of wave storms. The instrumental record presented in this work is very valuable in terms of the quantity and the quality of the observations: wave information was available at up to three instruments at increasing distance, and within 60km, from the coast. The study of the wave growth curves in such variable conditions, and their deviation from the theoretical growth laws, will be the first steps to assess the accuracy of the parameterizations of the wave models in this region. During the period of measure two typical storm events were recorded (northwest and east), which study in detail would provide valuable information about these kinds of energetic events. Bimodal storms were also observed and, although the instrumental records are less complete, valuable information can be obtained from its study in-depth. Unfortunately, wind data in the sea and more directional measurements are lacking. Nonetheless, the available data will be further studied to understand the wave generation and growth that took place during the period considered. To start with, additional data from a meteorological station at Tarragona's harbour will be analyzed, and directional wave information at A-dw(D) will be processed. These data were unfortunately not available at the time of this work, but will be included shortly.

Further data analysis will first focus on different aspects such as:

- To study the data along the instrumental transect and compare it with the theoretical wave-growth curves.
- To analyze and compare the directional spectra at E-iw(D) and A-dw(D) to identify the different wave trains registered during the wave record available.
- To study the directionality of each frequency and search for slanting fetch effects. The role of wind and wave direction on wave growth will also be considered in detail.

- To assess the role of bimodal spectra in regional wave growth.

Afterwards, and once the instrumental data is fully analyzed, the focus will be on assessing the performance of the high-resolution wind and wave models in predicting the fully-developed sea conditions recorded during storm 4. Because eastern swell are being generated as far as Corsica and Sardinia, for this future simulation the wave model will be nested into a coarser regional model of the western Mediterranean.

Besides, aiming to improve the deficiencies of the high resolution wave model identified in this work, and following the research line started, further work will include:

- To assess the effect of the different wind field on the predictions at all the instruments along the transect, and not only at the most offshore station.
- To better understand the generation and dissipation formulations in SWAN to identify the sources of error (i.e. convergence problems) of the implemented formulations, including the JAN formulation.
- To identify the reason the model is still under-predicting the integrated wave parameters, H_s and T_p , and to study the importance of changing WiD and the variability of the wind field on increasing H_s .
- To assess the performance of the model in predicting the directional spectra.
- To adapt the model code (parameterizations) to better predict wave generation and propagation in the region of study.
- To assess a pre-operational implementation of the high-resolution model in the study region.

Overall, further work aims to better understand generation and growth mechanisms in highly variable regions and to improve wave modelling in the NW Mediterranean.

8 Acknowledgements

This work was supported by the RIMA project (MEC, TRA2006-05132/TMAR), and the Marie – Modelling and Assimilation for Rofl Environments project (MTKD-CT-2004-014509). The buoy data available was provided by the XIOM network and Puertos del Estado. The energy spectra were obtained using WAFO Toolbox ((Brodtkorb et al. 2000). Coastal and modelled meteorological data were provided by the Meteorological service of Catalonia. The bathymetry data used as input to the wave model was the GEBCO One Minute Grid, version 2.0 (<http://www.gebco.net>). The predicted wind fields were provided by the Meteorological Service of Catalonia: many thanks to Abdel Sairouni for running the atmospheric models.

I would also like to thank my advisors, Agustín Sánchez-Arcilla and Rodolfo Bolaños, the advice and time of Francisco Ocampo Torres and Judith Wolf, and the research centres that received and supported me during this study: Laboratori d'Enginyeria Marítima (LIM/UPC), Proudman Oceanographic Laboratory (POL) and Centro de Investigación Científica y Educación Superior de Ensenada (CICESE).

Special thanks to my family and friends.

9 References

- Alves, J. H. G. M., and Banner, M. L. (2003). "Performance of a saturation-based dissipation-rate source term in modeling the fetch-limited evolution of wind waves." *J.Phys.Oceanogr.*, 33(6), 1274-1298.
- Ardhuin, F., Herbers, T. H. C., van Vledder, G. P., Watts, K. P., Jensen, R., and Graber, H. C. (2007). "Swell and Slanting-Fetch Effects on Wind Wave Growth." *J. Phys. Oceanogr.*, 37(4), 908-931.
- Bolaños, R. (2004). "Tormentas de oleaje en el Mediterráneo: Física y predicción." PhD thesis, ETSECCPB-UPC.
- Bolaños, R., Rotés, A., and Sanchez-Arcilla, A. (2005). "Spectral wave climate at northern Spain's Mediterranean coast." ASCE.
- Bolaños, R., and Sánchez-Arcilla, A. (2006). "A note on nearshore wave features: Implications for wave generation." *Prog. Oceanogr.*, 70(2-4), 168-180.
- Bolaños, R., Sanchez-Arcilla, A., and Cateura, J. (2007). "Evaluation of two atmospheric models for wind-wave modelling in the NW Mediterranean." *J. Marine Syst.*, 65(1-4), 336-353.
- Bolaños, R., Sánchez-Arcilla, A., Gómez, J., Vilaclara, E., and Sairouni, A. (2003). "Predicción de sucesos extremos en la costa catalana." *Libro de Resúmenes VII Jornadas Españolas de Ingeniería de Costas y Puertos*, Fundación para el Fomento de la Ingeniería del Agua, 523-525.
- Bolaños, R., Jorda, G., Cateura, J., Lopez, J., Puigdefabregas, J., Gomez, J., and Espino, M. (2009). "The XIOM: 20 years of a regional coastal observatory in the Spanish Catalan coast." *J.Mar.Syst.*, 77(3), 237-260.
- Brodtkorb, P. A., Johannesson, P., Lindgren, G., Rychlik, I., Ryden, J., and Sjö, E. (2000). "WAFO - a Matlab Toolbox for the analysis of random waves and loads." *Proc. 10th Int. Offshore and Polar Eng. Conf., ISOPE*, 343-350.
- Cavaleri, L., Alves, J. -. G. M., Ardhuin, F., Babanin, A., Banner, M., Belibassakis, K., Benoit, M., Donelan, M., Groeneweg, J., Herbers, T. H. C., Hwang, P., Janssen, P. A. E. M., Janssen, T., Lavrenov, I. V., Magne, R., Monbaliu, J., Onorato, M., Polnikov, V., Resio, D., Rogers, W. E., Sheremet, A., McKee Smith, J., Tolman, H. L., van Vledder, G., Wolf, J., and Young, I. (2007). "Wave modelling – The state of the art." *Prog.Oceanogr.*, 75(4), 603-674.
- Cavaleri, L., and Bertotti, L. (2004). "Accuracy of the modelled wind and wave fields in enclosed seas." *Tellus A*, 56 167-175(9).

- Cavaleri, L., and Bertotti, L. (1997). "In search of the correct wind and wave fields in a minor basin." *Mon. Weather Rev.*, 1964-1975.
- Cavaleri, L., and Scavo, M. (2006). "The calibration of wind and wave model data in the Mediterranean Sea." *Coast. Eng.*, 53(7), 613-627.
- Cavaleri, L., and Malanotte-Rizzoli, P. (1981). "Wind Wave Prediction in Shallow Water: Theory and Applications." *Journal of Geophysical Research*, 86(c11), 10961-10973.
- Codina, B., Aran, M., Young, S., and Redaño, A. (1997). "Prediction of a mesoscale convective system over Catalonia (Northeastern Spain) with a nested numerical model." *Meteorol. Atmos. Phys.*, 62(1), 9-22.
- Cortés, F. (2003). "El tráfico marítimo y la actividad portuaria." *Boletín Económico De ICE*, 2758 27-31.
- Coto, P., and Inglada, V. (1999). "Análisis del transporte marítimo en España (1974-1999): competencia y regulación." *Papeles De Economía Española*, 82 59-70.
- Flamant, C., Pelon, J., Hauser, D., Quentin, C., Drennan, W. M., Gohin, F., Chapron, B., and Gourrion, J. (2003). "Analysis of surface wind and roughness length evolution with fetch using a combination of airborne lidar and radar measurements." *J. Geophys. Res.*, 108.
- Gómez, J., Espino, M., Sánchez-Arcilla, A., Solano, M., and Vela, J. (2005). "Extreme wave conditions in a torrential climate. The Catalan case." *Book of Abstracts, Ocean Wave Measurement and Analysis, 5th International Symposium WAVES 2005*, CEDEX, 31.
- Grell, G. A., Dudhia, J., and Stauffer, D. R. (1994). "Description of the fifth-Generation Penn State/NCAR Mesoscale Model (MM5)." *NCAR Tech. Note, NCAR/TN-398-STR* .
- Grell, G. A. (1993). "Prognostic evaluation of assumptions used by cumulus parameterizations." *Mon. Weather Rev.*, 121(3), 764-787.
- Janssen, P. A. E. M. (1991). "Quasi-linear theory of wind-wave generation applied to wave forecasting." *J. Phys. Oceanogr.*, 21(11), 1631-1642.
- Janssen, P. A. E. M. (1989). "Wave-induced stress and the drag of air flow over sea waves." *J. Phys. Oceanogr.*, 19(6), 745-754.
- Jiménez, J. A., Sánchez-Arcilla, A., Valdemoro, H. I., Gracia, V., and Nieto, F. (1997). "Processes reshaping the Ebro delta." *Marine Geology*, 144(1-3), 59-79.
- Klein, R. J. T., Nicholls, R. J., Ragoonaden, S., Capobianco, M., Aston, J., and Buckley, E. N. (2001). "Technological Options for Adaptation to Climate Change in Coastal Zones." *J.Coast.Res.*, 531-543.
- Komen, G. J., Cavaleri, L., Donelan, M., Hasselmann, K., Hasselmann, S., and Janssen, P. A. E. M. (1994). *Dynamics and modelling of ocean waves*. Cambridge University Press, Cambridge, UK.
- Komen, G. J., Hasselmann, K., and Hasselmann, K. (1984). "On the existence of a fully developed wind-sea spectrum." *J. Phys. Oceanogr.*, 14(8), 1271-1285.

- Miles, J. W. (1957). "On the generation of surface waves by shear flows." *J.Fluid Mech.*, 3(2), 185-204.
- Monbaliu, J. (2003). "Spectral wave models in coastal areas." *Advances in Coastal Modeling*, C. Lakhani, ed., 133-158.
- Monbaliu, J., Padilla-Hernandez, R., Hargreaves, J. C., Albiach, J. C. C., Luo, W., Sclavo, M., and Günther, H. (2000). "The spectral wave model, WAM, adapted for applications with high spatial resolution." *Coast. Eng.*, 41 41-62.
- Niclasen, B. A. (2006). "An operational wave model for the Faroe Shelf." PhD thesis, University of the Faroe Islands, Torshavn, Denmark.
- Nombela, G., and Trujillo, L. (1999). "El sector portuario español: organización actual y perspectivas." *Papeles En Economía Española*, 82 71-85.
- Phillips, O. M. (1957). "On the generation of waves by turbulent wind." *J.Fluid Mech.*, 2(5), 417-445.
- Puertos del Estado. (2008). "Tráficos portuarios." *Revista De Puertos*, 144 78-82.
- Ris, R. C. (1997). "Spectral modelling of wind waves in coastal areas." PhD thesis, Delft University Press, Delft, Netherlands.
- Sánchez-Arcilla, A., González-Marco, D., and Bolaños, R. (2008). "A review of wave climate and prediction along the Spanish Mediterranean coast." *Nat. Hazard. Earth Sys.*, 8(6), 1217-1228.
- Schneggenburger, C., Günther, H., and Rosenthal, W. (2000). "Spectral wave modelling with non-linear dissipation: validation and applications in a coastal tidal environment." *Coastal Engineering*, 41(1-3), 201-235.
- Schultz, P. (1995). "An explicit cloud physics parameterization for operational numerical weather prediction." *Mon. Weather Rev.*, 123(11), 3331-3343.
- Siddons, L. A., Wyatt, L. R., and Wolf, J. (2009). "Assimilation of HF radar data into the SWAN wave model." *J.Mar.Syst.*, 77(3), 312-324.
- Signell, R. P., Carniel, S., Cavaleri, L., Chiggiato, J., Doyle, J. D., Pullen, J., and Sclavo, M. (2005). "Assessment of wind quality for oceanographic modelling in semi-enclosed basins." *J. Marine Syst.*, 53(1-4), 217-233.
- The SWAN team. (2008). "SWAN technical documentation." Delft University of Technology, Delft, Netherlands.
- Tucker, M. J., and Pitt, E. G. (2001). *Waves in ocean engineering*. Elsevier Science Ltd., Oxford, UK.
- van der Westhuysen, A. J., Zijlema, M., and Battjes, J. A. (2007). "Nonlinear saturation-based whitecapping dissipation in SWAN for deep and shallow water." *Coast.Eng.*, 54(2), 151-170.

- Van Rijn, L. C., Walstra, D. J. R., Grasmeyer, B., Sutherland, J., Pan, S., and Sierra, J. P. (2003). "The predictability of cross-shore bed evolution of sandy beaches at the time scale of storms and seasons using process-based Profile models." *Coast. Eng.*, 47 295-327.
- Yan, L. (1987). "An improved wind input source term for third generation ocean wave modelling." *Rep. No. WR-87-8*, De Bilt, The Netherlands.