



Title	PURIFICATION AND CHARACTERIZATION OF NEUROTOXIN PRODUCED BY CLOSTRIDIUM BOTULINUM TYPE C STRAIN 6813
Author(s)	TERAJIMA, Jun
Citation	Japanese Journal of Veterinary Research, 33(1-2), 97-97
Issue Date	1985-04-30
Doc URL	<a href="http://hdl.handle.net/2115/2352">http://hdl.handle.net/2115/2352</a>
Type	bulletin (article)
File Information	KJ00002374324.pdf



[Instructions for use](#)

PURIFICATION AND CHARACTERIZATION OF NEUROTOXIN  
PRODUCED BY *CLOSTRIDIUM BOTULINUM* TYPE C STRAIN 6813

Jun TERAJIMA

*Department of Biochemistry  
Faculty of Veterinary Medicine  
Hokkaido University, Sapporo 060, Japan*

The toxin produced by *Clostridium botulinum* type C strain 6813 (C-6813) was purified 1009-fold from the culture supernatant in an overall yield of 30%. The specific toxicity was  $1.1 \times 10^7$  mouse minimum lethal doses per mg of protein. The toxin had a molecular weight of 144,000 composed of light and heavy chains with molecular weights of 52,000 and 92,000, respectively, linked by one or two disulfide bond(s).

The purified C-6813 toxin heavy and light chains reacted strongly with anti-type D heavy chain IgG and anti-type C<sub>1</sub> (C-Stockholm) light chain IgG, respectively. The amino acid compositions of C-6813 toxin heavy and light chains were similar to those of type D heavy chain and type C<sub>1</sub> light chain than to those of type C<sub>1</sub> heavy chain and type D light chain, respectively. In neutralization test, anti-C-6813 toxin IgG had neutralizing activity of 99.7% against both C-6813 toxin and C<sub>1</sub> toxin. On the other hand, anti-type C<sub>1</sub> toxin IgG had neutralizing activity of 99.7% against C<sub>1</sub> toxin but only of 20% against C-6813 toxin.

These results indicate that in the neurotoxin produced by the type C strain at least two subtypes exist.