



Title	Molecular cloning of mouse MAIL cDNA and its expression and function during inflammation
Author(s)	KANEHIRA, Katsushi
Citation	Japanese Journal of Veterinary Research, 49(1), 57-57
Issue Date	2001-05-31
Doc URL	http://hdl.handle.net/2115/2885
Type	bulletin (article)
File Information	KJ00002400347.pdf



[Instructions for use](#)

Molecular cloning of mouse MAIL cDNA and its expression and function during inflammation

Katsushi Kanehira

Laboratory of Biochemistry
Department of Biomedical Sciences
School of Veterinary Medicine
Hokkaido University, Sapporo 060-0818, Japan

I have cloned the full length cDNA of a novel ankyrin-repeat protein from mice treated with lipopolysaccharide (LPS), and named it as "Molecule possessing Ankyrin-repeats Induced by Lipopolysaccharide" (MAIL). The deduced carboxyl-terminal portion of MAIL, including ankyrin-repeat domain (ARD), shared approximately 40% identity with the ARD of inhibitory kappa B ($I\kappa B$) family proteins. Intraperitoneal injection of LPS to mice rapidly induced MAIL mRNA in various tissues, particularly in the spleen, lymph node and lung. *In situ* hybridization study revealed that the induction of MAIL oc-

curred in macrophages and B lymphocytes. In concordance with these *in vivo* results, some cell lines of monocyte-macrophage and B lymphocyte origin abundantly expressed MAIL mRNA by LPS challenge *in vitro*. Ectopically expressed MAIL in Swiss/3T3 fibroblasts localized in the nucleus, and remarkably potentiated the LPS-induced mRNA expression and secretion of interleukin-6 (IL-6).

These findings indicate that MAIL is a new member of the nuclear $I\kappa B$ family, induced by LPS treatment, and an activator of IL-6 production.