



| | |
|------------------|---|
| Title | Effects of cycloheximide treatment on in-vitro development of porcine parthenotes and somatic cell nuclear transfer embryos |
| Author(s) | Diaz, Mario A. Martinez; Suzuki, Masami; Kagawa, Masumi; Ikeda, Koji; Takahashi, Yoshiyuki |
| Citation | Japanese Journal of Veterinary Research, 50(4), 147-155 |
| Issue Date | 2003-02-28 |
| DOI | 10.14943/jjvr.50.4.147 |
| Doc URL | http://hdl.handle.net/2115/2962 |
| Type | bulletin (article) |
| File Information | KJ00000699555.pdf |



[Instructions for use](#)

Effects of cycloheximide treatment on *in-vitro* development of porcine parthenotes and somatic cell nuclear transfer embryos

Mario A. Martinez Diaz^{1,2)}, Masami Suzuki¹⁾, Masumi Kagawa¹⁾, Koji Ikeda¹⁾
and Yoshiyuki Takahashi¹⁾*

(Accepted for publication : January 9, 2003)

Abstract

This study aimed to verify the beneficial effect of cycloheximide (CHX) treatment on the development of porcine somatic cell nuclear transfer (NT) embryos, which were constructed with enucleated oocytes and cumulus cells by using a single direct current (DC) pulse. In the first experiment, a single DC pulse applied to the induction of fusion and activation of NT embryos gave a high fusion rate. However, cleavage and subsequent development of fused couplets (NT embryos) to the blastocyst stage were poor. Experiment 2 was conducted to determine whether CHX treatment could enhance metaphase II (M II) oocyte activation and improve the subsequent parthenogenetic development. After giving the DC pulse and incubation with or without CHX, M II oocytes incubated with CHX showed higher cleavage and development to blastocysts compared with those incubated without CHX ($P < 0.05$). Experiment 3 was carried out to verify the beneficial effect of CHX on the development of NT embryos. The NT embryos treated with the DC pulse and CHX treatment showed higher cleavage and subsequent development compared with those treated with the DC pulse alone ($P < 0.05$). The present study demonstrates that CHX treatment enhances the electrical stimulus-induced activation of oocytes and NT embryos, and improves the subsequent development of parthenotes and NT embryos. The results indicate that protein synthesis inhibition treatment required for the induction of oocyte activation promotes the development of NT embryos.

¹⁾Laboratory of Theriogenology, Department of Veterinary Clinical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

²⁾Present address : Faculty of Veterinary Sciences, Universidad Austral de Chile, Valdivia, Chile

*Corresponding author : Yoshiyuki Takahashi

Laboratory of Theriogenology, Department of Veterinary Clinical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

Phone / Fax : +81-11-706-5231

E-mail : ytaka@vetmed.hokudai.ac.jp

Key words: activation; cloning; cycloheximide; nuclear transplantation; pig

Introduction

Since the birth of the famous cloned sheep "Dolly" was reported by Wilmut *et al.* (1997)³⁷⁾, cloned offspring has been produced by using somatic cell nuclear transfer (NT) in cattle¹³⁾, goats²⁾, pigs^{3,22,27)}, rabbits⁷⁾ and cats³⁰⁾. Somatic cell NT has also been successfully applied to the production of transgenic sheep²⁸⁾, cattle⁸⁾, goats¹⁴⁾ and pigs^{18,23,24)}. In the pig, transgenic technology is a valuable tool for the generation of animal models for human diseases and production of organs for xenotransplantation. In most successful porcine somatic cell NT protocols used for the production of cloned piglets, NT embryos at the 1- to 8-cell stages were transferred to the foster mothers (recipient animals)^{4,22,24,27)}. The transfer of transgenic NT embryos at the earlier developmental stages limits the monitoring of the expression of the foreign genes, which were transfected into the nuclear donor cells, compared with that at the morula and blastocyst stages. However, the *in-vitro* development of porcine somatic cell NT embryos to the blastocyst stage is very low^{4,16,17,33,34)}; therefore, there is a necessity to improve the *in-vitro* development of porcine somatic cell NT embryos to the blastocyst stage.

Many factors may affect the development of NT embryos such as genotype/cell line of nuclear donor cells, cell-cycle stages of nuclear donor cells and recipient oocytes, and methods of nuclear transfer including fusion/activation protocols¹⁰⁾. Cycloheximide (CHX), a protein synthesis inhibitor, was reported to be effective on the induction of activation of porcine metaphase II (M II) oocytes when combined with electric stimulation²¹⁾, calcium ionophore A23187 or ethanol⁵⁾. Our previous study¹²⁾ confirmed the effectiveness of the

CHX treatment combined with an electrical stimulation for the induction of activation (pronuclear formation) of porcine young M II oocytes. Porcine NT embryos treated with an electrical stimulus followed by a CHX treatment showed a higher development compared with those treated with an electrical stimulus and calcium ionophore A23187⁶⁾. However, a beneficial effect of CHX treatment in combination with an electrical stimulation on the development of porcine parthenotes and NT embryos to the blastocyst stage has not been confirmed.

In the present study, therefore, we treated porcine M II oocytes and NT embryos with an electrical stimulus combined with and without CHX to verify the beneficial effect of CHX treatment on the induction of activation of oocytes and NT embryos and on the subsequent development to the blastocyst stage.

Materials and Methods

Oocyte preparation

Abattoir-derived ovaries from prepubertal gilts were transported to the laboratory in a thermal flask with sterile physiological saline at 37°C within 2 hr of slaughter. Oocyte-cumulus-granulosa cell complexes (OCGCs) were collected from 4- to 6-mm antral follicles by a dissection method⁶⁾. Briefly, dissected non-atretic follicles with notable blood irrigation were kept in 25 mM Hepes-buffered TCM-199 (Sigma Chemical Co., St. Louis, MO, USA) supplemented with 0.1% polyvinyl alcohol (Sigma) at 30-35°C. Follicles were opened under a stereomicroscope, and OCGCs were released by scraping the inner surface of follicle wall with a pair of forceps. OCGCs were cultured in a 100- μ l drop (5 OCGCs in each drop) of bovine serum albumin (BSA)-free

North Carolina State University Medium 23 (NCSU-23)²⁶ supplemented with 0.57 mM cysteine (Sigma), 10% porcine follicular fluid, 10 IU/ml equine chorionic gonadotropin (Teikoku Hormone Mfg. Co. Ltd., Tokyo, Japan), 10 IU/ml human chorionic gonadotropin (Teikoku Hormone) and 50 µg/ml gentamicin sulfate (Sigma) covered with paraffin oil for 20 hr under a humidified atmosphere of 5% CO₂ in air at 39°C. This was followed by the culture in the same medium without hormones for additional 22-24 hr. After the maturation culture, oocytes were freed of granulosa/cumulus cells by vortexing the OCGCs for 4 min in Ca²⁺- and Mg²⁺-free Dulbecco's PBS (DPBS) containing 0.1% hyaluronidase (Sigma), and the oocytes with the first polar bodies (M II oocytes) were selected.

Oocyte enucleation

Matured-cumulus-free M II oocytes were mechanically enucleated as described previously¹. In brief, the first polar body and around 20% of the adjacent cytoplasm were removed using a pipette under an inverted microscope (Eclipse TE 300, Nikon, Tokyo, Japan) in HEPES-buffered TCM-199 supplemented with 5 µg/ml cytochalasin B (Sigma) and 10% fetal calf serum (FCS, Gibco Laboratories, Grand Island, NY, USA). After the enucleation, cytoplasts were incubated in Ca²⁺- and Mg²⁺-free DPBS supplemented with 5 µg/ml Hoechst 33342 and 3 mg/ml BSA for 20 min at 39°C. Enucleation was confirmed by exposing the oocytes to ultraviolet light for a few seconds under an inverted microscope equipped with an epifluorescence (TE-FM, Nikon) and UV-1A filter block (365 nm excitation and 400 nm emission).

Preparation of nuclear donor cells

Primary cultures of cumulus cells were established using OCGCs obtained after the

maturation culture as described previously¹. Briefly, cumulus cells were washed several times with Dulbecco's modified Eagle's medium: nutrient mixture F-12 (DMEM/F12, Gibco). They were then cultured for 3-4 days in DMEM/F12 supplemented with 10% FCS and 50 µg/ml gentamicin sulfate at a concentration of 1 x 10⁵ cells/ml in 4-well (16 mm diameter) tissue culture plate (Nalge Nunc International, Roskilde, Denmark). After establishing the primary cultures, the cells were further cultured in DMEM/F12 supplemented with 0.5% FCS and 50 µg/ml gentamicin sulfate for 4-5 days. A single-cell suspension was prepared by standard trypsinization, and middle-sized (20-24 µm) cumulus cells were used as nuclear donor cells for NT.

Production of nuclear transfer embryos

Nuclear donor cell was injected into the perivitelline space of an enucleated oocyte through the same slit in the zona pellucida as was made during enucleation. Couplets of enucleated oocytes and nuclear donor cells were placed and manually aligned between two wire electrodes, 1.0-mm apart, overlaid with 0.3M mannitol solution containing 0.1 mM MgSO₄, 0.1 mM CaCl₂ and 0.05 mg/ml BSA. Cell fusion and activation were induced simultaneously by applying a single DC pulse of 150 V/mm for 100 µsec using an Electro Cell Fusion instrument (LF 100, Life Tec Co., Tokyo, Japan). Fusion of the couplets was determined 30 min after the electrical stimulus.

Production of parthenotes

The oocytes at the M II stage received a single electrical stimulation which was the same DC pulse as that was used for the induction of fusion/activation of NT embryos. They were then incubated for 6 hr in NCSU-23 containing 5 µg/ml cytochalasin B and 50 µg/ml gentamicin sulfate.

In-vitro culture of NT embryos and parthenotes

NT embryos and parthenotes were cultured in 30- μ l drops of NCSU-23 containing 50 μ g/ml gentamicin sulfate covered with paraffin oil for 6 days under humidified air with 5% CO₂ at 39°C. The cleavage and subsequent development to the blastocyst stage were determined 48 and 144 hr after the electrical stimulus, respectively. At the end of the culture, all of NT embryos and parthenotes, which reached the blastocyst stage, were assigned for the cell count by using an air-dry method as described elsewhere³².

Experimental studies

In experiment 1, NT embryos were produced by using an electrical stimulus of a single DC pulse alone, and the rates of fusion, cleavage and development of NT embryos to the blastocyst stage were examined.

Experiment 2 was conducted to determine whether CHX treatment could enhance electrostimulation-induced oocyte activation, and could improve the subsequent parthenogenetic development to the blastocyst stage. In this experiment, *in vitro* mature M II oocytes were treated with a single DC pulse, and they were then incubated for 6 hr in NCSU-23 containing 5 μ g/ml cytochalasin B in the presence or absence of 10 μ g/ml CHX.

Experiment 3 was carried out to verify the beneficial effect of CHX on the develop-

ment of NT embryos. Fused couplets of enucleated oocytes and cumulus cells (NT embryos) produced by using a single DC pulse were incubated for 5 hr in NCSU-23 with or without 10 μ g/ml of CHX before assessing the *in vitro* development to the blastocyst stage.

Statistical analysis

Statistical differences between 2 means were analyzed by Student's *t*-test using Stat-View software (Abacus Concepts Inc., Berkeley, CA, USA).

Results

Experiment 1

After applying the electrical stimulation, 87.6 \pm 10.1% (mean \pm SD of 7 replicates, n=107) of couplets were fused. The proportions of fused couplets which cleaved and developed to the blastocyst stage were 28.9 \pm 7.3 and 5.5 \pm 6.1% (means \pm SD of 7 replicates, n=94), respectively.

Experiment 2

As shown in Table 1, the oocytes treated with CHX showed higher cleavage and blastocyst formation rates than those cultured without CHX ($P < 0.05$).

Experiment 3

Although there was no difference in the fusion rate between the two groups, cleavage and blastocyst rates of NT embryos were im-

Table 1. Effects of cycloheximide (CHX) treatment on parthenogenetic development of porcine oocytes

| Treatment | No. of oocytes examined | % of oocytes developed to | | Cell no. in blastocysts (n) |
|-----------|-------------------------|-----------------------------|-----------------------------|-----------------------------|
| | | ≥ 2 -cell | blastocyst | |
| No CHX | 114 | 31.4 \pm 9.8 ^a | 16.4 \pm 7.1 ^a | 24.3 \pm 5.8 (19) |
| With CHX | 116 | 63.0 \pm 6.9 ^b | 46.8 \pm 8.3 ^b | 22.7 \pm 4.4 (54) |

^{a,b}Values (means \pm SD of 4 replicates) with different superscripts in the same column differ significantly ($P < 0.05$).

Table 2 . Effects of cycloheximide (CHX) treatment on the *in vitro* development of porcine NT embryos

| Treatment | No. of couplets | % of fused couplets | % of couplets developed to ^a | | Cell no.in blastocysts (n) |
|-----------|-----------------|---------------------|---|-----------------------|----------------------------|
| | | | ≥ 2 -cell | blastocyst | |
| No CHX | 65 | 98.7±2.3 | 43.3±16.3 ^b | 5.4±4.8 ^b | 27.0±7.9 (3) |
| With CHX | 66 | 95.8±7.2 | 70.0±10.5 ^c | 16.9±3.5 ^c | 17.9±10.3 (11) |

^a% based on the number of fused couplets.

^{b,c}Values (means ±SD of 3 replicates) with different superscripts in the same column differ significantly ($P < 0.05$).

proved when they were treated with CHX ($P < 0.05$) as shown in Table 2.

Discussion

Excessive electrical stimuli for the induction of activation of porcine NT embryos was reported to reduce the development to the blastocyst stage¹⁶. To induce the fusion and activation of NT embryos, we used a single DC pulse of 150 V/mm for 100 μ sec, which was the lowest voltage capable of inducing a high activation (pronuclear formation) rate of porcine M II oocytes in our preliminary experiment ($64.3 \pm 6.0\%$, mean of 3 replicates, $n = 40$). In experiment 1, the single DC pulses efficiently induced the fusion between enucleated oocytes and nuclear donor cells; however, the cleavage and development of fused couplets (NT embryos) were poor. These results indicate that the single DC pulse used in the present study is insufficient to induce definitive activation of NT embryos.

Induction of activation of porcine young oocytes is more difficult than that of aging/aged oocytes^{12,15}. Although CHX alone could not induce the activation of porcine oocytes²¹, an electrical stimulus combined with CHX treatment was reported to enhance the activation (pronuclear formation) of porcine young M II oocytes¹². In experiments 2 and 3, therefore, CHX treatment was applied to the induction of activation of M II oocytes and NT embryos after giving the same electrical

stimulus as used in experiment 1. The higher rates of cleavage and development to blastocysts of parthenotes and NT embryos after the treatment with CHX clearly indicate that CHX treatment enhances the activation of M II oocytes and NT embryos, and leads the improvement in the subsequent development of parthenotes and NT embryos.

At the time of fertilization, penetration of sperm into the oocytes induces the oocyte activation and enables to exit from M II¹¹. To exit from the M II stage and to enter the first cleavage, the inactivation of both cytosstatic factor (CSF: c-mos/mitogen-activated protein kinase) and M-phase promoting factor (MPF: p34^{cdc2}/cyclin B) are required^{11,20}. As mentioned above, more than half (approximately 65%) of porcine M II oocytes, which received the single DC pulse, formed pronuclei in our preliminary experiment. Therefore, the single DC pulse may induce reduction of CSF and MPF activities; however, the CSF activity may increase again before definitive inactivation⁹. CHX could inhibit resynthesis of components of CSF and MPF^{19,31}. The present results indicate that protein synthesis inhibition treatment is required for the definitive inactivation of CSF and/or MPF, and to lead the initiation of cleavage and subsequent development of NT embryos when fusion and activation of NT embryos were induced simultaneously by using a single DC pulse.

Some groups have reported that the in-

terval between fusion and activation stimuli affected the development of somatic cell NT embryos in the pig¹⁶⁾, cattle^{29,36)} and mouse³⁵⁾: fusion several hours before activation was superior to the simultaneous fusion and activation treatment. In the present study, we induced fusion and activation of porcine NT embryos simultaneously by a single DC pulse before subjecting the NT embryos to the CHX treatment. Therefore, further studies are needed to clarify the role of CHX treatment in the development of somatic cell NT embryos with various intervals between fusion and activation stimuli.

Development of NT embryos to the blastocyst stage was poor compared with that of parthenotes even if they were treated with CHX. These results were in agreement with previous findings^{16,25)}; however, it was not clear why the NT embryos showed a poor development compared with the parthenotes. Several factors may affect the *in vitro* development of somatic cell NT embryos, and one of most important factors is the nuclear reprogramming¹⁰⁾. It remains to clarify the mechanisms involved and the effects of CHX on the inactivation of CSF/MPF and reprogramming of somatic cell nuclei after transfer to enucleated oocytes.

In conclusion, the present study demonstrates that CHX treatment enhances the electrical stimulus-induced activation of M II oocytes and NT embryos, and leads the improvement in the cleavage and subsequent development of parthenotes and NT embryos to the blastocyst stage. The results also indicate that protein synthesis inhibition treatment required for the induction of M II oocyte activation promotes the development of NT embryos when fusion and activation of NT embryos were induced by a single DC pulse.

Acknowledgments

The authors thank the staff members of the Ebetsu Meat Inspection Office (Hokkaido, Japan) for supplying the porcine ovaries. This study was supported by Grants-in-Aid for Scientific Research (Nos. 10556058 and 12480248) from the Japan Society for the Promotion of Science.

References

- 1) Atabay, E. C., Martinez Diaz, M.A., Dochi, O. and Takahashi, Y. 2001. Factors affecting enucleation rates of bovine and porcine oocytes after removal of cumulus cells by vortexing. *J. Reprod. Dev.*, 47 : 365-371.
- 2) Baguisi, A., Behboodi, E., Melican, D.T., Pollock, J.S., Destrempe, M.M., Cammuso, C., Williams, J.L., Nims, S.D., Porter, C.A., Midura, P., Palacios, M.J., Ayres, S.L., Denniston, R.S., Hayes, M.L., Ziomek, C.A., Meade, H.M., Godke, R.A., Gavin, W.G., Overstrom, E.W. and Echelard, Y. 1999. Production of goats by somatic cell nuclear transfer. *Nat. Biotechnol.*, 17 : 456-461.
- 3) Betthausen, J., Forsberg, E., Augenstein, M., Childs, L., Eilertsen, K., Enos, J., Forsythe, T., Golueke, P., Jurgella, G., Koppang, R., Lesmeister, T., Mallon, K., Mell, G., Misica, P., Pace, M., Pfister-Genskow, M., Strelchenko, N., Voelker, G., Watt, S., Thompson, S. and Bishop, M. 2000. Production of cloned pigs from *in-vitro* systems. *Nat. Biotechnol.*, 18 : 1055-1059.
- 4) Boquest, A.C., Grupen, C.G., Harrison, S.J., McIlpatrick, S.M., Ashman, R.J., d'Apice, A.J.F. and Nottle, M.B. 2002. Production of cloned pigs from cultured fetal fibroblast cells. *Biol. Reprod.*, 66 : 1283-1287.
- 5) Cha, S.K., Kim, N.H., Lee, S.M., Baik, C.S.,

- Lee, H.T. and Chung, K.S. 1997. Effect of cytochalasin B and cycloheximide on the activation rate, chromosome constituent and *in-vitro* development of porcine oocytes following parthenogenetic stimulation. *Reprod. Fertil. Dev.*, 9 : 441-444.
- 6) Cheong, H.T., Ikeda, K., Martinez Diaz, M. A., Katagiri, S. and Takahashi, Y. 2000. Development of reconstituted pig embryos by nuclear transfer of cultured cumulus cells. *Reprod. Fertil. Dev.*, 12 : 15-20.
- 7) Chesne, P., Adenot, P.G., Viglietta, C., Baratte, M., Boulanger, L. and Renard, J. P. 2002. Cloned rabbit produced by nuclear transfer from adult somatic cells. *Nat. Biotechnol.*, 20 : 366-369.
- 8) Cibelli, J.B., Stice, S.L., Golueke, P.J., Kane, J.J., Jerry, J., Blackwell, C., Ponce De Leon, F.A. and Robl, J.M. 1998. Transgenic bovine chimeric offspring produced from somatic cell-derived stem-like cells. *Nat. Biotechnol.*, 16 : 642-646.
- 9) Ciemerych, M.A. and Kubiack, J.Z. 1999. Transient reactivation of CSF in parthenogenetic one-cell mouse embryos. *Biol. Cell*, 91 : 641-647.
- 10) Dinnyes, A., De Souza, P., King, T. and Wilmut, I. 2002. Somatic cell nuclear transfer : Recent progress and challenges. *Cloning Stem Cells*, 4 : 81-90.
- 11) Fulka, J. Jr., Loi, P., Ledda, S., Moor, R. M. and Fulka, J. 2001. Nucleus transfer in mammals : how the oocyte cytoplasm modifies the transferred nucleus. *Theriogenology*, 55 : 1373-1380.
- 12) Ikeda, K. and Takahashi, Y. 2001. Effects of maturational age of porcine oocytes on the induction of activation and development *in-vitro* following somatic cell nuclear transfer. *J. Vet. Med. Sci.*, 63 : 1003-1008.
- 13) Kato, Y., Tani, T., Sotomaru, Y., Kurokawa, K., Kato, J., Doguchi, H., Yasue, H. and Tsunoda, Y. 1998. Eight calves cloned from somatic cells of a single adult. *Science*, 282 : 2095-2098.
- 14) Keefer, C.L., Baldassarre, H., Keyston, R., Wang, B., Bhatia, B., Bilodeau, A.S., Zhou, J.F., Leduc, M., Downey, B.R., Lazaris, A. and Karatzas, C.N. 2001. Generation of dwarf goat (*Capra hircus*) clones following nuclear transfer with transfected and nontransfected fetal fibroblasts and *in-vitro*-matured oocytes. *Biol. Reprod.*, 64 : 849-856.
- 15) Kikuchi, K., Izaïke, Y., Noguchi, J., Furukawa, T., Daen, F.P., Naito, K. and Toyoda, Y. 1995. Decrease of histone H 1 kinase activity in relation to parthenogenetic activation of pig follicular oocytes matured and aged *in vitro*. *J. Reprod. Fertil.*, 105 : 325-330.
- 16) Koo, D.B., Kang, Y.K., Choi, Y.H., Park, J. S., Han, S.K., Park, I.Y., Kim, S.U., Lee, K. K., Son, D.S., Chang, W.K. and Han, Y. M. 2000. *In-vitro* development of reconstructed porcine oocytes after somatic cell nuclear transfer. *Biol. Reprod.*, 63 : 986-992.
- 17) Koo, D.B., Kang, Y.K., Choi, Y.H., Park, J. S., Kim, H.N., Kim, T., Lee, K.K. and Han, Y.M. 2001. Developmental potential and transgene expression of porcine nuclear transfer embryos using somatic cells. *Mol. Reprod. Dev.*, 58 : 15-21.
- 18) Lai, L., Kolber-Simonds, D., Park, K.W., Cheong, H.T., Greenstein, J.L., Im, G.S., Samuel, M., Bonk, A., Rieke, A., Day, B.N., Murphy, C.N., Carter, D.B., Hawley, R.J. and Prather, R.S. 2002. Production of α -1, 3-galactosyltransferase knockout pigs by nuclear transfer cloning. *Science*, 295 : 1089-1092.
- 19) Levesque, J.T. and Sirard, M.A. 1996. Resumption of meiosis is initiated by the ac-

- cumulation of cyclin B in bovine oocytes. *Biol. Reprod.*, 55 : 1427-1436.
- 20) Masui, Y. and Markert, C.L. 1971. Cytoplasmic control of nuclear behavior during meiotic maturation of frog oocytes. *J. Exp. Zool.*, 177 : 129-145.
 - 21) Nussbaum, D.J. and Prather, R.S. 1995. Differential effects of protein synthesis inhibitors on porcine oocyte activation. *Mol. Reprod. Dev.*, 41 : 70-75.
 - 22) Onishi, A., Iwamoto, M., Akita, T., Mikawa, S., Takeda, K., Awata, T., Hanada, H. and Perry, A.C. 2000. Pig cloning by microinjection of fetal fibroblast nuclei. *Science*, 289 : 1188-1190.
 - 23) Park, K.W., Cheong, H.T., Lai, L., Im, G.S., Kuhholzer, B., Bonk, A., Samuel, M., Rieke, A., Day, B.N., Murphy, C.N., Carter, D.B. and Prather, R.S. 2001. Production of nuclear transfer-derived swine that express the enhanced green fluorescent protein. *Anim. Biotech.*, 12 : 173-181.
 - 24) Park, K.W., Lai, L., Cheong, H.T., Cabot, R., Sun, Q.Y., Wu, G., Rucker, E.B., Durtschi, D., Bonk, A., Samuel, M., Rieke, A., Day, B. N., Murphy, C.N., Carter, D.B. and Prather, R.S. 2002. Mosaic gene expression in nuclear transfer-derived embryos and the production of cloned transgenic pigs from ear-derived fibroblasts. *Biol. Reprod.*, 66 : 1001-1005.
 - 25) Park, K.W., Lai, L., Cheong, H.T., Im, G.S., Sun, Q.Y., Wu, G., Day, B.N. and Prather, R.S. 2001. Developmental potential of porcine nuclear transfer embryos derived from transgenic fetal fibroblasts infected with the gene for the green fluorescent protein : comparison of different fusion/activation conditions. *Biol. Reprod.*, 65 : 1681-1685.
 - 26) Petters, R.M. and Wells, K.D. 1993. Culture of pig embryos. *J. Reprod. Fertil. Suppl.*, 48 : 61-73.
 - 27) Polejaeva, I.A., Chen, S.H., Vaught, T.D., Page, R.L., Mullins, J., Ball, S., Dai, Y., Boone, J., Walker, S., Ayares, D.L., Colman, A. and Campbell, K.H.S. 2000. Cloned pigs produced by nuclear transfer from adult somatic cells. *Nature*, 407 : 86-90.
 - 28) Schnieke, A.E., King, A.J., Ritchie, W.A., Mycock, K., Scott, A.R., Ritchie, M., Wilmut, I., Colman, A. and Campbell, K. H. 1997. Human factor IX transgenic sheep produced by transfer of nuclei from transfected fetal fibroblasts. *Science*, 278 : 2130-2133.
 - 29) Shin, S.J., Lee, B.C., Park, J.I., Lim, J.M. and Hwang, W.S. 2001. A separate procedure of fusion and activation in an ear fibroblast nuclear transfer program improves preimplantation development of bovine reconstituted oocytes. *Theriogenology*, 55 : 1697-1704.
 - 30) Shin, T., Kraemer, D., Pryor, J., Liu, L., Rugila, J., Howe, L., Buck, S., Murphy, K., Lyons, L. and Westhusin, M. 2002. A cat cloned by nuclear transplantation. *Nature*, 415 : 859.
 - 31) Sun, Q. Y., Lai, L., Bonk, A., Prather, R. S. and Schatten, H. 2001. Cytoplasmic changes in relation to nuclear maturation and early embryo developmental potential of porcine oocytes : effects of gonadotropins, cumulus cells, follicular size, and protein synthesis inhibition. *Mol. Reprod. Dev.*, 59, 192-198.
 - 32) Takahashi, Y. and First, N.L. 1992. In vitro development of bovine one-cell embryos : influence of glucose, lactate, pyruvate, amino acids and vitamins. *Theriogenology*, 37 : 963-978.
 - 33) Tao, T., Machaty, Z., Boquest, A.C., Day, B. N. and Prather, R.S. 1999. Development of pig embryos reconstructed by microinjection of cultured fetal fibroblast cells

- into *in-vitro* matured oocytes. *Anim. Reprod. Sci.*, 56 : 133-141.
- 34) Uhm, S.J., Chung, H.M., Kim, C. Shim, H., Kim, N.H., Lee, H.T. and Chung, K.S. 2000. *In-vitro* development of porcine enucleated oocytes reconstructed by the transfer of porcine fetal fibroblasts and cumulus cells. *Theriogenology*, 54 : 559-570.
- 35) Wakayama, T., Perry, A.C., Zuccotti, M., Johnson, K.R. and Yanagimachi, R. 1998. Full-term development of mice from enucleated oocytes injected with cumulus cell nuclei. *Nature*, 394 : 369-374.
- 36) Wells, D.N., Misica, P.M., Tervit, H.R. and Vivanco, W.H. 1998. Adult somatic cell nuclear transfer is used to preserve the last surviving cow of the Enderby Island cattle breed. *Reprod. Fertil. Dev.*, 10 : 369-378.
- 37) Wilmut, I., Schnieke, A.E., McWhir, J., Kind, A.J. and Campbell, K.H. 1997. Viable offsprings derived from fetal and adult mammalian cells. *Nature*, 385 : 810-813.