



Title	Immunohistochemical localization of the estrogen receptor alpha (ER) and progesterone receptor (PR) in the uterus of sika deer (<i>Cervus nippon</i>) during pregnancy
Author(s)	Yanagawa, Yojiro; Matsuura, Yukiko; Suzuki, Masatsugu; Katagiri, Seiji; Tsubota, Toshio
Citation	Japanese Journal of Veterinary Research, 56(3), 139-149
Issue Date	2008-11
DOI	10.14943/jjvr.56.3.139
Doc URL	http://hdl.handle.net/2115/35114
Type	bulletin (article)
File Information	56-3_p139-149.pdf



[Instructions for use](#)

Immunohistochemical localization of the estrogen receptor alpha (ER α) and progesterone receptor (PR) in the uterus of sika deer (*Cervus nippon*) during pregnancy

Yojiro Yanagawa¹⁾, Yukiko Matsuura^{2,*}, Masatsugu Suzuki³⁾,
Seiji Katagiri⁴⁾ and Toshio Tsubota¹⁾

¹⁾Laboratory of Wildlife Biology, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

²⁾Field Science Center for Northern Biosphere, Hokkaido University, Sapporo, 060-0809, Japan

³⁾Laboratory of Zoo and Wildlife Medicine, Department of Veterinary Medicine, Faculty of Applied Biological Sciences, Gifu University, Gifu 501-1193, Japan

⁴⁾Laboratory of Theriogenology, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

Received for publication, October 14, 2008; accepted, October 28, 2008

Abstract

Information on steroid hormone receptor distribution in the uterus is essential to understand the roles of their ligands in pregnancy. This study examined the spatio-temporal localization of estrogen receptor alpha (ER α) and progesterone receptor (PR) in the uterus of sika deer (*Cervus nippon*) to determine the estrogen and progesterone action site during pregnancy. Ovaries and uteri were collected from 21 pregnant sika deer with single fetus and two corpora lutea, ranging from Day 20 to Day 207 of pregnancy. In addition, genital organs were also collected from three sika deer whose gestational status was unknown: one female had only one developing corpus luteum: \leq Day 4 (metestrus) and two females had two corpora lutea, one of which was at the developing stage equivalent to diestrus or early pregnancy: $>$ Day 7 (diestrus). Staining of ER α and PR was clear in all cell types during metestrus. During diestrus, the presence of ER α was also clear in deep glandular epithelium, stroma and myometrium, whereas it was suppressed in luminal epithelium and shallow glandular epithelium. Staining of PR was suppressed in luminal epithelium but was detectable in other cell types. Staining of ER α in all cell types and PR in luminal epithelium and glandular epithelium became undetectable by Day 28. PR was presented in stroma and myometrium throughout pregnancy. The distribution pattern of ER α and PR was different during diestrus from that in a ruminant. This could be attributed to estrogen secretion from the maturing and ovulating follicles in the presence of developed corpus luteum.

Key words: pregnancy, sika deer, steroid hormone receptor, uterus

*Corresponding author: Yukiko Matsuura, Field Science Center for Northern Biosphere, Hokkaido University, Sapporo 060-0809, Japan
Phone: +81-11-706-3653. Fax: +81-11-706-3450. E-mail: koyum@seagreen.ocn.ne.jp

Introduction

Sika deer (*Cervus nippon*) is a seasonal breeder, mating in autumn and fawning in early summer^{26,30} after 224-230 days' gestation^{29,30}. Ovarian steroid hormones should play a central role in the regulation of uterine function to establish and maintain pregnancy. Information on the steroid hormone receptor distribution in the uterus is, thus, essential to understand when and where steroid hormones (i.e., estrogen and progesterone) are required in the uterus during the estrous cycle and pregnancy³². Spatio-temporal distribution of estrogen receptor (ER) and progesterone receptor (PR) has been investigated in various domestic species, such as sheep³⁹, cattle²⁵, pigs^{16,17} and horses²¹, and has provided insights into uterine physiology. In ruminants, including cervid species, luteolysis, which is triggered by secretion of prostaglandin F_{2α} (PGF_{2α}) from the uterus, must be prevented to establish pregnancy^{19,28}. As PGF_{2α} secretion is triggered by oxytocin through an up-regulation of oxytocin receptor (OTR) which has been up-regulated by estrogen, the distribution pattern of ER is thought to be important in the onset of PGF_{2α} secretion in sheep^{39,40,43}. On the other hand, progesterone, the hormone of pregnancy, is essential to create a uterine environment that supports the development of the conceptus to term⁴⁴. It also influences uterine histotroph secretion, presumably aiding embryo survival to maintain pregnancy in sheep⁴¹. However, no study has documented the distribution of steroid hormone receptors in the uterus of cervid species.

This study was conducted to characterize the spatio-temporal distribution of endometrial and myometrial ER alpha (ERα) and PR protein during the gestation period to understand the physiological roles of estrogen and progesterone. It is known that over 80% of pregnant sika deer possess two steroidogenic corpora lutea (CLs), known to be formed at a different period, even though they have a single fetus^{31,47-49}. Therefore, uterine samples from sika deer with two CLs and single fetus were exclusively used in this study except for a

single sample (No. 1, Table 1), which represents the time between two ovulations.

Materials and Methods

Animals and uterine samples: Ovaries and uteri were collected in the wild from 21 pregnant sika deer (No.4-24, Table 1) and three additional females whose gestational status could not be determined (No.1-3, Table 1). The animals were shot legally on the island of Hokkaido, Japan, in January 2004, November 2005, January to May 2006 and November 2007 to January 2008. All pregnant females (No.4-24) had a single fetus and two CLs in the ovaries. Among three additional females (No.1-3), female No.1 had only one developing CL with a cavity and clot in it, indicating that she was most likely within four days after estrus²² (metestrus). Females No.2 and 3 had two CLs, one fully developed and the other at the developing stage. They were assumed to be seven days after estrus or later (diestrus), since development of CL takes about seven days in cattle and sheep^{34,36}. Since we failed to recover an embryo from these three animals, they were thus considered to be non-pregnant in this study. In sheep, it is known that there is no difference in uterine steroid hormone receptor distribution between pregnant and non-pregnant animals until Day 11³⁹. However it started to be different on Day 13: the period corresponded to the time of interferon tau (INFτ), the factor of maternal recognition of pregnancy, secretion by an embryo¹⁸. Since INFτ is known to be secreted by the embryo from Day 14¹² in cervid species, uterine steroid hormone receptor distribution is assumed to be the same until Day 14 regardless of the gestational status. Therefore, three non-pregnant animals are used to estimate the steroid hormone receptor distribution in early pregnancy.

Estimation of fetal age: Fetal weights were measured with an electronic balance or spring scale, and crown-rump lengths were measured with a caliper. Fetal age was estimated by crown-rump

length up to 60 days ($\hat{=}$ crown-rump length 80.6 mm) of pregnancy⁵²⁾ ($age = -2.08 + 14.15LnX$, X = crown-rump length) and fetuses over 60 days old (CRL > 80.6 mm) were estimated by fetal weight according to Suzuki *et al.*⁴⁸⁾ ($age = (\sqrt[3]{X} + 2.73)/0.091$, X = fetal body weight). The fetal ages ranged from Day 20 to day 207. Since the gestational length of sika deer is reported to be about 224-230 days^{29,30)}, samples covered almost the whole range of gestation periods.

Immunohistochemistry: Uteri were fixed within one hour after killing and preserved in 10% phosphate-buffered formalin for 3-18 months before embedding. Fixed tissues from the dorsal part of uterine horn in which the fetus was present were used as the sample for immunohistochemistry. In three animals (No.1-3), in which the presence of an embryo was not confirmed, tissue samples were obtained from the uterine horns ipsilateral to the ovary with the CL. After fixation, the specimens were dehydrated in an ethanol series and embedded in paraffin. Tissues were cut into 5- μ m-thick paraffin sections and mounted on silane-coated slide glass (MAS, S9226, Matsunami, Osaka, Japan).

To reveal the presence and distribution of ER α and PR in the uteri, the avidin-biotin immunoperoxidase technique (Vectastain[®] ABC kit; Vector Laboratories, Inc., Burlingame, U.K.) was used. After deparaffinization through an ethanol gradient and rehydration in phosphate-buffered salines, the sections were incubated briefly in an antigen-unmasking solution, Target Retrieval Solution (Dako Cytomation, Carpinteria, CA, USA) for 40 min at 99°C. The sections were then incubated in 3% hydrogen peroxide/methanol solution for 10 min to block the endogenous peroxidase activity, and treatment with blocking serum was applied to each section to prevent nonspecific reactions. Primary antibodies were added to sections overnight at 4°C for incubation. The rabbit polyclonal antibody (PA1-309, Affinity Bioreagents, Golden, CO, U.S.A.) to the synthetic peptide corresponds to amino acid residues 21-32 from human ER α ⁵⁾

which were completely conserved in several species, including the cervid species (accession X98007) and cross react with human, porcine, and rat ER α was used as primary antibody to ER α . Also the mouse monoclonal antibody (Ab-8, Neo Markers, Fremont, CA, U.S.A.) to human PR²⁰⁾ which cross reacts with human, horse, sheep and pig PR was used as primary antibody to PR. Negative controls were treated with normal rabbit serum or normal mouse serum. Colorization was performed by 3, 3'-diaminobenzidine-H₂O₂ solution (0.02% w/v) for 5 to 10 min. Steroid receptor staining intensities were scored visually referring to Spencer and Bazer³⁹⁾ (- absent, \pm scattering, + weak, ++ moderate, or +++ strong) in the luminal epithelium (LE), glandular epithelium (GE), endometrial stroma (ST) and myometrium (MYO). In addition, GE and ST were distinguished into shallow and deep parts when there was a difference in staining. We classify the area adjacent to epithelium as shallow and the area adjacent to myometrium as deep region.

Results

Estrogen receptor localization

Immunoreactive ER α were detected predominantly in the nuclei of uterine cells. Results of ER α staining in uteri are summarized in Table 1. In the uterus of No.1 (\leq Day 4), strong staining of ER α was observed in the deep GE, ST, and MYO, while the LE and shallow GE showed moderate staining (Fig. 1A, B). In deep GE, ST and MYO of uteri of female No. 2 and 3, ER α staining was similar to that in uterus of No. 1 (Fig. 1D), while the staining in the LE and shallow GE was weaker in No. 2 than in No. 1 (Fig. 1C). Staining of ER α in LE became undetectable by Day 20 (No. 4), and remained undetectable up to Day 207 (No. 24). In the deep GE, ST and MYO, weak to moderate ER α staining were observed on Day 20 (No. 4, Fig. 1E), but the staining became undetectable by Day 28 (No. 7) and thereafter (Fig. 1F). The staining became detectable during late pregnancy again (Day

164~, Fig. 1G). No appreciable staining was detected in uterine tissue sections incubated with normal rabbit serum substituted for primary antibody (Fig. 1H).

Progesterone receptor localization

Immunostaining for PR was localized predominantly in the nuclei of uterine cell. Results of PR staining in uteri are summarized in Table 1. The staining of PR in LE was only detected in No. 1 (\leq Day 4, Fig. 2A) and No. 4 (Day 20). In the GE, the staining of PR was detectable in No. 1-3 (Fig. 2 A-D) and No. 4 (Day 20, Fig. 2E), and became undetectable by Day 25 (No. 6). In contrast, moderate to strong staining was detectable in the ST and

MYO in all samples (Fig. 2F). From Day 164 to Day 207 (No. 21-24), they were consistently detected as strong (Fig. 2G). No appreciable staining was observed in sections incubated with normal mouse serum in place of primary antibody (Fig. 2 H).

Discussion

This is the first study to describe the distribution of steroid hormone receptors in the uterus of cervid species. The results indicate that distribution of uterine ER α and PR is differentially regulated in both a spatial and temporal manner dur-

Table 1. Summary of immunolocalization of steroid hormone receptors in the sika deer uterus

ID	Status	Estimated days of gestation ^{a)}	ER α							PR					
			GE			ST		MYO	GE			ST			
			LE	Shallow	Deep	Shallow	Deep		LE	Shallow	Deep	Shallow	Deep	MYO	
No. 1	metestrus	≤ 4	++	++	+++	+++	+++	+++	++	+++	+++	+++	+++	+++	
No. 2	diestrus	> 7	+	++	+++	+++	+++	++	-	+++	+++	++	++	+++	
No. 3	diestrus	> 7	\pm	+	+++	+++	+++	+++	-	\pm	++	++	+++	+++	
No. 4	Pregnant	20	-	+	+++	++	+	+	+	++	+++	++	++	+++	
No. 5	Pregnant	23	-	\pm	\pm	+	+	+	-	\pm	-	++	++	+++	
No. 6	Pregnant	25	-	-	-	+	-	-	-	-	-	++	++	++	
No. 7	Pregnant	28	-	-	-	-	-	-	-	-	-	++	++	++	
No. 8	Pregnant	45	-	-	-	-	-	-	-	-	-	++	++	++	
No. 9	Pregnant	50	-	-	-	-	-	-	-	-	-	++	++	++	
No. 10	Pregnant	51	-	-	-	-	-	-	-	-	-	++	++	++	
No. 11	Pregnant	57	-	-	-	-	-	-	-	-	-	++	++	++	
No. 12	Pregnant	58	-	-	-	-	-	-	-	-	-	+++	++	+++	
No. 13	Pregnant	60	-	-	-	-	-	-	-	-	-	+++	++	++	
No. 14	Pregnant	77	-	-	-	-	-	-	-	-	-	++	++	++	
No. 15	Pregnant	108	-	-	-	-	-	-	-	-	-	++	++	++	
No. 16	Pregnant	115	-	-	-	-	-	-	-	-	-	++	++	++	
No. 17	Pregnant	115	-	-	-	-	-	-	-	-	-	++	+++	+++	
No. 18	Pregnant	118	-	-	-	-	-	-	-	-	-	++	+++	+++	
No. 19	Pregnant	126	-	-	-	-	-	-	-	-	-	++	++	++	
No. 20	Pregnant	127	-	-	-	-	-	-	-	-	-	++	++	++	
No. 21	Pregnant	164	-	+	+	+	+	+	-	-	-	+++	+++	+++	
No. 22	Pregnant	197	-	\pm	+	+	+	+	-	-	-	+++	+++	+++	
No. 23	Pregnant	197	-	\pm	+	+	+	+	-	-	-	+++	+++	+++	
No. 24	Pregnant	207	-	+	+	+	+	+	-	-	-	+++	+++	++	

ER α , estrogen receptor alpha; PR, progesterone receptor; LE, luminal epithelium; GE, glandular epithelium; ST, stroma; MYO, myometrium; CL, corpus luteum;

^{a)}Fetal ages were estimated by crown-rump length up to 60 days according to Yanagawa *et al.*⁵²⁾, and by fetal body weight thereafter according to Suzuki *et al.*⁵⁰⁾. Days for No.1-3 are estimated days after estrus.

-, Absent; \pm , Scattering; +, Weak; ++, Moderate; +++, Strong




Fig. 1. Immunohistochemical localization of estrogen receptor alpha in the uterus of sika deer.

(A) Endometrium of female with one CL (No. 1). (B) Myometrium of female with one CL (No. 1). (C) Endometrium of female with two CLs, developed and developing CL (No. 3). (D) Myometrium of female with two CLs, developed and developing CL (No. 3). (E) Myometrium of female on Day 20 of pregnancy (No. 4). (F) Endometrium and myometrium of female in mid-pregnancy (No. 12). (G) Myometrium of female in late pregnancy (No. 21). (H) Negative control. LE, luminal epithelium; GE, glandular epithelium; ST, stroma; MYO, myometrium. Scale bar = 100 μ m

ing pregnancy. The distribution pattern of ER α and PR differed between the endometrium and myometrium, and also among cell types in the endometrium.

In the uterus with one developing CL assumed to be within several days after estrus (No. 1), the staining of ER α and PR were strong or moderate in all cell types. This is similar to that of a cyclic ruminant^{25,39}. The strong or moderate staining of ER α and PR may be due to the influence of a high concentration of circulating estrogen from the graafian follicle at the estrus, since estrogen is known to up-regulate the ER and PR⁵¹.

Although the present results show a trend in which ER staining decreases with days after estrus is similar to the change of ER in sheep, a domestic species in which steroid receptor distribution has been well documented, the time and extent of decline may differ between deer and sheep. In sheep, staining of ER in all cell types decreased dramatically by Day 11 of the estrous cycle or pregnancy and remained weak thereafter, when animals are pregnant, while staining of ER α delayed to decrease in sika deer and was detectable as late as Day 25 in the ST. The difference in the time of ER suppression between sheep and deer could be attributed to estrogen secretion by the follicle which is maturing and ovulating after formation of the first CL (>Day 7). This hypothesis could be verified when ER staining in pregnant deer, which does not form a second CL, decreased around Day 11 as in sheep. Alternatively, the delay in suppression of ER staining in shallow GE, ST and MYO up to Day 25 may suggest a potential role of estrogen in these cell types during early pregnancy in this species.

Lower staining of ER α in the LE and shallow GE than in the other part of the uterus is comparable to sheep. Since OTR appearance is regulated by estrogen, down-regulation of ER may have the advantage of preventing the luteolytic mechanism via OTR down-regulation as oxytocin stimulates the synthesis of luteolytic PGF $_{2\alpha}$ secretion. Down-regulation of ER in LE and shallow GE may be important since PGF $_{2\alpha}$ is thought to be secreted from

LE and shallow GE because of the specific localization of cyclooxygenase-2, an important enzyme for PGF $_{2\alpha}$ synthesis in these cell types in sheep⁸. The mechanism of ER appearance which is regulated differently between cell types is not clear, but down-regulation of ER in LE and shallow GE might be due to the effect of INF τ . INF τ produced by the conceptus and involved in the maternal pregnancy recognition^{2,40,41,45} is known to act in a paracrine fashion on LE and shallow GE to suppress transcription of ER and OTR genes^{14,33,42}, thereby preventing development of the endometrial luteolytic mechanism in sheep^{39,40}. In red deer, INF τ is reportedly produced by conceptus, and is known to have an anti-luteolytic effect^{1,11,12}. Therefore, INF τ may have been responsible for the reduced staining of ER α in LE and GE, which was similar to that in sheep, although the gestational status (i.e., the presence of an embryo) of No. 2 and 3 was unknown.

Staining of PR in uteri of No. 2 and 3 was relatively low in LE. Moreover, staining in LE and GE declined after Day 20 and became undetectable by Day 25. Decline of PR in LE and GE may be due to decline of PR up-regulation by estrogen since ER α appearance was suppressed at this period, or to prolonged exposure to progesterone action since progesterone has negative effects on PR appearance^{9,40,53}. On the other hand, high staining of PR in ST and MYO was detected during early pregnancy (Day 20-25) and might be important to establish and maintain pregnancy.

During Day 28 to Day 127 (No.7-20), ER α staining diminished in all cell types, possibly due to a high level of circulating progesterone as it down-regulated ER appearance^{13,40,41} and to prevent PGF $_{2\alpha}$ secretion and myometrial contraction⁶ for the maintenance of pregnancy. On the other hand, the reactivity of PR remained high in ST and MYO, consistent with that observed in sheep and cattle^{4,45,46}. The PR in ST may be required for proliferation of the endometrium which is stimulated by stromal cell-derived growth factors induced by progesterone⁴⁴. Presence of PR in MYO throughout pregnancy might also indicate a direct influ-




Fig. 2. Immunohistochemical localization of progesterone receptor in the uterus of sika deer.

(A) Endometrium of female with one CL (No. 1). (B) Myometrium of female with one CL (No. 1). (C) Endometrium of female with two CLs, developed and developing CL (No. 3). (D) Myometrium of female with two CLs, developed and developing CL (No. 3). (E) Myometrium of female Day 20 of pregnancy (No. 4). (F) Myometrium of female in mid-pregnancy (No. 16). (G) Myometrium of female in late pregnancy (No. 21). (H) Negative control. LE, luminal epithelium; GE, glandular epithelium; ST, stroma; MYO, myometrium. Scale bar = 100 μ m

ence of progesterone on contractile activity of this layer, finally resulting in myometrial quiescence⁶.

In late pregnancy, from Day 164 and later (No. 21-24), ER α staining was detected weakly in all uterine cell types except for LE. It may be due to secretion of estrogen toward parturition as reported in domestic animals²³. Increase in estrogen secretion toward term is also reported in cervid species; red deer²⁴, reindeer^{37,38} and white-tail deer³⁵. The presence of ER α in LE and GE is crucial in late pregnancy since ER up-regulation may increase OTR appearance^{3,15} which stimulates production of PGF_{2 α} ^{39,40,43}, a luteolytic factor. Also, the presence of ER α in ST might play an important role in luteolysis since estrogen induces epithelial proliferation through ER-positive ST¹⁰. The presence of ER in MYO may be important because oxytocin, which promotes myometrial contractility, coacted with PGF_{2 α} during late pregnancy and parturition^{6,7}. Interestingly, the timing of ER α up-regulation, which may be caused by estrogen, was about eight weeks prepartum in sika deer while estrogen secretion has been reported to increase about six weeks prepartum in other cervid species^{24,35,37,38}. Paired samples of uteri and peripheral blood of sika deer of known gestational age are needed to confirm the timing in increase of estrogen and up-regulation of ER in sika deer.

Acknowledgments

We would like to thank Professor S. Kondo, Dr. C. Yayota, Mr. K. Nishitani, Dr. M. Miyaji (Graduate School of Agriculture, Hokkaido University), Mr. K. Nishino (Field Science Center for Northern Biosphere, Hokkaido University) and Mr. Seino (School of Veterinary Medicine, Obihiro University of Agriculture and Veterinary Medicine) for their generous assistance in the field sampling. We are also indebted to Dr. H. Igota (Department of Biosphere and Environmental Science, Rakuno Gakuen University) and Mr. J. Igota (Nishiokoppe Wildlife Association) for providing us with samples. Our thanks also to Dr. M. Sasaki and Mr. D. Hay-

akawa (United Graduate School of Veterinary Science, Obihiro University of Agriculture and Veterinary Medicine) for their technical support for immunohistochemistry. This work was partially supported by a grant from JSPS (No. KAKENHI 19688013).

Reference

- 1) Bainbridge, D. R. and Jabbour, H. N. 1999. Source and site of action of anti-luteolytic interferon in red deer (*Cervus elaphus*): possible involvement of extra-ovarian oxytocin secretion in maternal recognition of pregnancy. *J. Reprod. Fertil.*, **116**: 305-313.
- 2) Bazer, F., Spencer, T. E., Ott, T. L. and Johnson, H. M. 1994. Cytokines and pregnancy recognition. In: *Immunobiology of Reproduction*. pp. 37-56, Hunt, J. ed., Springer-Verlag, New York.
- 3) Beard, A. P., Hunter, M. G. and Lamming, G. E. 1994. Quantitative control of oxytocin-induced PGF_{2 α} release by progesterone and oestradiol in ewes. *J. Reprod. Fertil.*, **100**: 143-150.
- 4) Boos, A., Kohtes, J., Janssen, V., Mulling, C., Stelljes, A., Zerbe, H., Hassig, M. and Thole, H. H. 2006. Pregnancy effects on distribution of progesterone receptors, oestrogen receptor alpha, glucocorticoid receptors, Ki-67 antigen and apoptosis in the bovine interplacentomal uterine wall and foetal membranes. *Anim. Reprod. Sci.*, **91**: 55-76.
- 5) Cardenas, H. and Pope, W. F. 2004. Attenuation of estrogenic effects by dihydrotestosterone in the pig uterus is associated with down-regulation of the estrogen receptors. *Biol. Reprod.*, **70**: 297-302.
- 6) Challis, J. R. G., Matthews, S. G., Gibb, W. and Lye, S. J. 2000. Endocrine and paracrine regulation of birth at term and preterm. *Endocr. Rev.*, **21**: 514-550.
- 7) Chard, T. 1989. Fetal and maternal oxytocin in human parturition. *Am. J. Perinatol.*, **6**: 145-152.
- 8) Charpigny, G., Reinaud, P., Tamby, J. P., Cre-

- minon, C., Martal, J., Maclouf, J. and Guillo-
mot, M. 1997. Expression of cyclooxygenase-1
and -2 in ovine endometrium during the es-
trous cycle and early pregnancy. *Endocrinol-
ogy*, **138**: 2163-2171.
- 9) Clarke, C. L. 1990. Cell-specific regulation of
progesterone receptor in the female reproduc-
tive system. *Mol. Cell. Endocrinol.*, **70**: C29-C
33.
 - 10) Cooke, P. S., Buchanan, D. L., Young, P.,
Setiawan, T., Brody, J., Korach, L. S., Taylor, J.,
Lubahn, D. B. and Cunha, G. R. 1997. Stromal
estrogen receptors mediate mitogenic effects
of estradiol on uterine epithelium. *Proc. Natl.
Acad. Sci. U.S.A.*, **94**: 6535-6540.
 - 11) Demmers, K. J., Kaluz, S., Deakin, D. W., Jab-
bour, H. N. and Flint, A. P. 1999. Production of
interferon by the conceptus in red deer *Cervus
elaphus*. *J. Reprod. Fertil.*, **115**: 59-65.
 - 12) Demmers, K. J., Jabbour, H. N., Deakin, D. W.
and Flint, A. P. F. 2000. Production of inter-
feron by red deer (*Cervus elaphus*) concep-
tuses and the effects of roIFN- τ on the timing
of luteolysis and the success of asynchronous
embryo transfer. *J. Reprod. Fertil.*, **118**: 387-
395.
 - 13) Evans, R. W. and Leavitt, W. W. 1980. Proges-
terone inhibition of uterine nuclear estrogen
receptor: dependence on RNA and protein syn-
thesis. *Proc. Natl. Acad. Sci. U.S.A.*, **77**: 5856-
5860.
 - 14) Fleming, J. A., Choi, Y., Johnson, G. A., Spencer,
T. E. and Bazer, F. W. 2001. Cloning of the
ovine estrogen receptor-alpha promoter and
functional regulation by ovine interferon-tau.
Endocrinology, **142**: 2879-2887.
 - 15) Fuchs, A. R., Helmer, H., Behrens, O., Liu, H.
C., Antonian, L., Chang, S. M. and Fields, M. J.
1992. Oxytocin and bovine parturition: a steep
rise in endometrial oxytocin receptors pre-
cedes onset of labor. *Biol. Reprod.*, **47**: 937-944.
 - 16) Geisert, R. D., Brenner, R. M., Moffatt, R. J.,
Harney, J. P., Yellin, T. and Bazer, F. W. 1993.
Changes in oestrogen receptor protein, mRNA
expression and localization in the en-
dometrium of cyclic and pregnant gilts. *Re-
prod. Fertil. Dev.*, **5**: 247-260.
 - 17) Geisert, R. D., Pratt, T. N., Bazer, F. W., Mayes,
J. S. and Watson, G. H. 1994. Immunocyto-
chemical localization and changes in en-
dometrial progesterin receptor protein during
the porcine oestrous cycle and early pregnancy.
Reprod. Fertil. Dev., **6**: 749-760.
 - 18) Godkin, J. D., Bazer, F. W., Moffatt, J., Sessions,
F. and Roberts, R. M. 1982. Purification and
properties of a major, low molecular weight
protein released by the trophoblast of seep
blastocysts at day 13-21. *J. Reprod. Fertil.*, **65**:
141-150
 - 19) Godkin, J. D., Smith, S. E., Johnson, R. D. and
Dore, J. J. 1997. The role of trophoblast inter-
ferons in the maintenance of early pregnancy
in ruminants. *Am. J. Reprod. Immunol.*, **37**:
137-143.
 - 20) Gray, G. O. and Satyaswaroop, P. G. 1988. Spe-
cies crossreactivity of human progesterone re-
ceptor monoclonal antibodies: Western blot
analysis. *Biochem. Biophys. Res. Commun.*,
157: 1067-1077.
 - 21) Hartt, L. S., Carlin, S. J., Joyce, M. M., Johnson,
G. A., Vanderwall, D. K. and Ott, T. L. 2005.
Temporal and spatial associations of oestrogen
receptor alpha and progesterone receptor in
the endometrium of cyclic and early pregnant
mares. *Reproduction*, **130**: 241-250.
 - 22) Ireland, J. J., Murphee, R. L. and Coulson, P. B.
1980. Accuracy of predicting stages of bovine
estrous cycle by gross appearance of the cor-
pus luteum. *J. Dairy Sci.*, **63**: 155-160.
 - 23) Jenkin, G. and Young, I. R. 2004. Mechanisms
responsible for parturition; the use of experi-
mental models. *Anim. Reprod. Sci.*, **82-83**: 567
-581.
 - 24) Kelly, R. W., McNatty, K. P., Moore, G. H., Ross,
D. and Gibb, M. 1982. Plasma concentrations
of LH, prolactin, oestradiol and progesterone
in female red deer (*Cervus Elaphus*) during
pregnancy. *J. Reprod. Fertil.*, **64**: 475-483.
 - 25) Kimmins, S. and MacLaren, L. A. 2001. Oes-
trous cycle and pregnancy effects on the distri-

- bution of oestrogen and progesterone receptors in bovine endometrium. *Placenta*, **22**: 742-748.
- 26) Koizumi, T. 1991. Reproductive characteristics of female sika deer, *Cervus nippon*, in Hyogo prefecture, Japan. In: *Ongulès/Ungulates*. pp. 561-563, Spitz, F., Janeau, G., Gonzalez, G. and Aulagnier, S. eds., Société Française pour l'Etude et la Protection des Mammifères-Institut de Recherche sur les Grands Mammifères, Paris.
- 27) Lazar, M. 2008. Mechanism of action of hormones that act on nuclear receptors. In: *Williams Textbook of Endocrinology*. 11th ed. pp. 37-46, Kronenberg, H. ed., Saunders/Elsevier, Philadelphia.
- 28) Martal, J., Chene, N., Camous, S., Huynh, L., Lantier, F., Hermier, P., L'Haridon, R., Charpigny, G., Charlier, M. and Chaouat, G. 1997. Recent developments and potentialities for reducing embryo mortality in ruminants: the role of IFN-tau and other cytokines in early pregnancy. *Reprod. Fertil. Dev.*, **9**: 355-380.
- 29) Matsuura, Y., Sato, K., Suzuki, M. and Ohtaishi, N. 2004. The effects of age, body weight and reproductive status on conception dates and gestation periods in captive sika deer. *Mammal Study*, **29**: 15-20.
- 30) Matsuura, Y., Sasamoto, Y., Sato, K., Takahashi, Y., Suzuki, M. and Ohtaishi, N. 2004. Monitoring ovarian cycle and conception by fecal progesterone analysis in sika deer. *Ecol. Res.*, **19**: 397-404.
- 31) Matsuura, Y., Suzuki, M., Hayakawa, D., Asano, M., Sasaki, M., Kitamura, N., Yamada, J., Tsubota, T. and Ohtaishi, N. 2004. Immunohistochemical localization of steroidogenic enzymes in corpus luteum of wild sika deer during early mating season. *Jpn. J. Vet. Res.*, **51**: 167-172.
- 32) Meikle, A., Tasende, C., Sosa, C. and Garofalo, E. G. 2004. The role of sex steroid receptors in sheep female reproductive physiology. *Reprod. Fertil. Dev.*, **16**: 385-94.
- 33) Mirando, M., Harney, J. P., Zhou, Y., Ogle, T. F., Ott, T. L., Moffatt, R. J. and Bazer, F. W. 1993. Changes in progesterone and oestrogen receptor mRNA and protein and oxytocin receptors in endometrium of ewes after intrauterine injection of ovine trophoblast interferon. *J. Mol. Endocrinol.*, **10**: 185-192.
- 34) Noakes, D. 1997. Ovarian changes during the cycle. In: *Fertility and Obstetrics in Cattle*. pp.4-5. Sutton, J. B. and Swift, S. T. eds, Blackwell Science, Cambridge.
- 35) Plotka, E. D., Seal, U. S., Verme, L. J. and Ozoga, J. J. 1977. Reproductive steroids in the white-tailed deer (*Odocoileus virginianus borealis*). 2. Progesterone and estrogen levels in peripheral plasma during pregnancy. *Biol. Reprod.*, **17**: 78-83.
- 36) Robertson, H. A. 1977. Reproduction in the ewe and the goat. In: *Reproduction in Domestic Animals*. pp. 475-498, Cole, A.A. and Cupps, P.T. eds., Academic Press, New York.
- 37) Ropstad, E., Veiberg, V., Sakkinen, H., Dahl, E., Kindahl, H., Holand, O., Beckers, J. F. and Eloranta, E. 2005. Endocrinology of pregnancy and early pregnancy detection by reproductive hormones in reindeer (*Rangifer tarandus tarandus*). *Theriogenology*, **63**: 1775-1788.
- 38) Shipka, M. P., Rowell, J. E. and Sousa, M. C. 2007. Steroid hormone secretion during the ovulatory cycle and pregnancy in farmed Alaskan reindeer. *J. Anim. Sci.*, **85**: 944-951.
- 39) Spencer, T. E. and Bazer, F. W. 1995. Temporal and spatial alterations in uterine estrogen receptor and progesterone receptor gene expression during the estrous cycle and early pregnancy in the ewe. *Biol. Reprod.*, **53**: 1527-1543.
- 40) Spencer, T. E., Becker, W. C., George, P., Mirando, M. A., Ogle, T. F. and Bazer, F. W. 1995. Ovine interferon- τ inhibits estrogen receptor up-regulation and estrogen-induced luteolysis in cyclic ewes. *Endocrinology*, **136**: 4932-4944.
- 41) Spencer, T. E., Becker, W. C., George, P., Mirando, M. A., Ogle, T. F. and Bazer, F. W. 1995. Ovine interferon- τ regulates expression

- of endometrial receptors for estrogen and oxytocin but not progesterone. *Biol. Reprod.*, **53**: 732-745.
- 42) Spencer, T. E. and Bazer, F. W. 1996. Ovine interferon tau suppresses transcription of the estrogen receptor and oxytocin receptor genes in the ovine endometrium. *Endocrinology*, **137**: 1144-1147.
- 43) Spencer, T. E., Gray, A., Johnson, G. A., Taylor, K. M., Gertler, A., Gootwine, E., Ott, T. L. and Bazer, F. W. 1999. Effects of recombinant ovine interferon tau, placental lactogen and growth hormone on the ovine uterus. *Biol. Reprod.*, **61**: 1409-1418.
- 44) Spencer, T. E. and Bazer, F. W. 2002. Biology of progesterone action during pregnancy recognition and maintenance of pregnancy. *Front. Biosci.*, **7**: 1879-1898.
- 45) Spencer, T. E., Burghardt, R. C., Johnson, G. A. and Bazer, F. W. 2004. Conceptus signals for establishment and maintenance of pregnancy. *Anim. Reprod. Sci.*, **82-83**: 537-550.
- 46) Spencer, T. E., Johnson, G. A., Burghardt, R. C. and Bazer, F. W. 2004. Progesterone and placental hormone actions on the uterus: insights from domestic animals. *Biol. Reprod.*, **71**: 2-10.
- 47) Suzuki, M., Koizumi, T. and Kobayashi, M. 1992. Reproductive characteristics and occurrence of accessory corpora lutea in sika deer *Cervus nippon centralis* in Hyogo prefecture, Japan. *J. Mammal. Soc. Jpn.*, **17**: 11-18.
- 48) Suzuki, M. 1993. Reproductive characteristics and morphology of accessory corpora lutea in sika deer in Hokkaido In: *Deer of China*. pp, 356-363, Ohtaishi, N. and Sheng, H.L. eds., Elsevier, Tokyo.
- 49) Suzuki, M. and Ohtaishi, N. 1993. Reproduction of female sika deer (*Cervus nippon yesoensis* Heude, 1884) in Ashoro district, Hokkaido. *J. Vet. Med. Sci.*, **55**: 833-836.
- 50) Suzuki, M., Kaji, K., Yamanaka, M. and Ohtaishi, N. 1996. Gestational age determination, variation of conception date, and external fetal development of sika deer (*Cervus nippon yesoensis* Heude, 1884) in eastern Hokkaido. *J. Vet. Med. Sci.*, **58**: 505-509.
- 51) Wathes, D. C. and Hamon, M. 1993. Localization of oestradiol, progesterone and oxytocin receptors in the uterus during the oestrous cycle and early pregnancy of the ewe. *J. Endocrinol.*, **138**: 479-492.
- 52) Yanagawa, Y., Matsuura, Y., Suzuki, M., Saga, S., Okuyama, H., Fukui, D., Bandou, G., Katagiri, S., Takahashi, Y. and Tsubota, T. 2009. Fetal age estimation of Hokkaido sika deer (*Cervus nippon yesoensis*) using ultrasonography during early pregnancy *J. Reprod. Dev.*, **55** (in press)
- 53) Zelinski, M. B., Hirota, N. A., Keenan, E. J. and Stormshak, F. 1980. Influence of exogenous estradiol-17beta on endometrial progesterone and estrogen receptors during the luteal phase of the ovine estrous cycle. *Biol. Reprod.*, **23**: 743-751.