



Title	Geochemical Characteristics of Tungsten in Miyako Granitoid and Scheelite-bearing Aplitic Veins at the Yamaguchi Cu-W Skarn Deposit, Iwate, Japan
Author(s)	Ogata, Takeyuki; Ishiyama, Daizo; Mizuta, Toshio; Satoh, Hinako; Matsueda, Hiroharu
Citation	Edited by Shunsuke F. Mawatari, Hisatake Okada., 137-142
Issue Date	2004
Doc URL	http://hdl.handle.net/2115/38518
Type	proceedings
Note	International Symposium on "Dawn of a New Natural History - Integration of Geoscience and Biodiversity Studies". 5-6 March 2004. Sapporo, Japan.
File Information	p137-142-neo-science.pdf



[Instructions for use](#)

Geochemical Characteristics of Tungsten in Miyako Granitoid and Scheelite-bearing Aplitic Veins at the Yamaguchi Cu-W Skarn Deposit, Iwate, Japan

Takeyuki Ogata¹, Daizo Ishiyama², Toshio Mizuta², Hinako Satoh² and Hiroharu Matsueda³

¹ COE for Neo-Science of Natural History, The Hokkaido University Museum, Sapporo 060-0810, Japan

² Faculty of Engineering and Resource Science, Akita University, Akita 010-8502, Japan

³ The Hokkaido University Museum, Sapporo 060-0810, Japan

ABSTRACT

Tungsten analyses were made on 16 samples collected from the North Miyako granitic body in Northeast Japan. Petrographic facies of the North granitic body vary from quartz diorite in the marginal zone (zone A), to tonalite and granodiorite (zone B), and to granite in the central zone (zone C). A large number of barren and scheelite-bearing aplitic veins are distributed around the Yamaguchi deposit which occurs in the contact aureole of zone C granite. The tungsten content of zone C granite is lower than that of the granitic rocks in zones A, B and the aplitic veins. It appears that tungsten in the differentiated granitic magma, which was associated with ore mineralization, was transported out of the magma chamber by magmatic fluids.

The tungsten content is generally low in the North Miyako granitic rocks but high in granitic rocks of Okinoshima zoned pluton, as well as in Otani granite and Busetsu granite from the Southwest Japan. In the case of magnetite-series, however, the behavior of tungsten in the Miyako granitic body from the tungsten metallogenic province is similar to that of the Okinoshima zoned pluton from molybdenum metallogenic province in Southwest Japan. Behavior of tungsten in granitic magma is affected by magmatic evolution during the process of saturation of granitic melt with magmatic fluid.

Keywords: Tungsten, Magmatic fluid, Devolatilization

INTRODUCTION

Ore elements such as tungsten, molybdenum and tin are important for understanding magmatic hydrothermal mineralization related to granitic magmatism. According to the computational models of magmatic devolatilization, behavior of ore metals such as tungsten, molybdenum and tin are affected by the generation of magmatic fluids separated from highly differentiated granitic magma [1].

Ishihara [2] divided the ore related granitoids into

three provinces: molybdenum metallogenic province, tungsten metallogenic province, and tin-tungsten metallogenic province (Fig. 1). The molybdenum and tin-tungsten metallogenic provinces are related to the magnetite-series granitoids and the ilmenite-series granitoids, respectively [2]. The difference between those two provinces is the result of different oxidation states of granitic magma between the two series [2]. On the other hand, magnetite-series granitoids are related to both molybdenum and tungsten metallogenic province. It is possible that difference




Fig. 1 Molybdenum and tungsten-tin metallogenetic zoning of the Japanese islands (after Ishihara [6]).

between tungsten and molybdenum mineralization related to magnetite-series granitic magmatism is affected by mechanism of fractional crystallization and generation of magmatic fluids, separated from highly differential granitic magma, rather than the physicochemical state of granitic magma. In order to understand tungsten mineralization related to granitic magmatism, it is necessary to concentrate on tungsten occurring in granitic rocks.

There are very few previous studies related to the concentration of tungsten in granitic rocks and the genetic relationship among concentrations of ore metals in the granitic rocks [3–7]. In this paper, the tungsten concentration in the Miyako granitic body associated with Yamaguchi W-Cu skarn deposit is described. Tungsten concentration data are applied to deduce the genetic relationship between the ore forming fluid and crystallizing granitic melt, as well as to characterize the granitic magma involved in the mineral-melt-vapor reaction.

OUTLINE OF GEOLOGY

Large numbers of the magnetite-series granitic bodies are distributed in Kitakami belt of the Outer

Zone in Northeast Japan. The Miyako granitic body is one of the Early Cretaceous magnetite-series zoned plutons in the North Kitakami belt. Ishihara [1] divided it into Mo and W metallogenetic provinces of Kitakami belt, based on the location of the Yamaguchi W-Cu skarn deposit.

The Miyako granitic body is distributed in the magnetite-series granitic terrain and tungsten metallogenetic province [1]. It has a size of about 50 km × 20 km in plan view with a balloon-like shape and a northwest-southeast extension (Fig. 2). The age of the emplacement of the Miyako granitoid was estimated by the K-Ar method to be 109–135 Ma [8] and by the Rb-Sr method to be 116–140 Ma [9]. The Miyako granitic body is divided into North and South bodies based on lithologic and compositional differences [10] (Fig. 1). Katada and Kanaya [11] reported that the South body intruded into the North body. The Yamaguchi Cu-W skarn deposit occurs in the contact aureole of the North Miyako granitic body.

On the basis of modal analysis, the North Miyako granitic body consists of quartz diorite (zone A), tonalite to hornblende-biotite grano-diorite (zone B) and hornblende-biotite granite (zone C) (Fig. 2). Ac-




Fig. 2 Simplified geological map of the Miyako granitic body adapted from Nishioka et al. [10]. Numerals are sample localities presented in Table 2.

According to the grain size of hornblende and biotite, the granitic rocks gradually change from fine-grained to coarse and non-equigranular from zone A through zone B to the boundary between zones B and C. The zone C granite frequently includes the country rocks as roof-pendant blocks. Aplite and pegmatite commonly occur in the zone C, close to the Yamaguchi deposit.

The Yamaguchi Cu-W skarn deposit occurs in the contact aureole of zone C granite. The distance between the Yamaguchi deposit and the granitic rocks in zone C is less than 100 m. Tungsten mineralization associated with scheelite (CaWO_4) occurs in aplitic veins. The scheelite-bearing aplitic veins cut massive skarn, which commonly contains the aggregates of fine-grained scheelite crystals mainly near the veins. The scheelite-bearing aplitic veins are abundant of albite, plagioclase, and titanite with subordinate amount of K-feldspar, quartz and scheelite.

ANALYTICAL PROCEDURE AND RESULTS

Major elements and tungsten analyses were made on about 16 samples taken from granitic rocks and

aplitic veins from the North Miyako granitic body. Locations of the analyzed samples are illustrated in Fig. 2. The concentration of SiO_2 was analyzed using XRF at Akita University (RIGAKU 3270) using fusion glasses made from a mixture of powdered sample and alkali fluxes in the proportion of 1 : 5. Tungsten was analyzed by ICP-MS (PQ-3, VG Elemental Co.) at Akita University. The methods of sample digestion and preparation of solution for ICP-MS analyses were the same as described by Sato et al. [12].

Results of analyses on SiO_2 and W are given in Table 1. The results from the duplicate analyses for tungsten concentration in the Japanese geological standards (JB-1 and JG-2) and the Miyako granitic rocks (MA-2, MB-2 and MC-5), used to calculate experimental error are shown in Table 2. Coefficients of variations (CV) of tungsten concentration were less than 12% for concentrations > 0.15 ppm in samples JB-1a, JG-2, MA-2 and MB-2. A larger CV value (34.1%) for MC-5 probably results from the smallest concentration of tungsten. The detection limit of tungsten concentration by ICP-MS, calculated after Yoshida et al. [14], is approximately 0.02

Table 1 Analytical results (contents of SiO₂ and W) for samples of the granitic rocks and aplitic veins

Sample No.	Rock type	SiO ₂ (wt.%)	W (ppm)
<i>North Miyako granitic rocks</i>			
<i>Zone A</i>			
MA-2	Biotite-homblende quartz diorite	61.64	0.40
MA-3	Biotite-homblende quartz diorite	54.55	0.15
MA-4	Biotite-homblende quartz diorite	55.92	0.18
MA-II	Biotite-homblende tonalite	64.67	0.46
<i>Zone B</i>			
MB-2	Biotite-homblende granodiorite	64.20	0.56
MB-4	Biotite-homblende granodiorite	65.69	0.34
MB-5	Biotite-homblende tonalite	62.88	0.63
MB-6	Biotite-homblende tonalite	62.68	0.39
<i>Zone C</i>			
MC-3	Biotite-homblende granite	72.16	0.19
MC-5	Biotite-homblende granite	72.34	0.15
MC-7	Biotite-homblende granite	71.46	0.28
<i>Aplitic veins</i>			
BA-1	Barren aplitic vein	76.42	0.63
BA-2	Barren aplitic vein	75.14	0.93
BA-4	Barren aplitic vein	76.59	1.10
MM-0	Scheelite-bearing aplitic vein	58.38	8000

ppm (2σ).

DISCUSSION

Figure 3 shows the variation diagram of tungsten against SiO₂ concentrations for the North Miyako granitic body in Northeast Japan. For comparison, data for similar plutons in Southwest Japan, such as Okinoyama zoned pluton (magnetite-series: molybdenum metallogenic provinces), the Busetsu granite (ilmenite-series: barren zone) and the Otani granite (ilmenite-series: tungsten metallogenic province) after Ishihara [6, 7] are also plotted in the same diagram.

The tungsten content in the North Miyako granitic body slightly increases from zone A (0.15–0.46 ppm) to zone B (0.34–0.63 ppm) and then decreases from zone B to zone C (0.15–0.28 ppm). However, the tungsten content drastically increases from zone C to the barren aplitic veins (0.63–47.67 ppm). Tungsten content in scheelite-bearing aplitic veins is ca. 0.8 wt% [15] that is much higher than in the granitic rocks from all zones.

The trends of concentration of tungsten in the North Miyako granitic body and aplitic veins are similar to those of copper [15]. Ogata et al. [15] suggested that behavior of Cu in the granitic rocks in the North Miyako granitic body was affected by a

Table 2 Duplicate analytical results (n = 5) of geochemical reference samples and the North Miyako granitic rocks

	R.V. (ppm)	mean (ppm)	σ (ppm)	CV (%)	R.E. (%)
JB-1	17.1	11.87	0.55	4.6	-31
JG-2	23	20.60	0.26	1.3	-10
MA-2	—	0.40	0.03	8.2	—
MB-2	—	0.56	0.06	11.4	—
MC-5	—	0.15	0.05	34.1	—


R.V.: recommended value^[13]

mean: arithmetic means of duplicate analyses

σ : standard deviations of duplicate analyses

CV: coefficients of variation

R.E.: relative error between R.V. and mean value.

**Fig. 3** Tungsten content (ppm) plotted against SiO₂ wt% in granitic rocks.

mechanism that differs from simple magmatic fractional differentiation and also transportation out of the granitic melt by fluid coexisting with the melt. Therefore, tungsten in granitic magma might have been transported out of the magma chamber by magmatic fluids.

The Tungsten concentration in the North Miyako granitic body is lower than that in the other plutons of Southwest Japan (Fig. 3). The tungsten concentration in granitoids is somewhat similar to tin contents, which are generally high in Southwest Japan and low in Northeast Japan [16]. Ishihara and Terashima [16] suggested that the low tin granitoids of Northeast Japan and high tin granitoids of Southwest Japan were formed by granitic magmatism which occurred in intra-oceanic island arc and continental margin, respectively.

Behavior of tungsten in the granitic melt is different between magnetite-series and ilmenite-series granitoids. In the magnetite-series granitoids, variation of tungsten content in the North Miyako granitic body is similar to that in the Okinoyama zoned pluton. On the other hand, tungsten contents in the Otani and Busetsu granites of ilmenite-series increase with increasing SiO₂ concentration. Therefore, it is possible that the Okinoyama granitic magma coexisted with magmatic fluid that transported tungsten out of granitic melt.

Tungsten and molybdenum species are contained mainly in OH complexes and in some cases in Na- and K- hydroxide complexes in the magmatic fluid [17, 18]. The solubility of tungsten in magmatic fluid coexisting with the granitic melt increases with increasing pressure and irrespective of the chloride concentration in magmatic fluid [19]. On the other hand, the solubility of molybdenum in magmatic fluid increases with increasing pressure and/or the chloride concentration in magmatic fluid [18, 19]. Shinohara et al. [20] suggested that the partition ratio of chloride (the ratio of Cl concentration in fluid to Cl concentration in melt) exhibits a strong positive pressure dependence. Therefore, W/Mo ratio of ore-forming magmatic fluid was probably determined by the depth of devolatilization of the granitic melt.

In the Okinoyama zoned pluton, high tungsten concentration among granitic rocks is observed in the granitic porphyry [7], which is the hypabyssal rock emplaced at shallow environment, rather than the North Miyako granitic body. Shimazaki [21] indicated that the depth of formation of the deposit in the inner zone of Southwest Japan is relatively shallower compared to the Kitakami belt of Northeast Japan. It is possible that the depth of emplacement and devolatilization of the North Miyako granitic body is relatively deeper than that of the Okinoyama zoned pluton. Chlorine concentration in apatite from magnetite-series granitoids distributed in the outer Southwest Japan is higher than in other similar granitoids in the Kitakami belt [22]. Therefore, chloride concentration of magmatic fluid coexisting with granitic magma of the North Miyako granitic body is relatively lower than those of the Okinoyama zoned pluton. Additionally, W/Mo ratio of the mineralization in the North Miyako granitic body is relatively higher than that in the Okinoyama zoned pluton, which is related to the high chloride bearing fluid. Ishihara [23] suggested that magnetite-series granitoids in W metallogenic province belonged to a different original magma and tectonic setting which

are characteristic of granitoids in Mo metallogenic province. However, it is emphasized that the W/Mo ratio of the magnetite-series granitoids may be controlled by factors such as the devolatilization depth related to the formation of ore metal complex.

Tungsten and molybdenum mineralization associated with magnetite-series granitic magmatism is affected mainly by magmatic evolution in the process of saturation of a granitic melt with magmatic fluid. Study of behavior of tungsten in granitic melt is important to understand its behavior as an ore metal during volatile exsolution and magmatic crystallization.

ACKNOWLEDGMENTS

The authors thank Dr. O. Fujikawa of the Hokkaido University Museum for his valuable comments. They are grateful to Dr. M.D.P.L. Francis, a COE PD-researcher, for his valuable advice and critical reading of the original manuscript. This research was supported by a 21st Century Center of Excellence (COE) Program on "Neo-Science of Natural History" (Program Leader: Hisatake Okada) at Hokkaido University financed by the Ministry of Education, Culture, Sports, Science and Technology, Japan.

REFERENCES

1. Candela, P.A. and Piccoli, P.M., 1995. Model ore-metal partitioning from melts into vapor and vapor/brine mixtures. *Mineral. Assoc. Canada Short Course*, 23, 101–122.
2. Ishihara, S., 1984. Granitoid series and Mo/W-Sn mineralization in East Asia. *Rept. Geol. Surv. Japan*, 263, 173–208.
3. Kuroda, R., 1956. Distribution of tungsten in granitic rocks in Japan. *J. Chem. Soc. Japan*, 77, 1142–1145.
4. Burrows, D.R. and Spooner, E.T.C., 1987. Generation of a magmatic H₂O-CO₂ fluid enriched in Mo, Au, and W within an Archean Sodic granodiorite stock, Mink Lake, Northwestern Ontario. *Econ. Geol.*, 82, 1931–1957.
5. Keith, J.D., van Middelbaar, W., Clark, A.H. and Hodgson, C.J., 1989. Granitoid texture, compositions, and volatile fugacities associated with the formation of tungsten-dominated skarn deposits. *Rev. Econ. Geology*, 4, 235–251.
6. Ishihara, S., 2002. Chemical characteristics of the mineralized granitoids (I): Mo and W provinces of the Inner Zone of Southwest Japan. *Bull. Geol. Surv. Japan*, 53, 657–672.
7. Ishihara, S., 2002. Chemical characteristics of the mineralized granitoids (II): Polymetallic province of the west-central Hyogo Prefecture, Japan. *Bull. Geol. Surv. Japan*, 53, 673–688.
8. Shibata, K., Matsumoto, T., Yanagi, Y. and Hamamoto, R., 1978. Isotopic ages and stratigraphic control of Mesozoic igneous rocks in Japan. *Contributions to the Geologic Time Scale, AAPG Studies in Geology*, 6, 143–164.
9. Maruyama, T., Miura, K. and Yamamoto, M., 1993. Initial Sr isotopic ratios of the some Late-Mesozoic igneous masses in

- the Kitakami Mountains, northeastern Japan. *Rept. Res. Inst. Nat. Resources Mining Coll. Akita Univ.*, 56, 29–52.
10. Nishioka, Y., Yoshida, T., Kanisawa, S. and Aoki, K., 1987. Geochemistry of the rocks from the Miyako granitic body in the Kitakami Mountains. *Res. Rep. Lap. Nucl. Sci. Tohoku Univ.*, 20, 340–350.
 11. Katada, M. and Kanaya, H., 1980. Potassium, rubidium, strontium, thorium and uranium of the Cretaceous plutons in the Kitakami Mountains. *J. Min. Petr. Econ. Geol.*, 75, 173–185.
 12. Sato, H., Ishiyama, D., Mizuta, T. and Ishikawa, Y., 1999. Rare earth element analysis of rock and thermal water samples by inductively coupled plasma mass spectrometry. *Sci. Tech. Rept. Fac. Eng. Res. Sci. Akita Univ.* No. 20, 1–8
 13. Imai N., Terashima S., Itoh S. and Ando A., 1995. 1994 compilation of analytical data for minor and trace elements in seventeen GSJ geochemical reference samples, "Igneous rock series". *Geostandards Newsletter*, 19, 135–213.
 14. Yoshida, T., Yamasaki, S. and Tsumura, A., 1992. Determination of trace and ultra-trace elements in 32 international geostandards by ICP-MS. *J. Min. Petr. Econ. Geol.*, 87, 107–122.
 15. Ogata, T., Ishiyama, D., Mizuta, T. and Nakashima, K., 2002. Coexistence of fluid phase with granitic magma: Geochemical characteristics of REE and Cu in Miyako granite and scheelite-bearing aplitic veins at the Yamaguchi Cu-W skarn deposit, Iwate, Japan. *Resource Geol.*, 52, 173–186.
 16. Ishihara, S. and Terashima, S., 1977. The tin content of the Japanese granitoids and its geological significance on the Cretaceous magmatism. *J. Geol. Soc. Japan*, 10, 657–664.
 17. Wood, S.A., 1992. Experimental determination of the solubility of WO_3 (s) and the thermodynamic properties of H_2WO_4 (aq) in the range 300–600°C at 1kbar: Calculation of scheelite solubility. *Geochim. Cosmochim. Acta*, 56, 1827–1836.
 18. Webster, J.D., 1999. Exsolution of magmatic volatile phases from Cl-enriched mineralizing granitic magmas and implications for ore metal transport. *Geochim. Cosmochim. Acta*, 61, 1017–1029.
 19. Bai, T.B. and Koster van Groos, A.F., 1999. The distribution of Na, K, Rb, Sr, Al, Ge, Cu, W, Mo, La, and Ce between granitic melts and coexisting aqueous fluids. *Geochim. Cosmochim. Acta*, 63, 1117–1131.
 20. Shinohara, H., Iiyama, T. and Matsuo, S., 1989. Partition of chlorine compounds between silicate melt and hydrothermal solutions: I. Partition of NaCl-KCl. *Geochim. Cosmochim. Acta*, 53, 2617–2630.
 21. Shimazaki, H., 1975. The ratios of Cu/Zn-Pb of pyrometasomatic deposits in Japan and their genetic implications. *Econ. Geology*, 70, 717–724.
 22. Imai, A., 2004. Variation of Cl and SO_3 contents of microphenocrystic apatite in intermediate to silicic igneous rocks in the Western Pacific island arcs and the implication to metallogenesis of porphyry Cu deposits. In: Abstract of *Japan-Swiss Seminar "Spatial and temporal relationships between deep magmatic, porphyry and epithermal environments, and significance for ore formation processes"*, Tsukuba.
 23. Ishihara, S., 1973. The Mo-W metallogenic provinces and the related granitic provinces. *Mining Geol.*, 23, 13–32.