



Title	Arthroscopically Assisted Reattachment of Avulsed Triangular Fibrocartilage Complex to the Fovea of the Ulnar Head
Author(s)	Iwasaki, Norimasa; Minami, Akio
Citation	The Journal of Hand Surgery, 34(7), 1323-1326 <a href="https://doi.org/10.1016/j.jhsa.2009.02.026">https://doi.org/10.1016/j.jhsa.2009.02.026</a>
Issue Date	2009-09
Doc URL	<a href="http://hdl.handle.net/2115/39285">http://hdl.handle.net/2115/39285</a>
Type	article (author version)
File Information	JHS34-7_p1323-1326.pdf



[Instructions for use](#)

**Arthroscopically Assisted Reattachment of Avulsed Triangular Fibrocartilage**

**Complex to the Fovea of the Ulnar Head**

*Surgical Technique*

**K2008-102R1, 2<sup>nd</sup> revision**

Norimasa Iwasaki, MD, PhD; Akio Minami, MD, PhD

Department of Orthopaedic Surgery, Hokkaido University School of Medicine,

Sapporo, Japan

**Running Title:** Reattachment of Avulsed TFCC to the Fovea

**Key words:** Arthroscopic reattachment, Fovea, TFCC, Wrist arthroscopy

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

**Acknowledgment:** The authors thank Dr. Yuichiro Abe for figure preparation and Dr. Tamotsu Kamishima for radiological evaluation.

**Correspondence to:** Norimasa Iwasaki, MD, PhD

Department of Orthopaedic Surgery, Hokkaido University School of Medicine, Kita 15,  
Nishi 7, Kita-Ku, Sapporo 060-8638, Japan.

Phone: 81-11-706-5937; Fax: 81-11-706-6054; E-mail: niwasaki@med.hokudai.ac.jp

1

2        **Arthroscopically Assisted Reattachment of Avulsed Triangular Fibrocartilage**

3                                **Complex to the Fovea of the Ulnar Head**

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

**Abstract**

19

20 Triangular fibrocartilage complex (TFCC) insertion into the fovea of the distal ulna  
21 plays a crucial role in stabilizing the distal radioulnar joint. Consequently, surgical  
22 reattachment against avulsion of the foveal TFCC insertion is required to stabilize the  
23 distal radioulnar joint. However, because of technical difficulties, no arthroscopic  
24 procedure for such a lesion has currently been established. We present a new  
25 technique for arthroscopic reattachment of the avulsed TFCC into the fovea. An  
26 osseous tunnel 2.9 mm in diameter is created from the ulnar neck to the foveal surface.  
27 Under arthroscopic guidance, a nonabsorbable suture passed into a 21 gauge needle is  
28 placed into the TFCC through the osseous tunnel. The avulsed portion of the TFCC is  
29 anchored to the fovea by means of a repair suture passed through the TFCC. To  
30 achieve normal tension of the TFCC, the suture is tied onto the periosteum around the  
31 proximal entrance of the osseous tunnel. Our arthroscopic technique is relatively  
32 simple and has significant advantages for progressive healing at the attachment site  
33 between the TFCC and the fovea.

34

35

36

37

## Introduction

38           The triangular fibrocartilage complex (TFCC) is a complex structure consisting  
39 of fibrocartilaginous and ligamentous tissues on the ulnar side of the wrist. This  
40 structure plays a crucial role in stabilizing the distal radioulnar joint (DRUJ) during  
41 forearm rotations. Traumatic or chronic disruption of the TFCC leads to pathologic  
42 conditions of the wrist due to DRUJ instability.

43           Recent biomechanical studies have clarified that the TFCC insertion into the  
44 fovea of the distal ulna has a greater effect on DRUJ stability than other insertion sites  
45 (1). Therefore, surgical reattachment against avulsion of the foveal TFCC insertion  
46 must be considered to provide stability for the DRUJ. Although open techniques of  
47 reattachment for avulsed TFCC to the fovea of the ulnar head have been introduced  
48 (2,3), no arthroscopic procedure has currently been developed due to technical  
49 difficulties.

50           Arthroscopic surgery has an advantage of accurately diagnosing and locating  
51 the lesions, which is often difficult to do in open procedures. Our aim was to suggest a  
52 novel technique of arthroscopic reattachment of the avulsed TFCC into the foveal  
53 region of the distal ulna.

54

## **Surgical Technique**

55  
56 All patients had a positive ulnar fovea sign, positive provocative manners  
57 described by Kleinman (4), and DRUJ instability. We defined the DRUJ instability as  
58 an asymmetry of the constraint of the radius translation relative to the distal ulna in  
59 forearm pronation and supination (5). Preoperative water-excited 3D double-echo  
60 steady state (DESS) magnetic resonance coronal T2-weighted images showed an area of  
61 high-signal-intensity at just distal to the fovea in all patients. Arthroscopic procedures  
62 are performed under general anesthesia with a tourniquet. A traction tower  
63 (CONMED, Largo, FL) maintains 3.2 to 4.5 kg of distraction throughout the procedure  
64 via finger traps placed on all the fingers except for the thumb. An arthroscope is  
65 introduced into the 3-4 portal using the standard technique. Outflow is established  
66 through the 6-U portal. An accessory portal includes the 6-R or 4-5 portal. The  
67 lunotriquetral interosseous ligaments and the TFCC are each inspected for wear or tear.  
68 The articular surface of the lunate and the triquetrum are also investigated. TFCC  
69 tension or resilience (trampoline effect) on ballottement with an arthroscopic probe is  
70 tested to make a diagnosis (3). The final diagnosis of avulsion of the foveal TFCC  
71 insertion is determined by a loss of the normal trampoline effect and a displacement of  
72 the TFCC in multiple directions by pulling on the TFCC using a probe (5).

73           Once the diagnosis has been confirmed, a 1.5-mm Kirschner wire is used as a  
74   guide pin and percutaneously inserted from the ulnar neck to the foveal region of the  
75   ulnar head under C-arm visualization (Fig. 1). Then, a 1.5 cm incision is made around  
76   the Kirschner wire and the ulna is exposed between the extensor carpai ulnaris and the  
77   flexor carpi ulnaris. A 2.9 mm cannulated drill (Depuy, Warsaw, IN) is driven into the  
78   just distal to the fovea over the inserted Kirschner wire. This procedure creates an  
79   osseous tunnel 2.9 mm in diameter from the ulnar neck to the foveal surface and  
80   debrides fibrous connective tissues at the foveal surface (Fig. 2A). Under arthroscopic  
81   guidance in the 3-4 portal, a 2-0 nonabsorbable suture (Prolene, ETHICON, Some Ville,  
82   NJ) passed into a 21 gauge needle is placed into the TFCC through the osseous tunnel.  
83   Then, a 2-0 nonabsorbable suture loop is advanced into the TFCC using the same  
84   manner. The suture end is captured by the loop and delivered out of the osseous tunnel  
85   by proximally withdrawing the loop (Fig. 2A, B). Then, the two free ends of the repair  
86   suture are pulled through the osseous tunnel to bring the suture onto the TFCC surface.  
87   The avulsed portion of the TFCC is anchored to the fovea by means of this manner.  
88   Near-normal tension of the TFCC is then reconstituted by tightening both ends of the  
89   suture. With the forearm in neutral rotation, the suture is tied onto the ulnar  
90   periosteum around the proximal entrance of the osseous tunnel (Fig. 2C).

91 Each patient is splinted with a long-arm cast in 45° of supination for 4 weeks  
92 postoperatively. Then, a removable wrist brace is applied for an additional 2 weeks.  
93 Vigorous rehabilitation of the wrist and forearm begins at 6 weeks postoperatively.

94

## 95 **Discussion**

96 The proximal portion of the TFCC consists of the ligamentous component,  
97 including the radioulnar ligament, which plays a crucial role in stabilizing the DRUJ  
98 during forearm rotations. Anatomical studies have emphasized that this ligament  
99 originates from the fovea as well as the base of the ulnar styloid process (6). In the  
100 wrist joint, the center of forearm rotation runs through the ulnar head close to the foveal  
101 region. Therefore, the foveal insertion has a greater effect on DRUJ stability during  
102 forearm rotations than the styloid insertion (1).

103 Sennwald et al (3) showed that an open surgical reattachment of the avulsed  
104 TFCC to its foveal insertion gave promising results after a mean follow-up of 3 years.  
105 Although arthroscopic suture techniques for ulnar-sided TFCC tear (Palmer class 1B)  
106 have been reported (7-9), no arthroscopic procedure for the reattachment of the TFCC to  
107 the fovea has been established yet. To biologically bond the avulsed TFCC to the  
108 ulnar insertion, fibrous connective tissue at the foveal surface is debrided and the



109 bleeding from bone marrow is enhanced by curettage to cancellous bone. Under the  
110 standard arthroscopic technique, these manners require direct visualization of the foveal  
111 surface by inserting an arthroscope beneath the avulsed TFCC. This makes it difficult  
112 to perform arthroscopic surgery for the avulsed TFCC from the fovea.

113 Our new technique has been developed to overcome the technical difficulties  
114 mentioned above. The creation of an osseous tunnel for an outside-in arthroscopic  
115 suture technique can enhance bleeding from cancellous bone. In previous  
116 experimental studies, anchoring of the meniscus into a bone tunnel resulted in  
117 progressive healing at the attachment site between the two tissues (10,11). These  
118 experimental data provide the rationale for our arthroscopic technique against avulsion  
119 of the foveal TFCC insertion.

120 From 4 to 8 weeks postoperatively, computed tomography images suggested  
121 the osseous tunnel had been filled with new bone formation (Fig. 3). Magnetic  
122 resonance images at 12 weeks postoperatively showed findings indicating attachment of  
123 the TFCC to the distal entry site of the tunnel. The technique we present allows an  
124 arthroscopic repair for avulsion of the TFCC at its foveal insertion. This technique is  
125 relatively simple for reattachment of avulsed TFCC to the fovea. We consider it a  
126 promising alternative to the procedures actually in use.

127

## References

- 128 1. Haugstvedt JR, Berger RA, Nakamura T, Neale P, Berglund L, An KN. Relative  
129 contributions of the ulnar attachments of the triangular fibrocartilage complex to the  
130 dynamic stability of the distal radioulnar joint. *J Hand Surg* 2006;31A:445-451.
- 131 2. Hermansdorfer JD, Kleinman WB. Management of chronic peripheral tears of the  
132 triangular fibrocartilage complex. *J Hand Surg* 1991;16A:340-346.
- 133 3. Sennwald GR, Lauterburg M, Zdravkovic V. A new technique of reattachment after  
134 traumatic avulsion of the TFCC at its ulnar insertion. *J Hand Surg*  
135 1995;20B:178-184.
- 136 4. Kleinman WB. Stability of the distal radioulna joint: biomechanics, pathophysiology,  
137 physical diagnosis, and restoration of function what we have learned in 25 years. *J*  
138 *Hand Surg* 2007;32A:1086-1106.
- 139 5. Tay SC, Tomita K, Berger RA. The “ulnar fovea sign” for defining ulnar wrist pain:  
140 an analysis of sensitivity and specificity. *J Hand Surg* 2007;32A:438-444.
- 141 6. Nakamura T, Makita A. The proximal ligamentous component of the triangular  
142 fibrocartilage complex. *J Hand Surg* 2000;25B:479-486.
- 143 7. Böhringer G, Schädel-Höpfner M, Petermann J, Gotzen L. A method for all-inside  
144 arthroscopic repair of Palmer 1B triangular fibrocartilage complex tears.

145 Arthroscopy 2002;18:211-213.

146 8. Pederzini LA, Tosi M, Prandini M, Botticella C. All-inside suture technique for  
147 Palmer class 1B triangular fibrocartilage repair. Arthroscopy 2007;23:1130.e1-4

148 9. Yao J, Dantuluri P, Osterman AL. A novel technique of all-inside arthroscopic  
149 triangular fibrocartilage complex repair. Arthroscopy 2007;23:1357.e1-4.

150 10. Gao J, Wei X, Messner K. Healing of the anterior attachment of the rabbit meniscus  
151 to bone. Clin Orthop Relat Res 1998;348:246-258.

152 11. Vail TP, Koukoubis TD, Schenkman D, Feagin JA. Graft-bone healing of a meniscus  
153 autograft anchored in bone tunnels. Knee Surg, Sports Traumatol, Arthroscopy  
154 1995;3:105-110.

155

156

157

158

159

160

161

162

### Figure Legends

163

164 **Figure 1.** Under C-arm visualization, a guide pin is inserted from the ulnar neck to the  
165 foveal region of the ulnar head to create an osseous tunnel.

166

167 **Figure 2. (A)** An osseous tunnel 2.9 mm in diameter from the ulnar neck to the foveal  
168 surface. Using a suture loop, the end of repair suture is delivered out of the osseous  
169 tunnel. **(B)** The two free ends of the repair suture are pulled through the osseous  
170 tunnel to bring the suture onto the TFCC surface. **(C)** The avulsed portion of the  
171 TFCC is anchored to the fovea with near normal tension. The suture is tied onto the  
172 ulnar periosteum around the proximal entrance of the osseous tunnel.

173

174 **Figure 3.** Computed tomography images suggest that the osseous tunnel is filled with  
175 new bone formation at 8 weeks postoperatively.

176

177

**Reference Citation from the Pub/Med/Medline Web Site**

1. [Haugstvedt JR, Berger RA, Nakamura T, Neale P, Berglund L, An KN.](#) Relative contributions of the ulnar attachments of the triangular fibrocartilage complex to the dynamic stability of the distal radioulnar joint. J Hand Surg [Am]. 2006 Mar;31(3):445-51. PMID: 16516740 [PubMed - indexed for MEDLINE]
2. [Hermansdorfer JD, Kleinman WB.](#) Management of chronic peripheral tears of the triangular fibrocartilage complex. J Hand Surg [Am]. 1991 Mar;16(2):340-6. PMID: 2022850 [PubMed - indexed for MEDLINE]
3. [Sennwald GR, Lauterburg M, Zdravkovic V.](#) A new technique of reattachment after traumatic avulsion of the TFCC at its ulnar insertion. J Hand Surg [Br]. 1995 Apr;20(2):178-84. PMID: 7797966 [PubMed - indexed for MEDLINE]
4. [Kleinman WB.](#) Stability of the distal radioulna joint: biomechanics, pathophysiology, physical diagnosis, and restoration of function what we have learned in 25 years. J Hand Surg [Am]. 2007 Sep;32(7):1086-106. Review. No abstract available. PMID: 17826566 [PubMed - indexed for MEDLINE]
5. [Tay SC, Tomita K, Berger RA.](#) The "ulnar fovea sign" for defining ulnar wrist pain: an analysis of sensitivity and specificity. J Hand Surg [Am]. 2007 Apr;32(4):438-44. PMID: 17398352 [PubMed - indexed for MEDLINE]

6. [Nakamura T, Makita A](#). The proximal ligamentous component of the triangular fibrocartilage complex. *J Hand Surg [Br]*. 2000 Oct;25(5):479-86. PMID: 10991819 [PubMed - indexed for MEDLINE]
7. [Böhringer G, Schädel-Höpfner M, Petermann J, Gotzen L](#). A method for all-inside arthroscopic repair of Palmer 1B triangular fibrocartilage complex tears. *Arthroscopy*. 2002 Feb;18(2):211-3. PMID: 11830818 [PubMed - indexed for MEDLINE]
8. [Pederzini LA, Tosi M, Prandini M, Botticella C](#). All-inside suture technique for Palmer class 1B triangular fibrocartilage repair. *Arthroscopy*. 2007 Oct;23(10):1130.e1-4. Epub 2007 Mar 13. PMID: 17916483 [PubMed - indexed for MEDLINE]
9. [Yao J, Dantuluri P, Osterman AL](#). A novel technique of all-inside arthroscopic triangular fibrocartilage complex repair. *Arthroscopy*. 2007 Dec;23(12):1357.e1-4. Epub 2007 May 7. PMID: 18063183 [PubMed - indexed for MEDLINE]
10. [Gao J, Wei X, Messner K](#). Healing of the anterior attachment of the rabbit meniscus to bone. *Clin Orthop Relat Res*. 1998 Mar;(348):246-58. PMID: 9553559 [PubMed - indexed for MEDLINE]

11. [Vail TP, Koukoubis TD, Schenkman D, Feagin JA](#). Graft-bone healing of a meniscus autograft anchored in bone tunnels. *Knee Surg Sports Traumatol Arthrosc.* 1995;3(2):105-10. PMID: 7553004 [PubMed - indexed for MEDLINE]

**Figure 1**  
[Click here to download high resolution image](#)





Figure 2A  
[Click here to download high resolution image](#)




Figure 2B  
[Click here to download high resolution image](#)




Figure 2C  
[Click here to download high resolution image](#)




Figure 3  
[Click here to download high resolution image](#)

