



Title	Purification and biochemical characterization of a cellulase from the digestive organs of the short-spined sea urchin <i>Strongylocentrotus intermedius</i>
Author(s)	Hasegawa, Syuto; Ura, Kazuhiro; Tanaka, Hiroyuki; Ojima, Takao; Takagi, Yasuaki
Citation	Fisheries Science, 78(5), 1107-1115 https://doi.org/10.1007/s12562-012-0528-y
Issue Date	2012-09
Doc URL	http://hdl.handle.net/2115/50293
Rights	© 2012 公益社団法人日本水産学会; © 2012 The Japanese Society of Fisheries Science
Type	article
File Information	FS78-5_1107-1115.pdf



[Instructions for use](#)

Purification and biochemical characterization of a cellulase from the digestive organs of the short-spined sea urchin *Strongylocentrotus intermedius*

Syuto Hasegawa · Kazuhiro Ura · Hiroyuki Tanaka · Takao Ojima · Yasuaki Takagi

Received: 23 September 2011 / Accepted: 5 June 2012 / Published online: 3 July 2012
© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract We isolated a cellulase from the digestive organs of the short-spined sea urchin *Strongylocentrotus intermedius* using a combination of ion-exchange chromatography and gel filtration together with an assay for carboxymethylcellulase activity. The isolated cellulase was stained as a single band by Congo red. The molecular weight of the isolated cellulase, as determined by sodium dodecyl sulfate-polyacrylamide gel electrophoresis under reducing conditions, was 59 kDa. The isolated cellulase exhibited hydrolytic activity toward carboxymethyl cellulose, with an optimum temperature and pH of 30 °C and pH 8.0, respectively. The thermal stability of the enzyme was characterized by determining the temperature at which activity decreased by 50 % with treatment for 30 min at pH 7.0 and found to be 32 °C. Cellulase activity remained at a high level at 5–20 °C, which is the growth temperature of the short-spined sea urchin. These results confirm that the short-spined sea urchin should preferably be reared at a water temperature of <20 °C.

Keywords Cellulase · Digestive organ · Enzymatic properties · Protein isolation · Sea urchin

Introduction

Cellulose, a major cell-wall component in plants, is a linear polysaccharide consisting of β -1,4-linked D-glucose residues that hydrolyzes the β -1,4-glycosyl linkages of the cellulose chain to produce cello-oligosaccharides, such as cellotriose and cellobiose [1]. This enzyme family is grouped into endolytic (endo-1,4- β -glucanase, EC 3.2.1.4) and exolytic (cellobiohydrolase, EC 3.2.1.91) enzymes. Complete depolymerization of cellulose to glucose requires another sort of cello-oligosaccharide-degrading enzyme, 1,4- β -glucosidase (EC 3.2.1.21). Thus far, cellulolytic enzymes have been isolated and characterized from bacteria [2], fungi [2], plants [3], molds [4], microbes [5], arthropods [6–11], nematodes [12], mollusks [13–19] and sea urchin [20].

The sea urchin is a well-known marine organism distributed widely from the rocky intertidal areas to the abyssal depths in all of the world's oceans. The gonads of sea urchins are a high value seafood product, and considerable research is being undertaken worldwide on the development of sea urchin aquaculture. More recently, considerable attention has been directed toward the development of artificial diets that enhance the quantity and quality of roe produced from sea urchin aquaculture [21, 22]. To develop such artificial diets and aquaculture systems, it is important to understand the digestive physiology of sea urchins. Although numerous edible sea urchins are kelp feeders, a surprisingly limited amount of information exists on the biochemical characterization of cellulases in the digestive system of sea urchins, although several reports have been published on the detection of cellulase activity in the digestive organs of sea urchins [13, 23–29]. In the respective studies, cellulase activity was analyzed using crude extracts from the digestive organs and, therefore, the exact

S. Hasegawa · Y. Takagi
Laboratory of Comparative Physiology,
Graduate School of Fisheries Science, Hokkaido University,
3-1-1 Minato-cho, Hakodate 041-8611, Japan

K. Ura (✉)
Graduate School of Fisheries Science, Hokkaido University,
3-1-1 Minato-cho, Hakodate 041-8611, Japan
e-mail: kazu@fish.hokudai.ac.jp

H. Tanaka · T. Ojima
Laboratory of Marine Molecular Biology,
Graduate School of Fisheries Science, Hokkaido University,
3-1-1 Minato-cho, Hakodate 041-8611, Japan

biochemical characterization of the cellulase has remained unclear. Nishida et al. [20] reported the isolation, biochemical characterization and cDNA cloning of cellulase from the digestive organs of the Northern sea urchin *Strongylocentrotus nudus*. However, these authors focused on the biochemical characterization of the cellulase at a temperature that is not considered to be the growth temperature of the Northern sea urchin. Thus, the isolation and biochemical characterization of digestive enzymes at the growth temperature of sea urchin is an important step toward understanding the digestive physiology of this organism and the development of efficient aquaculture systems. To the best of our knowledge, there has been no study on the biochemical characterization of cellulases from sea urchin at its growth temperature.

In the study reported here, we isolated and biochemically characterized a cellulase from the digestive organs of short-spined sea urchin, *Strongylocentrotus intermedius*. We also compared the biochemical characterization of the cellulases from the short-spined sea urchin and Northern sea urchin and discussed digestive physiology in terms of the biochemical characterizations of the cellulases at physiological growth temperatures.

Materials and methods

Materials

Digestive organs from the short-spined sea urchin were obtained at the local fish market in Hakodate, Hokkaido Prefecture, Japan. The pooled tissues (1.5 kg) were thoroughly rinsed in sea water to remove the intestinal contents and then dehydrated by the addition of 1.5 l acetone at -20°C . Precipitates were collected by centrifugation at 10,000 g for 15 min, rinsed five times in 0.5 l cold acetone, then rinsed again in 0.5 l acetone at room temperature and finally collected by centrifugation at 10,000 g for 15 min. The acetone-precipitated tissues were dried at room temperature, and the acetone-dried powder was stored at -20°C until use.

Purification of cellulase from the short-spined sea urchin

Crude enzyme was extracted from 2.5 g acetone-dried powder from the sea urchin digestive tracts with 100 ml sodium phosphate (10 mM; pH 7.0) at 0°C for 30 min with occasional stirring. The enzyme extract was centrifuged at 10,000 g for 15 min and the supernatant then dialyzed against 10 mM sodium phosphate (pH 7.0) at 4°C for 24 h. The supernatant was then applied to a TOYOPEARL-DEAE 650M column (2.0 × 26 cm) (Toyo Soda Mfg Co.,

Tokyo, Japan). After unadsorbed materials were washed out with 10 mM sodium phosphate (pH 7.0), proteins were eluted by a linear NaCl gradient from 0 to 300 mM (total volume 700 ml) in 10 mM sodium phosphate (pH 7.0) at a 90 ml/h flow rate. The eluate was collected in 8-ml fractions and carboxymethyl cellulose (CMCase) activity was determined. CMCase activity was detected in the fractions eluted at approximately 0.2 M NaCl (pooled as the DE-1 fraction). The DE-1 fraction was subjected to the same chromatography procedure after dialysis in 10 mM sodium phosphate (pH 7.0). The active fractions (pooled as the DE-2 fraction) were concentrated to 1.2 ml with a Vivaspin 20 ml centrifugal concentrator (Sartorius, Goettingen, Germany). The DE-2 fraction was subjected to ion exchange chromatography using a Mono Q 5/50 GL column (GE Healthcare UK, Buckinghamshire, UK) after dialysis in 10 mM sodium phosphate (pH 7.0). The retained proteins were eluted with a linear NaCl gradient from 0 to 300 mM at a 60 ml/h flow rate using a fast protein liquid chromatography system (GE Healthcare Biosciences). The active fractions were concentrated to 1.0 ml with a Vivaspin 5 ml centrifugal concentrator (Sartorius) and then subjected to gel filtration chromatography using a Superdex 75 column (GE Healthcare Biosciences) at a 60 ml/h flow rate. A 59-kDa protein with high CMCase activity was eluted as a single major peak.

Cellulase activity assay

A standard assay for cellulase activity was performed with a reaction mixture containing 0.52 % carboxymethyl cellulose (CMC; medium viscosity, Bio Medics, Orange, CA) in 10-mM sodium phosphate (pH 7.0) at 30°C . Reduced sugar produced by the reaction was determined using the method described by Park and Johnson [30]. One unit of cellulase is defined as the amount of enzyme that produces reduced sugar equivalent to 1 μmol glucose during 1 min. The temperature dependence of the cellulase was determined over a range of 5 to 60°C and at pH 7.0; pH dependence was determined at 30°C in the following reaction buffers: 10 mM HCl-glycine (pH 3–4), 10 mM $\text{CH}_3\text{COOH}-\text{CH}_3\text{COONa}$ (pH 4–5), 10 mM $\text{NaH}_2\text{PO}_4-\text{Na}_2\text{HPO}_4$ (pH 5–7), 10 mM Tris-HCl (pH 7–9) and 10 mM NaOH-glycine (pH 9–10). The thermal stability was characterized by incubating the isolated enzyme at different temperatures ($5-60^{\circ}\text{C}$) and pH 7.0 for 30 min followed by assaying the activity as described below.

Effects of metal ions on the cellulase activity

The effects of several metal ions and EDTA on the activity of the purified cellulase (1.5 units/ml) activity was determined in a reaction mixture containing 1 mM (final

concentration) of AgNO_3 , CaCl_2 , CdCl_2 , CoCl_2 , CuSO_4 , FeCl_2 , MgCl_2 , MnCl_2 , ZnCl_2 and EDTA in 10 mM sodium phosphate (pH 7.0) at 30 °C for 60 min, followed by an assay of the activity as described below.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was performed with 3 % stacking and 10 % separating gels as described by Laemmli et al. [31]. Samples were treated with equal volumes of sample buffer containing 4 % SDS, 14 % glycerol and 10 % 2-mercaptoethanol at 100 °C for 2 min or equal volumes of sample buffer containing 4 % SDS and 20 % glycerol. Gels were stained with 0.1 % Coomassie brilliant blue R-250 in ethanol, acetic acid and deionized water (DW) (40:10:50), then destained in ethanol, acetic acid, glycerol and DW (200:50:25:725). Molecular weights were estimated using Broad Range Standard Markers (New England Biolabs, Ipswich, MA).

Detection of cellulase activity in polyacrylamide gels

Cellulase activity in the polyacrylamide gels was detected using a modification of the method reported by Beguin [32]. Briefly, enzyme samples were treated with equal volumes of sample buffer containing 0.25 % SDS and 10 % glycerol. After SDS-PAGE, SDS was removed by washing the gel twice for 30 min in 10 mM sodium phosphate (pH 7.0) and 25 % isopropanol, followed by another two washes for 10 min in 10 mM sodium phosphate (pH 7.5). Agar gels (2 %; thickness 5 mm) containing 0.1 % CMC in 10 mM Tris-HCl (pH 7.5) were prepared. Excess liquid on the polyacrylamide gels was removed by wiping the gel surface, and the gels were then laid on top of the agar gels and incubated at 37 °C for 3 h. After incubation, the agar gels were stained with 0.1 % Congo red solution and then destained with 1 M NaCl. All procedures were performed at 4 °C except for the incubation.

Thin-layer chromatography

Thin-layer chromatography (TLC) for analysis of degradation products of cellulose and cello-oligosaccharides was performed on a TLC-60 plate (Merck, Darmstadt, Germany) and a developing solvent consisting of *n*-butanol/acetic acid/water (2:1:1, v/v/v). The sugar developed on the plate was detected by heating at 120 °C for 15 min after being sprayed with 10 % (v/v) sulfuric acid ethanol. Cellulose powders were purchased from Advantec (San Diego, CA) and Merck. The 0.1-ml reaction mixture (0.5 mg of phosphoric acid-swollen cellulose and crystalline cellulose, 5 mM sodium

phosphate, pH 7.0, 0.26 U of purified cellulase) was incubated at 30 °C for up to 24 h. The supernatant of the reaction mixture was subjected to TLC. The reaction mixture of the 0.05 ml containing 0.05 mg of G2 (cellobiose)–G6 (cellohexaose), 5 mM sodium phosphate (pH 7.0), and 0.13 U of purified cellulase was incubated at 30 °C for 20 h, then subjected to TLC.

Protein concentration

Protein concentrations were determined with bovine serum albumin fraction V (Wako Pure Chemicals, Osaka, Japan) as a standard protein using either the biuret method [33] or the method described elsewhere by Lowry et al. [34].

Amino acid sequencing

The N-terminal amino acid sequence of the intact protein was determined by electrophoretically blotting a sample onto a polyvinylidene difluoride membrane (Advantec) and performing the sequencing in an ABI 492HT protein sequencer (Applied Biosystems, Foster city, CA, USA) following the manufacturer's instructions.

Results

Cellulase isolation

A cellulase was isolated from the acetone-dried powder of the digestive organs of the short-spined sea urchin. Isolation was performed using a combination of ion-exchange chromatography and gel filtration together with an assay for CMCase activity (Fig. 1). The isolation procedures resulted in a target protein that was purified to a 1,156-fold concentration with a 5.4 % final yield and 28.9 U/mg specific activity (Table 1). The purified short-spined sea urchin cellulase showed a single band with an apparent molecular weight of 59 kDa, as determined by SDS-PAGE under reducing conditions (Fig. 2a), and was stained by Congo red (Fig. 2b).

Enzymatic properties of short-spined sea urchin cellulase

The basic properties of short-spined sea urchin cellulase were investigated and compared with those of cellulase from the Northern sea urchin [20]. As shown in Fig. 3, the optimal temperature and pH of short-spined cellulase was 30 °C and pH 8.0, respectively. These properties differ from those of the Northern sea urchin cellulase, which are 35 °C and pH 6.3, respectively (Fig. 3a, b). The thermal stability was characterized as the temperature at which




Fig. 1 Purification of short-spined sea urchin (*Strongylocentrotus intermedius*) cellulase. **a** First TOYOPEARL-DEAE 650M chromatography. **b** Second TOYOPEARL-DEAE 650M chromatography. **c** MonoQ column chromatography. **d** Superdex 75 gel filtration. The chromatography conditions are described in the [Materials and methods](#). The fractions indicated by *solid bars* were pooled and

designated as the DE-1 fraction (**a**; fractions eluted at approximately 0.2 M NaCl), DE-2 fraction (**b**; active fractions obtained after DE-1 fraction was subjected to the same chromatography procedure after dialysis in 10 mM sodium phosphate), MonoQ fraction (**c**) and Superdex 75 fraction (**d**)

Table 1 Purification of cellulase from the short-spined sea urchin (*Strongylocentrotus intermedius*)

Purification step ^a	Total protein (mg)	Specific activity (U/mg)	Total activity (U)	Purification yield	
				Fold	%
Crude extract	1,677.8	0.025	42.12	1	100
First DEAE fraction	24.9	0.743	18.53	29.7	43.9
Second DEAE fraction	5.08	2.27	11.53	90.8	27.3
Mono Q	0.20	16.9	3.39	676	8.0
Superdex 75	0.08	28.9	2.31	1,156	5.4

^a The chromatography conditions are described in the [Materials and methods](#)

activity decreased to 50 % with treatment for 30 min at pH 7.0; this temperature was found to be 32 and 37 °C for short-spined sea urchin and Northern sea urchin cellulase, respectively (Fig. 3c). Table 2 shows the effect of metal ions and EDTA on the activity of the short-spined sea urchin cellulase. The activity of cellulase was strongly inhibited by the addition of Ag⁺ and Cu²⁺. The TLC analysis showed that short-spined sea urchin cellulase has a higher degradation activity toward phosphoric acid-swollen cellulose than toward crystalline cellulose (Fig. 4a, b). The short-spined sea urchin cellulase degraded the phosphoric

acid-swollen cellulose to produce cellotetraose, cellotriose and cellobiose as major products (Fig. 4a). It also degraded the crystalline cellulose to produce cellotetraose, cellotriose and cellobiose; however, the degradation activity on this substrate was very low (Fig. 4b). Figure 4c shows the degradation of the cello-oligosaccharides. The short-spined sea urchin cellulase readily degraded both cellohexasae and cellopentaose to cellotetraose, cellotriose, cellobiose and glucose, and slowly degraded cellotetraose to cellotriose, cellobiose and glucose. However, it scarcely degraded cellotriose and cellobiose.




Fig. 2 **a** Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) of short-spined sea urchin cellulase from various purification steps. **b** Detection of cellulase activity in polyacrylamide gels. Gels were stained by Congo red. *Crude* Crude extract, *DE-1* DE-1 fraction, *DE-2* DE-2 fraction, *MonoQ* MonoQ fraction, *Superdex 75* Superdex 75 fraction, *MW* molecular-weight marker proteins

N-terminal amino-acid sequence

Using a protein sequencer, we determined the 15 residue N-terminal amino acid sequence of short-spined sea urchin cellulase to be SDYXDVLHKSILFFE. This sequence showed 86 % identity with the Northern sea urchin cellulase sequence at amino acid positions 18–32 (Table 3).

Discussion

To date, cellulases have been isolated and characterized from various organisms. However, biochemical characterization of the cellulases in marine organisms has been limited. In sea squirts and sea urchins, cellulase primary structures have been analyzed by means of genomic and expressed sequence tag analyses [35, 36]. In more recent studies, cellulases have been isolated from abalone and sea urchin for biochemical characterization and primary structure determination [18, 20]. In our study, short-spined sea urchin cellulase was purified using a combination of ion-exchange chromatography and gel filtration together with CMCase activity assays. The molecular weights of cellulases have been reported from various organisms, such as plants (50–70 kDa [3]), molds (42 kDa [4]), arthropods (40–47 kDa [6, 10, 11]), nematodes (49 kDa [12]), rotifer (62 kDa [37]), mollusks (44–70 kDa [15–18]) and sea

urchin (54 kDa [20]), and the range has been found to be quite wide. The molecular weight of purified short-spined sea urchin cellulase was estimated to be 59 kDa, a value which falls well within the range of previously reported cellulases from marine invertebrates, including abalone [18] and Northern sea urchin [20]. Furthermore, the N-terminal amino-acid sequence of purified short-spined sea urchin cellulase showed 86 % identity with that of Northern sea urchin [20], providing good proof that the purified protein is actually the cellulase of short-spined sea urchin.

In this study, we investigated the effects of metal ions and EDTA on the activity of short-spined sea urchin cellulase. To the best of our knowledge, this is the first such report in marine organisms. The activity of the enzyme was strongly inhibited with the addition of Ag^+ and Cu^{2+} . To date, there are very few published studies on the effect of metal ions on the activity of cellulase. In *Bacillus sphaericus*, the activity of cellulase was inhibited by adding Fe^{3+} and Hg^{2+} and stimulated by adding Co^{2+} and K^+ [38]. Working with various *Bacillus* strains, Mawadza et al. [39] isolated and characterized two types of cellulases (HR66 and CH43). The activity of HR66 and CH43 was stimulated by adding Co^{2+} and inhibited by adding Hg^{2+} . The activity of CH43 was also inhibited by adding Ag^+ , but that of HR66 was stimulated by adding Ag^+ . Other metal ions (K^+ , Mg^{2+} , Cu^{2+} , Ca^{2+} , Ni^{2+} , Zn^{2+} , Fe^{3+}) had no effects on the activity of either HR66 or CH43. Based on their results, the authors suggested that the effects of metal ions on the activity of cellulase was different depending on the type of cellulase and organisms. We found that the short-spined sea urchin cellulase showed a higher degradation activity toward phosphoric acid-swollen cellulose than to crystalline cellulose. Moreover, the degradation pattern was similar to that of the Northern sea urchin [20]. The short-spined sea urchin cellulase scarcely degraded cellotri-ose and cellobiose. β -Glucosidase activity was reported to be present in crude extracts from digestive organs of the sea urchin [40, 41], leading to the suggestion that in the sea urchin cellulose is first degraded to cellotri-ose and cellobiose by the cellulase, followed by the possible degradation of these cello-oligosaccharides to glucose by β -glucosidase. However, no data have been published on the isolation and biochemical characterization of β -glucosidase in sea urchin. In addition, in another study a cellulase was purified which raised a specific antibody against the enzyme from abalone; this antibody reacted with three protein bands by western blotting analysis, suggesting that there are several types of cellulase in marine organisms (Ojima, unpublished data). Further studies are being contemplated to determine the degradation mechanism of cellulose in sea urchin.

In echinoderms, less attention has been paid to the relationship between digestive enzymes and digestive physiology. Almost every study performed to date has




Fig. 3 Temperature, pH dependence and thermal stability activities of short-spined sea urchin and Northern sea urchin cellulases. **a** Activities of the cellulases were measured at various temperatures in a reaction mixture containing 0.52 % carboxymethyl cellulose (CMC) and 10 mM sodium phosphate (pH 7.0). **b** Activities were measured in the presence of 0.52 % CMC at 30 °C in various reaction buffers at varying pH, including 10 mM HCl-glycine (pH 3–4) (filled triangles), 10 mM $\text{CH}_3\text{COOH}-\text{CH}_3\text{COONa}$ (pH 4–5) (solid lines),

10 mM $\text{NaH}_2\text{PO}_4-\text{Na}_2\text{HPO}_4$ (pH 5–7) (filled diamonds), 10 mM Tris-HCl (pH 7–9) (filled squares), 10 mM NaOH-glycine (pH 9–10) (x symbols). **c** Thermal stability of the purified enzyme was measured by pre-incubation at various temperatures at a constant pH, and the activity was assayed in a reaction mixture containing 0.52 % CMC, 10 mM sodium phosphate (pH 7.0). Filled circles Short-spined sea urchin cellulase; open circles Northern sea urchin cellulase (Nishida et al. [20])

Table 2 N-terminal amino acid sequence of the cellulases

Organisms	N-terminal amino acid sequence of the cellulase
<i>Strongylocentrotus intermedius</i> (short-spined sea urchin)	SDYXDVLHKLSILFFE
<i>Strongylocentrotus nudus</i> (Northern sea urchin)	YDYXDVIHKLSILFFE

Table 3 Effects of metal ions and EDTA on the activity of cellulase of *Strongylocentrotus intermedius*


Compound	Relative activity (%)
None	100
AgNO_3	5.5
CaCl_2	97.0
CdCl_2	100.5
CoCl_2	95.2
CuSO_4	43.5
FeCl_2	83.6
MgCl	96.1
MnCl	106.7
ZnCl_2	101.7
EDTA	93.9

EDTA, Ethylenediaminetetraacetic acid

focused on the biochemical characterization of digestive enzymes at temperatures not considered to be the growth temperature of sea urchin. Consequently, enzyme activity has been assayed at temperatures ranging from 25 to 35 °C in various studies [28, 42–45], but few studies have focused on assaying digestive enzyme activity at the growth temperature of echinoderms. Klinger et al. [46] showed that there was no difference in digestive enzyme activity between polar and subtropical sea urchins at 15 and 25 °C after sea urchins had been reared at 16 or 23 °C. Thus far, there are several reports on the detection of cellulase activity in the digestive organs of sea urchin [13, 23–29], and the activity of digestive enzymes from sea urchins has been analyzed using crude extracts from the digestive organs. However, the physiological function of digestive enzymes in the digestive system of sea urchins remains unclear. The isolation and biochemical characterization of the digestive enzymes is an important step towards understanding the digestive physiology of the sea urchin. In echinoderms, the isolation and biochemical characterization of a cellulase from the digestive organs of Northern sea urchin was first reported by Nishida et al. [20]. However, this report also focused on the biochemical characterization of a cellulase.

Fig. 4 Degradation of phosphoric acid-swollen cellulose, crystalline cellulose and cello-oligosaccharides by the short-spined sea urchin cellulase. The reaction mixture of 0.1 ml containing 0.5 mg of phosphoric acid-swollen cellulose (a) or crystalline cellulose (b), 5 mM sodium phosphate (pH 7.0) and 0.26 U of purified cellulase was incubated at 30 °C for up to 24 h. The supernatant of the reaction mixture was subjected to thin-layer chromatography (TLC). The reaction mixture of 0.05 ml containing 0.05 mg of G2–G6, 5 mM sodium phosphate (pH 7.0) and 0.13 U of purified cellulase was incubated at 30 °C for 20 h, then subjected to TLC (c). G1 Glucose, G2 cellobiose, G3 cellotriose, G4 cellotetraose, G5 cellopentaose, G6 cellohexaose, M marker sugars

In our study, we isolated a cellulase from the digestive tract of the short-spined sea urchin and found that the optimum temperature (approximately 30 °C) for short-spined sea urchin cellulase activity was lower than that of Northern sea urchin cellulase (approximately 35 °C) [20]. Moreover, the enzyme activity remained at 20–40 % between 5 and 20 °C, which is the growth temperature for short-spined sea urchin. In our previous study, we isolated and biochemically characterized the proteolytic enzyme subtilase from the digestive tracts of the short-spined sea urchin, which is a major proteolytic enzyme in sea urchin, and determined that the optimum temperature is approximately 50 °C and that enzyme activity remains at 40–50 % between 10 and 20 °C [47]. The thermal stability of the short-spined sea urchin cellulase was also characterized, and although the enzyme activity remained at a high level between 5 and 20 °C, it decreased abruptly above 20 °C and was 50 % at 32 °C. In contrast, in the Northern sea urchin, the enzyme activity remained at a high level between 10 and 30 °C and was 50 % at 37 °C. Gonadal growth and growth rate are significantly suppressed in the short-spined sea urchin when rearing occurs at a water temperature of >21 °C versus <20 °C [48]. Also, the value of assimilation efficiency in the short-spined sea urchin was found to be 55 % at approximately 5 °C after kelp *Laminaria japonica* was supplied as food [49, 50]. In addition, gonadal weight in the short-spined sea urchin increased at approximately 5 °C after the kelp *L. japonica* had been supplied as food (T. Kayaba, personal communication), but it did not change in the Northern sea urchin under the same feeding conditions (K. Ura, unpublished data). In *S. droebachiens* and *Tripneustes esculentus*, both residents of low-temperature water zones, the value of assimilation efficiency is 50–60 % in winter [51, 52]. Other sea urchins (*S. droebachiens*) which live in water zones characterized by the same low temperature as the habitat of the short-spined sea urchin are known to undergo a decrease in survival rate at water temperatures of >19 °C, with the lethal temperature being approximately 24 °C [53]. In comparison, in the Northern sea urchin, which can live at higher water temperatures than the short-spined sea urchin, the value of assimilation efficiency is approximately 50 % at 10–25 °C;



however, assimilation efficiency is approximately 10 % at 5 °C after kelp is supplied as food [54]. From these results, we conclude that the activity of the short-spined sea urchin cellulose is weaker at high water temperatures than the Northern sea urchin cellulose.

The pH of the food canal of the sea urchin in an unfed state food is around pH 7.0 [55]. In the short-spined sea urchin, the optimal pH of purified cellulase was found to be pH 8.0; in contrast, that of the Northern sea urchin is pH 7.0, although enzyme activity was high at around pH 6.0–9.0. In our previous study, we isolated and biochemically characterized the proteolytic enzyme subtilase from the digestive tracts of the short-spined sea urchin and found that the optimum pH for subtilase activity is approximately pH 9.0. These data suggest that the digestive enzymes of sea urchin function under weak alkaline conditions. However, to date there have been no published studies on the changes in the pH of the food canal of sea urchin after feeding. Further work is needed to determine the physiological conditions of the food canal and the role of each of the enzymes in the food canal of sea urchin.

In Northern sea urchin, although cellulase activity has been detected in the esophagus, stomach, and intestine (T. Ojima, unpublished data), the localization of the cellulase in the digestive organs is still unclear. In our previous study, we isolated subtilase and raised a specific antibody against the enzyme [47]. Moreover, we found that subtilase is localized in granules in the anterior and posterior stomach and the anterior intestine [56]. From these data, it is likely that the localization of subtilase and cellulase may differ. Further work is needed to determine the localization of cellulase in the digestive organs of sea urchin.

In conclusion, we isolated and biochemically characterized a cellulase from the digestive organs of the short-spined sea urchin. We demonstrated that cellulase from the short-spined sea urchin possesses a physiological function at low temperatures, which is the growth temperature of this organism. Further work is in progress to confirm the localization of cellulase in the digestive organs in order to gain an understanding of the digestive system of sea urchin.

Acknowledgments This study was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture, Japan.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Bhat MK, Bhat S (1997) Cellulose degrading enzymes and their potential industrial applications. *Biotech Adv* 15:583–620
- Tomme P, Warren RAJ, Gilkes NR (1995) Cellulose hydrolysis by bacteria and fungi. *Adv Microbiol Physiol* 37:1–81
- Brummel DA, Lashbrook CC, Bennett AB (1994) Plant endo-1,4- β -D-glucanase. *ACS Symp Ser* 566:100–129
- Blume JE, Ennis HL (1991) A *Dictyostelium discoideum* cellulase is a member of a spore germination-specific gene family. *J Biol Chem* 266:15432–15437
- Moriya S, Ohkuma M, Kudo T (1998) Phylogenetic position of symbiotic protist *Dinemympha exilis* in the hindgut of the termite *Reticulitermes speratus* inferred from the protein phylogeny of elongation factor 1 α . *Gene* 210:221–227
- Watanabe H, Nakamura M, Tokuda G, Yamaoka I, Scrivener AM, Noda H (1997) Site of secretion and properties of endogenous endo- β -1,4-glucanase components from *Reticulitermes speratus* (Kolbe), a Japanese subterranean termite. *Insect Biochem Mol Biol* 27:305–313
- Watanabe H, Noda H, Tokuda G, Lo N (1998) A cellulase gene of termite origin. *Nature* 394:330–331
- Tokuda G, Watanabe H, Matsumoto T, Noda H (1997) Cellulose digestion in the wood-eating higher termite, *Nasutitermes takasagonesis* (Shiraki): distribution of cellulases and properties of endo- β -1,4-glucanase. *Zool Sci* 14:83–93
- Tokuda G, Lo N, Watanabe M, Slaytor M, Matsumoto T, Noda H (1999) Metazoan cellulase genes from termites: intron/exon structure and sites of expression. *Biochem Biophys Acta* 1447:146–159
- Xue XM, Anderson AJ, Richardson NA, Anderson AJ, Xue GP, Mather PB (1999) Characterization of cellulase activity in the digestive system of the redclaw cray-fish (*Cherax quadricarinatus*). *Aquaculture* 180:373–386
- Sugimura M, Watanabe H, Lo N, Saito H (2003) Purification characterization, cDNA cloning and nucleotide sequence of a cellulase from the yellow-spotted longicorn beetle, *Psacotheta hilaris*. *Eur J Biochem* 270:3455–3460
- Smant G, Stokkermans JPWG, Yan Y, De Boer JM, Baum TJ, Wang X, Hussey RS, Gommers FJ, Henriessat B, Davis EL, Helder J, Schots A, Bakker J (1998) Endogenous cellulases in animals: isolation of β -1,4-endoglucanases genes from two species of plant-parasitic cyst nematodes. *Proc Natl Acad Sci USA* 95:4906–4911
- Yokoe Y, Yasumasu I (1964) The distribution of cellulase in invertebrates. *Comp Biochem Physiol* 13:323–338
- Marshall JJ (1973) Purification of β -1,4-glucan hydrolase (cellulase) from the snail, *Helix pomatia*. *Comp Biochem Physiol* 44B:981–988
- Marshall JJ, Grand RJA (1976) Characterization of a β -1,4-glucan hydrolase from the snail, *Helix pomatia*. *Comp Biochem Physiol* 53B:231–237
- Anzai H, Nisizawa K, Matsuda K (1984) Purification and characterization of a cellulase from *Dolabella auricularia*. *J Biochem* 96:1381–1390
- Xu B, Hellman U, Ersson B, Jason J-C (2000) Purification, characterization and amino-acid sequence analysis of a thermostable, low molecular mass endo- β -1,4-glucanase from blue mussel, *Mytilus edulis*. *Eur J Biochem* 267:4970–4977
- Suzuki K, Ojima T, Nishita K (2003) Purification and cDNA cloning of a cellulase from abalone *Haliotis discus hannai*. *Eur J Biochem* 270:771–778
- Wang J, Ding M, Li Y, Chen Q, Xu G, Zhao F (2003) Isolation of a multifunctional enzogenous cellulase gene from mollusk, *Ampullaria crossean*. *Acta Biochem Biophys Sin* 35:941–946
- Nishida Y, Suzuki K, Kumagai Y, Tanaka H, Inoue A, Ojima T (2007) Isolation and primary structure of a cellulase from the Japanese sea urchin *Strongylocentrotus nudus*. *Biochimie* 89:1002–1011
- Kelly MS (2004) Echinoderms. In: Heinzeller T, Nebeisick JH (eds) *Munchen*. Taylor and Francis, London, pp 283–289

22. Lawrence JM (2007) Biology and ecology. In: Lawrence JM (ed) Edible sea urchins. Elsevier Science, Amsterdam, pp 1–9
23. Hultin E, Wanntorp I (1966) Viscosimetric determination of cellulase activity in the intestine of the sea urchin: reaction mechanism and equilibrium constant for cellulase stabilization with calcium. *Acta Chem Scand* 20:2667–2677
24. Elyakova LA (1972) Distribution of cellulases and chitinases in marine invertebrates. *Comp Biochem Physiol* 43B:67–70
25. Elyakova LA, Shevchenko NM, Avaeva SM (1981) A comparative study of carbohydrate activities in marine invertebrates. *Comp Biochem Physiol* 69B:905–908
26. Suzuki M, Nikuchi R, Onishi T (1984) The polysaccharide degradation activity in digestive tract of sea urchin *Strongylocentrotus nudus*. *Bull Jpn Soc Sci Fish* 50:1255–1260
27. Yamaguchi K, Araki I, Aoki T, Tseng C-H, Kitamikodo M (1989) Algal cell wall-degrading enzymes from viscera of marine animals. *Nippon Suisan Gakkaishi* 55:105–110
28. Obrietan K, Drinkwine M, Williams DC (1991) Amylase, cellulase and protease activities in surface and gut tissues of *Dendraster excentricus*, *Pisaster ochraceus* and *Strongylocentrotus droebachiensis* (Echinodermata). *Mar Biol* 109:53–57
29. Gomez-Pinchetti JL, Garcia-Reina G (1993) Enzymes from marine phycophages that degrade cell walls of seaweeds. *Mar Biol* 116:553–558
30. Park J, Johnson MJ (1949) A submicrodetermination of glucose. *Biol Chem* 181:149–151
31. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680–685
32. Beguin P (1983) Detection of cellulase activity in polyacrylamide gels using Congo red-stained agar replicas. *Anal Biochem* 131:333–336
33. Gornall AG, Bardawill CJ, David MM (1949) Determination of serum proteins by means of the biuret reaction. *J Biol Chem* 177:751–766
34. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265–275
35. Zhu XD, Mahairas G, Illies M, Cameron RA, Davidson EH, Etensohn CA (2001) A large-scale analysis of mRNA expressed by primary mesenchyme cells of the sea urchin embryo. *Development* 128:2615–2627
36. Dehal P, Satou Y, Campbell RK, Chapman J, Degnan B et al (2002) The draft genome of *Ciona intestinalis*: insights into chordate and vertebrate origins. *Science* 298:2157–2167
37. Chun CZ, Hur SB, Kim YT (1997) Purification and characterization of an endoglucanase from the marine rotifer, *Brachionus plicatilis*. *Mol Biol Int* 43:241–249
38. Singht J, Batra N, Sobti RC (2004) Purification and characterization of alkaline cellulase produced by a novel isolate, *Bacillus sphaericus* JS1. *J Microbiol Biotechnol* 31:51–56
39. Mawadza C, Hatti-Kaul R, Zvauya R, Mattiasson B (2000) Purification and characterization of cellulase produced by two *Bacillus* strains. *J Biotechnol* 83:177–187
40. Molodtsov NV, Vafina MG (1972) The distribution of β -N-acetylglucosaminidase in marine invertebrates. *Comp Biochem Physiol* 41B:113–120
41. Molodtsov NV, Vafina MG, Kim A, Sundukova EV, Artyukov AA, Bliknov YG (1974) Glycosidases of marine invertebrates from Poset Bay, Sea of Japan. *Comp Biochem Physiol* 48B:463–470
42. Klinger TS (1984) Activities and kinetics of digestive α - and β -glucosidase and β -galactosidase of five species of echinoids (Echinodermata). *Comp Biochem Physiol* 78A:597–600
43. Holzman TF, Russo SF, Williams DC (1985) Effects of feeding and starvation on proteolytic and tryptic activities in pyloric caecal tissues and duct fluids of the sea star *Pisaster ochraceus*. *Mar Biol* 90:55–59
44. Farrand AL, Williams DC (1988) Isolation, purification and partial characterization of four digestive proteases from the purple seastar *Pisaster ochraceus*. *Mar Biol* 97:231–236
45. Nakatani N, Kobayashi I (1996) Enzymatic properties of α -amylase from sea urchin, *Strongylocentrotus nudus*. *Comp Biochem Physiol* 113B:383–386
46. Klinger TS, Hsierh HL, Pangallo RA, Chen CP, Lawrence JM (1986) The effect of temperature on feeding, digestion, and absorption of *Lyechinus variegatus* (Lamarck) (Echinodermata: Echinoidea). *Physiol Zool* 59:332–336
47. Ura K, Tanaka E, Todo T, Gotoh T, Shimizu M, Ojima T, Takagi Y (2009) Purification of subtilase from short-spined sea urchin *Strongylocentrotus intermedius*. *Bull Fish Sci Hokkaido Univ* 58:21–28
48. Chang Y-Q, Lawrence JM, Cao X-B, Lawrence AL (2005) Food consumption, absorption, assimilation and growth of the sea urchin *Strongylocentrotus intermedius* fed a prepared food and the alga *Laminaria japonica*. *J World Aquac Soc* 36:68–75
49. Fuji A (1962) Studies on the biology of the sea urchin, V. Food consumption of *Strongylocentrotus intermedius*. *Jpn J Ecol* 12:181–186
50. Fuji A (1967) Ecological studies on the growth and food consumption of Japanese common littoral sea urchin, *Strongylocentrotus intermedius* (A.Agassiz). *Mem Fac Fish Hokkaido Univ* 12:83–160
51. Miller RJ, Mann KH (1973) Ecological energetics of the seaweed zone in marine bay on the Atlantic coast of Canada. III. Energy transformations by sea urchins. *Mar Biol* 18:99–114
52. Larson BR, Vadas RL, Keser M (1980) Feeding and nutritional ecology of the sea urchin *Strongylocentrotus droebachiensis* in Maine USA. *Mar Biol* 59:49–62
53. Pearce CM, Williams SW, Yuan F, Castell JD, Robinson SMC (2005) Effect of temperature on somatic growth and survivorship of early post-settled green sea urchin, *Strongylocentrotus droebachiensis* (Muller). *Aquac Res* 36:600–609
54. Agatsuma Y (2007) Ecology of *Strongylocentrotus nudus*. In: Lawrence JM (ed) Edible sea urchins. Elsevier Science, Amsterdam, pp 443–457
55. Stott FC (1995) The food canal of the sea-urchin *Echinus esculentus* L. and its functions. *Proc Zool Soc Lond* 125:63–86
56. Ura K, Hasegawa S, Tanaka E, Gotoh T, Ojima T, Takagi Y (2012) Immunohistochemical analysis of the proteolytic enzyme subtilase in the digestive organs of the sea urchin *Strongylocentrotus intermedius*. *Anat Rec* 295:73–77