



Title	Identification of the recognition sequence and target proteins for DJ-1 protease
Author(s)	Mitsugi, Hitomi; Niki, Takeshi; Takahashi-Niki, Kazuko; Tanimura, Kyoko; Yoshizawa-Kumagaye, Kumiko; Tsunemi, Masahiko; Iguchi-Ariga, Sanae M. M.; Ariga, Hiroyoshi
Citation	Febs letters, 587(16), 2493-2499 https://doi.org/10.1016/j.febslet.2013.06.032
Issue Date	2013-08-19
Doc URL	http://hdl.handle.net/2115/53264
Type	article (author version)
File Information	Mitsugi et al..pdf



[Instructions for use](#)

Identification of the recognition sequence and target proteins for DJ-1 protease

Hitomi Mitsugi^a, Takeshi Niki^b, Kazuko Takahashi-Niki^a, Kyoko Tanimura^c, Kumiko Yoshizawa-Kumagaye^c, Masahiko Tsunemi^c, Sanae M.M. Iguchi-Ariga^b, Hiroyoshi Ariga^{a,*}

^a*Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita-ku, Sapporo 060-0812, Japan*

^b*Graduate School of Agriculture, Hokkaido University, Kita-ku, Sapporo 060-8589, Japan*

^c*Peptide Institute, Inc. Minoh-shi, Osaka 562-8686, Japan*

*Corresponding author. Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita 12, Nishi 6, Kita-ku, Sapporo 060-0812, Japan. Tel: +81 11 706 3745, Fax: +81 11 706 4988.

E-mail address: hiro@pharm.hokudai.ac.jp (H. Ariga).

ABSTRACT

DJ-1 is a product for familial Parkinson's disease gene and for an oncogene and plays a role in anti-oxidative stress reaction. DJ-1 is also a cysteine protease, but the characteristics of DJ-1 protease remain unclear. In this study, we identified the recognition sequence for DJ-1 protease by using recombinant DJ-1 and a peptide library. Protease activity of DJ-1 lacking C-terminal α -helix (DJ-1 Δ H9) was stronger than that of full-sized DJ-1, and the most susceptible sequence digested by DJ-1 Δ H9 was valine-lysine-valine-alanine (VKVA) under the optimal conditions of pH 5.5 and 0 mM NaCl. Divalent ions, especially Cu²⁺, were inhibitory to DJ-1's protease activity. *c-abl* oncogene 1 product (ABL1) and kinesin family member 1B (KIF1B) containing VKVA were digested with DJ-1 Δ H9.

Keywords: DJ-1, protease, biochemistry

1. Introduction

DJ-1 was first identified by our group as a novel oncogene that transformed mouse NIH3T3 cells in cooperation with activated *ras* [1] and was later found to be a causative gene for a familial form of Parkinson's disease [2]. DJ-1 has multiple functions, including transcriptional regulation [3-6], anti-oxidative stress function [7-9], mitochondrial regulation [8, 10-12], and functions as a chaperone [13] and protease [14-16]. The structure of DJ-1 is similar to that of cysteine protease from *Pyrococcus horikoshii*, but C-terminal α -helix 9 blocks a catalytic domain of protease [17]. DJ-1 is cleaved at the C-terminal region under oxidative stress conditions *in vitro* and in cells [16, 18], and a C-terminally cleaved form of DJ-1 enhances cytoprotective action against oxidative stress-induced apoptosis [16]. Transthyretin, a causative protein in familial amyloidotic polyneuropathy, is degraded in cells transfected with full-sized DJ-1 and *in vitro* by recombinant DJ-1 lacking α -helix 9 (DJ-1 Δ H9), and mutation of C106 in DJ-1 results in loss of its protease activity [15]. These results suggest that α -helix 9 of DJ-1 is opened in cells by oxidative stress that causes pathological phenomena such as neurodegenerative diseases. Identification of the recognition sequence of DJ-1 protease and of the protein(s) that opens α -helix 9 of DJ-1 in cells will lead to elucidation of the physiological role of DJ-1 protease.

In this study, we identified the recognition sequence of DJ-1 by using recombinant DJ-1 Δ H9 and a peptide library, and *c-abl* oncogene 1 product (ABL1) and kinesin family member 1B (KIF1B) were digested by DJ-1 Δ H9.

2. Materials and methods

2.1. Plasmids and purification of DJ-1

Plasmids harboring GST-tagged wild-type DJ-1, C106S DJ-1, DJ-1 Δ H9 and C106S DJ-1 Δ H9 were described previously [15]. GST-tagged DJ-1s were expressed in and purified from *E. coli* and digested with PreScission protease (GE Bioscience, Little Chalfont, UK) to cleave off GST

as described previously [15] To construct deletion mutants lacking VKVA sequences of ABL1 and KIF1B, following PCR primers were used. ABL1 Δ VKVA-S:

5'-GGAGAACCACTTGGATTTTGGCCTGAGC-3', ABL1 Δ VKVA-AS:

5'-GCTCAGGCCAAAATCCAAGTGGTTCTCC-3', KIF1B Δ VKVA-S:

5'-GGAGCCTCAGTCCGGGTAAGGCC-3', and KIF1B Δ VKVA-AS:

5'-GGCCTTACCCGGACTGAGGCTCC-3'. PCR was carried out using pTOPO-ABL1 and

pTOPO-KIF1B as templates and primers above. Resultant PCR products were digested with

DpnI and used for transformation into *E. coli*.

2.2. Screening of the peptides digested with DJ-1

First screening: Ten μ M fluorescence-quenching substrate library FRETs-25Xaa Series (Peptide Institute, Osaka, Japan) was reacted with 15 μ g of recombinant DJ-1 in a buffer containing 0.5 mM acetate (pH 5.5) and 0.2 mM CaCl₂ at 37°C for 8-21 hrs and its fluorescence intensity was measured using a fluorometer (GloMax, Promega, Madison, WI, USA). Second screening: Three candidates of DJ-1 target sequences that had been identified in the first screening (FRETs-25Val, FRETs-25Ile and FRETs-25Ala) were reacted with 100 μ M FRETs-25Xaa Series and 67.5 μ g of DJ-1 Δ H9 in the same buffer as above at 37°C for 8 or 7 hrs. Then, a final concentration of 0.5 M EDTA (pH 8.1) was added to each reaction mixture, and the reaction mixtures were frozen in liquid nitrogen and subjected to LC-MS analysis to determine amino acids corresponding to Yaa and Zaa sites as described previously [19].

2.3. Digestion of target protein candidates for DJ-1 protease *in vitro*

cDNAs encoding target proteins for DJ-1 protease were cloned into pcDNA3. ³⁵S-labeled proteins were then synthesized *in vitro* using reticulocyte lysate of the TNT transcription-translation coupled system (Promega) and reacted with 1 μ g of recombinant DJ-1 in a buffer containing 0.5 mM acetate (pH 5.5) and 0.2 mM CaCl₂ at 37°C for 4 hrs. Bovine serum albumin (BSA) as a negative control was also reacted with ³⁵S-labeled proteins. Labeled

proteins were then separated on SDS-containing polyacrylamide gels and visualized by fluorography.

2.7. Statistical analyses

Data are expressed as means \pm S.E. Statistical analyses were performed using the Tukey-Kramer test.

3. Results

3.1. Identification of amino acid sequences digested by DJ-1

Since a C-terminal α -helix 9 in DJ-1 blocks the putative catalytic region of DJ-1 protease and since we have shown that recombinant DJ-1 lacking α -helix 9 (DJ-1 Δ H9) digested transthyretin *in vitro* and that this activity was diminished by substitution of an amino acid from cysteine to serine at 106 (C106S) [15], 4 recombinant DJ-1s were expressed in and purified from *E. coli* (Fig. 1A) and used in this study. BSA was also used as a negative control.

Fluorescence-quenching substrate library FRETs-25Xaa Series possesses a fluorescence group (Nma) and a quenching group (Dnp), and Nma's emission of fluorescence light is not activated by Dnp (Fig. 1B). When peptides in the library are digested with protease, fluorescence intensity of Nma is increased, enabling measurement of protease activity. The Xaa (P1) site contains 19 different amino acids other than cysteine, and Yaa (P2) and Zaa (P3) sites contain 5 kinds of amino acids with different characteristics. Each Xaa series is therefore a mixture of 25 kinds of peptides, and the whole library contains 475 kinds of peptides (Fig. 1B).

To determine amino acids in an Xaa site, FRETs-25Xaa Series was reacted with 4 recombinant DJ-1s and BSA for 21 hrs and its fluorescence was measured. Results for valine (Val), isoleucine (Ile), alanine (Ala) and proline (Pro) in the Xaa site are shown as examples, in which Pro in the Xaa site of the peptide was not digested with DJ-1 (Fig. 2A). While wild-type DJ-1, C106S DJ-1, C106S Δ H9 DJ-1 and BSA showed no or little protease activity against FRETs-25Xaa Series, DJ-1 Δ H9 digested FRETs-25Xaa Series with Val, Ile and Ala in the Xaa

sites in a time-dependent manner. A summary of DJ-1 protease activities using various amino acids in the Xaa sites in the library is shown in Fig. 2B. Of 19 amino acids, Val, Ile and Ala, which are categorized into nonpolar amino acids, in the Xaa sites of the peptide were found to be good substrates for protease activity of DJ-1 Δ H9. As the second screening, FRETs-25Val, -25Ile and -25Ala were reacted with DJ-1 Δ H9 for 4 hrs, and amino acids in Yaa and Zaa of FRETs-25Val, -25Ile and -25Ala were determined by using LC-MS. Ratio of peak area for FRETs-25Val in MS profiles versus enzyme digestibility was plotted (Fig. 2C), and a summary of all of the results is shown in Table 1. In the case of FRETs-25Val, for instance, the order of sequence digestibility to DJ-1 Δ H9 is VKV > VPV > RYV > FKV > RKV. DJ-1 protease digested the peptide bond between P1 amino acids and Ala.

3.2. Characteristics of DJ-1 protease

Biochemical characterization of DJ-1 protease was then carried out using VKVA peptide (D-A2pr(Nma)-GVKVAFPK(Dnp)rr) as a substrate and recombinant DJ-1s. Protease activity of DJ-1 Δ H9 appeared at pH 5.5-6.0 and disappeared at pH 7.4 and 9.5 (Fig. 3A), indicating that a weak acidic condition is optimal. DJ-1 protease activity was decreased in a salt-dependent manner and the concentration of 0 mM NaCl was maximal (Fig. 3B). Also, DJ-1 protease did not require divalent ions such as Ca⁺⁺, Mg⁺⁺ and Cu⁺⁺, and even a low concentration of Cu⁺⁺ diminished protease activity (Fig. 3C). Kinetic analysis of the enzymatic reaction showed that the reaction of DJ-1 Δ H9 protease activity obeyed the Michaelis-Mentene formula in which substrate concentration and reaction rate were placed on horizontal and longitudinal axes, respectively (Fig. 3D). Molecular activity and Michaelis-Menthen constant (*k*_{cat} and *K*_m values, respectively) were then calculated. The *k*_{cat} value is represented as substrate concentration/enzyme concentration x seconds, and *k*_{cat} of DJ-1 Δ H9 protease was calculated to be 1.02 x 10⁻⁴ (S⁻¹) (Fig. 3F). After the Lineweaver-Burk plot was made, *K*_m was calculated to be 40.1 μ M, which is 1/10⁵ of that of trypsin, a strong protease (Figs. 3E and 3F, respectively). These results indicate that although protease activity of DJ-1 Δ H9 is not strong, DJ-1 Δ H9 has

biochemical characteristics for protease.

Since Pfp1, whose crystal structure is similar to that of DJ-1, is a class of the cysteine protease and since a cysteine substitution mutant of DJ-1, C106S DJ-1, lost protease activity (Fig. 2A), it is thought that DJ-1 Δ H9 is also a cysteine protease. To ascertain this, two inhibitors for cysteine protease were reacted with DJ-1 Δ H9. E-64 is a specific inhibitor of cysteine protease and inhibits protease activity of papain. N-ethylmaleimide (NEM) nonselectively inhibits cysteine protease. As shown in Figs. 3G and 3H, protease activity of DJ-1 Δ H9 was inhibited by NEM, suggesting that, like Pfp1, DJ-1 Δ H9 is a cysteine protease.

3.3. Identification of candidate proteins for DJ-1 Δ H9 protease

A search for proteins that contain recognition sequences of DJ-1 protease was made by using the BLAST search against 3 amino acids shown in Table 1, and more than 2000 proteins were found. Since it is not possible to examine digestibility of all of these proteins, we focused on recognition sequence-containing proteins that interacted with DJ-1 such as DJBP, HIPK1 α , HIPK1 β , Topors, PTEN and tyrosine hydroxylase or that play roles in pathogenesis of cancer such as ABL1 and of Parkinson's disease (PD) such as LRRK2 and in mitochondrial regulation such as KIF1B, POMC and UCP1 [3, 4, 7, 20-27]. These proteins were synthesized *in vitro* using reticulocyte lysates and reacted with DJ-1 Δ H9, wild-type DJ-1 and C106S Δ H9. Of 14 proteins examined, only ABL1 and KIF1B were cleaved by DJ-1 Δ H9 but not by wild-type DJ-1 and C106S Δ H9, and the results for ABL1, KIF1B and HIPK1 α as a negative example are shown in Fig. 4A. When ABL1 and KIF1B were reacted with various amounts of DJ-1 Δ H9, they were digested with DJ-1 Δ H9 in a dose-dependent manner (Fig. 4B). ABL1 and KIF1B possess a VKVA sequence, the most digestible sequence for DJ-1 Δ H9 protease, at amino acid numbers 398 and 6 in the kinase domain and in the motor domain, respectively (Fig. 4D). When these sites were digested with DJ-1 Δ H9, ABL1 and KIF1B should be cleaved to two fragments with approximately 44 and 82 kDa and to those with approximately 0.7 and 126 kDa, respectively, and sizes of cleaved bands observed in Figs. 4A and 4B are well matched. A 126-kDa band of KIF1B cleaved by

DJ-1 Δ H9 was observed on bands that appeared in all the lanes. These bands may be translated from internal methionine of KIF1B or bands cleaved by protease present in reticulocyte lysates. The KIF1B fragment of 0.7 kDa was run through gels and only a band with 126 kDa was observed. To confirm whether a VKVA sequence in ABL1 and KIF1B is a target sequence for DJ-1 protease, ABL1 and KIF1B mutants lacking the VKVA sequence were constructed and reacted with wild-type DJ-1, DJ-1 Δ H9 and C106S Δ H9. As shown in Fig. 4C, these mutants were not cleaved by DJ-1 Δ H9. The reduced level of full-sized KIF1B mutants lacking the VKVA sequence was reproducibly observed. This may be due to conformational change of KIF1B mutant that resulted in low translational efficiency. These results suggest that ABL1 and KIF1B are candidate proteins for DJ-1 protease.

4. Discussion

In this study, we first identified recognition sequences for DJ-1 protease and found that VKVA, VKIA and VYAA were preferentially cleaved by DJ-1 lacking C-terminal α -helix 9 (DJ-1 Δ H9) *in vitro*, and wild-type, C10S DJ-1 and C106S Δ H9 DJ-1 had no or little protease activity (Figs. 1 and 2, Table 1). DJ-1 Δ H9 protease has biochemical characteristics of protease (Fig. 3), and ABL1 and KIF1B were identified to be candidate proteins for DJ-1 protease (Fig. 4). Preferred amino acids of the P1 site for DJ-1 protease are Val, Ile and Ala, which are non-polar amino acids, and these phenomena are similar to those for Pfp1 [28]. Optimal pH for DJ-1 protease was 5.5-6.0 and its activity decreased at more alkaline pH (Fig. 3). These results imply two possibilities: One is that DJ-1 protease works under oxidative stress conditions that induce acidic regions locally in cells, and the other possibility is that DJ-1 protease works in some organelles in which relatively low pH is maintained. Such organelles include the endosome, lysosome, Golgi apparatus and synaptic vesicle. Of these organelles, the lysosome contains many proteases whose optimal pH is around 5. It is possible that DJ-1 translocates to the lysosome to explore its protease activity under some stress conditions. Cu⁺⁺ strongly inhibited DJ-1 Δ H9 protease activity (Fig. 3C). Cu⁺⁺ is known to be a risk factor for Parkinson's disease and has been reported to bind

to DJ-1 [29]. It is therefore thought that Cu^{++} binds to a near active site of DJ-1 protease, thereby inhibiting its activity.

ABL1 and KIF1B were identified as target proteins for DJ-1 protease (Fig. 4). ABL1 is a proto-oncogene product and works as a non-receptor-type tyrosine kinase in cell differentiation, cell growth and cell adhesion, and its overexpression or structural abnormality results in onset of cancer [30]. Augmented expression and activity have also been reported in patients with Alzheimer's disease and amyotrophic lateral sclerosis [31-33], and overexpression of ABL1 leads to apoptosis induction [33]. Furthermore, it has been reported that ABL1 phosphorylates Parkin, a causative gene product of familial Parkinson's disease (Park 2), resulting in inactivation of its ubiquitin ligase activity [34]. It is therefore possible that DJ-1 degrades ABL1 to inhibit its kinase activity, thereby maintaining neuroprotective activity of Parkin. KIF1B is a motor protein belonging to the Kinesin family of proteins and has two splicing isoforms, KIF1B α and KIF1B β [25, 35, 36]. KIF1B α and KIF1B β transport mitochondria in nerve cells and the endoplasmic reticulum, respectively, and possess a VKVA sequence. It is possible that DJ-1 inhibits mitochondrial transport through cleavage of KIF1B α .

We have shown that transthyretin is cleaved by DJ-1 ΔH9 *in vitro* and that transthyretin is cleaved by transfection of full-sized DJ-1 in cells [15]. Chen et al. reported that DJ-1 in dopaminergic cells undergoes C-terminal cleavage in response to mild oxidative stress, and a C-terminally cleaved form of DJ-1 with activated protease activity enhances cytoprotective action against oxidative stress-induced apoptosis [16]. These results suggest that the C-terminal sequence of DJ-1 is opened under some stress conditions in cells. Although the physiological function of DJ-1 protease is still not known, the present findings will contribute to elucidation of DJ-1 function against stresses.

Acknowledgements

This work was supported by grants-in-aid from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) and by the Program for Promotion of Fundamental Studies in

Health Sciences of the National Institute of Biomedical Innovation (NIBIO) in Japan.

References

- [1] Nagakubo, D., Taira, T., Kitaura, H., Ikeda, M., Tamai, K., Iguchi-Ariga, S.M.M. and Ariga, H. (1997) DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras. *Biochem. Biophys. Res. Commun.* 231, 509-513.
- [2] Bonifati, V. et al. (2003) Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science* 299, 256-259.
- [3] Takahashi, K., Taira, T., Niki, T., Seino, C., Iguchi-Ariga, S.M.M. and Ariga, H. (2001) DJ-1 positively regulates the androgen receptor by impairing the binding of PIASx alpha to the receptor. *J. Biol. Chem.* 276, 37556-37563.
- [4] Niki, T., Takahashi-Niki, K., Taira, T., Iguchi-Ariga, S.M.M. and Ariga, H. (2003) DJBP: a novel DJ-1-binding protein, negatively regulates the androgen receptor by recruiting histone deacetylase complex, and DJ-1 antagonizes this inhibition by abrogation of this complex. *Mol. Cancer Res.* 1, 247-261.
- [5] Zhong, N. et al. (2006) DJ-1 transcriptionally up-regulates the human tyrosine hydroxylase by inhibiting the sumoylation of pyrimidine tract-binding protein-associated splicing factor. *J. Biol. Chem.* 281, 20940-20948.
- [6] Clements, C.M., McNally, R.S., Conti, B.J., Mak, T.W. and Ting, J.P.-Y. (2006) DJ-1, a cancer- and Parkinson's disease-associated protein, stabilizes the antioxidant transcriptional master regulator Nrf2. *Proc. Natl. Acad. Sci. USA* 103, 15091-15096.
- [7] Taira, T., Saito, Y., Niki, T., Iguchi-Ariga, S.M.M., Takahashi, K. and Ariga, H. (2004) DJ-1 has a role in antioxidative stress to prevent cell death. *EMBO Rep.* 5, 213-218.
- [8] Canet-Aviles, R.M. et al. (2004) The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfinic acid-driven mitochondrial localization. *Proc. Natl. Acad. Sci. USA* 101, 9103-9108.
- [9] Martinat, C., Shendelman, S., Jonason, A., Leete, T., Beal, M.F., Yang, L. and Floss, T.

- (2004) Abeliovich, sensitivity to oxidative stress in DJ-1-deficient dopamine neurons: an ES-derived cell model of primary parkinsonism. *PLoS Biol.* 2, e327.
- [10] Hao, L.Y., Giasson, B.I. and Bonini, N.M. (2010). DJ-1 is critical for mitochondrial function and rescues PINK1 loss of function. *Proc. Natl. Acad. Sci. USA* 107, 9747-9752.
- [11] Krebiehl, G. et al. (2010). Reduced basal autophagy and impaired mitochondrial dynamics due to loss of Parkinson's disease-associated protein DJ-1. *PLoS One* 5, e9367.
- [12] Thomas, K.J. et al. (2011). DJ-1 acts in parallel to the PINK1/parkin pathway to control mitochondrial function and autophagy. *Hum. Mol. Genet.* 20, 40-50.
- [13] Zhou, W., Zhu, M., Wilson, M.A., Petsko, G.A. and Fink, A.L. (2006) The oxidation state of DJ-1 regulates its chaperone activity toward alpha-synuclein. *J. Mol. Biol.* 356, 1036–1048.
- [14] Olzmann, J.A. et al. (2004) Familial Parkinson's disease-associated L166P mutation disrupts DJ-1 protein folding and function. *J. Biol. Chem.* 279, 8506-8515.
- [15] Koide-Yoshida, S. et al. (2007) DJ-1 degrades transthyretin and an inactive form of DJ-1 is secreted in familial amyloidotic polyneuropathy. *Int. J. Mol. Med.* **19**, 885-893.
- [16] Chen, J., Li, L. and Chin, L.S. (2010) Parkinson disease protein DJ-1 converts from a zymogen to a protease by carboxyl-terminal cleavage. *Hum. Mol. Genet.* 19, 2395-2408.
- [17] Honbou, K., Suzuki, N.N., Horiuchi, M., Niki, T., Taira, T., Ariga, H. and Inagaki, F. (2003) The crystal structure of DJ-1, a protein related to male fertility and Parkinson's disease. *J. Biol. Chem.* 278, 31380-31384.
- [18] Ooe, H., Maita, C., Maita, H., Iguchi-Ariga, S.M.M. and Ariga, H. (2006) Specific cleavage of DJ-1 under an oxidative condition. *Neuroscience Lett.* 406, 165-168.
- [19] Tanskul, S., Oda, K., Oyama, H., Noparatnaraporn, N., Tsunemi, M. and Takada, K. (2003) Substrate specificity of alkaline serine proteinase isolated from photosynthetic bacterium, *Rubrivivax gelatinosus* KDD51. *Biochem. Biophys. Res. Commun.* 309, 547-551.

- [20] Sekito, A., Koide-Yoshida, S., Niki, T., Taira, T., Iguchi-Arigo, S. M. and Arigo, H. (2006) DJ-1 interacts with HIPK1 and affects H₂O₂-induced cell death. *Free Radic. Res.* 40, 155-116.
- [21] Kim, R.H. et al. (2005) DJ-1, a novel regulator of the tumor suppressor PTEN. *Cancer Cell* 7, 263-273.
- [22] Ishikawa, S., Taira, T., Takahashi-Niki, K., Niki, T., Arigo, H. and Iguchi-Arigo, S.M.M. (2010) Human DJ-1-specific transcriptional activation of tyrosine hydroxylase gene. *J. Biol. Chem.* 285, 39718-39731.
- [23] Paisán-Ruíz, C. et al. (2004) Cloning of the gene containing mutations that cause PARK8-linked Parkinson's disease. *Neuron* 44, 595-600.
- [24] Wang, J. Y., Ledley, F., Goff, S., Lee, R., Groner, Y. and Baltimore, D. (1984) The mouse c-abl locus: molecular cloning and characterization. *Cell* 36, 349-356.
- [25] Nangaku, M., Sato-Yoshitake, R., Okada, Y., Noda, Y., Takemura, R., Yamazaki, H. and Hirokawa, N. (1994) KIF1B, a novel microtubule plus end-directed monomeric motor protein for transport of mitochondria. *Cell* 79, 1209-1220.
- [26] Kimura, T. (1986) Transduction of ACTH signal from plasma membrane to mitochondria in adrenocortical steroidogenesis. Effects of peptide, phospholipid, and calcium. *J. Steroid. Biochem.* 25, 711-716.
- [27] Kozak, L.P. and Koza, R.A. (1999) Mitochondria uncoupling proteins and obesity: molecular and genetic aspects of UCP1. *Int. J. Obes. Relat. Metab. Disord.* 23 Suppl 6, S33-S37.
- [28] Halio, S.B., Bauer, M. W., Mukund, S., Adams, M. and Kelly, R.M. (1997) Purification and Characterization of Two Functional Forms of Intracellular Protease PfpI from the Hyperthermophilic Archaeon *Pyrococcus furiosus*. *Appl. Environ. Microbiol.* 63, 289-295.
- [29] Xu, X. M., Lin, H., Maple, J., Björkblom, B., Alves, G., Larsen, J. P. and Møller, S.G. (2010) The Arabidopsis DJ-1a protein confers stress protection through cytosolic SOD activation. *J. Cell Sci.* 123, 1644-1651.
- [30] Koretzky, G.A. (2007) The legacy of the Philadelphia chromosome. *J. Clin. Invest.* 117,

2030-2032.

- [31] Cancino, G.I., Toledo, E.M., Leal, N.R., Hernandez, D.E., Yévenes, L.F., Inestrosa, N.C. and Alvarez, A.R. (2008) STI571 prevents apoptosis, tau phosphorylation and behavioural impairments induced by Alzheimer's beta-amyloid deposits. *Brain* 131, 2425-2442.
- [32] Alvarez, A.R., Sandoval, P.C., Leal, N.R., Castro, P.U. and Kosik, K.S. (2004) Activation of the neuronal c-Abl tyrosine kinase by amyloid-beta-peptide and reactive oxygen species. *Neurobiol. Dis.* 17, 326-336.
- [33] Katsumata, R. et al. (2012) c-Abl inhibition delays motor neuron degeneration in the G93A mouse, an animal model of amyotrophic lateral sclerosis. *PLoS One* 7, e46185.
- [34] Ko, H.S. et al. (2010) Phosphorylation by the c-Abl protein tyrosine kinase inhibits parkin's ubiquitination and protective function. *Proc. Natl. Acad. Sci. USA* 107, 16691-16696.
- [35] Gong, T.W., Winnicki, R.S., Kohrman, D.C. and Lomax, M.I. (1999) A novel mouse kinesin of the UNC-104/KIF1 subfamily encoded by the Kif1b gene. *Gene* 239, 117-127.
- [36] Conforti, L., Buckmaster, E.A., Tarlton, A., Brown, M.C., Lyon, M.F., Perry, V.H. and Coleman, M.P. (1999) The major brain isoform of kif1b lacks the putative mitochondria-binding domain. *Mamm. Genome.* 10, 617-622.

Legends of figures

Fig. 1. DJ-1 mutants and FRETs-25Xaa Series.

A. Schematic drawing of recombinant DJ-1s used in this study (left panel). Purified DJ-1 proteins were separated on 12% polyacrylamide gel and stained with coomassie brilliant blue (right panel).

B. Schematic drawing of FRETs-25Xaa Series.

Fig. 2. Identification of recognition sequences for DJ-1 protease.

A. Amino acids in the P1 site of FRETs-25Xaa Series were identified using various DJ-1s shown in Fig. 1A and BSA. Results for Val, Ile, Ala and Pro in the P1 site are shown.

B. Summary of the results of experiments described in the legend for Fig. 2A.

C. Amino acids in P2 and P3 sites of FRETTS-25Val were identified as described in Materials and methods.

Fig. 3. Biochemical characterization of DJ-1 protease.

A-C. Optimal conditions of pH (A), NaCl concentration (B) and concentration of divalent ions (C) were examined for valine, lysine and valine (VKV) in P1, P2 and P3 sites of FRETTS-25Xaa using DJ-1 Δ H9.

D, E. Michaelis-Menten kinetics (D) and Lineweaver-Burk plot (E) were drawn after reaction of DJ-1 Δ H9 with VKV in P1, P2 and P3 sites of FRETTS-25Xaa.

F. *k*_{cat} and *K*_m were calculated from the results shown in Figs. 3D and 3E.

G, H. DJ-1 Δ H9 was reacted with VKV in P1, P2 and P3 sites of FRETTS-25Xaa and various concentrations of E-64 (G) and N-ethylmaleimide (H), and protease activity of DJ-1 Δ H9 is shown.

Fig. 4. Identification of ABL1 and KIF1B as target proteins for DJ-1 protease.

A. ³⁵S-labeled ABL1, KIF1B and HIPK1 α were synthesized *in vitro* using reticulocyte lysates, reacted with wild-type, Δ H9 and C106S Δ H9 DJ-1 and BSA, and separated on polyacrylamide gels. Proteins were then visualized by fluorography.

B. ³⁵S-labeled ABL1 and KIF1B were reacted with various concentrations of wild-type and Δ H9 DJ-1 and visualized by fluorography.

C. ³⁵S-labeled ABL1, ABL1 Δ VKVA, KIF1B and KIF1B Δ VKVA were reacted with wild-type, Δ H9 and C106S Δ H9 DJ-1 and subjected to fluorography. Full-sized ABL1 and KIF1B, and large cleaved products of ABL1 and KIF1B are shown by arrows.

D. Schematic drawing of structures of ABL1 and KIF1B.

Table 1. Relative hydrolysis rate of DJ-1 Δ H9 for FRETTS substrates.

FRETS-25Val		FRETS-25Ile		FRETS-25Ala	
P3P2P1	Relative rate	P3P2P1	Relative rate	P3P2P1	Relative rate
VKV	100 %	VKI	100 %	VYA	100 %
VPV	64	RYI	64	VKA	59
RYV	54	VPI	50	RYA	50
FKV	52	RKI	49	VPA	34
RKV	51	FKI	47	RIA	33

Digestibility of amino acids in P1, P2 and P3 sites for DJ-1 Δ H9 protease is shown.




Fig. 1







Fig. 2



D. Michaelis-Menten kinetics




E. Lineweaver-Burk plot



F

	k_{cat} (S^{-1})	K_m (μM)	k_{cat}/K_m ($M^{-1} S^{-1}$)
DJ-1 Δ H9	1.02×10^{-4}	40.1	2.5
Trypsin			2.5×10^5

G. E-64



H. N-Ethylmaleimide




Fig. 3




Fig. 4