

Pyrroloquinoline Quinone Prevents Oxidative Stress-Induced Neuronal Death Probably through Changes in Oxidative Status of DJ-1

Kana NUNOME,^a Shin MIYAZAKI,^a Masahiko NAKANO,^b Sanae IGUCHI-ARIGA,^c and Hiroyoshi ARIGA^{*,a}

^aGraduate School of Pharmaceutical Sciences, Hokkaido University; ^cGraduate School of Agriculture, Hokkaido University; Kita-ku, Sapporo 060–8589, Japan; and ^bMitsubishi Gas Chemical Co., Inc., Niigata Research Laboratory; Kita-ku, Niigata 950–3112, Japan. Received March 22, 2008; accepted May 1, 2008; published online May 2, 2008

Pyrroloquinoline quinone (PQQ) has been shown to play a role as an anti-oxidant in neuronal cells and prevent neuronal cell death in a rodent stroke model. DJ-1, a causative gene product for a familial form of Parkinson's disease, plays a role in anti-oxidative stress function by self-oxidation of DJ-1. In this study, the expression level and oxidation status of DJ-1 were examined in SHSY-5Y cells and primary cultured neurons treated with 6-hydroxydopamine (6-OHDA) or H₂O₂ in the presence or absence of PQQ. The pI shift of DJ-1 to an acidic point, which was observed in SHSY-5Y cells treated with 6-OHDA, was inhibited by PQQ. TOF-MS analyses showed that while the level of a reduced form of DJ-1, one of the active forms of DJ-1, was decreased in SHSY-5Y cells treated with 6-OHDA or H₂O₂, PQQ increased the level of the reduced form of DJ-1. These results suggest that PQQ prevents oxidative stress-induced changes in oxidative status of DJ-1. Therefore, the neuroprotective effects of PQQ on oxidative stress-induced neuronal death may be at least in part involved in increased level of an active form of DJ-1.

Key words DJ-1; pyrroloquinoline quinone; oxidative stress; neurotoxicity

Pyrroloquinoline quinone (PQQ) was identified as a redox cofactor of bacterial dehydrogenases¹⁾ and later found to be present in various organisms, including mammals.^{2–6)} PQQ has been shown to act as an anti-oxidant by scavenging superoxide radicals and to protect mitochondria from oxidative stress-induced damage.⁷⁾ PQQ has also been reported to prevent neuronal cell death in a rodent stroke model^{8–14)} and to prevent fibril formation of α -synuclein *in vitro*.¹⁵⁾ The mechanism underlying PQQ activity, however, has not been elucidated.

DJ-1 was identified by us as a novel oncogene¹⁶⁾ and later found to be a causative gene for a familial form Parkinson's disease (PD), *PARK7*.¹⁷⁾ DJ-1 is a multi-functional protein and has been shown to play roles in anti-oxidative stress reaction,^{18–21)} transcriptional regulation^{22–26)} and chaperone reaction.²⁷⁾ DJ-1 knockdown cells or DJ-1 knockout mice that had been treated with H₂O₂ or with several neurotoxins such as MPP+, 6-hydroxydopamine (6-OHDA), paraquat, and rotenone became highly susceptible to cell death,^{18–21,28–33)} and administration of DJ-1 into the substantia nigra of PD model rats that had been treated with 6-OHDA protected the rat against dopaminergic cell death and decreased their locomotion defects.³⁴⁾ It has also been shown that activity of mitochondrial complex I was reduced in DJ-1 knockdown cells^{19,35)} or in DJ-1 knockout mice,^{28,29)} suggesting that DJ-1 also regulates activity of mitochondrial complex I.

DJ-1 has three cysteines located at amino acid numbers 46, 53 and 106 (C46, C53 and C106, respectively). Of these three cysteines, C106 is the most susceptible to oxidation and is oxidized as SOH, SO₂H and SO₃H forms.³⁶⁾ Although oxidation at C106 is necessary for exerting activities of DJ-1,^{18–20,37)} it has been reported that highly oxidized DJ-1 at C106 lost its activity³⁸⁾ and that abnormally oxidized forms of DJ-1 were observed in patients with sporadic forms of PD and Alzheimer's disease.^{39,40)}

In this study, the expression levels and oxidative status at

C106 of DJ-1 were examined in human neuroblastoma cell line SH-SY5Y cells treated with H₂O₂ or 6-OHDA in the presence or absence of PQQ. The results showed that although the expression level of DJ-1 was not changed, the levels of reduced DJ-1 and oxidized DJ-1 at C106 with SOH were increased in cells treated with 6-OHDA and PQQ compared to the level in cells treated with 6-OHDA alone, suggesting that anti-oxidative activity of PQQ is, at least in part, catalyzed by DJ-1 activity.

MATERIALS AND METHODS

Materials PQQ was provided by Mitsubishi Gas Chemical Co., Inc. Vitamin C (ascorbic acid) and vitamin E (α -tocopherol) were purchased from Wako Pure Chemical, Japan.

Cells and Cell Viability Assay Human SH-SY5Y cells were cultured in Dulbecco's modified Eagle's medium supplemented with 10% calf serum. Cells in 96-well plates (5000 cells/well) were cultured with or without various amounts of PQQ, vitamin C or vitamin E for 24 h and then treated with various amounts of H₂O₂ for 3 h or with 6-OHDA for 48 h. The cells were harvested and their viability was examined by a methyl thiazolyl tetrazolium (MTT) assay using a cell counting kit 8 (Dojindo).

Primary Neuronal Culture of the Ventral Mesencephalon Cultures of the rat mesencephalon were established according to methods described previously.⁴¹⁾ The ventral two-thirds of the mesencephalon was dissected from rat embryos on the 17–19th days of gestation. The dissected regions included dopaminergic neurons from the substantia nigra and the ventral tegmental area but not noradrenergic neurons from the locus ceruleus. Neurons were dissociated mechanically and plated onto 0.1% polyethyleneimine-coated 96-well plates at a density of 2.5×10^6 cells/well. The cells were cultured in Dulbecco's modified Eagle's medium containing 10% fetal calf serum for 2 d, and then in the same medium containing 2% B-27 supplement (Invitrogen) and

* To whom correspondence should be addressed. e-mail: hiro@pharm.hokudai.ac.jp

2 mg/ml aphidicolin (Sigma) without fetal calf serum from the 3rd day onwards. The animals were treated in accordance with guidelines published in the NIH Guide for the Care and Use of Laboratory Animals.

To examine the effects of PQQ, vitamin C and vitamin E on oxidative stress-induced cell death, the cells were cultured in the presence or absence of 15 μM of each compound for 24 h and then treated with 40 μM 6-OHDA for 24 h or with 300 μM H_2O_2 for 1.5 h. Cell viability was then examined by an MTT assay.

Western Blotting To examine the expression of DJ-1 in SH-SY5Y cells that had been treated with 100 μM 6-OHDA for 4 h in the presence or absence of 15 μM PQQ, proteins were extracted from SH-SY5Y cells after incubation in a buffer containing 150 mM NaCl, 1 mM EDTA, 20 mM Tris (pH 8.0), and 0.5% NP-40. Proteins were then separated on a 12.5% polyacrylamide gel containing SDS and subjected to Western blotting with a rabbit anti-DJ-1 (1/2000 dilution) or mouse anti-actin antibody (1/4000 dilution) (Chemicon). Proteins on the membrane were reacted with an IRDye800 (Rockland) or Alexa680-conjugated secondary antibody (Molecular Probe) and visualized by using an infrared imaging system (Odyssey, LI-COR). For isoelectric focusing analysis, proteins were then separated in an isoelectric focusing gel of pH 5–8 and subjected to Western blotting as described above. The rabbit anti-DJ-1 antibody was established by us as described previously.¹⁶⁾

Identification of Oxidative Status of C106 of DJ-1 by MALDI-TOF/TOF-MS Analysis SH-SY5Y cells were treated with H_2O_2 or 6-OHDA in the presence or absence of PQQ as described above. Cell extracts were prepared and immunoprecipitated with a rabbit anti-DJ-1 polyclonal antibody (final concentration of 4 $\mu\text{g}/\text{ml}$), and immunoprecipitates were separated on a 12.5% polyacrylamide gel. After the gel had been stained with Coomassie brilliant blue (CBB), a band corresponding to DJ-1 was cut out, reduced, alkylated with a buffer containing iodoacetamide, and digested with trypsin. The peptide solutions were desalted, mixed with α -cyano-4-hydroxycinnamic acid, and applied onto a target plate. MS/MS spectra of the C106-containing peptide spanning 100–122 amino acids were obtained by using an Ultraflex (Brucker Daltonics) in reflector mode and analyzed with flexanalysis software (Brucker Daltonics). Protein identification was carried out with Mascot software against the database of NIBI.

Binding of PQQ to the C106 Region by a Quartz Crystal Microbalance (QCM) Fixation of PQQ and DJ-1-binding compounds on a sensor chip of QCM (Affinix Q, Initiium) was carried out as follows. In the case of compound A, the sensor chip was incubated with 100 μl of 5 mM 3,3'-dithiodipropionic acid for 1 h and then with 50 μl each of 100 mg/ml 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) and 100 mg/ml *N*-hydroxysuccinimide (NHS) for 20 min. After washing the chip with H_2O , the chip was incubated with 4 μl of 1 mM compound A dissolved in H_2O for 1 h. In the case of compound C, the sensor chip was washed with a solution containing H_2O_2 and sulfonic acid (molar ratio=1:3), and then it was incubated with 4 μl of 1 mM compound C or PQQ dissolved in chloroform until the solution had evaporated. In the case of compound PQQ, the sensor chip was incubated with 20 mM 2-aminoethanethiol

for 1 h, and then it was incubated with 4 μl of 1 mM PQQ dissolved in 50 μl each of 100 mg/ml EDC and 100 mg/ml NHS for 20 min. After washing the chip, it was incubated with 100 μl of 1 M Tris-HCl (pH 8.8). To the sensor chips fixed with compounds in Affinix Q, 8 μl of 1 mg/ml wild-type DJ-1 was applied, and their frequency was measured according to the manufacturer's protocol.

Statistical Analysis Data are expressed as means \pm S.D. Statistical analyses were performed using analysis of variance (one-way ANOVA) followed by unpaired Student's *t*-test.

RESULTS

Effect of PQQ on Oxidative Stress-Induced Cell Death

The effect of PQQ on cell viability was first examined. Human dopaminergic neuroblastoma cell line SH-SY5Y cells were incubated with various concentrations of PQQ for 24 h and cell viability was measured by an MTT assay (Fig. 1A). Although PQQ at concentrations up to 20 μM had little effect on cell viability, PQQ at concentration of more than 30 μM significantly reduced cell viability. We therefore used 15 μM PQQ in this study. The effect of PQQ on oxidative stress-induced cell death was then examined. Vitamins C and E, well known anti-oxidants, were used as positive controls. SH-SY5Y cells were incubated with 15 μM each of PQQ, vi-




Fig. 1. Effect of PQQ on Oxidative Stress-Induced Death in SHSY-5Y Cells

(A) SH-SY5Y cells were pretreated with various concentrations of PQQ for 24 h, and cell viability was measured by an MTT assay. ** p <0.01 vs. cell viability without PQQ. (B, C) SH-SY5Y cells were pretreated with 15 μM of PQQ, vitamin C or vitamin E for 24 h and then treated with 6-OHDA for 48 h (B) or with H_2O_2 for 3 h (C) and cell viability was measured by an MTT assay. Significance: * p <0.05, ** p <0.01.

tamin C and vitamin E for 24 h and then treated with 6-OHDA for 48 h or with H_2O_2 for 3 h, and cell viability was measured by the MTT assay (Figs. 2B, C, respectively). Without PQQ, vitamin C and vitamin E, cell death occurred in a dose-dependent manner. With PQQ, vitamin C and vitamin E, on the other hand, cell death induced by addition of H_2O_2 or 6-OHDA was inhibited, and the level of inhibitory activity of PQQ was much higher than that of vitamin C and vitamin E at the concentration used.

Primary neuronal cells of the ventral mesencephalon were prepared from rat embryos on the 17–19th days of gestation, and cells were pretreated with compounds for 24 h and then treated with $300 \mu M H_2O_2$ for 1.5 h or with $40 \mu M$ 6-OHDA for 24 h. While neither vitamin C nor vitamin E showed significant protective activity, PQQ significantly abrogated cell death (Fig. 2A or B, respectively).

Effect of PQQ on the Expression and Oxidative Status of DJ-1 SH-SY5Y cells were cultured with $15 \mu M$ PQQ for 24 h and then cultured with or without $100 \mu M$ 6-OHDA for 48 h. Protein extract was prepared from cells and analyzed by Western blotting with anti-DJ-1 and anti-actin antibodies, and the expression level of DJ-1 in cells was normalized against that of actin (Fig. 3A). The results showed that there was no change in DJ-1 expression in cells after treatment with PQQ.

Since the oxidative status of DJ-1 at C106 is important for its activity, proteins in cells were separated on two-dimension (2D) gels and subjected to Western blotting with an anti-DJ-1 antibody (Fig. 3B). Without PQQ and 6-OHDA treatment, major and minor spots with isoelectric points of pH 6.2 and 5.8, respectively, were observed, and a pattern similar to

that without PQQ treatment was observed after cells treated with PQQ (Figs. 3B-a, B-b, respectively). After treatment of cells with 6-OHDA, on the other hand, a band with a more acidic point appeared and this band disappeared in cells treated with 6-OHDA and PQQ (Figs. 3B-c, B-d, respectively). These results suggest that PQQ prevents oxidation of DJ-1 that is induced by 6-OHDA.

To verify the oxidative status of DJ-1 precisely, SH-SY5Y cells were treated with 6-OHDA or with H_2O_2 and PQQ as described above. The cell extract was then immunoprecipitated with an anti-DJ-1 antibody and precipitates were separated on polyacrylamide gel. A protein band corresponding to DJ-1 was cut out after staining the gel, digested with trypsin, and subjected to MALDI-TOF/TOF-MS analysis (Fig. 4A). A segment containing C106 had seven peaks that correspond to reduced and oxidized C106 as SOH, SO_2H and SO_3H . Since some of cysteine residues that had been reduced and oxidized as SOH and SO_2H , but not as SO_3H , were modified with iodoacetamide, molecular weight 56 was added to molecular weight of C106-containing peptides with unmodified and modified forms (Fig. 4A).

Without H_2O_2 or 6-OHDA treatment, the reduced form of C106 was 90% of total forms (Fig. 4B). After treatment with 6-OHDA, the reduced form of C106 decreased to 75%. Pretreatment with PQQ increased the reduced form of C106 to about 90%, but pretreatment with vitamin C decreased the reduced form of C106 to 65% (Fig. 4B, left parts). After treatment with H_2O_2 , on the other hand, the reduced form of C106 decreased to 42%. Pretreatment with PQQ and with vitamin C increased the reduced form of C106 to about 76% and to 46%, respectively (Fig. 4B, right parts). The oxidized form of C106 as SO_3H that had appeared after treatment with H_2O_2 also decreased in cells pretreated with PQQ. These results indicate that DJ-1 at least with the reduced form of




Fig. 2. Effect of PQQ on Oxidative Stress-Induced Death of Primary Neurons

Rat mesencephalic cultured neurons were pretreated with $15 \mu M$ of PQQ, vitamin C or vitamin E for 24 h and then treated with $40 \mu M$ 6-OHDA for 24 h (A) or with $300 \mu M H_2O_2$ for 1.5 h (B) and cell viability was measured by an MTT assay. Significance: $**p < 0.01$ vs. cell viability with 6-OHDA or H_2O_2 in the absence of compounds.




Fig. 3. Abrogation of Superfluous Oxidation of DJ-1 at C106 by PQQ in SH-SY5Y Cells

(A) SH-SY5Y cells were cultured with or without $15 \mu M$ of PQQ for 24 h and then treated with $100 \mu M$ 6-OHDA for 4 h. Cell extracts were then prepared and analyzed by Western blotting with anti-DJ-1 and anti-actin antibodies as described in Materials and Methods. After quantification of intensities of bands corresponding to DJ-1 and actin using an infrared imaging system (Odyssey, LI-COR), relative expression (DJ-1/actin) was determined. (B) SH-SY5Y cells were treated with PQQ and 6-OHDA as described in the legend of Fig. 3A, proteins in cell extracts were separated on 2D gels and subjected to Western blotting with anti-DJ-1 and anti-actin antibodies as described in Materials and Methods.




Fig. 4. Effect of PQQ on the Oxidative Status of DJ-1 in SHSY-5Y Cells

(A) SH-SY5Y cells were pretreated with 15 μ M of PQQ or vitamin C for 24 h and then treated with 100 μ M 6-OHDA for 4 h or with 400 μ M H₂O₂ for 90 min. Cell extracts were immunoprecipitated with an anti-DJ-1 antibody and the precipitates were separated on a 12% polyacrylamide gel. A stained band corresponding to DJ-1 was analyzed by MALDI-TOF/TOF-MS as described in Materials and Methods. Since cysteine residues that had been reduced and oxidized C106 as SOH and SO₂H, but not as SO₃H, were modified with iodoacetamide, a molecular weight of 56 was added to the molecular weights of C106-containing peptides spanning 100–122 amino acids. (B) Ratios of reduced and oxidized forms of C106 are presented.

C106 is an active form and that DJ-1 with the SO₃H form of C106 is an inactive form. It is therefore thought that PQQ prevents superfluous oxidation of DJ-1.

No Binding of PQQ to DJ-1 We identified compounds that bind to the C106 region and found that these compounds prevented oxidative stress-induced death of cells.⁴²⁾ We therefore examined whether PQQ also binds to DJ-1 by using a quartz crystal microbalance in which PQQ, compound A or compound C was fixed on a sensor chip and recombinant

wild-type DJ-1 was applied. In this case, compounds A and C were used as DJ-1-binding and non-binding compounds, respectively. As shown in Fig. 5, compound A but not C bound to DJ-1, and PQQ hardly bound to DJ-1.

DISCUSSION

6-OHDA is a neurotoxin that compromises mitochondrial complex I, resulting in production of reactive oxygen species




Fig. 5. No Binding of PQQ to DJ-1

Binding of PQQ to DJ-1 was examined by using a quartz crystal microbalance as described in Materials and Methods. Compounds A and C were used as DJ-1-binding and nonbinding compounds, respectively. ΔF indicates decreased frequency of a sensor chip.

(ROS) such as superoxide, hydroxyl radical and H_2O_2 . The ROS produced then attacks mitochondria, resulting in the damage in mitochondrial complex I. Reactive nitrogen species such as nitrogen oxide (NO) and peroxynitrite also cause damage to cells, including neuronal cells. It has been reported that PQQ scavenges superoxide anion and hydroxyl radicals in an *in vitro* system⁷) and that PQQ suppresses peroxynitrite formation and NO-induced neurotoxicity.⁹) The molecular mechanism underlying the preventive effect of PQQ, however, has not been elucidated. In this study, we showed that PQQ prevented H_2O_2 - and 6-OHDA-induced death of the dopaminergic neuroblastoma cell line SH-SY5Y and primary rat neurons and that its preventive effect was stronger than that of vitamins C and E, well-known antioxidants, at the concentration used in this study. Under these conditions in which SH-SY5Y cells were treated with H_2O_2 or 6-OHDA in the presence or absence of PQQ, we found that the level of oxidation of DJ-1 at C106 was reduced by PQQ addition compared to that without PQQ, though the expression level and localization (data not shown) of DJ-1 were not changed. Although it is not clear which forms of oxidized C106, SH (reduced), SOH, SO_2H and SO_3H , of DJ-1 are active forms, the level of C106 with SH was clearly changed. It has been reported that DJ-1 has a chaperone activity inhibiting aggregation of α -synuclein *in vitro* and that DJ-1 at C106 with SO_2H is the active form in this reaction.³⁸) In SH-SY5Y cells treated with H_2O_2 or 6-OHDA, however, the level of C106 with SO_2H increased, suggesting that other forms such as SH and SOH forms of C106 are active. In cells treated with H_2O_2 , the level of C106 with SO_3H increased and then decreased after addition of PQQ. Since DJ-1 at C106 with SO_3H was inactive in terms of chaperone activity³⁸) and a highly oxidized form of DJ-1, perhaps DJ-1 at C106 with SO_3H , was observed in brains of patients with Parkinson's disease and Alzheimer's disease,^{39,40}) DJ-1 at C106 with SO_3H is thought to be inactive against oxidative stress-induced cell death.

The level of DJ-1 at C106 with SO_2H was increased after SH-SY5Y cells were treated with 6-OHDA and vitamin C compared to that in cells treated with 6-OHDA alone (Fig. 4). This phenomenon was different from that in cells treated with 6-OHDA and PQQ, though both PQQ and vitamin C exerted protective effects against 6-OHDA-induced cell death (Fig. 1), suggesting that there is a distinct mechanism of anti-oxidative stress function between PQQ and vitamin C.

The cysteine residue at amino acid number 106 (C106) is an essential amino acid for DJ-1 to exert its activity.^{18–20,37}) We have identified compounds that bind to the C106 region of DJ-1, and these compounds prevented oxidative stress-in-

duced neuronal cell death and movement defect in cultured cells and in 6-OHDA-injected PD rat models.⁴²) These compounds A (UCP0045037) and B (UCP0054278), which are novel and uncharacterized compounds, respectively, prevent superficial oxidation of C106 by binding to the C106 region, thereby keeping active forms of DJ-1. Of DJ-1-binding compounds, compound A was used as a positive control in a DJ-1-binding experiment of PQQ (Fig. 5). Since PQQ was found not to bind to DJ-1, there are at least two possibilities. One possibility is that elimination of ROS by PQQ leads to prevention of DJ-1 oxidation. Another possibility is that PQQ affects some signaling pathways that regulate oxidation of DJ-1. Although pathways in oxidative stress reactions that are affected by PQQ have not been identified, it has recently been reported that PQQ activated *ras* signaling pathways such as the Raf/ERK pathway, leading to stimulation of growth of *ras*-transformed NIH3T3 but not untransformed NIH3T3 cells.⁴³) Since *DJ-1* was identified as an oncogene in cooperation with activated *ras*,¹⁶) factors involved in *ras* signaling pathways may affect the oxidation status of DJ-1.

Acknowledgments We thank Yoko Misawa and Kiyomi Takaya for their technical assistance. We also thank Kazuko Takahashi-Niki for preparation of rat primary neurons. This work was supported by Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology and by the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBIO) in Japan.

REFERENCES

- 1) Salisbury S. A., Forrest H. S., Cruse W. B., Kennard O., *Nature* (London), **280**, 843–844 (1979).
- 2) Gallop P. M., Paz M. A., Flückiger R., Kagan H. M., *Trends Biochem. Sci.*, **14**, 343–346 (1989).
- 3) Goodwin P. M., Anthony C., *Adv. Microb. Physiol.*, **40**, 1–80 (1998).
- 4) Smidt C. R., Steinberg F. M., Rucker R. B., *Proc. Soc. Exp. Biol. Med.*, **197**, 19–26 (1991).
- 5) Bishop A., Gallop P. M., Karnovsky M. L., *Nutr. Rev.*, **56**, 287–293 (1998).
- 6) McIntire W. S., *Annu. Rev. Nutr.*, **18**, 145–177 (1998).
- 7) Misra H. S., Khairnar N. P., Barik A., Indira Priyadarini K., Mohan H., Apte S. K., *FEBS Lett.*, **578**, 26–30 (2004).
- 8) Jensen F. E., Gardner G. J., Williams A. P., Gallop P. M., Aizenman E., Rosenberg P. A., *Neuroscience*, **62**, 399–406 (1994).
- 9) Zhang Y., Rosenberg P. A., *Eur. J. Neurosci.*, **16**, 1015–1024 (2002).
- 10) He K., Nukada H., Urakami T., Murphy M. P., *Biochem. Pharmacol.*, **65**, 67–74 (2003).
- 11) Zhang Y., Feustel P. J., Kimelberg H. K., *Brain Res.*, **1094**, 200–206 (2006).
- 12) Hara H., Hiramoto H., Adachi T., *Neurochem. Res.*, **32**, 489–495 (2007).
- 13) Tao R., Karliner J. S., Simonis U., Zheng J., Zhang J., Honbo N., Alano C. C., *Biochem. Biophys. Res. Commun.*, **363**, 257–262 (2007).
- 14) Ohwada K., Takeda H., Yamazaki M., Isogai H., Nakano M., Shimomura M., Fukui K., Urano S., *J. Clin. Biochem. Nutr.*, **42**, 29–34 (2008).
- 15) Kobayashi M., Kim J., Kobayashi N., Han S., Nakamura C., Ikebukuro K., Sode K., *Biochem. Biophys. Res. Commun.*, **349**, 1139–1144 (2006).
- 16) Nagakubo D., Taira T., Kitaura H., Ikeda M., Tamai K., Iguchi-Ariga S. M. M., Ariga H., *Biochem. Biophys. Res. Commun.*, **31**, 509–513 (1997).
- 17) Bonifati V., Rizzo P., van Baren M. J., Schaap O., Breedveld G. J., Krieger E., Dekker M. C., Squitieri F., Ibanez P., Joosse M., van Dongen J. W., Vanacore N., van Swieten J. C., Brice A., Meco G., van

- Duijn C. M., Oostra B. A., Heutink P., *Science*, **299**, 256—259 (2003).
- 18) Taira T., Saito Y., Niki T., Iguchi-Ariga S. M. M., Takahashi K., Ariga H., *EMBO Rep.*, **5**, 213—218 (2004).
 - 19) Canet-Aviles R. M., Wilson M. A., Miller D. W., Ahmad R., McLendon C., Bandyopadhyay S., Baptista M. J., Ringe D., Petsko G. A., Cookson M. R., *Proc. Natl. Acad. Sci. U.S.A.*, **101**, 9103—9108 (2004).
 - 20) Martinat C., Shendelman S., Jonason A., Leete T., Beal M. F., Yang L., Floss T., Abeliovich A., *PLoS Biol.*, **2**, e327 (2004).
 - 21) Yokota T., Sugawara K., Ito K., Takahashi R., Ariga H., Mizusawa H., *Biochem. Biophys. Res. Commun.*, **312**, 1342—1348 (2003).
 - 22) Takahashi K., Taira T., Niki T., Seino C., Iguchi-Ariga S. M. M., Ariga H., *J. Biol. Chem.*, **276**, 37556—37563 (2001).
 - 23) Niki T., Takahashi-Niki K., Taira T., Iguchi-Ariga S. M. M., Ariga H., *Mol. Cancer Res.*, **1**, 247—261 (2003).
 - 24) Shinbo Y., Taira T., Niki T., Iguchi-Ariga S. M. M., Ariga H., *Int. J. Oncol.*, **26**, 641—648 (2005).
 - 25) Xu J., Zhong N., Wang H., Elias J. E., Kim C. Y., Woldman I., Pifl C., Gygi S. P., Geula C., Yankner B. A., *Hum. Mol. Genet.*, **14**, 1231—1241 (2005).
 - 26) Zhong N., Kim C. Y., Rizzu P., Geula C., Porter D. R., Pothos E. N., Squitieri F., Heutink P., Xu J., *J. Biol. Chem.*, **281**, 20940—20948 (2006).
 - 27) Shendelman S., Jonason A., Martinat C., Leete T., Abeliovich A., *PLoS Biol.*, **2**, e362 (2004).
 - 28) Goldberg M. S., Pisani A., Haburcak M., Vortherms T. A., Kitada T., Costa C., Tong Y., Martella G., Tschertner A., Martins A., Bernardi G., Roth B. L., Pothos E. N., Calabresi P., Shen J., *Neuron*, **45**, 489—496 (2005).
 - 29) Kim R. H., Smith P. D., Aleyasin H., Hayley S., Mount M. P., Pownall S., Wakeham A., You-Ten A. J., Kalia S. K., Horne P., Westaway D., Lozano A. M., Anisman H., Park D. S., Mak T. W., *Proc. Natl. Acad. Sci. U.S.A.*, **102**, 5215—5220 (2005).
 - 30) Chen L., Cagniard B., Mathews T., Jones S., Koh H. C., Ding Y., Carvey P.M., Ling Z., Kang U. J., Zhuang X., *J. Biol. Chem.*, **280**, 21418—21426 (2005).
 - 31) Manning-Bo A. B., Caudle W. M., Perez X. A., Reaney S. H., Paletzki R., Isla M. Z., Chou V. P., McCormack A. L., Miller G. W., Langston J. W., Gerfen C. R., Dimonte D. A., *Neurobiol. Dis.*, **27**, 141—150 (2007).
 - 32) Andres-Mateos E., Perier C., Zhang L., Blanchard-Fillion B., Greco T. M., Thomas B., Ko H. S., Sasaki M., Ischiropoulos H., Przedborski S., Dawson T. M., Dawson V. L., *Proc. Natl. Acad. Sci. U.S.A.*, **104**, 14807—14812 (2007).
 - 33) Chandran J. S., Lin X., Zapata A., Höke A., Shimoji M., Moore S. O., Galloway M. P., Laird F. M., Wong P. C., Price D. L., Bailey K. R., Crawley J. N., Shippenberg T., Cai H., *Neurobiol. Dis.*, **29**, 505—514 (2008).
 - 34) Inden M., Taira T., Kitamura Y., Yanagida T., Tsuchiya D., Takata K., Yanagisawa D., Nishimura K., Taniguchi T., Kiso Y., Yoshimoto K., Agatsuma T., Koide-Yoshida S., Iguchi-Ariga S. M. M., Shimohama S., Ariga H., *Neurobiol. Dis.*, **24**, 144—158 (2006).
 - 35) Ooe H., Taira T., Iguchi-Ariga S. M. M., Ariga H., *Toxicol. Sci.*, **88**, 114—126 (2005).
 - 36) Kinumi T., Kimata J., Taira T., Ariga H., Niki E., *Biochem. Biophys. Res. Commun.*, **317**, 722—728 (2004).
 - 37) Takahashi-Niki K., Niki T., Taira T., Iguchi-Ariga S. M. M., Ariga H., *Biochem. Biophys. Res. Commun.*, **320**, 389—397 (2004).
 - 38) Zhou W., Zhu M., Wilson M. A., Petsko G. A., Fink A. L., *J. Mol. Biol.*, **356**, 1036—1048 (2006).
 - 39) Bandopadhyay R., Kingsbury A. E., Cookson M. R., Reid A. R., Evans I. M., Hope A. D., Pittman A. M., Lashley T., Canet-Aviles R., Miller D. W., McLendon C., Strand C., Leonard A. J., Abou-Sleiman P. M., Healy D. G., Ariga H., Wood N. W., de Silva R., Revesz T., Hardy J. A., Lees A. J., *Brain*, **127**, 420—430 (2004).
 - 40) Choi J., Sullards M. C., Olzmann J. A., Rees H. D., Weintraub S. T., Bostwick D. E., Gearing M., Levey A. I., Chin L.-S., Li L., *J. Biol. Chem.*, **281**, 10816—10824 (2006).
 - 41) Sawada H., Kohno R., Kihara T., Izumi Y., Sakka N., Ibi M., Nakanishi M., Nakamizo T., Yamakawa K., Shibasaki H., Yamamoto N., Akaike A., Inden M., Kitamura Y., Taniguchi T., Shimohama S., *J. Biol. Chem.*, **279**, 10710—10719 (2004).
 - 42) Miyazaki S., Yanagida T., Nunome K., Ishikawa S., Inden M., Kitamura K., Nakagawa S., Taira T., Hirota K., Niwa M., Iguchi-Ariga S. M. M., Ariga H., *J. Neurochem.*, **105**, 2418—2434 (2008).
 - 43) Kumazawa T., Hiwasa T., Takiguchi M., Suzuki O., Sato K., *Int. J. Mol. Med.*, **19**, 765—770 (2007).