



Title	Mitochondrial activity during pre-maturational culture in in vitro-grown bovine oocytes is related to maturational and developmental competences.
Author(s)	Huang, Weiping; Kang, Sung-Sik; Nagai, Katsuhisa; Yanagawa, Yojiro; Takahashi, Yoshiyuki; Nagano, Masashi
Citation	Reproduction, fertility, and development, 28(3), 349-356 https://doi.org/10.1071/RD14023
Issue Date	2014-07-17
Doc URL	http://hdl.handle.net/2115/59542
Rights	Disclaimer. This is not the definitive version of record of this article. This manuscript has been accepted for publication in Reproduction, but the version presented here has not yet been copy edited, formatted or proofed. Consequently, the journal accepts no responsibility for any errors or omissions it may contain. The definitive version is available at http://dx.doi.org/10.1071/RD14023 © 2014 Society for Reproduction and Fertility.
Type	article (author version)
File Information	Huang et al. RFD.pdf



[Instructions for use](#)

1 Full paper

2

3 Title: Mitochondrial activity during pre-maturational culture in *in vitro*-grown bovine oocytes
4 related to maturational and developmental competences

5

6 Authors: Weiping HUANG, Sung-Sik KANG, Katsuhisa NAGAI, Yojiro YANAGAWA, Yoshiyuki
7 TAKAHASHI, Masashi NAGANO*

8

9 Institution: Laboratory of Theriogenology, Department of Veterinary Clinical Sciences,
10 Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

11

12

13 Running head: Mitochondrial activity and IVG oocyte development

14

15 *Corresponding author: M. Nagano

16 Laboratory of Theriogenology, Department of Veterinary Clinical Sciences, Graduate School of
17 Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

18 Tel.: +81-11-706-5232, Fax: +81-11-706-5232

19 e-mail: mnaga@vetmed.hokudai.ac.jp

20

21 **Abstract**

22 The objective of this study was to investigate the dynamics of mitochondrial activity in *in*
23 *vitro*-grown (IVG) bovine oocytes during pre-maturational culture (pre-IVM) and its relationship
24 to their developmental competence upon being subjected to different pre-IVM durations. After
25 12-day IVG culture, oocytes were cultured for 0, 10 or 20 h with 3-isobutyl-1-methylxanthine
26 (IBMX) as pre-IVM. Mitochondrial activity in IVG oocytes after 10 h pre-IVM was the highest
27 among all the pre-IVM durations ($P < 0.05$). In addition, cleavage (79.4%) and blastocyst rates
28 (38.9%) of embryos derived from IVG oocytes with 10 h pre-IVM were higher than those with
29 20 h pre-IVM (63.0 and 25.8%, respectively; $P < 0.05$) and similar to those of *in vivo*-grown
30 oocytes (82.7 and 36.7%, respectively). To confirm the developmental ability of IVG oocytes
31 with 10 h pre-IVM beyond the blastocyst stage *in vivo*, embryo transfer was attempted.
32 Transferred embryos developed to the elongated embryonic stage (63.6%, 7/11) in the
33 recipient uterus at Day 16 of oestrus, and a male calf was delivered (50%, 1/2). In conclusion,
34 it was indicated that the mitochondrial activity of bovine IVG oocytes peaked at 10 h pre-IVM
35 and was closely correlated with the nuclear maturation and developmental competences of
36 IVG oocytes

37

38

39 **Additional Keywords:** early antral follicle, embryo transfer, IVG, mitochondria, ROS

40

41 **Introduction**

42 The development to blastocysts of bovine oocytes derived from in vitro-growth (IVG)
43 culture was found to be markedly lower than that of in vivo-grown oocytes (Hirao et al. 2004).
44 In our previous studies, we showed that the maturational and developmental competences of
45 IVG oocytes were improved by shortening the IVG duration from 14 to 12 days (Huang et al.
46 2013, 2014) and by performing pre-maturational culture (pre-IVM) for 20 h in medium
47 supplemented with phosphodiesterase (PDE) inhibitor (3-isobutyl-1-methylxanthine (IBMX))
48 between IVG and in vitro maturational (IVM) cultures (Huang et al. 2013, 2014). In one
49 previous study (Huang et al. 2014), the normal fertilisation rates of IVG oocytes were similar
50 (around 70%) regardless of pre-IVM, despite the low nuclear maturation rate of IVG oocytes
51 without pre-IVM. Therefore, the authors speculated that IVG oocytes without pre-IVM achieved
52 nuclear maturation during the culture for fertilisation, and that the improvement of maturational
53 and developmental competences of IVG oocytes with pre-IVM is probably caused by
54 synchronisation of meiotic progression because most IVG oocytes may be fertilised at an
55 appropriate time for embryonic development. On the other hand, it was reported that in
56 vivo-grown bovine oocytes with 18 to 21 h pre-IVM showed higher developmental competence
57 to blastocysts than those without pre-IVM, regardless of the timing of fertilisation (Hashimoto
58 et al. 2002). Hashimoto et al. (2002) concluded that immature bovine oocytes required a
59 certain time to acquire developmental competence during meiotic arrest. For this reason, we
60 speculate that cytoplasmic changes of oocytes occur during pre-IVM and this contributes to
61 the acquisition of developmental competence. Therefore, it is necessary to investigate the
62 cytoplasmic changes in IVG oocytes during pre-IVM culture.

63 Mitochondria play important roles in oocyte maturation and subsequent embryonic

64 development by being a steady source of adenosine triphosphate (ATP) through oxidative
65 phosphorylation (Nagano et al. 2006; Tarazona et al. 2006; Van Blerkom 2011). In mammalian
66 oocytes, mitochondrial activity increased with follicular development (Kanaya et al. 2007;
67 Machatkova et al. 2012) and the higher mitochondrial activity before the resumption of meiosis
68 was associated with higher competences in nuclear maturation (Egerszegi et al. 2010;
69 Machatkova et al. 2012) and development to blastocysts (Kanaya et al. 2007). On the other
70 hand, it was reported that reactive oxygen species (ROS) such as hydroxyl radical and
71 hydrogen peroxide were generated by the mitochondrial respiratory chain (Fleury et al. 2002)
72 and ROS suppressed the developmental competence of oocytes (Hashimoto et al. 2000). For
73 this reason, there is a possibility that mitochondrial activity and ROS production change during
74 pre-IVM culture and affect the nuclear maturation and embryonic development of IVG oocytes.

75 In the present study, in order to clarify the cause of the improvement of maturational and
76 developmental competences of IVG oocytes by pre-IVM, we examined mitochondrial activity
77 and ROS production during pre-IVM culture and the competences in nuclear maturation and
78 development to blastocysts of IVG oocytes with different pre-IVM durations. Moreover, to
79 confirm the developmental ability of IVG oocytes beyond the blastocyst stage, we also
80 attempted to transfer the embryos derived from IVG oocytes into recipient cows.

81

82 **Materials and Methods**

83 **Chemicals**

84 All the chemicals used for this study were purchased from Sigma-Aldrich (St. Louis, MO,
85 USA), unless otherwise stated.

86

87 **Collection of early antral follicles and IVG culture of oocyte-granulosa cell complexes**

88 Bovine ovaries obtained at a local abattoir were kept in plastic bags at 20°C and were
89 transported to the laboratory within 6 to 10 h of their collection, as described previously (Huang
90 et al. 2013, 2014). The oocyte–cumulus–granulosa complexes (OCGCs) with normal
91 appearance were cultured individually in a 96-well culture plate (Becton, Dickinson and Co.,
92 Franklin Lakes, NJ, USA) with 200 µL of the growth medium for 12 days at 39°C in humidified
93 air with 5% CO₂, as described previously (Huang et al. 2013, 2014). The growth medium was
94 HEPES-buffered TCM-199 (Invitrogen, Grand Island, NY, USA) supplemented with 0.91 mM
95 sodium pyruvate, 1 µg mL⁻¹ oestradiol-17β, 5% fetal calf serum (FCS; Invitrogen), 4 mM
96 hypoxanthine, 4% polyvinylpyrrolidone (MW 36000) and 50 µg mL⁻¹ gentamicin sulfate.
97 Hypoxanthine was added to prevent premature meiotic resumption as described previously
98 (Eppig and Downs 1987; Harada et al. 1997; Hirao et al. 2004). Every 4 days of IVG culture,
99 half (100 µL) of the growth medium was replaced with the same amount of fresh medium.

100

101 **Pre-IVM and IVM of IVG oocytes**

102 After 12-day IVG culture, oocytes surrounded by several layers of cumulus cells were
103 collected from morphologically normal OCGCs and submitted to IVM with or without pre-IVM,
104 as described previously (Huang et al. 2013, 2014). Briefly, pre-IVM culture of IVG oocytes was
105 performed as they were cultured individually in each well of micro-well plates (NUNC, Roskilde,
106 Denmark) filled with 6 mL of HEPES-buffered TCM-199 supplemented with 0.2 mM sodium
107 pyruvate, 2×10^{-6} units mL⁻¹ follicle-stimulating hormone (FSH, from porcine pituitary), 0.5
108 mM IBMX, 1 µg mL⁻¹ oestradiol-17β, 10% FCS and 50 µg mL⁻¹ gentamicin sulfate at 39°C

109 under 5% CO₂ in air for 0, 10 or 20 h. For IVM, oocytes were cultured individually in each well
110 of micro-well plates filled with 6 mL of IVM medium, which was HEPES-buffered TCM-199
111 supplemented with 0.2 mM sodium pyruvate, 0.02 units mL⁻¹ FSH, 1 µg mL⁻¹ oestradiol-17β,
112 10% FCS and 50 µg mL⁻¹ gentamicin sulfate, at 39°C under 5% CO₂ in air for 22 h (Nagano
113 et al. 2013).

114 Oocytes collected from antral follicles of 2 to 8 mm in diameter (in vivo-grown oocytes)
115 served as controls. IVM of in vivo-grown oocytes was conducted as described previously
116 (Takahashi et al. 1996). In brief, cumulus–oocyte complexes (COCs) were incubated in
117 droplets of IVM medium that was the same as used for IVG oocytes (~10 COCs per 50 µL)
118 covered with paraffin oil at 39°C under 5% CO₂ in air for 22 h.

119

120 **IVF and IVC of inseminated oocytes**

121 IVF (Day 0) was performed using frozen semen from a Holstein bull according to a
122 procedure described previously (Takahashi and Kanagawa 1998) with slight modification. In
123 brief, motile spermatozoa (5 × 10⁶ spermatozoa mL⁻¹) separated from thawed semen using a
124 Percoll gradient (45 and 90%) were co-incubated with COCs in a 100-µL droplet (~10 COCs
125 per droplet) of modified Brackett and Oliphant isotonic medium (Brackett and Oliphant 1975)
126 containing 3 mg mL⁻¹ fatty-acid-free bovine serum albumin (BSA) and 2.5 mM theophylline
127 (Takahashi and First 1992) for 18 h at 39°C in a humidified atmosphere of 5% CO₂, 5% O₂
128 and 90% N₂.

129 IVC of inseminated oocytes (presumptive zygotes) was performed as described
130 previously (Takahashi and Kanagawa 1998). Briefly, after co-incubation with spermatozoa,

131 presumptive zygotes were freed from cumulus cells by vortexing and washing three times in
132 the culture medium. Cumulus-free zygotes were cultured for ~150 h in a 30- μ L droplet of the
133 culture medium at 39°C under 5% CO₂, 5% O₂ and 90% N₂. The culture medium was a
134 modified synthetic oviduct fluid containing 12 essential amino acids for basal medium Eagle, 7
135 non-essential amino acids for minimum essential medium, and further supplementation with 1
136 mM glutamine, 5 mM glycine, 5 mM taurine, 1 mM glucose, 10 μ g mL⁻¹ insulin and 3 mg
137 mL⁻¹ fatty-acid-free BSA instead of polyvinyl alcohol (Takahashi and Kanagawa 1998). After
138 30 h (Day 2) and ~150 h (Day 7) of IVC, cleavage and development of presumptive zygotes to
139 the blastocyst stage were assessed, respectively. Total cell numbers of blastocysts obtained
140 after ~150 h of IVC were counted using an air-drying method (Takahashi and First 1992).

141

142 **Evaluation of oocyte nuclear status**

143 After pre-IVM and IVM, oocytes were denuded from cumulus cells by vortexing. Denuded
144 oocytes were stained with 1% aceto-orcein and examined for their nuclear status as described
145 elsewhere (Nagano et al. 2006). Nuclear statuses were divided into germinal vesicle (GV),
146 germinal vesicle breakdown (GVBD), chromosomal condensation (CC), metaphase I (MI),
147 anaphase I–telophase I (AI–TI) and metaphase II (MII) by observation under a phase-contrast
148 microscope.

149

150 **Evaluation of mitochondrial activity in IVG oocytes during pre-IVM culture**

151 IVG oocytes after pre-IVM and IVM were treated with 500 IU mL⁻¹ hyaluronidase in
152 Dulbecco's phosphate-buffered saline (DPBS) for 10 min and then the oocytes were denuded

153 from cumulus cells by repeat pipetting with a fine pipette. Denuded oocytes were incubated for
154 15 min in the dark at 37°C in DPBS supplemented with 1 μM
155 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolyl carbocyanine iodide (JC-1), 1 μg
156 mL⁻¹ Hoechst 33342 and 10% FCS, as described previously (Ge et al. 2012) with slight
157 modification. Images of oocytes were acquired using a digital fluorescence microscope
158 (BZ-9000; Keyence, Osaka, Japan) and the mean fluorescence intensity of the images was
159 calculated using analysis software (BZ-H2A; Keyence). Nuclear status of IVG oocytes was
160 evaluated by blue fluorescence of Hoechst 33342 and oocytes having GV before IVM and
161 metaphase plate with polar body after IVM were subjected to the evaluation of mitochondrial
162 activity. Membrane potentials of mitochondria in GV and MII oocytes were calculated as the
163 ratio of fluorescence intensity of activated mitochondria, expressed as red fluorescence of JC-1
164 staining, to less-activated mitochondria, expressed as green fluorescence of JC-1 staining
165 ($\Delta\psi_m$, red/green fluorescence intensity).

166

167 **Assessment of ROS generation in IVG oocytes during pre-IVM culture**

168 IVG oocytes after pre-IVM were denuded from cumulus cells by vortexing. Denuded
169 oocytes were incubated for 15 min in the dark at 37°C in DPBS supplemented with 10 μM
170 2',7'-dichlorodihydrofluorescein diacetate (DCHFDA) and 10% FCS, as described previously
171 (Hashimoto et al. 2000) with slight modification. Images of oocytes were acquired using a
172 digital fluorescence microscope and the mean fluorescence intensity of the images was
173 calculated using analysis software. ROS in oocytes were defined as the mean fluorescence
174 intensity of oocytes.

175

176 **Experimental design**

177 **Experiment 1**

178 After 0, 10 and 20 h pre-IVM, a total of 83 IVG oocytes derived from 129 OCGCs were
179 used for the evaluation of mitochondrial activity in oocytes (six to seven oocytes per replicate).

180 After IVM with 0, 10 and 20 h pre-IVM durations, a total 90 IVG oocytes at MII stage from 263
181 OCGCs were also used for the evaluation of mitochondrial activity (five to twelve oocytes per
182 replicate). In addition, the ROS generation in IVG oocytes was assessed during pre-IVM
183 culture using a total of 61 IVG oocytes derived from 94 OCGCs (six to seven oocytes per
184 replicate).

185

186 **Experiment 2**

187 To examine the relationship between mitochondrial activity and the maturational and
188 developmental competences of IVG oocytes with different pre-IVM durations, a total of 419
189 IVG oocytes derived from 635 OCGCs (10 to 30 oocytes per replicate) were subjected to IVM
190 with 0, 10 or 20 h pre-IVM culture. After 10 and 22 h IVM culture, their nuclear statuses were
191 investigated. Before the examination of nuclear status, some of the IVG oocytes after 22 h IVM
192 (with 0, 10 and 20 h pre-IVM: n = 53, 71 and 37, respectively) were provided for the
193 measurement of their diameter, as described previously (Huang et al. 2013, 2014). To confirm
194 meiotic arrest immediately before IVM culture, nuclear statuses of IVG oocytes after 0, 10 or
195 20 h pre-IVM were evaluated by using a total of 157 IVG oocytes derived from 254 OCGCs (11
196 to 27 oocytes per replicate). Moreover, IVG oocytes with 10 or 20 h pre-IVM were subjected to

197 IVM–IVF–IVC and examined for their cleavage, development to blastocyst stage and cell
198 numbers in blastocysts using a total of 357 IVG oocytes derived from 503 OCGCs (25 to 40
199 oocytes per replicate). In vivo-grown oocytes after IVM served as controls.

200

201 **Experiment 3**

202 For evaluating embryonic development beyond the blastocyst stage, eleven fresh
203 blastocysts (Day 7) obtained after IVC were transferred transcervically into a recipient heifer
204 synchronised at Day 8 of oestrus by EAZI-BREED CIDR (Inter Ag, Te Rapa Road, Hamilton,
205 NZ) synchronisation (Katagiri and Takahashi 2006). Transcervical uterine flushing using a
206 two-way Foley catheter (Fujihira Industry, Tokyo, Japan) was performed at Day 16 of oestrus
207 for embryo collection, and collected embryos were evaluated for their developmental stage. In
208 addition, two blastocysts were also transferred into another heifer synchronised by the same
209 protocol, and her pregnancy was diagnosed at Days 29 and 62 of oestrus by ultrasonography
210 (5 MHz, HS101V; Honda Electronics, Tokyo, Japan).

211

212 **Statistical analysis**

213 All percentage data were subjected to arcsine transformation before statistical analysis.
214 All data were analysed by one-way ANOVA or two-way ANOVA followed by Tukey-Kramer's
215 honestly significant difference test. All statistical analyses were performed using software
216 (JMP Version 10; SAS Institute, Cary, NC, USA).

217

218 **Results**

219 **Experiment 1**

220 There was no significant interaction by two-way ANOVA between pre-IVM duration and
221 with or without IVM culture in terms of mitochondrial activity. Mitochondrial activity in IVG
222 oocytes at 10 h pre-IVM was higher than at 0 and 20 h pre-IVM ($P < 0.05$; Fig. 1). After IVM,
223 the mitochondrial activities of IVG oocytes at MII stage were similar regardless of pre-IVM
224 durations (Fig. 1). Also, great variability of mitochondrial activity among each oocyte type was
225 observed compared with before IVM culture. Mitochondrial activity of IVG oocytes with 0 and
226 20 h pre-IVM was higher after IVM than before IVM ($P < 0.05$), although it was similar in IVG
227 oocytes with 10 h pre-IVM both before and after IVM. ROS generation in IVG oocytes with 0 h
228 pre-IVM was higher than in those with 10 and 20 h pre-IVM ($P < 0.05$; Fig. 2).

229

230 **Experiment 2**

231 After 10 h IVM, the MI rate (82.7%) of IVG oocytes with 10 h pre-IVM tended to be higher
232 than in those with 0 h pre-IVM (48.3%, $P = 0.17$) and their GVBD/CC rate (1.5%) was lower
233 than that of the 20 h pre-IVM group (34.2%, $P < 0.05$) and tended to be lower than the 0 h
234 pre-IVM group (37.3%, $P = 0.08$; Table 1). After 22 h IVM, the MII rate of IVG oocytes with 10 h
235 (91.5%) was higher than in those with 0 h (50.5%, $P < 0.01$) and 20 h pre-IVM (79.0%, $P <$
236 0.05) and similar to that of in vivo-grown oocytes (88.6%; Table 2). The percentage of MI-stage
237 oocytes was significantly lower in IVG oocytes with 10 h pre-IVM (5.5%) than in the 0 h (25.4%,
238 $P < 0.01$) and 20 h pre-IVM groups (19.8%, $P < 0.05$). The mean diameter of IVG oocytes after
239 22 h IVM was $115.2 \pm 3.4 \mu\text{m}$ (range: 105.9 to 122.7 μm , $n = 161$). Before IVM culture, all IVG
240 oocytes with 0 and 10 h pre-IVM were at GV stage (Table 3). The percentages (100%) of

241 GV-stage oocytes after 0 and 10 h pre-IVM were higher than after 20 h pre-IVM (90.0%, P <
242 0.01).

243 As shown in Table 4, cleavage (79.4%) and blastocyst rates (38.9%) of IVG oocytes with
244 10 h pre-IVM were higher than those with 20 h pre-IVM (63.0 and 25.8%, respectively; P <
245 0.05) and similar to those of in vivo-grown oocytes (82.7 and 36.7%, respectively). However,
246 blastocyst rates based on cleaved oocytes were similar regardless of pre-IVM duration. In
247 addition, total cell numbers in blastocysts of IVG oocytes with 10 h pre-IVM were similar to
248 those of the control (P = 0.38).

249

250 **Experiment 3**

251 Seven embryos (63.6%, 7/11) were recovered from recipient heifers at Day 16 of oestrus.
252 Six embryos were elongated, ranging from 40 to 100 mm (70.0 ± 21.9 mm; Fig. 3), and the
253 other one showed retardation in terms of length (30 mm). Another heifer subjected to embryo
254 transfer was confirmed as pregnant by detection of an embryo (50%, 1/2) with a heartbeat on
255 Days 29 and 62 of oestrus. After 276 days of oestrus, the heifer delivered a healthy male
256 offspring with 46 kg bodyweight.

257

258 **Discussion**

259 In the present study, mitochondrial activity in IVG oocytes was high after 10 h pre-IVM,
260 and the oocytes showed higher maturational and developmental competence to blastocyst
261 stage. Furthermore, most of the IVG oocytes (more than 80%) with 10 h pre-IVM and in
262 vivo-grown oocytes reached the M1 stage after 10 h IVM culture, and around 90% of them

263 reached the MII stage after 22 h IVM culture, indicating that the nuclear maturation of IVG
264 oocytes was enhanced and synchronised by 10 h pre-IVM. On the other hand, when the
265 pre-IVM duration was extended to 20 h, IVG oocytes at MII stage were decreased to around
266 80%, and also 10% of IVG oocytes started meiotic resumption spontaneously and some of the
267 IVG oocytes were already degenerated. These results indicate that the extension of pre-IVM to
268 20 h induced the degradation of oocytes or cumulus cells, resulting in the lost of the gap
269 junction between the IVG oocyte and cumulus cells. It was reported that reducing nuclear
270 asynchrony during IVM by temporary meiotic arrest using dibutyryl cyclic AMP (dbcAMP)
271 appeared to enhance the nuclear maturation and developmental competence of porcine
272 oocytes (Somfai et al. 2003). It was also reported that increasing mitochondrial activity and
273 ATP content during pre-IVM by using IBMX and forskolin improved the nuclear maturation and
274 developmental competence of mouse oocytes (Zeng et al. 2013). It is well known that cAMP
275 can be increased in oocytes by cAMP analogues (e.g. dbcAMP), activators of adenylate
276 cyclase (e.g. FSH, forskolin) and PDE inhibitors (e.g. IBMX; Gilchrist and Thompson 2007).
277 The intracellular second messenger, cAMP, plays a significant role in the regulation of
278 mitochondrial activity in mammalian cells (Papa 2006; Carlucci et al. 2008) and further meiosis
279 progression in oocytes (Thomas et al. 2002). It was reported that bovine oocytes derived from
280 larger follicles (>6 mm in diameter) showed higher mitochondrial activity (Machatkova et al.
281 2012) and a higher proportion of blastocyst development (Lonergan et al. 1994) than those
282 from small follicles (<6 mm in diameter). These reports suggest that mitochondrial activity in
283 bovine oocytes increases during follicular development and that higher mitochondrial activity is
284 associated with greater developmental competence of immature oocytes. The present pre-IVM

285 culture system may have mimicked follicular development in terms of the dynamics of
286 mitochondrial activity. By adding IBMX and FSH to pre-IVM medium, the cAMP concentration
287 in IVG oocytes may increase, resulting in the improvement of mitochondrial activity before IVM
288 culture, and may have led to the improved maturational and developmental competences of
289 IVG oocytes with 10 h pre-IVM in the present study.

290 After IVM culture, the mitochondrial activity in IVG oocytes at MII stage became similar
291 regardless of pre-IVM duration, because mitochondrial activities in IVG oocytes with 0 and 20
292 h were enhanced. It was reported that bovine IVM oocytes exhibited higher mitochondrial
293 activity than immature oocytes (Tarazona et al. 2006). This corresponded to our results of IVG
294 oocytes with 0 and 20 h pre-IVM. However, in vivo-matured porcine oocytes showed similar
295 mitochondrial activity to immature oocytes (Romek et al. 2011). This corresponded to our
296 results of IVG oocytes with 10 h pre-IVM. These results may indicate that IVG oocytes with 10
297 h pre-IVM have similar characteristics to in vivo-matured oocytes. In our previous study
298 (Koyama et al. 2014), we showed that the mitochondrial activity and ATP content were
299 increasing in parallel as the duration of IVM culture was extended, and that the developmental
300 competence of bovine oocytes did not correlate with them directly. It was also reported that
301 ATP content in ovum pick up-derived oocytes did not increase during IVM culture, but ATP
302 content in abattoir-derived oocytes did increase (Tamassia et al. 2004). From these results, we
303 speculate that IVG oocytes with 10 h pre-IVM may have similar competence of nuclear
304 maturation and development to blastocysts from in vivo-grown oocytes immediately collected
305 from live cows. On the other hand, IVG oocytes with 0 and 20 h pre-IVM may have not
306 acquired these competences yet and may have already lost them, respectively. In further

307 studies we should confirm the mitochondrial activity of in vivo-matured bovine oocytes.

308 ROS in IVG oocytes with 0 h pre-IVM were higher than in those with 10 and 20 h pre-IVM.

309 A previous study indicated that, when goat granulosa cells were treated with FSH in vitro,

310 catalase activity was stimulated significantly (Behl and Pandey 2002). In cumulus

311 cell-enclosed bovine oocytes, enzymatic antioxidant (superoxide dismutase, glutathione

312 peroxidase and catalase) systems can attenuate the effect of oxidative stress by scavenging

313 ROS (Cetica et al. 2001). In the present study, pre-IVM culture medium supplemented with

314 FSH may directly or indirectly neutralise ROS generation in oocytes. Our previous results

315 showed that the developmental rate to blastocysts (12.7%) without pre-IVM was significantly

316 lower than with 20 h pre-IVM (26.1%), despite the similar normal fertilisation rate (Huang et al.

317 2014). It is speculated that high ROS content in IVG oocytes without pre-IVM may result in a

318 lower developmental rate to blastocysts than with pre-IVM.

319 The mitochondrial activity and blastocyst development of IVG oocytes with 10 h pre-IVM

320 were decreased by extending the pre-IVM duration to 20 h in the present study, although there

321 was no significant difference in ROS content between IVG oocytes with 10 and 20 h pre-IVM.

322 This result indicates that the decreases of mitochondrial activity and developmental

323 competence in IVG oocytes with 20 h pre-IVM were probably not due to ROS production.

324 During apoptosis, mitochondria rapidly lose their transmembrane potential (Ly et al. 2003). It

325 was reported that the pro-apoptotic pathway can be promoted by the cAMP signalling pathway

326 in most cells (Insel et al. 2012) and that granulosa cells stimulated with forskolin and IBMX

327 exhibited a longer-lasting rise in intracellular cAMP, resulting in cell apoptosis (Aharoni et al.

328 1995). It was indicated that the duration of the intracellular rise in cAMP plays an important role

329 in controlling the extent of apoptosis in granulosa cells (Aharoni et al. 1995). Therefore,
330 investigation of cAMP concentration and apoptotic program induction during pre-IVM culture
331 should be conducted in future studies.

332 Following the embryo transfer of blastocysts derived from IVG oocytes with 10 h pre-IVM,
333 six out of seven elongated embryos showed a normal length (40 to 100 mm), similar to the rate
334 of in vivo-derived bovine embryos (Betteridge et al. 1980; Gustafsson and Ploen 1986), and
335 one out of two embryos was maintained in the pregnancy. After 276 days of oestrus, the heifer
336 delivered a male calf with normal bodyweight (46 kg). Large calf syndrome was not observed
337 in the present study although the oocytes were cultured for 3 weeks in vitro. These results
338 suggest that IVG oocytes with 10 h pre-IVM acquired full developmental competence the same
339 as in vivo-grown IVM oocytes and probably in vivo-grown in vivo-matured oocytes.

340 In conclusion, we clarified that the mitochondrial activity of bovine IVG oocytes peaked at
341 10 h pre-IVM culture with IBMX and was closely correlated with the nuclear maturation and
342 developmental competences of IVG oocytes. The IVG oocytes with 10 h pre-IVM can acquire
343 high maturational and developmental competences, the same as in vivo-grown oocytes.
344 However, the extension of pre-IVM duration to 20 h induced oocyte or cumulus cell
345 degradation and reduced the mitochondrial activity before IVM culture and the developmental
346 competence of IVG oocytes.

347

348 **Acknowledgments**

349 This study was financially supported by a Grant-in-Aid for Scientific Research (No.
350 25450441) from the Japan Society for the Promotion of Science to M. Nagano. Weiping

351 Huang was sponsored by the China Scholarship Council. We thank Genetics Hokkaido
352 Association for the donation of frozen bull sperm.

353

354 **References**

355

356 Aharoni, D., Dantes, A., Oren, M., and Amsterdam, A. (1995) Camp-mediated signals as
357 determinants for apoptosis in primary granulosa-cells. *Exp. Cell Res.* **218**, 271-282.

358

359 Behl, R., and Pandey, R.S. (2002) FSH induced stimulation of catalase activity in goat
360 granulosa cells *in vitro*. *Anim. Reprod. Sci.* **70**, 215-221.

361

362 Betteridge, K.J., Eaglesome, M.D., Randall, G.C., and Mitchell, D. (1980) Collection,
363 description and transfer of embryos from cattle 10-16 days after oestrus. *J. Reprod. Fertil.* **59**,
364 205-216.

365

366 Brackett, B.G., and Oliphant, G. (1975) Capacitation of rabbit spermatozoa *in vitro*. *Biol.*
367 *Reprod.* **12**, 260-274.

368

369 Carlucci, A., Lignitto, L., and Feliciello, A. (2008) Control of mitochondria dynamics and
370 oxidative metabolism by cAMP, AKAPs and the proteasome. *Trends Cell Biol.* **18**, 604-613.

371

372 Cetica, P.D., Pintos, L.N., Dalvit, G.C., and Beconi, M.T. (2001) Antioxidant enzyme activity
373 and oxidative stress in bovine oocyte *in vitro* maturation. *IUBMB Life* **51**, 57-64.

374

375 Egerszegi, I., Alm, H., Ratky, J., Heleil, B., Brussow, K.P., and Torner, H. (2010) Meiotic
376 progression, mitochondrial features and fertilisation characteristics of porcine oocytes with
377 different G6PDH activities. *Reprod. Fertil. Dev.* **22**, 830-838.

378

379 Eppig, J.J., and Downs, S.M. (1987) The effect of hypoxanthine on mouse oocyte growth and
380 development *in vitro*: maintenance of meiotic arrest and gonadotropin-induced oocyte
381 maturation. *Dev. Biol.* **119**, 313-321.

382

383 Fleury, C., Mignotte, B., and Vayssiere, J.L. (2002) Mitochondrial reactive oxygen species in
384 cell death signaling. *Biochimie* **84**, 131-141.

385

386 Ge, H., Tollner, T.L., Hu, Z., Dai, M., Li, X., Guan, H., Shan, D., Zhang, X., Lv, J., Huang, C.,
387 and Dong, Q. (2012) The importance of mitochondrial metabolic activity and mitochondrial
388 DNA replication during oocyte maturation *in vitro* on oocyte quality and subsequent embryo
389 developmental competence. *Mol. Reprod. Dev.* **79**, 392-401.

390

391 Gilchrist, R.B., and Thompson, J.G. (2007) Oocyte maturation: emerging concepts and
392 technologies to improve developmental potential *in vitro*. *Theriogenology* **67**, 6-15.
393

394 Gustafsson, H., and Ploen, L. (1986) The morphology of 16 and 17 day old bovine blastocysts
395 from virgin and repeat breeder heifers. *Anat. Histol. Embryol.* **15**, 277-287.
396

397 Harada, M., Miyano, T., Matsumura, K., Osaki, S., Miyake, M., and Kato, S. (1997) Bovine
398 oocytes from early antral follicles grow to meiotic competence *in vitro*: effect of FSH and
399 hypoxanthine. *Theriogenology* **48**, 743-755.
400

401 Hashimoto, S., Minami, N., Takakura, R., and Imai, H. (2002) Bovine immature oocytes
402 acquire developmental competence during meiotic arrest *in vitro*. *Biol. Reprod.* **66**, 1696-1701.
403

404 Hashimoto, S., Minami, N., Yamada, M., and Imai, H. (2000) Excessive concentration of
405 glucose during *in vitro* maturation impairs the developmental competence of bovine oocytes
406 after *in vitro* fertilization: relevance to intracellular reactive oxygen species and glutathione
407 contents. *Mol. Reprod. Dev.* **56**, 520-526.
408

409 Hirao, Y., Itoh, T., Shimizu, M., Iga, K., Aoyagi, K., Kobayashi, M., Kacchi, M., Hoshi, H., and
410 Takenouchi, N. (2004) *In vitro* growth and development of bovine oocyte-granulosa cell
411 complexes on the flat substratum: effects of high polyvinylpyrrolidone concentration in culture
412 medium. *Biol. Reprod.* **70**, 83-91.
413

414 Huang, W., Nagano, M., Kang, S.S., Yanagawa, Y., and Takahashi, Y. (2013a) Effects of *in*
415 *vitro* growth culture duration and prematuration culture on maturational and developmental
416 competences of bovine oocytes derived from early antral follicles. *Theriogenology* **80**,
417 793-799.
418

419 Huang, W., Nagano, M., Kang, S.S., Yanagawa, Y., and Takahashi, Y. (2013b) Prematurational
420 culture with IBMX synchronizes meiotic progression of the germinal vesicle stage and
421 improves nuclear maturation and embryonic development in *in vitro*-grown bovine oocytes. *J.*
422 *Reprod. Dev.* DOI: 10.1262/jrd.2013-082
423

424 Insel, P.A., Zhang, L., Murray, F., Yokouchi, H., and Zambon, A.C. (2012) Cyclic AMP is both a
425 pro-apoptotic and anti-apoptotic second messenger. *Acta Physiologica* **204**, 277-287.
426

427 Kanaya, H., Hashimoto, S., Teramura, T., Morimoto, Y., Matsumoto, K., Saeki, K., Iritani, A.,
428 and Hosoi, Y. (2007) Mitochondrial dysfunction of *in vitro* grown rabbit oocytes results in
429 preimplantation embryo arrest after activation. *J. Reprod. Dev.* **53**, 631-637.
430

431 Katagiri, S., and Takahashi, Y. (2006) Potential relationship between normalization of
432 endometrial epidermal growth factor profile and restoration of fertility in repeat breeder cows.
433 *Anim. Reprod. Sci.* **95**, 54-66.
434

435 Koyama, K., Kang, S.S., Huang, W., Yanagawa, Y., Takahashi, Y., and Nagano, M. (2014)
436 Aging-related changes in *in vitro*-matured bovine oocytes: oxidative stress, mitochondrial
437 activity and ATP content after nuclear maturation. *J. Reprod. Dev.* **60**, 136-142.
438

439 Lonergan, P., Monaghan, P., Rizos, D., Boland, M.P., and Gordon, I. (1994) Effect of follicle
440 size on bovine oocyte quality and developmental competence following maturation, fertilization,
441 and culture *in vitro*. *Mol. Reprod. Dev.* **37**, 48-53.
442

443 Ly, J.D., Grubb, D.R., and Lawen, A. (2003) The mitochondrial membrane potential ($\Delta\psi_m$) in
444 apoptosis; an update. *Apoptosis* **8**, 115-128.
445

446 Machatkova, M., Jeseta, M., Hulinska, P., Knitlova, D., Nemcova, L., and Kanka, J. (2012)
447 Characteristics of bovine oocytes with different meiotic competence in terms of their
448 mitochondrial status and expression of nuclear-encoded factors. *Reprod. Domest. Anim.* **47**,
449 806-814.
450

451 Nagano, M., Kang, S.S., Koyama, K., Huang, W., Yanagawa, Y., and Takahashi, Y. (2013) *In*
452 *vitro* maturation system for individual culture of bovine oocytes using micro-volume multi-well
453 plate. *Jpn. J. Vet. Res.* **61**, 149-154.
454

455 Nagano, M., Katagiri, S., and Takahashi, Y. (2006) ATP content and
456 maturational/developmental ability of bovine oocytes with various cytoplasmic morphologies.
457 *Zygote* **14**, 299-304.
458

459 Papa, S. (2006) Does cAMP play a part in the regulation of the mitochondrial electron
460 transport chain in mammalian cells? *IUBMB Life* **58**, 173-175.
461

462 Romek, M., Gajda, B., Rolka, M., and Smoraż, Z. (2011) Mitochondrial activity and morphology
463 in developing porcine oocytes and pre-implantation non-cultured and cultured embryos.
464 *Reprod. Domest. Anim.* **46**, 471-480.
465

466 Somfai, T., Kikuchi, K., Onishi, A., Iwamoto, M., Fuchimoto, D., Papp, A.B., Sato, E., and Nagai,
467 T. (2003) Meiotic arrest maintained by cAMP during the initiation of maturation enhances
468 meiotic potential and developmental competence and reduces polyspermy of IVM/IVF porcine
469 oocytes. *Zygote* **11**, 199-206.
470

471 Takahashi, Y., and First, N.L. (1992) *In vitro* development of bovine one-cell embryos:
472 Influence of glucose, lactate, pyruvate, amino acids and vitamins. *Theriogenology* **37**,
473 963-978.
474

475 Takahashi, Y., Hishinuma, M., Matsui, M., Tanaka, H., and Kanagawa, H. (1996) Development
476 of *in vitro* matured/fertilized bovine embryos in a chemically defined medium: influence of
477 oxygen concentration in the gas atmosphere. *J. Vet. Med. Sci.* **58**, 897-902.
478

479 Takahashi, Y., and Kanagawa, H. (1998) Effect of oxygen concentration in the gas atmosphere
480 during *in vitro* insemination of bovine oocytes on the subsequent embryonic development *in*
481 *vitro*. *J. Vet. Med. Sci.* **60**, 365-367.
482

483 Tamassia, M., Nuttinck, F., May-Panloup, P., Reynier, P., Heyman, Y., Charpigny, G., Stojkovic,
484 M., Hiendleder, S., Renard, J.P., and Chastant-Maillard, S. (2004) *In vitro* embryo production
485 efficiency in cattle and its association with oocyte adenosine triphosphate content, quantity of
486 mitochondrial DNA, and mitochondrial DNA haplogroup. *Biol. Reprod.* **71**, 697-704.
487

488 Tarazona, A.M., Rodriguez, J.I., Restrepo, L.F., and Olivera-Angel, M. (2006) Mitochondrial
489 activity, distribution and segregation in bovine oocytes and in embryos produced *in vitro*.
490 *Reprod. Domest. Anim.* **41**, 5-11.
491

492 Thomas, R.E., Armstrong, D.T., and Gilchrist, R.B. (2002) Differential effects of specific
493 phosphodiesterase isoenzyme inhibitors on bovine oocyte meiotic maturation. *Dev. Biol.* **244**,
494 215-225.
495

496 Van Blerkom, J. (2011) Mitochondrial function in the human oocyte and embryo and their role
497 in developmental competence. *Mitochondrion* **11**, 797-813.
498

499 Zeng, H.T., Ren, Z., Guzman, L., Wang, X., Sutton-McDowall, M.L., Ritter, L.J., De Vos, M.,
500 Smitz, J., Thompson, J.G., and Gilchrist, R.B. (2013) Heparin and cAMP modulators interact
501 during pre-*in vitro* maturation to affect mouse and human oocyte meiosis and developmental
502 competence. *Hum. Reprod.* **28**, 1536-1545.
503
504
505
506

507 **Figure Legends**

508 **Fig. 1.** Mitochondrial activity ($\Delta\psi_m$) of IVG oocytes with different pre-IVM durations before and
509 after 22 h IVM culture.

510 The ratio of red to green fluorescence was quantified as an indicator of mitochondrial activity.

511 Lines of the boxes delineate the 25th, 50th and 75th percentiles and the whiskers depict the
512 10th and 90th percentiles of a population. The circles represent the outliers.

513 ^{a, b} Different characters indicate a significant difference between different pre-IVM duration ($P <$
514 0.05).

515 * Asterisk indicates a significant difference between same pre-IVM duration before IVM and
516 after IVM ($P < 0.05$).

517 Numbers in parentheses are the numbers of IVG oocytes used (5 replicates before IVM and 3
518 replicates after IVM).

519

520 **Fig. 2.** Reactive oxygen species (ROS) in IVG oocytes at different pre-IVM durations.

521 Lines of the boxes delineate the 25th, 50th and 75th percentiles and the whiskers depict the
522 10th and 90th percentiles of a population. The circles represent the outliers.

523 ^{a, b} Different characters indicate a significant difference ($P < 0.05$).

524 Numbers in parentheses are the numbers of IVG oocytes used (3 replicates).

525

526 **Fig. 3.** The embryo (length 100 mm) recovered from cattle at 16 days of estrus was elongated.

527 Bar represents 5 mm.

528

Fig. 1.




Fig. 2.





Fig. 3.



1 **Table 1. Effect of pre-IVM duration on meiotic resumption of IVG oocytes after 10 h IVM**

2 ^{a,b} Values (mean ± SD) with different superscripts within columns are significantly different (P <
3 0.05).

4 GV, germinal vesicle; GVBD/CC, germinal vesicle breakdown/chromosomal condensation; M I,
5 metaphase I; M II, metaphase II; Deg, degenerate

6

Pre-IVM duration (h)	No. of IVM (replicates)	% of oocytes at each stage				
		GV	GVBD/CC	M I	M II	Deg
0	33 (3)	14.4 ± 15.5	37.3 ± 27.7 ^{ab}	48.3 ± 20.2	0	0
10	49 (3)	13.5 ± 13.7	1.5 ± 2.6 ^b	82.7 ± 16.1	0	2.2 ± 3.9
20	30 (3)	7.5 ± 6.6	34.2 ± 8.0 ^a	60.0 ± 8.7	2.7 ± 4.8	0
Control*	40 (3)	0	14.0 ± 13.4 ^{ab}	86.0 ± 13.4	0	0

7 *The oocytes collected from antral follicles (2 to 8 mm in diameter) served as *in vivo*-grown
8 controls.

9

10

11 **Table 2. Effect of pre-IVM duration on nuclear maturation of IVG oocytes after 22 h IVM**

12 Values are mean ± SD.

13 GV, germinal vesicle; GVBD, germinal vesicle breakdown; M I, metaphase I; A I/T I, anaphase

14 I/telophase I; M II, metaphase II

15

Pre-IVM duration (h)	No. of IVM (replicates)	% of oocytes at each stage				
		GV	GVBD	M I	A I	M II
0	94 (5)	5.0 ± 7.2	3.9 ± 3.9	25.4 ± 7.2 ^a	14.2 ± 10.7	50.5 ± 10.1 ^c
10	117 (5)	0	0	5.5 ± 5.2 ^b	3.8 ± 4.8	91.5 ± 2.3 ^a
20	96 (5)	1.3 ± 2.8	0	19.8 ± 12.6 ^a	0	79.0 ± 12.0 ^b
Control*	103 (5)	0	0	9.6 ± 2.7 ^{ab}	1.8 ± 2.5	88.6 ± 2.6 ^{ab}

16 *The oocytes collected from antral follicles (2 to 8 mm in diameter) served as *in vivo*-grown
17 controls.

18 **Table 3. Effect of pre-IVM duration on meiotic arrest of IVG oocytes**

19 ^{a,b} Values (mean \pm SD) with different superscripts within columns are significantly different (P <
20 0.05).

21 GV, germinal vesicle; GVBD, germinal vesicle breakdown; M I, metaphase I

22

Pre-IVM duration (h)	No. of IVM (replicates)	% of oocytes at each stage			
		GV	GVBD	M I	Degeneration
0	42 (3)	100.0 \pm 0.0 ^a	0	0	0
10	60 (3)	100.0 \pm 0.0 ^a	0	0	0
20	55 (3)	90.0 \pm 5.5 ^b	5.7 \pm 1.7	1.7 \pm 2.9	2.6 \pm 4.4

23

24

25

26

27 **Table 4. Effect of pre-IVM duration on embryonic development of IVG oocytes**

28 ^{a,b} Values (mean \pm SD) with different superscripts within columns are significantly different ($P < 0.05$).

29

Pre-IVM duration (h)	No. of oocytes (replicates)	% of cleaved oocytes	% of blastocysts based on		Cell no. in blastocysts (n)
			Inseminated oocytes	Cleaved oocytes	
10	164 (5)	79.4 \pm 2.4 ^a	38.9 \pm 4.2 ^a	49.0 \pm 5.0	141 \pm 62 (49 ^{**})
20	193 (5)	63.0 \pm 5.6 ^b	25.8 \pm 7.3 ^b	41.2 \pm 12.0	135 \pm 59 (51)
Control*	150 (5)	82.7 \pm 7.6 ^a	36.7 \pm 3.4 ^a	44.7 \pm 6.4	156 \pm 55 (55)

30 *The oocytes collected from antral follicles (2 to 8 mm in diameter) served as *in vivo*-grown controls.

31 ** Total cell numbers in blastocysts were evaluated in four replicates because some of the blastocysts after IVC were transferred

32