



Title	p62/SQSTM1 Plays a Protective Role in Oxidative Injury of Steatotic Liver in a Mouse Hepatectomy Model
Author(s)	Haga, Sanae; Ozawa, Takeaki; Yamada, Yuma; Morita, Naoki; Nagashima, Izuru; Inoue, Hiroshi; Inaba, Yuka; Noda, Natsumi; Abe, Riichiro; Umezawa, Kazuo; Ozaki, Michitaka
Citation	Antioxidants & redox signaling, 21(18), 2515-2530 https://doi.org/10.1089/ars.2013.5391
Issue Date	2014-12-20
Doc URL	http://hdl.handle.net/2115/60349
Rights	The final published version is available from Mary Ann Liebert, Inc., publishers at http://dx.doi.org/10.1089/ars.2013.5391
Type	article (author version)
File Information	Antioxid Redox Signal_21(18)_2515-2530.pdf



[Instructions for use](#)

1 Original Research Communication

2 p62/SQSTM1 plays a protective role in oxidative injury

3 of steatotic liver in a mouse hepatectomy model.

4
5 Sanae Haga¹, Takeaki Ozawa², Yuma Yamada³, Naoki Morita⁴,
6 Izuru Nagashima⁴, Hiroshi Inoue⁵, Yuka Inaba⁵, Natsumi Noda¹, Riichiro Abe⁶,
7 Kazuo Umezawa⁷, Michitaka Ozaki¹†

8
9 ¹ Laboratory of Molecular and Functional Bio-imaging, Faculty of Health
10 Sciences, Hokkaido University, Sapporo, Japan

11 ² Department of Chemistry, School of Science, The University of Tokyo, Tokyo,
12 Japan

13 ³ Laboratory for Molecular Design of Pharmaceuticals, Faculty of
14 Pharmaceutical Sciences, Hokkaido University, Sapporo, Japan

15 ⁴ Bioproduction Research Institute, National Institute of Advanced Industrial
16 Science and Technology (AIST), Sapporo, Japan

17 ⁵ Department of physiology and metabolism, Brain/Liver Interface Medicine
18 Research Center, College of Medical, Pharmaceutical and Health Sciences,
19 Kanazawa University, Kanazawa, Japan

20 ⁶ Department of Dermatology, Hokkaido University Graduate School of
21 Medicine, Sapporo, Japan

22 ⁷ Department of Molecular Target Medicine Screening, Aichi Medical University
23 School of Medicine, Nagakute, Japan

24
25 *Abbreviated title (Running title):*

26 p62/SQSTM1 & oxidative injury in steatotic liver

27
28
29
30

1 *[†]Corresponding author:*

2 Michitaka Ozaki M.D., Ph.D.

3 Address: Laboratory of Molecular and Functional Bio-imaging, Faculty of Health
4 Sciences, Hokkaido University

5 N-12, W-5, Kita-ku, Sapporo, Hokkaido, 060-0812 Japan

6 Tel.: +81-11-706-3337/+81-11-706-4753

7 Fax: +81-11-706-3337/ +81-11-717-4753

8 E-mail address: ozaki-m@med.hokudai.ac.jp

9

10 *Word count:*

11 Abstract-abbreviation: 5,619 words

12

13 *Number of references:*

14 40 references

15

16 *Number of illustrations:*

17 4 grayscale illustrations (Figs. 4, 5, 6, 7 and 8) and 4 color illustrations (online
18 only, Figs. 1, 2, 3 and 9)

19

20 *Conflicts of interest:*

21 The authors declare that no conflicts of interest exist.

22

23 *Financial support:*

24 This work was supported by Grants-in-Aid for Scientific Research from the
25 Ministry of Education, Culture, Sports, Science and Technology of Japan
26 (#23659631, #23249066 and #20249060 to MO), a JSPS Grant for Young
27 Scientists (#10J06373 to SH) and the heartfelt donation from Mr. & Mrs.
28 Fujikawa (to MO).

29

30

1 ABSTRACT

2 *Aims:* Liver injury and regeneration involve complicated processes and are
3 affected by various physio-pathological factors. We investigated the
4 mechanisms of steatosis-associated liver injury and delayed regeneration in a
5 mouse model of partial hepatectomy.

6 *Results:* Initial regeneration of the steatotic liver was significantly delayed
7 after hepatectomy. Although hepatocyte proliferation was not significantly
8 suppressed, severe liver injury with oxidative stress occurred immediately
9 after hepatectomy in the steatotic liver. FasL/Fas expression was up-regulated
10 in the steatotic liver, whereas the expression of anti-oxidant and anti-apoptotic
11 molecules (catalase/Mn-SOD/Ref-1 and Bcl-2/Bcl-xL/FLIP, respectively) and
12 p62/SQSTM1, a steatosis-associated protein, were down-regulated.
13 Interestingly, pro-survival Akt was not activated in response to hepatectomy,
14 although it was sufficiently expressed even before hepatectomy. Suppression of
15 p62/SQSTM1 increased FasL/Fas expression and reduced Nrf-2-dependent ARE
16 activity and anti-oxidant responses in steatotic and non-steatotic hepatocytes.
17 Exogenously added FasL induced severe cellular oxidative stress and
18 necrosis/apoptosis in steatotic hepatocytes, with only the necrosis being

1 inhibited by pre-treatment with anti-oxidants, suggesting that
2 FasL/Fas-induced oxidative stress mainly leads to necrosis. Furthermore,
3 p62/SQSTM1 re-expression in the steatotic liver markedly reduced liver injury
4 and improved liver regeneration.

5 *Innovation:* The present study is the first to demonstrate that reduced
6 expression of p62/SQSTM1 plays a crucial role in post-hepatectomy acute injury
7 and delayed regeneration of steatotic liver, mainly via redox-dependent
8 mechanisms.

9 *Conclusion:* In the steatotic liver, reduced expression of p62/SQSTM1 induced
10 FasL/Fas overexpression and suppressed anti-oxidant genes mainly through
11 Nrf-2 inactivation, which, together with the hypo-responsiveness of Akt,
12 caused post-hepatectomy necrotic/apoptotic liver injury and delayed
13 regeneration, both mainly via a redox-dependent mechanism.

14

15

16

17

18

1 Introduction

2

3 Liver regeneration occurs as a series of physio-pathological events that result in
4 quantitative recovery from loss of liver mass, compensating for decreased
5 hepatic volume and impaired function. The liver has the ability to restore lost
6 volume, a phenomenon that is rarely seen in other organs (10, 22). It is well
7 established that normal adult hepatocytes usually are quiescent but have the
8 potential to replicate. After surgical procedures that reduce liver mass, such as
9 partial hepatectomy (PH) or live-donor liver transplantation, a rapid
10 enlargement of the residual or grafted liver commonly occurs to restore liver
11 mass and function (9). Poor or insufficient regeneration of diseased liver, for
12 example in non-alcoholic steatohepatitis (NASH) and liver cirrhosis, or of aged
13 liver, is potentially fatal for these patients (1, 25, 32, 33). Therefore, a better
14 understanding of the molecular mechanisms of liver injury and delayed
15 regeneration in various pathological conditions may lead to clinical benefits.

16 Hepatic steatosis (fatty liver) is a commonly encountered hepatic
17 disorder that may be caused by various factors, such as obesity, diabetes
18 mellitus and alcohol consumption (33). It is often considered a benign condition

1 because it does not usually cause severe clinical symptoms. However,
2 living-donor liver transplantation from a donor with a steatotic liver is
3 problematic, because steatosis often causes immediate graft failure (primary
4 non-function) (33).

5 In the normal liver, various mitotic factors and cytokines promptly
6 activate various cellular signals and events, eventually leading to sufficient
7 liver regeneration after PH (9, 10, 22, 36). In the steatotic liver, impairment of
8 signaling mechanisms due to adaptation to chronic metabolic abnormalities and
9 decreased adenosine triphosphate (ATP) production has been reported to be
10 the likely cause of increased mortality and impaired regeneration after PH (24,
11 33, 35). In addition, hepatic steatosis is considered to reduce tolerance to
12 ischemic injury and oxidative stress (OS) (2, 33).

13 This study was designed to explore the root causes of post-PH liver
14 injury and delayed regeneration of the steatotic liver, and to identify the
15 molecules responsible for these phenomena. Here, we report that the
16 reduction of steatosis-associated p62/sequestosome1 (p62/SQSTM1)
17 accelerates post-PH liver injury with OS and is involved in impairment of the
18 initial regeneration of steatotic liver following PH.

1 Results

2

3 *Initial liver regeneration was disturbed in the steatotic liver after PH, despite*
4 *sufficient hepatocyte proliferation.*

5

6 First, we monitored liver recovery in db/db steatotic mice and lean
7 littermate controls after two-thirds PH (Fig. 1). The liver lipid content was
8 determined by Sudan III stain in the lean controls and the db/db steatotic mice
9 (Fig. 1A), which showed evident lipid accumulation only in the db/db mouse
10 liver. Liver recovery occurred immediately after PH in the non-steatotic liver of
11 the control mice, whereas it was markedly disturbed in the steatotic liver of
12 the db/db mice, at least until 72 hr post-PH (Fig. 1B). The recovery of the
13 steatotic liver, however, was equivalent to that of the control liver at 7 days
14 post-PH. This observation suggests that the regeneration of the steatotic liver
15 may be initially disturbed by a reduced mitotic response after PH.

16 To investigate the mitotic response in the steatotic liver, we examined
17 and counted proliferating cell nuclear antigen (PCNA)-positive hepatocytes (Fig.
18 1C) and mitotic hepatocytes (Fig. 1D) in the regenerating liver after PH. These

1 histological analyses showed that PCNA positivity and mitosis in the
2 regenerating liver peaked at 36 hr and 48 hr post-PH, respectively, and
3 thereafter decreased. Western blot analysis showed that cyclinD1 and PCNA
4 similarly peaked at 36 hr post-PH and thereafter decreased. Signal transducer
5 and activator of transcription 3 (STAT3), which is mainly associated with cell
6 proliferation, was found to be expressed at equivalent levels in both livers and
7 to be phosphorylated at 4 hr post-PH in both livers (Fig. 1E). Although post-PH
8 expression of CyclinD1 and PCNA was more evident in the db/db mouse liver
9 and the control liver, respectively, the counts of mitotic cells were almost
10 equal in both groups (Fig. 1D).

11 In light of the above, we can say that hepatocyte proliferation in the
12 db/db mouse liver was equivalent to that in the control liver, suggesting that a
13 mechanism other than delayed or disturbed mitotic response was the main
14 cause of the delayed regeneration of the steatotic liver.

15

16 *OS and injury occurred in the steatotic liver of db/db mice immediately after*
17 *PH.*

18

1 Next, we monitored post-PH liver injury in the db/db mice to elucidate the
2 mechanism underlying delayed liver regeneration. Marked liver injury occurred
3 within 12 hr of PH in the db/db mice (Fig. 2A). The serum levels of aspartate
4 aminotransferase (AST), alanine aminotransferase (ALT) and lactate
5 dehydrogenase (LDH) were markedly elevated at 12 to 24 hr post-PH in the
6 db/db mice but had recovered to baseline levels at 3 days post-PH, suggesting
7 that the harmful cellular events associated with PH occurred immediately.
8 Histological examination also confirmed liver injury with sporadic necrosis and
9 neutrophil infiltration observed especially in the peri-portal areas of the
10 steatotic liver 24 hr after PH (Fig. 2A).

11 To determine whether OS is induced in the steatotic liver in the early
12 post-PH period, we studied hepatic OS using a reduction-oxidation-sensitive
13 green fluorescent protein (roGFP) probe (Fig. 2B) (7, 15). This redox-sensitive
14 roGFP probe showed the immediate generation of robust OS only in the
15 steatotic liver, which peaked within 30 min of PH and continued for at least 8
16 hr.

17 We next examined if and when apoptotic cell death occurred in the
18 post-PH steatotic liver. Bio-imaging of the post-PH remnant liver showed

1 immediate activation of caspase-3, peaking at 4 hr post-PH and continuing for
2 at least 24 hr (Fig. 2C). Hepatic apoptosis in the steatotic liver also began to
3 increase immediately after PH and peaked at 12 hr post-PH (Fig. 2D). These
4 analyses reveal that apoptotic cell death occurred at 4 hr post-PH in the
5 steatotic liver, but did not occur in the control liver.

6 Given that hepatic OS and apoptosis occurred sequentially and that
7 some caspases are redox-sensitive (17), we considered that the severe post-PH
8 OS might have activated redox-sensitive caspases and thereby induced
9 apoptosis-mediated injury in the db/db mice.

10

11 *Analysis of hepatic signals associated with cell proliferation, cell*
12 *survival/death and OS.*

13

14 To elucidate the mechanisms of hepatic OS, injury and delayed
15 regeneration in the db/db mice, we investigated the hepatic signal molecules
16 involved.

17 Fas and Fas-ligand (FasL) were both expressed at higher levels in the
18 steatotic liver before and after PH as compared with the control liver (Fig. 3A).

1 Akt, mainly associated with cell survival, was expressed at equivalent levels in
2 both livers, but was transiently phosphorylated after PH with a peak at 24 hr in
3 the control liver. Interestingly, in the steatotic liver, it was activated even
4 before PH and there was no increase in activation after PH.

5 Anti-oxidant proteins, catalase, manganese superoxide dismutase
6 (Mn-SOD) and redox factor-1 (Ref-1), were expressed at lower levels in the
7 steatotic liver than in the control liver, and only copper/zinc-superoxide
8 dismutase (Cu/Zn-SOD) was expressed at equivalent levels (Fig. 3B).
9 Anti-apoptotic proteins such as Bcl-2/-xL and FLICE-like inhibitory protein
10 (FLIP) were expressed at lower levels in the steatotic liver. Together with the
11 hypo-responsiveness of Akt and the decreased levels of anti-apoptotic and
12 anti-oxidant molecules, these harmful signals mediated by highly expressed
13 FasL/Fas may play a central role in post-PH liver injury and delayed liver
14 regeneration.

15

16 *FasL/Fas signals induce cellular ROS and death in steatotic hepatocytes.*

17

18 Next, we determined whether the induction of steatosis in hepatocytes

1 leads to FasL/Fas overexpression and whether FasL stimuli induce cellular
2 reactive oxygen species (ROS) and more cell death (apoptosis/necrosis) in
3 steatotic hepatocytes. Treatment of AML12 liver cells with T0901317, an
4 agonist for Liver X Receptor- α (4), induced a marked steatotic change in
5 hepatocytes within 5 days (Fig. 4A). Similar to the *in vivo* observations, the
6 steatotic hepatocytes showed overexpression of both FasL and Fas, but less
7 expression of p62/SQSTM1.

8 Steatotic hepatocytes were more prone to die than non-steatotic
9 hepatocytes when challenged with exogenous FasL (Jo2 antibody) (Fig. 4B).
10 Steatotic hepatocytes stimulated by FasL showed more cell death and
11 caspase-3 activity as compared with non-steatotic hepatocytes. Although cell
12 death continued for 20-24 hr after Jo2 administration, caspase-3 was
13 transiently activated with a peak at 4 hr. These facts indicate that
14 FasL/Fas-mediated death signals induce hepatocyte death through
15 caspase-3-mediated apoptosis and/or non-apoptotic cell death, and that
16 steatotic hepatocytes are more susceptible than control hepatocytes to
17 FasL/Fas-induced cell death.

18 To examine the involvement of ROS in Fas-induced cell death and to

1 exclude neutrophil-derived ROS in post-PH liver injury, we next examined ROS
2 generation in hepatocytes with and without steatosis (Fig. 4C). Steatotic
3 hepatocytes expressing greater amounts of FasL and Fas molecules, as
4 compared with non-steatotic hepatocytes, clearly generated cellular ROS after
5 stimulation of FasL. Hepatocytes in which Fas was depleted by small interfering
6 RNAs (siRNA), however, generated little cellular ROS, even when stimulated
7 with Jo2, which confirmed that FasL/Fas signals induce more cellular ROS in
8 steatotic hepatocytes.

9 To investigate the involvement of FasL/Fas-mediated cellular ROS in
10 the death of steatotic cells, we studied the effects of anti-oxidants on cell
11 death (apoptosis and necrosis) (Fig. 4D). Unexpectedly, FasL/Fas-induced
12 necrotic cell death was significantly inhibited by pre-treatment with the
13 anti-oxidants ascorbic acid, Trolox and N-acetyl cysteine (NAC), whereas
14 apoptotic cell death was not inhibited at all. These data indicate that
15 FasL/Fas-induced cell death occurred through apoptosis and necrosis in a
16 caspase-dependent manner and a ROS-dependent manner, respectively.

17

18 *Reduced expression of p62/SQSTM1 in the steatotic liver and hepatocytes is*

1 *responsible for increased FasL/Fas expression and OS.*

2

3 To examine the molecular mechanism underlying the increased
4 FasL/Fas signaling and OS in the steatotic liver, we focused on p62/SQSTM1, a
5 molecule associated with steatosis (23), whose expression was reduced only in
6 the steatotic liver and hepatocytes (Fig. 3A, 3C and 4A). It has been reported
7 that p62/SQSTM1 is associated negatively with hepatic steatosis (23, 28) and
8 that its reduced expression may impair the nuclear factor erythroid 2-related
9 factor 2 (Nrf-2)-dependent cellular anti-oxidant system (16, 19).

10 Acute reduction of p62/SQSTM1 by siRNA did not affect the cellular
11 triglyceride content of hepatocytes (Fig. 5A), suggesting that p62/SQSTM1 does
12 not primarily induce steatotic changes in hepatocytes and that its reduced
13 expression occurs secondary to the steatotic changes in hepatocytes. In the
14 steatotic liver, p62/SQSTM1 was expressed at much lower levels before PH,
15 although it was induced transiently at 24 hr post-PH in both the normal and the
16 steatotic liver (Fig. 3A). The expression of *p62/SQSTM1* mRNA and protein (Fig.
17 3A, 4A, 5B) was down-regulated in the steatotic liver and hepatocytes.
18 Conversely, the expression of FasL and Fas was up-regulated in the steatotic

1 liver and hepatocytes (Fig. 3A, 4A). Interestingly, even non-steatotic
2 hepatocytes strongly expressed Fas and FasL proteins when depleted of
3 p62/SQSTM1 (Fig. 5C). In contrast, induction of p62/SQSTM1 in normal and
4 steatotic hepatocytes reduced FasL and Fas expression (Fig. 5D). It is
5 interesting that the induction of p62/SQSTM1 was less efficient in steatotic
6 hepatocytes than in non-steatotic hepatocytes, resulting in an insufficient
7 reduction of FasL/Fas molecules in steatotic hepatocytes (Fig. 5D). This
8 suggests that p62/SQSTM1 negatively regulates these harmful molecules.

9

10 *p62/SQSTM1 is responsible for the anti-oxidant property of hepatocytes.*

11

12 p62/SQSTM1 is known to positively regulate the DNA-binding activity of
13 Nrf-2 by physical association with Kelch-like ECH-associated protein-1 (Keap-1),
14 which binds directly to Nrf-2 and inactivates its transcriptional activity on
15 anti-oxidant response elements (ARE) (16, 19). Therefore, we examined the
16 regulatory role of p62/SQSTM1 in Nrf-2-mediated ARE-transcriptional activity
17 in hepatocytes. As expected, reduction of *p62/SQSTM1* expression by siRNA
18 decreased ARE-transcription significantly, but did not decrease *Nrf-2* reporter

1 activity (Fig. 6A). Although the transcription activity of Nrf-2 was increased by
2 *p62/SQSTM1* depletion, possibly by a negative feedback mechanism, its protein
3 expression was slightly decreased (Fig. 6B). These data suggest that free
4 Keap-1, increased by the reduced expression of *p62/SQSTM1*, alternatively
5 bound to Nrf-2, eventually leading to the suppression of *ARE* transcription.

6 The expression of a series of anti-oxidant proteins (catalase, Mn-SOD,
7 Ref-1, HO-1, TRX and GPx) in hepatocytes was reduced by depletion of
8 *p62/SQSTM1* (Fig. 6B), probably through the suppression of *ARE* activity (Fig.
9 6A). Among these proteins, catalase, Mn-SOD, Ref-1 and Nrf-2 were increased
10 significantly by overexpression of *p62/SQSTM1*. These data clearly indicate that
11 *p62/SQSTM1* is responsible for the expression of anti-oxidants both in cultured
12 hepatocytes and in the liver. Overall, *p62/SQSTM1* is critically involved in the
13 increased FasL/Fas expression and the decreased anti-oxidant properties of
14 steatotic hepatocytes, its expression being suppressed by cellular steatosis.

15 Because hypoxia/reoxygenation (H/R) of hepatocytes induces cellular
16 ROS and injury, we examined whether H/R-induced ROS is exacerbated in
17 hepatocytes by previous ablation of *p62/SQSTM1* (Fig. 6C). Four hours of
18 hypoxia followed by reoxygenation induced cellular ROS in control hepatocytes,

1 and this was significantly exacerbated in hepatocytes without p62/SQSTM1.
2 These data confirm the general role of p62/SQSTM1 in protection from various
3 kinds of OS in hepatocytes. Furthermore, the induction of p62/SQSTM1 in
4 non-steatotic and steatotic hepatocytes equally and sufficiently suppressed
5 Jo2-induced necrotic and apoptotic cell death, although Jo2 induced cell death
6 more in steatotic hepatocytes than in non-steatotic hepatocytes (Fig. 6D).
7 These data demonstrate the anti-oxidant and cytoprotective capacities
8 (against both necrosis and apoptosis) of p62/SQSTM1 in hepatocytes with or
9 without steatosis.

10

11 *Induction of p62/SQSTM1 in the db/db steatotic liver reduced liver injury after*
12 *PH and improved liver regeneration.*

13

14 In order to examine the *in vivo* effect of p62/SQSTM1, we determined
15 whether or not re-expression of p62/SQSTM1 in the steatotic liver of db/db
16 mice improves liver regeneration after PH by reducing injury. PH was
17 performed 36 hr after hydrodynamic injection of the *p62/SQSTM1* gene.
18 p62/SQSTM1 protein was robustly expressed in the liver of db/db mice at the

1 time of liver resection (Fig. 7A). Induction of *p62/SQSTM1* in the steatotic liver
2 clearly protected it from PH-induced injury (Fig. 7B), although some liver
3 injury remained. The expression levels of PCNA and cyclinD1 proteins peaked at
4 48 hr post-PH similarly in both the control and *p62/SQSTM1*-expressing liver
5 (Fig. 7C). In addition, there was no significant difference in the mitotic index
6 and the counts of PCNA-positive nuclei in hepatocytes between the two livers.
7 Induction of *p62/SQSTM1* gene did not affect post-PH cell proliferation in the
8 db/db mouse liver.

9 By re-introducing *p62/SQSTM1* into the db/db mouse liver and clearly
10 reducing liver injury, post-PH liver regeneration was partially but consistently
11 improved at 72 hr post-PH (Fig. 7D).

12

13 *A different model of liver steatosis (HFD mice) also shows reduced*
14 *p62/SQSTM1 in liver, leading to increased FasL/Fas and decreased*
15 *anti-oxidant/anti-apoptotic molecules.*

16

17 In order to determine whether or not the reduction of *p62/SQSTM1* and
18 its effects are limited to the liver in db/db mice, we studied a different model

1 of liver steatosis (HFD model). In this model, mice were fed on chow containing
2 35% fat by weight from 4 to 16 weeks of age, which made hepatocytes
3 sufficiently steatotic (Fig. 8A). As shown in Fig. 8B, the protein level of
4 p62/SQSTM1 was reduced in the steatotic liver tissue. FasL and Fas were
5 increased, and anti-oxidant molecules such as catalase, MnSOD and Ref-1 and
6 anti-apoptotic molecules such as Bcl-2, Bcl-xL and FLIP were all reduced. These
7 changes were the same as those observed in the db/db mouse liver. Thus,
8 reduction of p62/SQSTM1 and p62/SQSTM1-dependent regulation of these
9 pathways/signals are not confined to the db/db mouse model, but occur in the
10 steatotic liver of various origins.

11

12

1 Discussion

2

3 Fas expression and apoptosis are well-recognized features of human
4 NASH (11) and other steatotic liver diseases (6, 38), suggesting the potential
5 role of Fas in injury of the steatotic liver. However, the precise mechanism of
6 injury of the steatotic liver has not been thoroughly elucidated. In this study,
7 using a mouse model of hepatectomy, we first demonstrated that lack of
8 p62/SQSTM1 plays a pivotal role in post-PH acute injury of the steatotic liver
9 (Fig. 9). Initial regeneration of the steatotic liver after PH was delayed not by a
10 suppressed mitotic response but by an immediate post-PH liver injury
11 consisting of Fas/caspase-mediated apoptosis and Fas/OS-dependent necrosis.

12 In terms of the mechanism underlying acute injury of the steatotic liver,
13 we demonstrated that steatosis-induced reduction of p62/SQSTM1, by
14 enhancing the cellular FasL/Fas pathway, causes post-PH necrotic and
15 apoptotic acute liver injury in a redox-dependent and redox-independent
16 manner, respectively. Of particular note, reduction of p62/SQSTM1 lowered the
17 expression of anti-oxidant molecules (catalase, MnSOD, Ref-1, HO-1, TRX and
18 GPx) by decreasing Keap-1/Nrf-2 binding (i.e., *ARE* activity), making the liver

1 susceptible to OS. Additionally, we demonstrated that OS induced by FasL/Fas
2 in the steatotic liver mainly caused the necrotic cell death of hepatocytes. Just
3 as important was the finding that the hypo-responsiveness of Akt, a pro-survival
4 and anti-oxidant molecule, enhanced/promoted OS-induced necrotic injury
5 and caspase-mediated apoptotic liver injury (14, 30), together with reduced
6 expression of anti-oxidant molecules and anti-apoptotic molecules,
7 respectively.

8 In the present study, steatotic changes in the liver or hepatocytes
9 reduced the expression of p62/SQSTM1, which may have induced mild but
10 continuous liver injury by causing overexpression of FasL/Fas even under basal
11 conditions. Serum levels of AST, ALT and LDH without PH showed a slight but
12 significant elevation of these enzymes [AST (IU/L): 13.0 \pm 0.0 and 70.0 \pm 24.0;
13 ALT (IU/L): 35.5 \pm 5.0 and 54.5 \pm 5.0; LDH (IU/L): 349.5 \pm 12.0 and
14 635.5 \pm 132.2, in control and db/db mice, respectively; $p < 0.05$] (Fig. 2A),
15 indicating continuous mild liver injury even without PH. Increased expression
16 and activation of Akt was also observed without PH, which again may have been
17 induced by continuous liver injury under basal conditions. Surgical stress such
18 as PH might exacerbate the liver injury or trigger marked liver injury, because

1 surgery induces a strong inflammatory reaction and oxidative stress. In
2 particular, the steatotic liver may be more vulnerable to such insults, because
3 Fas-receptor expression is elevated and anti-oxidative capacity is reduced by
4 p62-associated Nrf2/ARE inactivation. These mechanisms may underlie
5 steatosis-associated acute liver injury in the mouse. The compensatory
6 regeneration (mitotic response) was not clearly observed under basal
7 conditions (Fig. 1A); however, phosphorylation of STAT3 (Supplementary Fig.)
8 and Akt (Fig. 3A) was observed even without PH, as previously reported (24, 37).
9 This may explain the cellular response for proliferation (p-STAT3) and survival
10 (p-Akt) against mild and continuous hepatocyte injury. After PH, acute liver
11 injury may be enhanced by the PH-associated rapid increase in exogenous FasL,
12 and the ensuing apoptosis and OS-mediated necrosis in the steatotic liver.
13 Hepatic OS might also be induced/enhanced by ROS generated from
14 mitochondria and the NOX system after PH more in the steatotic liver than in
15 the normal liver (3, 13).

16 Although expression of p62/SQSTM1 was markedly decreased in the
17 steatotic liver and hepatocytes (Fig. 3A, 3C, 4A, 5B), depletion of *p62/SQSTM1*
18 did not induce steatotic changes in hepatocytes (Fig. 5A), suggesting that

1 hepatic steatosis is the primary cause of the reduced expression of p62/SQSTM1.
2 Also, the db/db mice showed moderate hyperglycemia in addition to steatotic
3 liver, raising the question of whether hyperglycemia might cause the reduction
4 of p62/SQSTM1. As shown by the *in vitro* assay in Fig. 4, however, chemically
5 induced steatosis in hepatocytes with a normal glucose level consistently
6 decreased p62/SQSTM1 and increased FasL/Fas. Additionally, in an alternative
7 *in vivo* steatotic liver model, in which steatotic liver was induced by high fat
8 feeding, the p62/SQSTM1 level in the liver was similarly reduced (Fig. 8). Thus,
9 we conclude that it is steatosis, and not hyperglycemia, that mainly leads to
10 the reduction of p62/SQSTM1 in hepatocytes.

11 The exact mechanism underlying the reduced expression of
12 p62/SQSTM1 and overexpression of FasL/Fas was not elucidated in this study;
13 however, a study on p62/SQSTM1 and its anti-apoptotic capacity showed that
14 p62/SQSTM1 exerts anti-apoptotic capacity by inhibiting BID and caspase-3
15 activities (20). Generally, it seems that p62/SQSTM1, when it is present, serves
16 as a cell-protection molecule by providing an anti-apoptotic function. In the
17 present study, when p62/SQSTM1 was not present, the expression of assaultive
18 FasL and Fas genes/proteins were both up-regulated. p62/SQSTM1, when

1 present, may also “positively” suppress FasL/Fas expression at the
2 transcriptional level as a “gatekeeper molecule”. Furthermore, the negative
3 regulation of FasL/Fas expression by p62/SQSTM1 was observed in hepatocytes
4 without steatosis (Fig. 5C, 5D), suggesting that p62/SQSTM1 is a key molecule
5 in the protection of hepatocytes and other types of cells that potentially
6 express the Fas molecule.

7 Deterioration of the anti-oxidant defense mechanism in the steatotic
8 liver may have enhanced OS-induced acute liver injury after PH. If this is the
9 case, then the steatotic liver with reduced levels of p62/SQSTM1 may be very
10 vulnerable to OS, even in a static condition. p62/SQSTM1 itself is involved in
11 the regulation of cellular OS through a Keap-1/Nrf-2-dependent anti-oxidant
12 mechanism involving glutathione S-transferase, cytochrome p450, HO-1, SOD,
13 catalase or p62/STSQM1 (8, 16, 19, 23, 28, 31, 40). Indeed, in the present study,
14 these anti-oxidant molecules were found to be down-regulated in the steatotic
15 liver (Fig. 3B) and the ablation of *p62/SQSTM1* was found to reduce both *ARE*
16 reporter activity and their expression, even in non-steatotic hepatocytes (Fig.
17 6A, B). These facts strongly suggest that p62/SQSTM1 confers anti-oxidant
18 properties on hepatocytes through the Keap-1/Nrf-2 system. Therefore, a

1 decrease in the p62/SQSTM1 level of hepatocytes, whatever the cause, might
2 make the liver and hepatocytes vulnerable to OS.

3 Our previous paper (24) analyzed delayed regeneration of steatotic
4 liver from the aspect of cell cycle progression and proliferation. It revealed
5 that hepatocyte proliferation was mildly reduced by the suppressed expression
6 of Wee1/Myt1 at 1-3 days post-PH, but that it continued even after 3 days
7 post-PH, until the lost liver mass was made up. In the control liver, however,
8 hepatocyte proliferation ceased after 3 days post-PH. These observations
9 indicate that post-PH hepatocytes in the db/db mouse liver can proliferate
10 until 7 days post-PH, despite the severe injury early after PH. We consider that
11 transient but severe liver injury early after PH in the steatotic liver, together
12 with somewhat slow cell proliferation, causes the delayed liver regeneration
13 but that hepatocyte proliferation continues until the lost liver mass is made up
14 (3-7 days post-PH). In the present study, re-expression of p62/SQSTM1 in the
15 steatotic liver did not affect cell proliferation after PH (Fig. 7C).

16 In an acute stress model such as PH, the increase in FasL/Fas caused by
17 reduction of p62/SQSTM1 seems to be highly relevant in inducing acute liver
18 injury, because the increased FasL/Fas signal induced post-PH liver injury in a

1 positive manner (caspase-dependent apoptosis and OS-dependent necrosis). By
2 contrast, the reduced Nrf-2/Keap-1 signal accelerated the above injurious
3 signals by reducing the expression of anti-oxidant molecules. Previous reports
4 have described the pathological relevance of p62/SQSTM1 in acute and chronic
5 situations in humans, such as neuronal, musculo-skeletal and infectious
6 diseases (5, 12, 27, 34). Many studies focused on analyses of p62/SQSTM1 with
7 respect to autophagy, because p62/SQSTM1 has been known to be a good in
8 vivo marker of autophagy. However, one report indicated the relevance of
9 p62/SQSTM1 and the FasL/Fas pathway in the acute lung injury that occurs in
10 acute respiratory distress syndrome (ARDS) (20). Thus, p62/SQSTM1 might also
11 play an important role in human diseases other than steatotic liver.

12 This study is the first to show the critical role of p62/SQSTM1 in
13 redox-dependent steatotic liver injury and disturbed early regeneration after
14 PH. The lack of p62/SQSTM1 in some pathological conditions such as steatosis
15 induces acute liver injury. Further studies are required to elucidate thoroughly
16 the mechanism of liver protection and injury, but the present data clearly
17 demonstrate the critical role of p62/SQSTM1 in the liver anti-oxidant system
18 and provide important clues toward the development of new therapies for

1 various pathological liver conditions.

2

1 Innovation

2

3 Clinically, steatotic liver has been considered to be the likely cause both of
4 increased mortality after hepatectomy, mainly as a result of reduced tolerance
5 to ischemic injury and OS, and of impaired mitotic response. The present study
6 is the first to demonstrate that the reduced expression of p62/SQSTM1 in
7 steatotic liver directly causes post-PH liver injury, mainly in a redox-dependent
8 manner, and that restoration of p62/SQSTM1 reduces injury and restores the
9 regeneration of steatotic liver. Furthermore, the study importantly suggests
10 the potential role of p62/SQSTM1 in protecting the liver from all kinds of injury.

11

12

13

1 Materials and methods

2

3 *Animal experiments*

4

5 Male homozygous db/db mice (38-42 g body weight, 9-10 weeks old) and lean
6 littermate controls (20-25 g body weight, 9-10 weeks old) were obtained from
7 CLEA Japan (Tokyo, Japan) and used for the 2/3 PH experiment. In the high fat
8 diet (HFD) mouse model, mice (C57BL6/J, male) were fed on chow containing
9 35% fat by weight from 4 to 16 weeks of age (#D12492, Research Diets, Inc., NJ).
10 The mice were fasted overnight prior to the experiments and were
11 anesthetized with 1.5-2.0 % isoflurane (Forane©, Abbott, Tokyo, Japan). The
12 percentage of the whole liver constituted by each lobe was similar and surgical
13 resection of the middle and left lobes resulted in 2/3 PH in both mice. Sudan III
14 stained lipid droplets in more than 90% of the hepatocytes in the liver of db/db
15 mice but very few droplets in the hepatocytes of the control mice. After
16 laparotomy, the left and median liver lobes were surgically resected. The mice
17 were sacrificed for the collection of liver specimens at the time points
18 indicated before or after hepatectomy, and the liver/body weight ratios were

1 calculated to estimate the recovery of liver mass. The animals were
2 maintained under standard conditions and treated according to the Guidelines
3 for the Care and Use of Laboratory Animals of Hokkaido University.

4

5 *Histological analysis*

6

7 To visualize lipid accumulation in the liver and hepatocytes, frozen sections of
8 formalin-fixed liver tissue and hepatocytes were stained with Sudan III (Fig. 1A)
9 and Oil red O dye (Fig. 4A), respectively. To evaluate the proliferation of
10 hepatocytes after PH, PCNA-positive and mitotic hepatocytes were examined
11 histologically in liver tissues at 36, 48 and 72 hr post-PH. To assess the
12 expression of p62/SQSTM1 in mouse liver, immunohistochemical analysis of
13 p62/SQSTM1 was performed in the untreated liver tissues. These tissues were
14 fixed in 10% buffered formalin and paraffin-embedded, and subjected to
15 hematoxylin and eosin (H&E) staining and immunohistochemical staining with
16 anti-PCNA and anti-p62/SQSTM1. At least 500 hepatocytes were counted for
17 mitotic index or PCNA-positivity at least in triplicate in different sections in
18 each group.

1

2 *Western blot analysis*

3

4 Whole-liver protein extracts (30 µg) were separated by 10% SDS-PAGE and
5 transferred to a PVDF membrane. The following primary antibodies were used:
6 Akt/phospho-Akt, signal STAT3/phospho-STAT3, CyclinD1, p62/SQSTM1 (Cell
7 Signaling Technology, Boston, MA), Fas, Ref-1, CuZn-SOD, Bcl-2/xL, Nrf-2, HO-1
8 (Santa Cruz Biotechnology, Santa Cruz, CA), FasL, GPx (Abcam, Cambridge, UK),
9 Catalase, FLIP (EMD Biosciences, Darmstadt, Germany), Mn-SOD and TRX (BD
10 Transduction Laboratories, Franklin Lakes, NJ).

11

12 *In vivo imaging of mouse liver redox states (oxidative stress) and caspase-3*
13 *activity (apoptosis)*

14

15 We developed a recombinant adenovirus encoding roGFP, which allows
16 real-time visualization of the oxidation/reduction potentials of various cells in
17 vitro (7). Three days before each experiment, *AdroGFP* was administered
18 intravenously via the tail vein in a volume of 100 µl (5×10^7 pfu/mouse). For *in*

1 *in vivo* liver imaging, the liver was exposed under anesthesia to enable a CCD
2 camera to record images directly using a Photon Imager (Biospace Co., Paris,
3 France). Fluorescence intensities at 530 nm from excitation at 480 nm were
4 measured and plotted reciprocally [$1/ (em. 530 \text{ at ex. } 480 \text{ nm})$]. This allowed
5 the signal to increase with oxidation and to decrease with reduction and was
6 used as an index of *in vivo* redox states.

7 An optical probe termed pcFirefly luciferase (pcFluc)-DEVD was used to detect
8 caspase-3 activity in real-time (18, 29). We also developed a recombinant
9 adenovirus encoding pcFluc-DEVD and used it to infect mice (5×10^7
10 pfu/mouse). D-Luciferin, a luciferase substrate, was injected intraperitoneally
11 at 1 mg in 100 μ l of PBS. *In vivo* imaging of the mouse liver was performed using
12 an *in vivo* imager for 5 minutes at 5-10 minutes after injection.

13

14 *Biochemical measurement of liver injury (apoptosis/necrosis)*

15

16 For evaluation of apoptotic cell death, an ELISA kit (Cell Death Detection
17 ELISA^{PLUS}; Roche, Basel, Switzerland) was used according to the manufacturer's
18 instructions. Biochemical analyses, such as serum AST, ALT and LDH levels, as

1 indices of liver injury, were performed at the time points indicated before and
2 after PH.

3

4 *In vitro and in vivo gene silencing with RNA interference and gene transfer*
5 *with cDNA*

6

7 For *in vitro* and *in vivo* experiments, small interfering RNAs (siRNAs) for mouse
8 *p62/SQSTM1* (sense 5'-GGAACUCGCUAUAAGUGCATT-3', antisense
9 5'-UGCACUUAUAGCGAGUCCCA) and *GAPDH* used as the control were
10 purchased from Ambion, Inc. (Austin, TX) and cDNA for mouse *p62/SQSTM1* was
11 purchased from Origene, Inc. (Rockville, MD). Luciferase cDNA was used as a
12 control. For the *in vitro* experiments, transfection of siRNAs and cDNA into
13 AML12 liver cells was accomplished using Lipofectamine 2000 (Invitrogen,
14 Rockville, MD) according to the manufacturer's instructions. *p62/SQSTM1* and
15 *GAPDH* expression was evaluated by PCR and Western blot analyses. In the *in*
16 *vivo* experiments, hydrodynamic injections were used to introduce the DNA
17 into mouse liver. Plasmid DNA of *luciferase* (*pLuc* as a control) or *p62/SQSTM1*
18 gene (*p-p62*) was injected rapidly into the tail veins of the db/db mice (10 µg in

1 2 ml of saline). The livers were harvested 48 hr after injection, and the
2 luciferase activity and mRNA and protein levels of p62/SQSTM1 were measured
3 to detect expression. The supernatants of liver tissue homogenates were
4 examined for luciferase activity by a Luciferase Assay Systems kit (Promega,
5 Madison, WI).

6

7 *RT-PCR assay*

8

9 First-strand cDNA synthesis used 5 µg of total RNA, Superscript III reverse
10 transcriptase and oligo (dT) 20 primers (Invitrogen), according to the
11 manufacturer's instructions. The cDNA was amplified by PCR with specific
12 primers for mouse p62/SQSTM1 (225 bp): sense
13 *5'-GATGTGGAACATGGAGGGAAGAG-3'*, antisense
14 *5'-AGTCATCGTCTCCTCCTGAGCA-3'*. PCR was performed by 30-25 cycles of
15 denaturation at 94°C for 30 s, annealing at 55°C for 30 s, and extension at 72°C
16 for 30 s.

17

18 *Reporter probes for ARE/Nrf-2 transcriptional activation*

1

2 The ARE-luciferase reporter probe was constructed from the ARE sequences of
3 human *NQO1* in the pGL4.17 firefly luciferase reporter vector (Promega) (26).

4 The secreting type of metridia luciferase reporter probe was used for Nrf-2
5 transcriptional activity assay. The promoter region of *nrf2* (21, 39) was isolated
6 by PCR amplification from genomic DNA of human gastric cells and ligated into
7 pMetLuc2-Reporter (Clontech, Mountain View, CA). Next, the CMV
8 promoter-driven Renilla luciferase gene from pGL4.75 was introduced into this
9 reporter plasmid for normalization of the reporter gene assay. After 48 hr of
10 incubation, the cell lysate or the culture medium was subjected to
11 ARE-luciferase reporter assay (Dual-luciferase reporter assay system, Promega)
12 or Nrf-2-metridia luciferase reporter assay (Ready-To-Glow secreted luciferase
13 reporter assay, Clontech), respectively.

14

15 *Monitoring of viable, apoptotic and necrotic cells*

16

17 Cells at 70%-80 % confluence were plated in a 24-well plate. Growth curves
18 were determined by plating the cells in the IncuCyte system (Essen Instruments,

1 Ann Arbor, MI), which allows the automated non-invasive monitoring of live
2 cells in culture. Cell apoptosis and necrosis were determined using Caspase-Glo
3 3/7 (Essen BioScience, Ann Arbor, MI) and YOYO-1 (Molecular Probe, Eugene,
4 Oregon), respectively.

5

6 *Evaluation of cellular redox states in vitro*

7

8 The redox-sensitive roGFP was adenovirally transduced into AML12 cells (5 moi)
9 48 hr prior to the experiment. The cellular redox potentials were imaged using
10 fluorescence microscopy with an emission wavelength of 530 nm and an
11 excitation wavelength of 400/480 nm. Fluorescence intensities resulting from
12 excitation at 400/480 nm were quantified using BZ-analyzer software (Keyence
13 Corp., Osaka, Japan). At least 20 cells were measured in each experiment, and
14 the ratio ex 400/480 nm was used as a cellular redox index.

15

16 *Statistical analysis*

17

18 All results were expressed as means \pm standard error of the mean (SEM). Data

1 were compared by Fisher's test, and p values of less than 0.05 were considered
2 to be statistically significant.

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

1 Acknowledgments

2

3 The authors are grateful to Ms. Y. Yasuzaki for excellent technical support.

4

1 Abbreviations Used

2 ALT = alanine aminotransferase

3 ARE = anti-oxidant response elements

4 AST = aspartate aminotransferase

5 ATP = adenosine triphosphate

6 Cu/Zn-SOD = copper/zinc-superoxide dismutase

7 FasL = Fas-ligand

8 FLIP = FILICE-like inhibitory protein

9 GPx = glutathione peroxidase

10 H&E = hematoxylin and eosin

11 HFD = high fat diet

12 HO-1 = heme oxygenase 1

13 Keap-1 = Kelch-like ECH-associated protein 1

14 LDH = lactate dehydrogenase

15 Mn-SOD = manganese superoxide dismutase

16 NAC = N-acetyl cysteine

17 NASH = non-alcoholic steatohepatitis

18 Nrf-2 = nuclear factor erythroid 2-related factor 2

- 1 OS = oxidative stress
- 2 p62/SQSTM1 = p62/sequestosome1
- 3 pcFluc = pc firefly luciferase
- 4 PCNA = proliferating cell nuclear antigen
- 5 PH = partial hepatectomy
- 6 Ref-1 = redox factor-1
- 7 roGFP = reduction-oxidation-sensitive green fluorescent protein
- 8 ROS = reactive oxygen species
- 9 SEM = standard error of the mean
- 10 siRNA(s) = small interfering RNA(s)
- 11 STAT3 = signal transducer and activator of transcription 3
- 12 TRX = thioredoxin
- 13

1 References

2

- 3 1. Aoyama T, Ikejima K, Kon K, Okumura K, Arai K and Watanabe S.
4 Pioglitazone promotes survival and prevents hepatic regeneration failure
5 after partial hepatectomy in obese and diabetic KK-A(y) mice. *Hepatology*
6 49: 1636-1644, 2009.
- 7 2. Caraceni P, Domenicali M, Vendemiale G, Grattagliano I, Pertosa A, Nardo B,
8 Morselli-Labate AM, Trevisani F, Palasciano G, Altomare E and Bernardi M.
9 The reduced tolerance of rat steatotic liver to ischemia reperfusion is
10 associated with mitochondrial oxidative injury. *J Surg Res* 124: 160-168,
11 2005.
- 12 3. Cardoso AR, Kakimoto PA, Kowaltowski AJ. Diet-sensitive sources of
13 reactive oxygen species in liver mitochondria: role of very long chain
14 acyl-CoA dehydrogenases. *PLoS One* 8: e77088, 2013.
- 15 4. Cha JY and Repa JJ. The liver X receptor (LXR) and hepatic lipogenesis. The
16 carbohydrate-response element-binding protein is a target gene of LXR. *J*
17 *Biol Chem* 282: 743-751, 2007.
- 18 5. Deretic V. Autophagy in infection. *Curr Opin Cell Biol* 22: 252-262, 2010.

- 1 6. Donohue TM Jr, Clemens DL, Galli A, Crabb D, Nieto N, Kato J and Barve SS.
2 Use of cultured cells in assessing ethanol toxicity and ethanol-related
3 metabolism. *Alcohol Clin Exp Res* 25: 875-935, 2001.
- 4 7. Dooley CT, Dore TM, Hanson GT, Jackson WC, Remington SJ and Tsien RY.
5 Imaging dynamic redox changes in mammalian cells with green fluorescent
6 protein indicators. *J Biol Chem* 279: 22284-22293, 2004.
- 7 8. Faraonio R, Vergara P, Di Marzo D, Pierantoni MG, Napolitano M, Russo T and
8 Cimino F. p53 suppresses the Nrf2-dependent transcription of antioxidant
9 response genes. *J Biol Chem* 281: 39776-39784, 2006.
- 10 9. Fausto N. Liver regeneration. *J Hepatol* 32: 19-31, 2000.
- 11 10. Fausto N. Liver regeneration: from laboratory to clinic. *Liver Transpl* 7:
12 835-844, 2001.
- 13 11. Feldstein AE, Canbay A, Angulo P, Tanai M, Burgart LJ, Lindor KD and Gores
14 GJ. Hepatocyte apoptosis and fas expression are prominent features of
15 human nonalcoholic steatohepatitis. *Gastroenterology* 125: 437-443, 2003.
- 16 12. Garner TP, Long J, Layfield R, Searle MS. Impact of p62/SQSTM1 UBA
17 domain mutations linked to Paget's disease of bone on ubiquitin recognition.
18 *Biochemistry* 50: 4665-4674, 2011.

- 1 13. Guichard C, Moreau R, Pessayre D, Epperson TK, Krause KH. NOX family
2 NADPH oxidases in liver and in pancreatic islets: a role in the metabolic
3 syndrome and diabetes? *Biochem Soc Trans* 36: 920-929, 2008.
- 4 14. Haga S, Morita N, Irani K, Fujiyoshi M, Ogino T, Ozawa T and Ozaki M.
5 p66(Shc) has a pivotal function in impaired liver regeneration in aged mice
6 by a redox-dependent mechanism. *Lab Invest* 90: 1718-1726, 2010.
- 7 15. Haga S, Remington SJ, Morita N, Terui K and Ozaki M. Hepatic ischemia
8 induced immediate oxidative stress after reperfusion and determined the
9 severity of the reperfusion-induced damage. *Antioxid Redox Signal* 11:
10 2563-2572, 2009.
- 11 16. Jain A, Lamark T, Sjøttem E, Larsen KB, Awuh JA, Øvervatn A, McMahon M,
12 Hayes JD and Johansen T. p62/SQSTM1 is a target gene for transcription
13 factor NRF2 and creates a positive feedback loop by inducing antioxidant
14 response element-driven gene transcription. *J Biol Chem* 285: 22576-22591,
15 2010.
- 16 17. Kannan K and Jain SK. Oxidative stress and apoptosis. *Pathophysiology* 7:
17 153-163, 2000.
- 18 18. Kanno A, Yamanaka Y, Hirano H, Umezawa Y and Ozawa T. Cyclic luciferase

- 1 for real-time sensing of caspase-3 activities in living mammals. *Angew*
2 *Chem Int Ed Engl* 46: 7595-7599, 2007.
- 3 19. Komatsu M, Kurokawa H, Waguri S, Taguchi K, Kobayashi A, Ichimura Y, Sou
4 YS, Ueno I, Sakamoto A, Tong KI, Kim M, Nishito Y, Iemura S, Natsume T,
5 Ueno T, Kominami E, Motohashi H, Tanaka K and Yamamoto M. The selective
6 autophagy substrate p62 activates the stress responsive transcription factor
7 Nrf2 through inactivation of Keap1. *Nat Cell Biol* 12: 213-223, 2010.
- 8 20. Liang X, Wei SQ, Lee SJ, Fung JK, Zhang M, Tanaka A, Choi AM, Jin Y. p62
9 sequestosome 1/light chain 3b complex confers cytoprotection on lung
10 epithelial cells after hyperoxia. *Am J Respir Cell Mol Biol* 48:489-496, 2013.
- 11 21. Marzec JM, Christie JD, Reddy SP, Jedlicka AE, Vuong H, Lanken PN, Aplenc
12 R, Yamamoto T, Yamamoto M, Cho HY, Kleeberger SR. Functional
13 polymorphisms in the transcription factor NRF2 in humans increase the risk
14 of acute lung injury. *FASEB J* 21: 2237-2246, 2007.
- 15 22. Michalopoulos GK and DeFrances MC. Liver regeneration. *Science* 276:
16 60-66, 1997.
- 17 23. Moscat J and Diaz-Meco MT. Feedback on fat: p62-mTORC1-autophagy
18 connections. *Cell* 147: 724-727, 2011.

- 1 24. Murata H, Yagi T, Iwagaki H, Ogino T, Sadamori H, Matsukawa H, Umeda Y,
2 Haga S, Takaka N and Ozaki M. Mechanism of impaired regeneration of
3 steatotic liver in mouse partial hepatectomy model. *J Gastroenterol*
4 *Hepatol* 22: 2173-2180, 2007.
- 5 25. Murata S, Hashimoto I, Nakano Y, Myronovych A, Watanabe M and Ohkohchi
6 N. Single administration of thrombopoietin prevents progression of liver
7 fibrosis and promotes liver regeneration after partial hepatectomy in
8 cirrhotic rats. *Ann Surg* 248: 821-828, 2008.
- 9 26. Niitsu Y, Hakamata M, Goto Y, Higashi T, Shoji M, Sugai T, Umezawa K.
10 Chemoenzymatic synthesis of
11 (2R,3R,4R)-dehydroxymethylepoxyquinomicin (DHMEQ), a new activator of
12 antioxidant transcription factor Nrf2. *Org Biomol Chem* 9: 4635-4641, 2011.
- 13 27. Nogalska A, Terracciano C, D'Agostino C, King Engel W, Askanas V.
14 p62/SQSTM1 is overexpressed and prominently accumulated in inclusions of
15 sporadic inclusion-body myositis muscle fibers, and can help differentiating
16 it from polymyositis and dermatomyositis. *Acta Neuropathol* 118:407-413,
17 2009.
- 18 28. Okada K, Yanagawa T, Warabi E, Yamastu K, Uwayama J, Takeda K,

- 1 Utsunomiya H, Yoshida H, Shoda J and Ishii T. The alpha-glucosidase
2 inhibitor acarbose prevents obesity and simple steatosis in sequestosome
3 1/A170/p62 deficient mice. *Hepatol Res* 39: 490-500, 2009.
- 4 29. Ozaki M, Haga S and Ozawa T. In vivo monitoring of liver damage using
5 caspase-3 probe. *Theranostics* 2: 207-214, 2012.
- 6 30. Ozaki M, Haga S, Zhang HQ, Irani K and Suzuki S. Inhibition of
7 hypoxia/reoxygenation-induced oxidative stress in HGF-stimulated
8 antiapoptotic signaling: role of PI3-K and Akt kinase upon rac1. *Cell Death*
9 *Differ* 10: 508-515, 2003.
- 10 31. Ramos-Gomez M, Dolan PM, Itoh K, Yamamoto M and Kensler TW.
11 Interactive effects of nrf2 genotype and oltipraz on benzo[a]pyrene-DNA
12 adducts and tumor yield in mice. *Carcinogenesis* 24: 461-467, 2003.
- 13 32. Schmucker DL. Age-related changes in liver structure and function:
14 Implications for disease ? *Exp Gerontol* 40: 650-659, 2005.
- 15 33. Selzner M and Clavien PA. Steatotic liver in liver transplantation and surgery.
16 *Semin Liver Dis* 21: 105-113, 2001.
- 17 34. Shen YF, Tang Y, Zhang XJ, Huang KX, Le WD. Adaptive changes in autophagy
18 after UPS impairment in Parkinson's disease. *Acta Pharmacol Sin* 34:

- 1 667-673, 2013.
- 2 35. Tanoue S, Uto H, Kumamoto R, Arima S, Hashimoto S, Nasu Y, Takami Y,
3 Moriuchi A, Sakiyama T, Oketani M, Ido A and Tsubouchi H. Liver
4 regeneration after partial hepatectomy in rat is more impaired in a
5 steatotic liver induced by dietary fructose compared to dietary fat.
6 *Biochem Biophys Res Commun* 407: 163-168, 2011.
- 7 36. Taub R. Liver regeneration: from myth to mechanism. *Nat Rev Mol Cell Biol*
8 5: 836-847, 2004.
- 9 37. Torbenson M, Yang SQ, Liu HZ, Huang J, Gage W, Diehl AM. STAT-3
10 overexpression and p21 up-regulation accompany impaired regeneration of
11 fatty livers. *Am J Pathol* 161: 155-161, 2002.
- 12 38. Wedemeyer I, Bechmann LP, Odenthal M, Jochum C, Marquitan G, Drebber
13 U, Gerken G, Gieseler RK, Dienes HP and Canbay A. Adiponectin inhibits
14 steatotic CD95/Fas up-regulation by hepatocytes: therapeutic implications
15 for hepatitis C. *J Hepatol* 50: 140-149, 2009.
- 16 39. Yamamoto T, Yoh K, Kobayashi A, Ishii Y, Kure S, Koyama A, Sakamoto T,
17 Sekizawa K, Motohashi H, Yamamoto M. Identification of polymorphisms in
18 the promoter region of the human NRF2 gene. *Biochem Biophys Res*

1 *Commun* 321: 72-79, 2004.

2 40. Zou X, Feng Z, Li Y, Wang Y, Wertz K, Weber P, Fu Y and Liu J. Stimulation of
3 GSH synthesis to prevent oxidative stress-induced apoptosis by
4 hydroxytyrosol in human retinal pigment epithelial cells: activation of Nrf2
5 and JNK-p62/SQSTM1 pathways. *J Nutr Biochem* 23: 994-1006, 2012.

6

1 Figure Legends

2

3 FIG. 1. Although post-PH initial liver regeneration is disturbed in db/db mice,
4 mitotic responses occur equally in lean control and db/db steatotic mice.

5 (A) Lipid content was confirmed by Sudan III stain in the liver of control and

6 db/db mice before PH. Lipid accumulation was more evident in the central

7 area than in the portal area of the db/db mouse liver. (B) Initial liver

8 regeneration was disturbed in the db/db mice after PH. (C) Counts of

9 PCNA-positive hepatocytes in the regenerating liver showed the mitotic

10 response peaked at 36 hr post-PH in lean control and db/db mouse liver. (D)

11 Histological examination (H&E staining) revealed similar mitotic responses at

12 36, 48 and 72 hr post-PH in the liver of lean control and db/db steatotic mice.

13 Scale bar: 50 μ m. Scale bar: 50 μ m. At least 5 mice were used for each

14 experiment (B, C and D). Data are expressed as mean \pm SEM. (E) Expression of

15 cell cycle-associated proteins was examined in the liver. Each blot represents

16 at least three independent experiments. The intensity of each band was

17 quantified by densitometry, showing the chronological relative changes for

18 each protein, and is plotted on the right. The results are expressed as mean \pm

1 SEM of at least three independent experiments; $p < 0.05$ was considered
2 statistically significant (B, C, D and E). Groups without an asterisk (*) were not
3 statistically different (E). (To see this illustration in color, the reader is referred
4 to the Web version of this article at www.liebertonline.com/ars)

5

6 FIG. 2. Steatotic liver shows acute OS and injury immediately after PH.

7 (A) Blood biochemistry shows immediate liver injury after PH in the db/db
8 steatotic mouse liver. Histologically, sporadic necrosis in the peri-portal areas
9 (area surrounded by arrows) and neutrophil infiltration (inset) are observed 24
10 hr after PH. (B) Bio-imaging of liver OS after PH. Emission from the roGFP
11 fluorescent probe was measured directly at the exposed liver surface, and
12 imaged and quantified. Photographs are representative images of the dynamic
13 changes in hepatic redox states (oxidation: green to blue; reduction: orange to
14 red). Early post-PH, OS is observed in the db/db steatotic liver. For each
15 experiment, the intensities of the hepatic roGFP signals are plotted relative to
16 the pre-PH values. (C) Bio-imaging of liver caspase-3 activity after PH. The
17 pcFluc-DEVD probe emitted signals at 4-24 hr post-PH in the remnant liver of
18 db/db mice, with no evident signal in the control liver. Data in the graph are

1 expressed as mean \pm SEM and are expressed relative to the pre-PH control. (D)
2 Apoptotic cell death is markedly increased at 4 hr post-PH in the steatotic liver.
3 Results are expressed as mean \pm SEM of five independent experiments; $p < 0.05$
4 was considered statistically significant (A to D). Each experiment was
5 performed five times and representative photographs are shown (B and C). (To
6 see this illustration in color, the reader is referred to the Web version of this
7 article at www.liebertonline.com/ars)

8

9 FIG. 3. Analysis of signals associated with cell proliferation and survival,
10 apoptosis and oxidative stress in hepatocytes.

11 (A) Protein expression of phospho-Akt, Akt, Fas, FasL and p62/SQSTM1 was
12 analyzed by Western blotting. (B) Anti-oxidant/anti-apoptotic proteins were
13 analyzed in the liver of the control and db/db mice. Each blot represents at
14 least three independent experiments. The intensity of each band was
15 quantified by densitometry and is plotted on the right. The data are expressed
16 as mean \pm SEM relative to the lean control values; $p < 0.05$ was considered
17 statistically significant (A and B). (C) Immunohistochemical study of the
18 p62/SQSTM1 expression in mouse liver, demonstrating the reduced expression

1 of p62/SQSTM1 in the db/db steatotic liver. (P: portal vein; C: central vein) (To
2 see this illustration in color, the reader is referred to the Web version of this
3 article at www.liebertonline.com/ars)

4

5 FIG. 4. Fas/FasL expression, ROS generation and cell death
6 (necrosis/apoptosis) in steatotic hepatocytes.

7 (A) Steatosis was induced in AML12 liver cells by treatment with 1 μ M T0901317
8 for 5 days. Oil red O stain showed a marked and homogenous accumulation of
9 lipid in the treated hepatocytes. Fas and FasL expression was up-regulated in
10 the T0901317-treated steatotic hepatocytes. Scale bar: 50 μ m. Photographs are
11 representative of three independent experiments. The intensities of each band
12 are expressed relative to the intensities of the “Non-treated” band; $p < 0.05$ was
13 considered significant. (B) Cell survival and caspase-3 activity were monitored
14 in the control and steatotic hepatocytes with or without Fas after anti-Fas
15 antibody challenge (0.75 μ g/ml of Jo2). (C) Anti-Fas-induced OS in the
16 steatotic hepatocytes was evaluated using a redox-sensitive roGFP probe. For
17 each group, the experiment was performed three times and the data are
18 expressed relative to values before anti-Fas treatment. (D) The effect of

1 anti-oxidants on necrosis and apoptosis was examined in the steatotic
2 hepatocytes. Anti-oxidants (ascorbic acid, Trolox and NAC) suppressed
3 anti-Fas-induced necrotic death but not apoptotic death in steatotic
4 hepatocytes. Cell apoptosis and necrosis were determined using the
5 fluorogenic probes Caspase-Glo 3/7 and YOYO-1, respectively. For each group,
6 the experiment was performed three times and the data are expressed relative
7 to the values of the untreated cells.

8

9 FIG. 5. Cellular steatosis reduces the expression of p62/SQSTM1, and the
10 expression of FasL and Fas is negatively regulated by p62/SQSTM1,
11 regardless of steatosis.

12 (A) Adipogenesis assay of p62/SQSTM1-ablated AML12 cells. Ablation of
13 p62/SQSTM1 alone did not induce steatosis in AML12 liver cells. Untreated
14 AML12 cells were used as a negative control, T0901317-treated AML12 cells as a
15 positive control. (B) mRNA and protein expression levels of p62/SQSTM1 are
16 reduced in steatotic hepatocytes and steatotic liver. The intensity of each band
17 was quantified and normalized to control hepatocytes by densitometry and is
18 plotted on the right; $p < 0.05$ was considered statistically significant. (C)

1 Ablation of p62/SQSTM1 alone induced FasL and Fas expression in AML12
2 hepatocytes, even without steatosis. (D) Overexpression of p62/SQSTM1 in both
3 non-steatotic and steatotic hepatocytes led to a reduction of FasL and Fas.
4 Each experiment was performed at least three times independently, and
5 representative data are shown.

6

7 FIG. 6. Reduced *p62/SQSTM1* leads to a decrease in cellular anti-oxidant
8 properties by suppressing ARE activity.

9 (A) Ablation of *p62/SQSTM1* suppressed *ARE* reporter activity in AML12 cells
10 (left panel), but not *Nrf-2* reporter activity (right panel). ARE reporter activity
11 is expressed relative to the ARE activity against *GAPDH* siRNA.
12 Tert-butylhydroquinone (tBHQ) was used as a positive control of the *ARE*
13 reporter probe. (B) Ablation of *p62/SQSTM1* reduced the expression of
14 anti-oxidant molecules, whereas its overexpression increased the levels of
15 these molecules. Photographs are representative of three independent
16 experiments. The intensity of each band was quantified, normalized against
17 tubulin, and expressed relative to the “no treatment” band (* $p < 0.05$ vs.
18 *GAPDH*-siRNA; ** $p < 0.05$ vs. *GAPDH*-siRNA and vs. *p62*-siRNA). (C) A roGFP

1 redox-sensitive probe showed that ablation of p62/SQSTM1 enhanced
2 hypoxia/reoxygenation-induced ROS generation in AML12 liver cells (4-hr
3 hypoxia, 10-min reoxygenation). (D) Induction of p62/SQSTM1 suppressed
4 Jo2-induced cell death (LDH release and apoptosis) in both control and
5 steatotic hepatocytes (0.75 µg/ml of Jo2, anti-Fas antibody). In each group,
6 the data are expressed as mean ± SEM values relative to the corresponding
7 pre-hypoxic cells; $p < 0.05$ was considered significant.

8

9 FIG. 7. Hepatic transduction of the p62/SQSTM1 gene markedly reduces
10 post-PH liver injury and improves regeneration of the steatotic liver of
11 db/db mice.

12 Post-PH liver injury and regeneration were evaluated in db/db mouse liver
13 transduced with the *p62/SQSTM1* gene. (A) Western blot analysis shows the
14 robust induction of p62/SQSTM1 protein in the liver of db/db mice by
15 hydrodynamic injection of the *p62/SQSTM1* gene (*p-p62*). Plasmid DNA of
16 *luciferase* gene was injected as a control (*p-Luc*). (B) Serum levels of AST, ALT
17 and LDH were reduced at 12 hr post-PH in the *p62/SQSTM1*-transduced
18 steatotic liver, as compared with the luciferase-transduced control liver. (C)

1 Expression of PCNA and cyclinD1 protein peaked at 48 hr post-PH similarly in
2 both the control and p62/SQSTM1-expressing liver; also, there was no
3 significant difference in the mitotic index and the counts of PCNA-positive
4 hepatocytes between the two types of liver. (D) Liver regeneration recovered
5 partially, but not completely, after transduction of p62/SQSTM1 into the
6 steatotic liver. At least 5 mice were used for each experiment. Data are
7 expressed as mean \pm SEM; $p < 0.05$ was considered statistically significant.

8

9 FIG. 8. High fat diet (HFD) induces liver steatosis and p62/SQSTM1 reduction
10 in mouse liver, leading to an increase in FasL/Fas and a decrease in
11 anti-oxidant/-apoptotic molecules.

12 (A) HFD feeding made hepatocytes sufficiently steatotic in the liver tissue (H&E
13 staining). Photographs are representative of three independent experiments.


14 NC: Normal chow. (B) Western blot analysis showed reduced expression of
15 p62/SQSTM1, increased FasL/Fas, and decreased anti-oxidant (catalase,
16 MnSOD, Ref-1) and anti-apoptotic (Bcl-2, Bcl-xL, FLIP) molecules. Each blot
17 represents at least three independent experiments. The intensity of each band
18 was quantified, normalized against GAPDH, and expressed relative to the “NC

1 mouse liver” band.


2

3 FIG. 9. Schematic showing the pivotal role of p62/SQSTM1 in injury of the
4 steatotic liver.


5 Steatosis-associated reduction of p62/SQSTM1 induces FasL/Fas in the steatotic
6 liver. Together with exogenous FasL, this directly causes post-PH necrotic and
7 apoptotic acute liver injury in, respectively, a redox-dependent and
8 redox-independent manner. Reduction of p62/SQSTM1 reduces Keap-1/Nrf-2
9 binding, which suppresses the expression of anti-oxidant molecules (catalase,
10 MnSOD, Ref-1, HO-1, TRX, GPx) and therefore makes the liver susceptible to OS.
11 Furthermore, the hypo-responsiveness of Akt enhances necrotic and apoptotic
12 injury, together with the reduced expression of anti-oxidant molecules and
13 anti-apoptotic molecules (Bcl-2, Bcl-xL, FLIP), respectively. These mechanisms
14 may collectively underlie steatosis-associated liver injury in the mouse. (To see
15 this illustration in color, the reader is referred to the Web version of this article,
16 at www.liebertonline.com/ars)




190x254mm (300 x 300 DPI)




190x254mm (300 x 300 DPI)




190x254mm (300 x 300 DPI)




190x254mm (300 x 300 DPI)





190x254mm (300 x 300 DPI)




190x254mm (300 x 300 DPI)



190x254mm (300 x 300 DPI)




190x254mm (300 x 300 DPI)



190x254mm (300 x 300 DPI)

ARS R3
SUPPLEMENTARY FIG.



SUPPLEMENTARY FIG. Phosphorylation of STAT3 in the steatotic liver was observed even without PH. The intensity of each band was quantified by densitometry. The results are expressed as mean \pm SEM of at least five independent experiments, with a p value of <0.05 considered significant.

254x190mm (300 x 300 DPI)