



Title	Draft Genome Sequence of High-Temperature-Adapted Protochlamydia sp. HS-T3, an Amoebal Endosymbiotic Bacterium Found in Acanthamoeba Isolated from a Hot Spring in Japan
Author(s)	Yamaguchi, Hiroyuki; Matsuo, Junji; Yamazaki, Tomohiro; Ishida, Kasumi; Yagita, Kenji
Citation	Genome Announcements, 3(1), e01507-14 https://doi.org/10.1128/genomeA.01507-14
Issue Date	2015
Doc URL	http://hdl.handle.net/2115/57999
Rights(URL)	http://creativecommons.org/licenses/by/3.0/
Type	article
File Information	Genome Announc-2015-Yamaguchi.pdf



[Instructions for use](#)

Draft Genome Sequence of High-Temperature-Adapted *Protochlamydia* sp. HS-T3, an Amoebal Endosymbiotic Bacterium Found in *Acanthamoeba* Isolated from a Hot Spring in Japan

Hiroyuki Yamaguchi,^a Junji Matsuo,^a Tomohiro Yamazaki,^{a,b} Kasumi Ishida,^{a,b} Kenji Yagita^c

Department of Medical Laboratory Science, Faculty of Health Sciences, Hokkaido University, Kita-ku, Sapporo, Japan^a; Japan Society for the Promotion of Science, Chiyoda-ku, Tokyo, Japan^b; National Institute of Infectious Diseases, Shinjuku-ku, Toyama, Tokyo, Japan^c

Here, we report the draft genome sequence of high-temperature-adapted *Protochlamydia* sp. strain HS-T3, an environmental chlamydia. This bacterium is an amoebal endosymbiont, found in *Acanthamoeba* isolated from a hot spring in Japan. Strain HS-T3 readily grew in mammalian cells at 37°C, a characteristic not previously reported for environmental chlamydiae.

Received 12 December 2014 Accepted 29 December 2014 Published 5 February 2015

Citation Yamaguchi H, Matsuo J, Yamazaki T, Ishida K, Yagita K. 2015. Draft genome sequence of high-temperature-adapted *Protochlamydia* sp. HS-T3, an amoebal endosymbiotic bacterium found in *Acanthamoeba* isolated from a hot spring in Japan. *Genome Announc* 3(1):e01507-14. doi:10.1128/genomeA.01507-14.

Copyright © 2015 Yamaguchi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Hiroyuki Yamaguchi, hiroyuki@med.hokudai.ac.jp.

Chlamydiae is a bacterial phylum and class comprising obligate intracellular bacteria. Two distinct lineages exist, pathogenic and environmental chlamydiae, and this divergence likely occurred 0.7 to 1.4 billion years ago (1). Pathogenic chlamydiae, including *Chlamydia trachomatis* and *Chlamydia pneumoniae*, are well-known human pathogens causing ocular infection resulting in trachoma (preventable blindness) (2), sexually transmitted diseases with an estimated five million new cases annually worldwide (3), and community-acquired pneumonia (4). However, evidence showing an association between environmental chlamydiae and pathogenesis is limited. Pathogenic chlamydiae have successfully adapted to growth in mammalian cells, including human cells (which provide a stable environment with minimal temperature changes) through genome reduction, with an average genomic size of around 1.0 Mb (5). In contrast, environmental chlamydiae have co-evolved with their invertebrate hosts as endosymbionts of lower eukaryotes, free-living amoebae such as *Acanthamoeba*, adapting to a less hospitable environment with a relatively low temperature, likely without genome reduction (1, 6). Most environmental chlamydiae therefore have not successfully been cultured in mammalian cells at the general culture temperature of 37°C (7, 8). However, we recently identified an environmental chlamydia, high-temperature-adapted *Protochlamydia* sp. strain HS-T3, which can readily grow in human immortal HEp-2 epithelial cells under normal culture conditions. This strain was identified in *Acanthamoeba* isolated from a hot spring in Kanagawa prefecture, Japan (9).

The draft genome of *Protochlamydia* HS-T3 was obtained using an Illumina HiSeq 2000 sequencer (Illumina, San Diego, CA, USA), with sequencing runs for paired-end sequences. The bacterial DNA libraries were prepared using a TruSeq DNA sample kit (Illumina). The genome was assembled into 34 contigs, ranging in size from 189 to 363,061 bp, using *de novo* sequence assembler software (Velvet, EMBL-EBI) (10). Gene prediction was performed with Rapid Annotation using Subsystem Technology (RAST) (<http://rast.nmpdr.org/>) (11), and functional annotation

was performed using the Kyoto Encyclopedia of Genes and Genomes (KEGG) (<http://www.genome.jp/kegg/>) (12). The sequencing runs and read assembly of the libraries were carried out by Hokkaido System Science (Sapporo, Japan).

The draft genome sequence of *Protochlamydia* sp. HS-T3 was 2,308,589 bp (2.3 Mb) with a G+C content of 38.74% and 950-fold genome coverage. The genome sequence contained 2,259 coding sequences, 40 tRNAs, and 2 ribosomal RNAs. Neighbor analysis using RAST revealed that the closest neighbor of *Protochlamydia* HS-T3 was *Protochlamydia amoebophila* UWE25 (NC_005861.1) (6), which is unable to grow in cells derived from higher forms of life, such as insects (13). Similarly to *Protochlamydia* UWE25, *Protochlamydia* HS-T3 was found by KEGG analysis to possess central carbon metabolic pathways (e.g., the Embden-Meyerhof pathway, tricarboxylic acid cycle, pentose phosphate pathway, Entner-Doudoroff pathway, and fatty acid biosynthesis) and type III general secretion gene clusters but lacked the genes encoding the type IV secretion system. In addition, 1,139 hypothetical proteins and 8 unknown proteins were identified.

Nucleotide sequence accession numbers. The draft genome sequence of *Protochlamydia* HS-T3 has been deposited at DDBJ/EMBL/GenBank under the accession numbers **BBPT01000001** through **BBPT01000034** (34 entries). The version described in this paper is the first version.

ACKNOWLEDGMENTS

This study was supported by grants-in-aid from the Japan Society for the Promotion of Science (JSPS) Fellows (13J04219) and Scientific Research from KAKENHI, grant numbers 21590474, 24659194, and 24117501 (Innovation Areas: Matryoshka-type evolution). Those funding this project played no role in study design, data collection or analysis, the decision to publish, or preparation of the manuscript.

We thank the staff at the Department of Medical Laboratory Science, Faculty of Health Sciences, Hokkaido University, for their assistance throughout this study.

REFERENCES

- Horn M. 2008. Chlamydiae as symbionts in eukaryotes. *Annu Rev Microbiol* 62:113–131. <http://dx.doi.org/10.1146/annurev.micro.62.081307.162818>.
- Bhosai SJ, Bailey RL, Gaynor BD, Lietman TM. 2012. Trachoma: an update on prevention, diagnosis, and treatment. *Curr Opin Ophthalmol* 23:288–295. <http://dx.doi.org/10.1097/ICU.0b013e32835438fc>.
- Da Ros CT, Schmitt Cda S. 2008. Global epidemiology of sexually transmitted diseases. *Asian J Androl* 10:110–114. <http://dx.doi.org/10.1111/j.1745-7262.2008.00367.x>.
- Burillo A, Bouza E. 2010. *Chlamydophila pneumoniae*. *Infect Dis Clin North Am* 24:61–71. <http://dx.doi.org/10.1016/j.idc.2009.10.002>.
- Subtil A, Dautry-Varsat A. 2004. *Chlamydia*: five years A.G. (after genome). *Curr Opin Microbiol* 7:85–92. <http://dx.doi.org/10.1016/j.mib.2003.12.012>.
- Horn M, Collingro A, Schmitz-Esser S, Beier CL, Purkhold U, Fartmann B, Brandt P, Nyakatura GJ, Droege M, Frishman D, Rattai T, Mewes HW, Wagner M. 2004. Illuminating the evolutionary history of chlamydiae. *Science* 304:728–730. <http://dx.doi.org/10.1126/science.1096330>.
- Hayashi Y, Nakamura S, Matsuo J, Fukumoto T, Yoshida M, Takahashi K, Mizutani Y, Yao T, Yamaguchi H. 2010. Host range of obligate intracellular bacterium *Parachlamydia acanthamoebae*. *Microbiol Immunol* 54:707–713. <http://dx.doi.org/10.1111/j.1348-0421.2010.00265.x>.
- Matsuo J, Hayashi Y, Nakamura S, Sato M, Mizutani Y, Asaka M, Yamaguchi H. 2008. Novel *Parachlamydia acanthamoebae* quantification method based on coculture with amoebae. *Appl Environ Microbiol* 74:6397–6404. <http://dx.doi.org/10.1128/AEM.00841-08>.
- Sampo A, Matsuo J, Yamane C, Yagita K, Nakamura S, Shouji N, Hayashi Y, Yamazaki T, Yoshida M, Kobayashi M, Ishida K, Yamaguchi H. 2014. High-temperature adapted primitive *Protochlamydia* found in *Acanthamoeba* isolated from a hot spring can grow in immortalized human epithelial HEp-2 cells. *Environ Microbiol* 16:486–497. <http://dx.doi.org/10.1111/1462-2920.12266>.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
- Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. 1999. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* 27:29–34. <http://dx.doi.org/10.1093/nar/27.1.29>.
- Sixt BS, Hiess B, König L, Horn M. 2012. Lack of effective anti-apoptotic activities restricts growth of *Parachlamydiaceae* in insect cells. *PLoS One* 7:e29565. <http://dx.doi.org/10.1371/journal.pone.0029565>.