




Title	Aflibercept Traps Galectin-1, an Angiogenic Factor Associated with Diabetic Retinopathy
Author(s)	Kanda, Atsuhiro; Noda, Kousuke; Saito, Wataru; Ishida, Susumu
Citation	Scientific reports, 5, 17946 <a href="https://doi.org/10.1038/srep17946">https://doi.org/10.1038/srep17946</a>
Issue Date	2015-12-09
Doc URL	<a href="http://hdl.handle.net/2115/60583">http://hdl.handle.net/2115/60583</a>
Rights(URL)	<a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a>
Type	article
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	srep17946-s1.pdf (Supplementary Information)



[Instructions for use](#)

# Aflibercept Traps Galectin-1, an Angiogenic Factor Associated with Diabetic Retinopathy

Atsuhiko Kanda, Kousuke Noda, Wataru Saito and Susumu Ishida



**B**


Identified protein	Gene name	Molecular mass (Da)	No. of peptides	Score	Sequence coverage (%)	Calculated pI value	p value	Accession number
Galectin-1	<i>Lgals1</i>	14,857	16	249	80	5.14	< 0.05	P11762

## Supplementary Figure S1. Molecular binding of aflibercept with galectin-1.

Rat retinal endothelial cell (TR-iBRB2) extracts were applied with aflibercept- or normal IgG-immobilized protein G beads. **A**, The eluted proteins were separated by SDS-PAGE and visualized with silver staining. A single protein band around 14 kDa was detected (red arrow). **B**, Summary of identified aflibercept-interacting protein.

## Aflibercept Traps Galectin-1, an Angiogenic Factor Associated with Diabetic Retinopathy

Atsuhiko Kanda, Kousuke Noda, Wataru Saito and Susumu Ishida



### Supplementary Figure S2. Binding affinity of deglycosylated aflibercept with galectin-1.

Sensorgrams obtained using biosensors loaded with PNGase F-treated aflibercept and incubated with different concentrations of galectin-1.