

Biochemical properties of *Paracoccus denitrificans* FnrP: reactions with molecular oxygen and nitric oxide

Jason C. Crack¹ · Matthew I. Hutchings² · Andrew J. Thomson¹ · Nick E. Le Brun¹

Received: 21 October 2015 / Accepted: 23 December 2015
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract In *Paracoccus denitrificans*, three CRP/FNR family regulatory proteins, NarR, NnrR and FnrP, control the switch between aerobic and anaerobic (denitrification) respiration. FnrP is a [4Fe–4S] cluster-containing homologue of the archetypal O₂ sensor FNR from *E. coli* and accordingly regulates genes encoding aerobic and anaerobic respiratory enzymes in response to O₂, and also NO, availability. Here we show that FnrP undergoes O₂-driven [4Fe–4S] to [2Fe–2S] cluster conversion that involves up to 2 O₂ per cluster, with significant oxidation of released cluster sulfide to sulfane observed at higher O₂ concentrations. The rate of the cluster reaction was found to be ~sixfold lower than that of *E. coli* FNR, suggesting that FnrP can remain transcriptionally active under microaerobic conditions. This is consistent with a role for FnrP in activating expression of the high O₂ affinity cytochrome *c* oxidase under microaerobic conditions. Cluster conversion resulted in dissociation of the transcriptionally active FnrP dimer into monomers. Therefore, along with *E. coli* FNR, FnrP belongs to the subset of FNR proteins in which cluster type is correlated with association state. Interestingly, two

key charged residues, Arg140 and Asp154, that have been shown to play key roles in the monomer–dimer equilibrium in *E. coli* FNR are not conserved in FnrP, indicating that different protomer interactions are important for this equilibrium. Finally, the FnrP [4Fe–4S] cluster is shown to undergo reaction with multiple NO molecules, resulting in iron nitrosyl species and dissociation into monomers.

Keywords Fumarate–nitrate reduction regulator · Gene regulation · Iron–sulfur cluster · Oxygen · Nitric oxide

Abbreviations

| | |
|---------------|---|
| DTT | Dithiothreitol |
| GST | Glutathione S-transferase |
| FNR | Fumarate nitrate reduction |
| HEPES | 4-(2-Hydroxyethyl)-1-piperazine-ethanesulfonic acid |
| IPTG | Isopropyl β-D-1-thiogalactopyranoside |
| LC–MS | Liquid chromatography mass spectrometry |
| NO | Nitric oxide |
| PROLI-NONOate | 1-(Hydroxy-NNO-azoxy)-L-proline, disodium salt |
| SOD | Superoxide dismutase |

Electronic supplementary material The online version of this article (doi:10.1007/s00775-015-1326-7) contains supplementary material, which is available to authorized users.

✉ Jason C. Crack
j.crack@uea.ac.uk

✉ Nick E. Le Brun
n.le-brun@uea.ac.uk

¹ Centre for Molecular and Structural Biochemistry, School of Chemistry, University of East Anglia, Norwich NR4 7TJ, UK

² School of Biological Sciences, University of East Anglia, Norwich NR4 7TJ, UK

Introduction

Paracoccus denitrificans is a popular model organism and one of the best studied prokaryotes with respect to respiration. It has a remarkable metabolic versatility allowing it to thrive in aerobic or anaerobic environments [1, 2]. Under anaerobic conditions nitrogen oxides can be utilized as terminal electron acceptors, in place of oxygen, and *P. denitrificans* is one of several organisms for which

the denitrification pathway is well understood. It expresses four essential reductases which sequentially reduce nitrate (*nar*), nitrite (*nir*), nitric oxide (*nor*) and nitrous oxide (*nos*) to dinitrogen [3]. Optimal switching from aerobic respiration to the denitrification pathway is thus a key requirement for this flexibility, and the coordination of the denitrification enzymes is tightly controlled at the transcriptional level [4].

In *E. coli*, the fumarate and nitrate reduction (FNR) transcriptional regulator is responsible for sensing environmental levels of O₂ and controlling the switch to anaerobic nitrate respiration [5]. FNR proteins represent a major sub-group of the cyclic-AMP receptor protein (CRP) family of bacterial transcriptional regulators, and, like CRP, consist of two distinct domains that provide DNA-binding and sensory functions [6–8]. However, unlike *E. coli*, some bacterial species possess multiple members of the FNR protein family [4, 9]; *P. denitrificans* has three major FNR paralogues which coordinate the regulation of the denitrification enzymes [4]. One of these paralogues, NarR, is a nitrate sensor involved in regulating nitrate reductase (*nar*) expression. The second, NnrR, is a heme-based nitric oxide sensor involved in the control of expression of nitrite (*nir*), nitric oxide (*nor*) and nitrous oxide reductases (*nos*) [4]. The third, FnrP, is a true orthologue of *E. coli* FNR as it regulates genes encoding aerobic and anaerobic respiratory enzymes in response to O₂ availability [4, 10–12].

FnrP becomes activated under anaerobic conditions by the insertion of an O₂-labile [4Fe–4S]²⁺ cluster, coordinated by four cysteine residues (Cys14, 17, 25 and 113) present in the N-terminal sensory domain [13]. Cluster incorporation enables the C-terminal DNA-binding domain to recognize specific binding sites located in FnrP regulated promoters. Inactivation of FnrP by O₂ very likely involves oxidative disassembly of the [4Fe–4S] cluster, causing FnrP to adopt a transcriptionally inactive form [14]. In this way, FnrP regulates most of the genes associated with adaptation to the anaerobic environment, including nitrate reductase (*nar*) and nitrous oxide reductases (*nos*) [4, 10, 11], see Fig. 1. For *E. coli* FNR, exposure to O₂ drives a [4Fe–4S]²⁺ to [2Fe–2S]²⁺ cluster conversion, causing the FNR dimer to dissociate into transcriptionally inactive monomers [5]. In contrast, FNR orthologues from *P. putida* (ANR) and *B. subtilis* (FnrB) both form a stable dimer under aerobic and anaerobic conditions, independent of the nature of the iron–sulfur cluster [9, 15].

Many bacteria that carry out anaerobic respiration with nitrate/nitrite as a terminal electron acceptor can produce nitric oxide (NO) endogenously [16, 17]. In *P. denitrificans*, NO production represents a key intermediary step during reduction of nitrate to dinitrogen [3]. NO is a gaseous lipophilic radical that can function as both a signaling molecule and a cytotoxic agent [18]. The latter arises from




Fig. 1 Regulation of denitrification in *P. denitrificans*. Nitrate reductase (*nar*) is co-regulated by NarR and FnrP. Nitrite reductase (*nir*) and nitric oxide reductase (*nor*) are regulated solely by NnrR. Nitrous oxide reductase (*nos*) is co-regulated by FnrP and NnrR. Summarized from references [4, 10, 11, 14]. Small molecule effectors required to activate or inhibit transcription are marked a (+) or (-), respectively. Note: under certain circumstances *nar* will reduce nitrite (NO₂⁻) to nitric oxide (NO)

the reactivity of NO with a variety of important cellular targets [19–21], including iron–sulfur cluster proteins [22]. Several regulatory proteins known to respond to NO contain iron–sulfur clusters as the sensory module [5, 23], and significant progress has been made recently in understanding the reactions of iron–sulfur clusters with NO in regulatory proteins [24, 25]. In the case of FnrP, there is growing in vivo evidence to suggest that, in addition to its primary function as an O₂ sensor, it also plays a role in modulating gene expression in response to NO [10] in a similar way to *E. coli* FNR [25, 26].

Here we report investigations of the biochemical properties of FnrP. We present information on the nature of the iron–sulfur cluster, its reaction with O₂ and NO, as well as the effect of these reactions on its association state. We compare our findings to those reported for *E. coli* FNR as well other FNR orthologues.

Materials and methods

Purification of FnrP

A GST-FnrP fusion protein (Fig. S1) was overproduced in aerobically grown *E. coli* BL21λDE3 harboring pSAD105, induced by the addition of 1 mM IPTG at 37 °C, as previously described [14]. For the in vivo assembly of [4Fe–4S] FnrP, the aeration of cultures harboring pSAD105 was reduced after induction, as previously described [27] to mimic semi-aerobic conditions [11]. FnrP was purified under anaerobic conditions using assay buffer (25 mM HEPES, 2.5 mM CaCl₂, 100 mM NaCl, 100 mM NaNO₃, pH 7.5) as previously described for *E. coli* FNR [28]. FnrP was cleaved from the fusion protein using thrombin (Fig. S1), and, where necessary, the [4Fe–4S] cluster

reconstituted, *in vitro*, as previously described [14, 27, 29], except that a 1 ml Q Sepharose column was used to concentrate the protein and assay buffer containing 500 mM KCl was used to elute the protein. Protein concentration was determined using the method of Bradford (BioRad), with bovine serum albumin as the standard [30]. FnrP iron and acid-labile sulfide content were determined as previously described [31, 32].

Spectroscopy

UV–visible absorbance measurements were made with a Jasco V550 spectrometer. The extinction coefficient for the *E. coli* [4Fe–4S] FNR ($\epsilon_{406\text{ nm}} = 16,200\text{ M}^{-1}\text{ cm}^{-1}$ [28]) was used to calculate the amount of [4Fe–4S] cluster present in FnrP samples. CD spectra were measured with a Jasco J810 spectropolarimeter. For liquid chromatography–mass spectrometry (LC–MS) an aliquot of FnrP (100 μL , 46 μM [4Fe–4S]) was combined with varying aliquots of aerobic (229 μM O_2 , 20 °C) or anaerobic assay buffer (200 μl final volume), and allowed to react for 15 min. Samples were diluted to $\sim 2\text{ }\mu\text{M}$ final concentration, with an aqueous mixture of 1 % (v/v) acetonitrile, 0.3 % (v/v) formic acid, sealed, removed from the anaerobic cabinet and analyzed by an LC–MS instrument consisting of an Ultimate 3000 UHPLC system (Dionex, Leeds, UK), a ProSwift RP-1S column (4.6 \times 50 mm) (Thermo Scientific), and a Bruker microQTOF-QIII mass spectrometer, running Hystar (Bruker Daltonics, Coventry, UK), as previously described [9].

Gel filtration

FnrP samples (loaded at $\sim 28\text{ }\mu\text{M}$ [4Fe–4S]) before and after exposure to O_2 were analyzed by gel filtration under anaerobic conditions using assay buffer and a calibrated Sephacryl S-100HR 16/50 column (GE Healthcare), at a flow rate of 1 ml min^{-1} .

Kinetic measurements

Kinetic measurements were performed under pseudo-first-order conditions (162 μM O_2) at 25 °C by combining varying ratios of aerobic and anaerobic assay buffer (2 ml total volume) with FnrP (8.5 μM [4Fe–4S]). Changes in $A_{406\text{ nm}}$ were used to track cluster conversion. A single or double exponential function, as necessary, was fitted to the data, as previously described [9, 28, 33]. Observed rate constants (k_{obs}) obtained from the fits (in the case of double exponential fits, the rate constant for the first reaction phase was used) were divided by the O_2 concentration, providing an estimate of the apparent second-order rate constant. Kinetic data fitting was performed using Origin (version 8, Origin

Labs). Estimates of errors for rate constants are represented as \pm the standard deviation.

Other analytical methods

FnrP (2 ml) was titrated against varying aliquots of O_2 ($\sim 220\text{ }\mu\text{M}$ dissolved in assay buffer) or NO (as NONOate; Cayman chemicals), using anaerobic cuvettes and gas tight syringes. Stock solutions of the NO donor PROLI-NONOate ($t_{1/2} = 1.5\text{ s}$) were prepared in 50 mM NaOH and quantified optically ($\epsilon_{252\text{ nm}} 8400\text{ M}^{-1}\text{ cm}^{-1}$).

Results and discussion

In vivo purified and *in vitro* reconstituted FnrP binds an identical [4Fe–4S] cluster

Cluster reconstitution of apo-FnrP yielded a straw brown-colored solution with a UV-visible absorbance spectrum (see Fig. 2a) containing a broad shoulder at 420 nm, together with a less well-resolved feature at $\sim 310\text{ nm}$, indicative of the presence of a [4Fe–4S] cluster, as previously described [14]. A cluster-containing form of FnrP was also expressed in *E. coli* cultures (hereafter referred to as native) and isolated under anaerobic conditions. Reconstituted FnrP samples used in this work contained 0.9 (± 0.1) clusters per monomer, based on absorbance measurements, and protein, iron, and acid-labile sulfide determinations. Native samples were found to contain variable amount of the [4Fe–4S] cluster, up to a maximum of ~ 0.7 per monomer.

As FnrP contains seven cysteine residues (Cys8, 14, 17, 25, 28, 113 and 144), it is possible that differing methods (in *vitro* versus *in vivo*) of iron–sulfur cluster assembly might lead to significant differences in the ligation pattern and hence the local environment of the cluster. Since iron–sulfur clusters derive their optical activity from the fold of the protein to which they are ligated, the CD spectrum provides information about the cluster environment. The anaerobic CD spectra of both native and reconstituted [4Fe–4S] FnrP displays positive (+) features at 335 and 440 nm, together with negative (–) features at 300, 375, and 520 nm (see Fig. 2b). The $\Delta\epsilon$ values for the native and reconstituted forms of FnrP are very similar, indicating that the [4Fe–4S] clusters are in essentially identical environments. Hence, reconstituted FnrP was used in subsequent experiments.

Interestingly, the shape of the CD spectrum of FnrP is quite distinct from that of *E. coli* FNR [28]. The major bands at (–)375 nm and (+)440 nm, in FnrP, which originate primarily from S \rightarrow Fe charge transitions, are equivalent to bands observed at (+)380 nm and (+)420 nm for




Fig. 2 Spectroscopic properties of [4Fe-4S] FnrP. **a.** Absorbance spectrum of reconstituted FnrP (48 μ M [4Fe-4S], 81 % cluster loaded), **b** CD spectrum of the same reconstituted FnrP (black) and

[4Fe-4S] FnrP assembled in vivo (gray), for comparison. The buffer was 25 mM HEPES 2.5 mM CaCl_2 100 mM NaCl 100 mM NaNO_3 , 500 mM KCl, pH 7.5

[4Fe-4S] FNR and other [4Fe-4S] containing proteins, such as HiPIP, WhiD and ANR [9, 34, 35]. However, there is no strict correlation between the cluster type and the shape or sign of bands in the CD spectrum; presumably, differences in the cluster binding cavity and/or the geometry of the cluster lead to variation in the CD spectrum.

Reaction of [4Fe-4S] FnrP with O_2 resembles that of *E. coli* FNR

Titration of reconstituted FnrP with aerobic buffer (229 μ M dissolved O_2 at 20 $^\circ\text{C}$) revealed a progressive decrease in $A_{406\text{ nm}}$ and concomitant increase at 530 nm (see Fig. 3a), features typically associated with an FNR-like [4Fe-4S] to [2Fe-2S] conversion [5, 36]. Clear end points to the titration were not observed at $A_{406\text{ nm}}$, possibly due to the slow degradation of the [2Fe-2S] cluster, which contributes to the $\Delta A_{406\text{ nm}}$ readings. Therefore, an identical titration was followed by CD spectroscopy (see Fig. 3b). The well-resolved features of the [4Fe-4S] cluster were gradually replaced with a broad spectrum reminiscent of *E. coli* [2Fe-2S] FNR, containing positive features at ~ 360 and ~ 510 nm, together with a single negative feature at ~ 450 nm, and isosbestic points at 408 and 480 nm. The spectral features of native [4Fe-4S] FnrP behaved in a comparable manner (not shown) during equivalent titrations. A plot of $\text{CD}_{380\text{ nm}} - \text{CD}_{480\text{ nm}}$ versus $[\text{O}_2]:[\text{4Fe-4S}]$ (Fig. 3c) revealed that the reaction reached completion at an $[\text{O}_2]:[\text{4Fe-4S}]$ ratio of ~ 2 ; this behavior is notably different to that of *E. coli* FNR [36], which showed that the reaction was 85 % complete at a ratio of 1 $[\text{O}_2]:[\text{4Fe-4S}]$. By contrast, plotting $\text{CD}_{440\text{ nm}} - \text{CD}_{480\text{ nm}}$ versus $[\text{O}_2]:[\text{4Fe-4S}]$ revealed clear inflection points at both 1 and 2 $[\text{O}_2]:[\text{4Fe-4S}]$ cluster

(see Fig. 3d), suggesting the presence of a meta-stable intermediate.

Analysis of iron-sulfur proteins via liquid chromatography-mass spectroscopy (LC-MS) invariably leads to the loss of the cluster, although previous studies have shown that cysteine persulfides formed during the reaction of the cluster with O_2 remain intact [37]. It was, therefore, of interest to determine if O_2 -mediated cluster conversion in FnrP resulted in retention of cluster sulfides as covalent adducts. [4Fe-4S] FnrP was treated with increasing amounts of O_2 (up to 5 molar equivalents) for 15 min, and analyzed by LC-MS, see Fig. 4a. Even prior to the introduction of O_2 , a small amount of sulfur adduct (at +32 Da of the 28,969 Da peak) was observed. As O_2 was introduced, the relative abundance of these adducts increased, with 1-4 persulfide adducts observed and with the single persulfide adduct at +32 Da the most abundant. Furthermore, the mass of the main protein peak decreased by 1-2 Da, indicating formation of disulfide bonds (Fig. 4b). Treatment with 2 mM DTT, post- O_2 exposure, resulted in the loss of all sulfur adducts and reduction of disulfides.

The [4Fe-4S] cluster of *E. coli* FNR has been shown to react with O_2 via a two-step mechanism (see Scheme 1). Step 1 involves the one electron oxidation of the $[\text{4Fe-4S}]^{2+}$ cluster leading to release of Fe^{2+} to generate a $[\text{3Fe-4S}]^{1+}$ intermediate along with superoxide, which may be recycled back to O_2 by the combined actions of superoxide dismutase (SOD) and catalase [33]. Step 2 involves the conversion of the $[\text{3Fe-4S}]^+$ cluster to the $[\text{2Fe-2S}]^{2+}$ cluster, with the concomitant release of a further iron ion and oxidation of cluster sulfides, which can be stored as cysteine persulfide [5, 33, 37]. In the case of FnrP, the two inflection points observed in the CD intensity plot (Fig. 3d) together with the LC-MS data (Fig. 4) indicate that the availability




Fig. 3 Titration of [4Fe-4S] FnrP with oxygen. **a** Absorbance spectra following addition of O_2 to [4Fe-4S] FnrP (15 μM in cluster). *Black lines* represent an $[O_2]:[4Fe-4S]$ ratio of 0 and 5, respectively. **b** CD spectra following addition of O_2 to [4Fe-4S] FnrP (16 μM in cluster) monitored by circular dichroism. *Black lines* represent $[O_2]:[4Fe-$

4S] ratios of 0, 1, 2 and 3.5, respectively. *Arrows* indicate the direction of movement in response to O_2 . Spectra recorded at intervening $[O_2]:[4Fe-4S]$ ratios are shown in *gray*. **c** $CD_{380-480\text{ nm}}$ and **d** $CD_{440-480\text{ nm}}$ values were normalized and plotted versus the $[O_2]:[4Fe-4S]$ ratio

of O_2 beyond a ratio of 1 per cluster results in increased sulfide oxidation/persulfide formation. This, causes further changes in the CD spectrum, up to a ratio of 2 $[O_2]:[4Fe-4S]$, with little further effect beyond this, suggesting that persulfidation of cluster-coordinating Cys residues occurs mostly in the 1–2 $[O_2]:[4Fe-4S]$ range.

FnrP reacts more slowly than *E. coli* FNR with O_2 in vitro




The first clue that [4Fe-4S] FnrP may be less sensitive to O_2 was the isolation of a straw brown protein from semi-aerobic cultures. Equivalent preparations with *E. coli* FNR typically yield apo-FNR or occasionally [2Fe-2S] FNR [38, 39]; anaerobic cultures are required to obtain [4Fe-4S] FNR [27, 40]. Hence, it was of interest to investigate the kinetics of O_2 -induced [4Fe-4S] FnrP cluster conversion. The $A_{406\text{ nm}}$ decay for FnrP was measured under

pseudo-first-order reaction conditions ($[O_2]:[4Fe-4S]$ ratio of ~ 19). The data were fitted best by a double exponential function (consistent with a two-step reaction [33]) with an observed rate constant (k_{obs}) for the first reaction of $0.0047 (\pm 0.0001) s^{-1}$ (see Fig. 5). Division of k_{obs} by the O_2 concentration (161 μM) provides an estimate of the apparent second-order rate constant, $k = 29 (\pm 1) M^{-1} s^{-1}$. This value establishes that the [4Fe-4S] cluster of FnrP is significantly less reactive with O_2 in vitro than the archetypal *E. coli* FNR cluster [28], and that it more closely resembles the *Pseudomonas putida* FNR proteins PP_3233 and PP_3287 and the previously characterized variant of *E. coli* FNR, FNR-S24F, in its reactivity (see Table 1) [9, 41].

Previous work with *E. coli* FNR showed that replacement of Ser24, located immediately adjacent to the cluster ligand Cys23, by Pro results in significant aerobic FNR activity, indicative of FeS cluster stabilization [41]. Interestingly, FnrP has Pro in the position equivalent to Ser24

Fig. 4 Detection of persulfide species of FnrP by LC-MS.

a ESI-TOF mass spectra of [4Fe-4S] FnrP (2.3 μ M) before and after the addition of increasing amounts of O₂ are shown, as indicated. The peak at 28,969 Da corresponds to the monomer molecular ion peak of FnrP, and the peaks at +32, +64, +96 and +128 Da correspond to the addition of one, two, three and four covalently bound sulfur atoms, respectively, as indicated. Treatment of an aerobic ([O₂]:[4Fe-4S] = 5) sample, post-O₂ exposure, with DTT led to loss of persulfide adducts. **b** During the titration, the mass of the FnrP decreased from 28,969 to 28,967 Da, corresponding to the loss of 2 protons and indicative of O₂-induced disulfide bond formation



Scheme 1 Reaction of *E. coli* [4Fe-4S] FNR with O₂




Fig. 5 Kinetics of O₂-mediated [4Fe-4S] cluster conversion. Kinetics were recorded under pseudo-first-order conditions (8.5 μ M [4Fe-4S], 161.6 μ M O₂) at an [O₂]:[4Fe-4S] ratio of 19 at 25 °C. A double exponential function (black line) was used to fit the data (gray). The rate constants reported in the text from these experiments are mean values with standard errors from two repeats

in FNR (Fig. 6a). Thus, this amino acid residue substitution could, at least partially, account for the lower O₂ reactivity.

We note that amino acid substitutions at other positions next to cluster-coordinating Cys residues are also known to influence the aerobic reactivity of the *E. coli* FNR cluster with O₂. For example, substitutions of Asp22 with Ser or Gly resulted in increased activity under aerobic conditions [42, 43]. The equivalent position in FnrP is occupied by Ile (Fig. 6a) and this difference likely alters the redox properties of the cluster, resulting in a lower reactivity towards O₂.

Under O₂-limiting conditions many bacteria induce high-affinity oxidases, as the ATP yield from oxygen respiration is significantly higher than anaerobic respiration [2]. *P. denitrificans* is no different in this respect, with FnrP activating the expression of the *cbb*₃-type cytochrome *c* oxidase (*cco*) in vivo [2, 11]. We note that *cco* promoter activity increases by eight times during the switch from aerobic to semi-aerobic growth conditions [11]. In contrast, the high-affinity *E. coli* cytochrome *bd*-I (*cydAB*) oxidase (a quinol:O₂ oxidoreductase) is repressed by [4Fe-4S] FNR under anaerobic conditions [44, 45]. Thus, *P. denitrificans* may begin to utilize its high-affinity oxidases at significantly higher environmental O₂ concentrations than its rivals to gain a competitive advantage.

The oligomeric state of FnrP is dependent on the [4Fe-4S] cluster

Under anaerobic conditions *E. coli* FNR acquires a [4Fe-4S] cluster, triggering a conformational change at the dimerization interface that leads to the formation of

Table 1 Kinetic data for the reaction of FNR homologues with O₂

| Species | Protein ID | Second-order rate constant (M ⁻¹ s ⁻¹) | References |
|-------------------------|------------|---|------------|
| <i>E. coli</i> | FNR | 180–200 | [5] |
| | FNR-D154A | 167–172 | [5] |
| | FNR-I151A | 130 | [5] |
| | FNR-S24F | 80 | [41] |
| <i>N. meningitidis</i> | NmFNR | 105 | [55] |
| <i>P. putida</i> | ANR | 280 | [9] |
| | PP_3287 | 55 | [9] |
| | PP_3233 | 38 | [9] |
| <i>P. denitrificans</i> | FnrP | 29 | This study |

homo-dimers (~60 kDa) and site-specific DNA binding [5]. Upon exposure to O₂, a [4Fe–4S] to [2Fe–2S] cluster conversion results in a rearrangement of the dimer interface, leading to monomerization (~30 kDa) [46]. To date, only *E. coli* FNR has been reported to undergo this monomer/dimer transition; with other FNR orthologues tending to remain dimeric irrespective of the cluster [9, 15]. Therefore, the oligomeric state of reconstituted [4Fe–4S] FnrP (~90 % loaded) was examined by analytical gel filtration (see Fig. 7). FnrP has a mass of ~29 kDa, per monomer. In the absence of O₂, the majority of the protein eluted with a relative molecular mass of 51.0 (±1.0) kDa, somewhat lower than, but close to, the expected mass for a homodimer. We note that a slight shoulder, corresponding to a relative molecular mass of ~32 kDa, was also observed. In the presence of O₂, the protein eluted with a relative molecular mass of 28.8 (±0.2) kDa, as expected for monomeric FnrP. These results indicate that cluster conversion drives a rearrangement of the dimer interface, leading to monomerization, as observed for *E. coli* FNR.

Moore and Kiley [46] showed that subunit interactions in *E. coli* FNR arise from a predominantly hydrophobic interface (see Fig. 6b), and that the negatively charged side chain of Asp154 is oriented towards this interface, where inter-subunit charge repulsion inhibits dimerization before cluster acquisition. Insertion of the [4Fe–4S]²⁺ cluster apparently causes shielding of the negative charge, by Ile-151, thereby facilitating dimerization. The recent structure of the *E. coli* FNR-like FNR from *Aliivibrio fischeri* [47] was broadly in agreement with this model, but indicated that hydrophobic contacts made between Ile-151 residues of the two subunits, rather than screening electrostatic repulsion, is key to stabilizing the dimer. Removal of the negatively charged side chain by substitution, such as in FNR-D154A, alleviates the repulsion even in the absence of a cluster, leading to a predominantly dimeric form whether the cluster is present or not. The positive charge of Arg140 is also important for FNR function. Substitution

of this positively charged side chain, such as in FNR-R140A, resulted in an FNR variant with little anaerobic activity, implying a defect in dimerization. In the *A. fischeri* FNR structure, Arg-140 is located near the N terminus of the dimerization helix where no dimerization helix interactions occur. Instead, the Arg-140 sidechain forms a salt bridge with Asp-130 of the other subunit (B helix) and it was suggested that this interaction plays a key role in dimer stability. Interestingly, the double mutant FNR-R140A/D154A regained ≥70 % activity under anaerobic conditions [46]. Assuming residues 130 to 149 participate in subunit interactions, analogous to *E. coli* FNR residues 140 to 159 (Fig. 6a), then Leu134, 141 and 148 presumably form the core of the hydrophobic interface in FnrP (see Fig. 6b). Moreover, charged residues Arg140 and Asp154, found to be important to the monomer/dimer equilibrium in *E. coli* FNR, appear to be replaced by Ala130 and Ala144 in FnrP. Thus, the ability of FnrP to undergo a cluster-induced monomer/dimer transition may depend on the lack of strong electrostatic interactions involving the side chains of Ala130 and Ala144 (*cf* the *E. coli* FNR variant R140A/D154A [46]). The remainder of the residues in FnrP appears to preserve the general nature of the *E. coli* FNR dimerization helix (see Fig. 6). We note that *Bradyrhizobium japonicum* FixK2 also contains Ala residues at equivalent positions to FnrP Ala130 and 144, and that other CRP/FNR family paralogues tend to contain an Ala residue in place of the Arg residue and a hydrophobic/non-charged residue in place of the negatively charged Asp residue (see Fig. S2).

FnrP is a nitric oxide (NO) sensor

In *P. denitrificans* at least three FNR-like transcriptional regulators are involved in the regulation of the four denitrification reductases (see Fig. 1), in response to nitrate, NO or O₂ [4]. In vivo studies have shown that FnrP is involved in the dual control of both the nitrate reductase (*nar*) and the nitrous oxide reductase (*nos*) with NarR and NnrR, respectively, and that factors besides O₂ influence the activity of FnrP [10, 12, 48]. In this respect, we note that the *cco* promoter activity drops to one-third of the semi-aerobic level when cultures are shifted from semi-aerobic to denitrifying conditions and that cultures experience a transitory accumulation of NO under anaerobic conditions prior to the establishment of denitrification. There is also increasing evidence to suggest that FnrP, like *E. coli* FNR, also responds to NO in vivo [10, 11]. Therefore, we investigated the stoichiometry of the reaction of the FnrP cluster with NO by measuring changes in CD and absorbance spectral properties following sequential additions of NO to anaerobic FnrP. A progressive decrease in A_{406 nm} and an increase in A_{360 nm} was observed. The final UV–visible spectrum,





Fig. 6 Sequence alignments of FNR proteins and comparison of the dimerization helices of FNR and FnrP. **a** Sequence alignment of FNR proteins. The dimerization helix is indicated. Cluster-coordinating residues are indicated by *blue arrowheads*. Residues next to cluster-coordinating Cys residues that are important for controlling cluster reactivity are indicated by *green arrowheads*. Two key residues within the helix that are important for *E. coli* FNR association state, Arg140 and Asp154, are indicated by *orange arrowheads*. Proteins are *E. coli* FNR (*EcFNR*), *Paracoccus denitrificans* FnrP (*PdFnrP*), *Pseudomonas aeruginosa* FNR (*PaNnrR*) and *Neisseria meningitidis* FNR (*NmCRP*). The alignment was generated using Clustal Omega [53] and annotated using Genedoc [54]. **b** Helical wheel projection of the dimerization helix of **a** *E. coli* FNR and **b** *P. denitrificans* FnrP, assuming the standard 3.5 residues per turn for a coiled coil. Residues that when substituted by alanine result in a significantly altered activity in *E. coli* FNR, and predicted equivalents in FnrP, occupy positions *a* and *d* of the helical wheel. Adapted from Moore et al. [46]

with a principal absorption band at 360 nm and a shoulder at 430 nm are consistent with previous observations of reaction of an iron–sulfur cluster with NO and indicate the formation of iron–nitrosyl species (see Fig. 8a) [24, 25]. CD signals arising from the [4Fe–4S] cluster decreased almost to zero as the reaction with NO proceeded (Fig. 8b).




Fig. 7 Association state properties of FnrP probed by gel filtration. **a** Chromatogram for reconstituted [4Fe–4S] FnrP in the absence (*upper trace*) or presence of O₂ (*lower trace*). In the absence of O₂ FnrP had an apparent molecular mass of ~51 kDa, indicative of a dimer. A small shoulder, corresponding to a mass of ~32 kDa (monomer) is also observed. In the presence of O₂, FnrP had an apparent mass of ~29 kDa, consistent with monomeric protein. **b** Standard calibration curve for a Sephacryl S100HR column. *Open circles* correspond to standard proteins (BSA, carbonic anhydrase, cytochrome *c*), *gray circles* correspond to the two states of *E. coli* FNR. *Black triangles* correspond to the two states of FnrP. $\ln M_w$, natural log of molecular weight in kDa




Fig. 8 Titration of [4Fe–4S] FnrP with NO. **a** Absorbance spectra of [4Fe–4S] FnrP (18 μ M [4Fe–4S]) following exposure to NO. After each addition of Proli-NONOate, the sample was incubated at an ambient temperature for 5 min prior to spectroscopic measurements. *Black lines* correspond to [NO]:[4Fe–4S] ratios of 0 and 18. Intervening spectra (*gray lines*) correspond to ratios of 4 and 8. **b** CD spectra of an equivalent titration (18 μ M [4Fe–4S]). *Black lines* correspond to [NO]:[4Fe–4S] ratios of 0 and 18, intervening spectra are *gray*. **c** CD_{380 nm} values were normalized and plotted versus the [NO]:[4Fe–4S] ratio. **d** Gel filtration analysis of nitrosylated FnrP. In the presence of NO, FnrP had a molecular mass of ~33 kDa (monomer). A broad shoulder, corresponding to a mass of 50–100 kDa, was also observed. The shape of this shoulder decreased in the presence of DTT, implying that it may arise from a disulfide-bonded FnrP aggregate

A plot of intensity at 380 or 430 nm versus [NO]:[4Fe–4S] (Fig. 8C) revealed that the reaction was complete at a stoichiometry of ≥ 7 NO molecules per cluster, with a clear inflection point at ~ 6 NO, similar to the previously reported nitrosylation of *E. coli* FNR [25].

Analysis by gel filtration of molecular mass changes upon nitrosylation of [4Fe–4S] FnrP revealed a decrease in mass from 51 to 33 kDa (Fig. 8d) for the bulk of the sample. A broad shoulder, corresponding to a mass of ~ 50 –100 kDa, was also observed. The shoulder decreased following treatment with DTT, implying that a small proportion of the sample was in the form of a disulfide-bonded FnrP aggregate. Moreover, LC–MS analysis of nitrosylated FnrP revealed the presence of sulfur adducts (not shown), as previously observed for *E. coli* FNR [25]. We conclude that nitrosylation of FnrP is likely to proceed in a manner similar to that previously reported for *E. coli* FNR, and that, like its reaction with O₂, nitrosylation of FnrP results in dissociation of the dimer into monomers, again similar to *E. coli* FNR.

Conclusion

FNR proteins are global transcription factors that respond to changes in environmental O₂ through the assembly and disassembly of an O₂-sensitive [4Fe–4S] cluster. In the archetypal FNR protein *E. coli* FNR, molecular O₂ brings about conversion of the [4Fe–4S] cluster into a [2Fe–2S] form, thereby triggering conformational changes that initiate monomerization and concomitant loss of sequence-specific DNA binding. In this respect, the *P. denitrificans* regulator FnrP is similar to its *E. coli* counterpart [5]. However, sequence alignment revealed that the residues important for the monomer–dimer equilibrium for FnrP are different to those in *E. coli* FNR and involve fewer charged side chain interactions. Furthermore, we have found that the FnrP cluster is at least 6 times less sensitive to O₂ than *E. coli* FNR. This finding is consistent with the observation of FnrP-activated expression of *cbb*₃-type cytochrome *c* oxidase, nitrate reductase and nitrous oxide reductase under semi-aerobic conditions in vivo [11, 48].

Many transcriptional regulators are known to respond to NO and in *P. denitrificans* the principal regulators are NsrR, NnrR, and FnrP. In other bacterial species NsrR regulates, amongst others, genes involved in detoxification, such as *hmp*, for which NO is a substrate [49, 50]. NnrR principally activates the expression of the nitrite, nitric oxide, and nitrous oxide reductases under anaerobic conditions in response to NO, thereby facilitating denitrification [13, 51, 52]. With regard to denitrification, FnrP co-regulates the expression of the nitrate and nitrous oxide reductases, ensuring their production under

semi-aerobic conditions. We note that nitrate reductase is the most important source of endogenously derived NO during nitrate/nitrite respiration [16]. It is suggested that if the NO detoxification systems are overwhelmed, FnrP will become nitrosylated leading to lowered expression of *cco*, and modulation of the *nar* and *nos* operons (that require [4Fe–4S] FnrP for activation). The concomitant detection of NO by NnrR would then ensure the timely expression of the *nir*, *nor* and *nos* operons, minimizing the transitory nitrosative stress as metabolic modes are switched over in favor of denitrification. Fig. S3 shows a summary of these regulatory systems. Here we have shown that [4Fe–4S] FnrP undergoes a nitrosylation reaction involving multiple NO molecules. This leads to dissociation of FnrP, containing iron–nitrosyl products similar to those observed for other NO-sensing iron–sulfur regulatory proteins, into monomers, providing a mechanistic basis for NO regulation of FnrP.

Acknowledgments This work was supported by the UK's Biotechnology and Biological Sciences Research Council grant BB/L007673/1 to NLB, AJT and JCC. We thank Nick Cull for technical assistance, Dr. Myles Cheesman for access to instrumentation and Prof. Stephen Spiro for pSAD105 encoding GST-FnrP.

Compliance with ethical standards

Dedication The authors dedicate this article to the memory of Bob Williams. They have been strongly influenced by his scientific work and inspired by his spirit. Andrew Thomson and Geoff Moore (see accompanying article on ferritins) studied in Oxford with Bob for their doctoral degrees. At UEA, together with the late Colin Greenwood, they established a multi-disciplinary research unit, the Centre for Metalloprotein Spectroscopy and Biology (CMSB). Nick Le Brun and Jason Crack carried out their doctoral studies in the CMSB under the supervision of Andrew and Geoff. Bob was an adviser to, and a strong supporter of, the CMSB.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Bakken LR, Bergaust L, Liu B, Frostegard A (2012) Regulation of denitrification at the cellular level: a clue to the understanding of N₂O emissions from soils. *Philos Trans R Soc Lond B Biol Sci* 367:1226–1234
- Richardson DJ (2000) Bacterial respiration: a flexible process for a changing environment. *Microbiology* 146:551–571
- Zumft WG (1997) Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev* 61:533–616
- Veldman R, Reijnders WN, van Spanning RJ (2006) Specificity of FNR-type regulators in *Paracoccus denitrificans*. *Biochem Soc Trans* 34:94–96

5. Crack JC, Green J, Thomson AJ, Le Brun NE (2014) Iron-sulfur clusters as biological sensors: the chemistry of reactions with molecular oxygen and nitric oxide. *Acc Chem Res* 47:3196–3205
6. Green J, Scott C, Guest JR (2001) Functional versatility in the CRP-FNR superfamily of transcription factors: FNR and FLP. *Adv Microb Physiol* 44:1–34
7. Matsui M, Tomita M, Kanai A (2013) Comprehensive computational analysis of bacterial CRP/FNR superfamily and its target motifs reveals stepwise evolution of transcriptional networks. *Genome Biol Evol* 5:267–282
8. Shaw DJ, Rice DW, Guest JR (1983) Homology between CAP and Fnr, a regulator of anaerobic respiration in *Escherichia coli*. *J Mol Biol* 166:241–247
9. Ibrahim SA, Crack JC, Rolfe MD, Borrero-de Acuna JM, Thomson AJ, Le Brun NE, Schobert M, Stapleton MR, Green J (2015) Three *Pseudomonas putida* FNR family proteins with different sensitivities to O₂. *J Biol Chem* 290:16812–16823
10. Bergaust L, van Spanning RJ, Frostegard A, Bakken LR (2012) Expression of nitrous oxide reductase in *Paracoccus denitrificans* is regulated by oxygen and nitric oxide through FnrP and NNR. *Microbiology* 158:826–834
11. Otten MF, Stork DM, Reijnders WN, Westerhoff HV, Van Spanning RJ (2001) Regulation of expression of terminal oxidases in *Paracoccus denitrificans*. *Eur J Biochem* 268:2486–2497
12. Van Spanning RJ, De Boer AP, Reijnders WN, Westerhoff HV, Stouthamer AH, Van Der Oost J (1997) FnrP and NNR of *Paracoccus denitrificans* are both members of the FNR family of transcriptional activators but have distinct roles in respiratory adaptation in response to oxygen limitation. *Mol Microbiol* 23:893–907
13. Hutchings MI, Spiro S (2000) The nitric oxide regulated nor promoter of *Paracoccus denitrificans*. *Microbiology* 146:2635–2641
14. Hutchings MI, Crack JC, Shearer N, Thompson BJ, Thomson AJ, Spiro S (2002) Transcription factor FnrP from *Paracoccus denitrificans* contains an iron-sulfur cluster and is activated by anoxia: identification of essential cysteine residues. *J Bacteriol* 184:503–508
15. Reents H, Gruner I, Harmening U, Bottger LH, Layer G, Heathcote P, Trautwein AX, Jahn D, Hartig E (2006) *Bacillus subtilis* Fnr senses oxygen via a [4Fe–4S] cluster coordinated by three cysteine residues without change in the oligomeric state. *Mol Microbiol* 60:1432–1445
16. Ralt D, Wishnok JS, Fitts R, Tannenbaum SR (1988) Bacterial catalysis of nitrosation: involvement of the nar operon of *Escherichia coli*. *J Bacteriol* 170:359–364
17. Watmough NJ, Butland G, Cheesman MR, Moir JW, Richardson DJ, Spiro S (1999) Nitric oxide in bacteria: synthesis and consumption. *Biochim Biophys Acta* 1411:456–474
18. Bruckdorfer R (2005) The basics about nitric oxide. *Mol Aspects Med* 26:3–31
19. Broillet MC (1999) S-nitrosylation of proteins. *Cell Mol Life Sci* 55:1036–1042
20. Kwon YM, Weiss B (2009) Production of 3-nitrosoindole derivatives by *Escherichia coli* during anaerobic growth. *J Bacteriol* 191:5369–5376
21. Weiss B (2006) Evidence for mutagenesis by nitric oxide during nitrate metabolism in *Escherichia coli*. *J Bacteriol* 188:829–833
22. Drapier JC (1997) Interplay between NO and [Fe–S] clusters: relevance to biological systems. *Methods* 11:319–329
23. Fleischhacker AS, Kiley PJ (2011) Iron-containing transcription factors and their roles as sensors. *Curr Opin Chem Biol* 15:335–341
24. Crack JC, Smith LJ, Stapleton MR, Peck J, Watmough NJ, Buttner MJ, Buxton RS, Green J, Oganesyan VS, Thomson AJ, Le Brun NE (2011) Mechanistic insight into the nitrosylation of the [4Fe–4S] cluster of WhiB-like proteins. *J Am Chem Soc* 133:1112–1121
25. Crack JC, Stapleton MR, Green J, Thomson AJ, Le Brun NE (2013) Mechanism of [4Fe–4S](Cys)₄ cluster nitrosylation is conserved amongst NO-responsive regulators. *J Biol Chem* 288:11492–11502
26. Pullan ST, Gidley MD, Jones RA, Barrett J, Stevanin TM, Read RC, Green J, Poole RK (2007) Nitric oxide in chemostat-cultured *Escherichia coli* is sensed by Fnr and other global regulators: unaltered methionine biosynthesis indicates lack of S nitrosation. *J Bacteriol* 189:1845–1855
27. Crack JC, Green J, Thomson AJ, Le Brun NE (2014) Techniques for the production, isolation, and analysis of iron-sulfur proteins. *Methods Mol Biol* 1122:33–48
28. Crack JC, Gaskell AA, Green J, Cheesman MR, Le Brun NE, Thomson AJ (2008) Influence of the environment on the [4Fe–4S]²⁺ to [2Fe–2S]²⁺ cluster switch in the transcriptional regulator FNR. *J Am Chem Soc* 130:1749–1758
29. Bates DM, Popescu CV, Khoroshilova N, Vogt K, Beinert H, Munck E, Kiley PJ (2000) Substitution of leucine 28 with histidine in the *Escherichia coli* transcription factor FNR results in increased stability of the [4Fe–4S]²⁺ cluster to oxygen. *J Biol Chem* 275:6234–6240
30. Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254
31. Beinert H (1983) Semi-micro methods for analysis of labile sulfide and of labile sulfide plus sulfane sulfur in unusually stable iron-sulfur proteins. *Anal Biochem* 131:373–378
32. Crack JC, Green J, Le Brun NE, Thomson AJ (2006) Detection of sulfide release from the oxygen-sensing [4Fe–4S] cluster of FNR. *J Biol Chem* 281:18909–18913
33. Crack JC, Green J, Cheesman MR, Le Brun NE, Thomson AJ (2007) Superoxide-mediated amplification of the oxygen-induced switch from [4Fe–4S] to [2Fe–2S] clusters in the transcriptional regulator FNR. *Proc Natl Acad Sci U S A* 104:2092–2097
34. Przysiecki CT, Meyer TE, Cusanovich MA (1985) Circular dichroism and redox properties of high redox potential ferredoxins. *Biochemistry* 24:2542–2549
35. Crack JC, den Hengst CD, Jakimowicz P, Subramanian S, Johnson MK, Buttner MJ, Thomson AJ, Le Brun NE (2009) Characterization of [4Fe–4S]-containing and cluster-free forms of *Streptomyces* WhiD. *Biochemistry* 48:12252–12264
36. Crack J, Green J, Thomson AJ (2004) Mechanism of oxygen sensing by the bacterial transcription factor Fumarate-Nitrate Reduction (FNR). *J Biol Chem* 279:9278–9286
37. Zhang B, Crack JC, Subramanian S, Green J, Thomson AJ, Le Brun NE, Johnson MK (2012) Reversible cycling between cysteine persulfide-ligated [2Fe–2S] and cysteine-ligated [4Fe–4S] clusters in the FNR regulatory protein. *Proc Natl Acad Sci USA* 109:15734–15739
38. Popescu CV, Bates DM, Beinert H, Munck E, Kiley PJ (1998) Mössbauer spectroscopy as a tool for the study of activation/inactivation of the transcription regulator FNR in whole cells of *Escherichia coli*. *Proc Natl Acad Sci USA* 95:13431–13435
39. Sutton VR, Stubna A, Patschkowski T, Munck E, Beinert H, Kiley PJ (2004) Superoxide destroys the [2Fe–2S]²⁺ cluster of FNR from *Escherichia coli*. *Biochemistry* 43:791–798
40. Yan A, Kiley PJ (2009) Techniques to isolate O₂-sensitive proteins: [4Fe–4S]-FNR as an example. *Meth Enzymol* 463:787–805
41. Jervis AJ, Crack JC, White G, Artymiuk PJ, Cheesman MR, Thomson AJ, Le Brun NE, Green J (2009) The O₂ sensitivity of the transcription factor FNR is controlled by Ser24 modulating

- the kinetics of [4Fe–4S] to [2Fe–2S] conversion. *Proc Natl Acad Sci USA* 106:4659–4664
42. Melville SB, Gunsalus RP (1990) Mutations in *fnr* that alter anaerobic regulation of electron transport-associated genes in *Escherichia coli*. *J Biol Chem* 265:18733–18736
 43. Kiley PJ, Reznikoff WS (1991) *Fnr* mutants that activate gene expression in the presence of oxygen. *J Bacteriol* 173:16–22
 44. Sawers G (1999) The aerobic/anaerobic interface. *Curr Opin Microbiol* 2:181–187
 45. Tseng CP, Albrecht J, Gunsalus RP (1996) Effect of microaerophilic cell growth conditions on expression of the aerobic (*cyoABCDE* and *cydAB*) and anaerobic (*narGHJI*, *frdABCD*, and *dmsABC*) respiratory pathway genes in *Escherichia coli*. *J Bacteriol* 178:1094–1098
 46. Moore LJ, Kiley PJ (2001) Characterization of the dimerization domain in the FNR transcription factor. *J Biol Chem* 276:45744–45750
 47. Volbeda A, Darnault C, Renoux O, Nicolet Y, Fontecilla-Camps JC (2015) The crystal structure of the global anaerobic transcriptional regulator FNR explains its extremely fine-tuned monomer-dimer equilibrium. *Sci Adv* 1:e1501086
 48. Bouchal P, Struharova I, Budinska E, Sedo O, Vyhliadalova T, Zdrahal Z, van Spanning R, Kucera I (2010) Unraveling an FNR based regulatory circuit in *Paracoccus denitrificans* using a proteomics-based approach. *Biochim Biophys Acta* 1804:1350–1358
 49. Crack JC, Munnoch J, Dodd EL, Knowles F, Al Bassam MM, Kamali S, Holland AA, Cramer SP, Hamilton CJ, Johnson MK, Thomson AJ, Hutchings MI, Le Brun NE (2015) NsrR from *Streptomyces coelicolor* is a nitric oxide-sensing [4Fe–4S] cluster protein with a specialized regulatory function. *J Biol Chem* 290:12689–12704
 50. Filenko N, Spiro S, Browning DF, Squire D, Overton TW, Cole J, Constantinidou C (2007) The NsrR regulon of *Escherichia coli* K-12 includes genes encoding the hybrid cluster protein and the periplasmic, respiratory nitrite reductase. *J Bacteriol* 189:4410–4417
 51. Lee YY (2006) Transcription factor NNR from *Paracoccus denitrificans* is a sensor of both nitric oxide and oxygen: isolation of *nnr** alleles encoding effector-independent proteins and evidence for a haem-based sensing mechanism. *Microbiology* 152:1461–1470
 52. van Spanning RJ, De Boer AP, Reijnders WN, Spiro S, Westerhoff HV, Stouthamer AH, van der Oost J (1995) Nitrite and nitric oxide reduction in *Paracoccus denitrificans* is under the control of NNR, a regulatory protein that belongs to the FNR family of transcriptional activators. *FEBS Lett* 360:151–154
 53. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Soding J, Thompson JD, Higgins DG (2011) Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Sys Biol* 7:539
 54. Nicholas KB, Nicholas HBJ (1997) Genedoc: a tool for editing and annotating multiple sequence alignments (**Distributed by the authors**)
 55. Edwards J, Cole LJ, Green JB, Thomson MJ, Wood AJ, Whittingham JL, Moir JW (2010) Binding to DNA protects *Neisseria meningitidis* fumarate and nitrate reductase regulator (FNR) from oxygen. *J Biol Chem* 285:1105–1112