

# SEPARATING CLUB-GUESSING PRINCIPLES IN THE PRESENCE OF FAT FORCING AXIOMS

DAVID ASPERÓ AND MIGUEL ANGEL MOTA

ABSTRACT. We separate various weak forms of Club Guessing at  $\omega_1$  in the presence of  $2^{\aleph_0}$  large, Martin's Axiom, and related forcing axioms.

We also answer a question of Abraham and Cummings concerning the consistency of the failure of a certain polychromatic Ramsey statement together with the continuum large.

All these models are generic extensions via finite support iterations with symmetric systems of structures as side conditions, possibly enhanced with  $\omega$ -sequences of predicates, and in which the iterands are taken from a relatively small class of forcing notions.

We also prove that the natural forcing for adding a large symmetric system of structures (the first member in all our iterations) adds  $\aleph_1$ -many reals but preserves CH.

## 1. INTRODUCTION

One is sometimes faced with the problem of building a model of set theory satisfying the following two requirements.

- (1)  $2^{\aleph_0} > \aleph_2$  holds in the model.
- (2) Some particular combinatorial principle  $P$  of the form “For all  $x$  there is some  $y$  such that  $Q(x, y)$ ”, where  $Q(x, y)$  is sufficiently absolute, holds in the model. Furthermore, for each  $x$  there is a natural proper forcing adding a  $y$  such that  $Q(x, y)$ . Hence,  $P$  can be forced by means of a countable support iteration of proper forcings, but in the corresponding extension  $2^{\aleph_0} \leq \aleph_2$  necessarily holds.

The method of iterated forcing with finite supports and symmetric systems of submodels as side conditions was developed in [6] in order to resolve the tension between (1) and (2) in various situations (see

---

2010 *Mathematics Subject Classification.* 03E50, 03E57, 03E35, 03E05.

Part of the work on this article was done during the Thematic Program “Forcing and its Applications” at the Fields Institute, Toronto, in 2012. Both authors acknowledge financial support from the Fields Institute.

[6] and [7] for background information). Variants of this method have been subsequently investigated in [7] and [3].

One of the central themes of the present article is the construction of iterations as in [6] where the iterands are chosen from some relatively small class of posets (let us call these constructions of the first type). The other central theme is a new variation of the general method from [6] obtained from associating sequences of predicates to the submodels in the side conditions of the iteration (this gives rise to the constructions of the second type). The main focus in this article is the separation of club-guessing principles at  $\omega_1$  in the presence of forcing axioms for relatively small classes of posets but with respect to large collections of dense sets. The corresponding models are obtained as forcing extensions via constructions of either the first or the second type. Also, using a construction of the first type we answer a question of Abraham–Cummings in the context of polychromatic Ramsey theory ([1]).

The rest of the paper is structured as follows. In the next subsection we prove several implications and non-implications between weak forms of Club Guessing at  $\omega_1$ , and present our main theorems (Theorems 1.13 and 1.14). In Section 2 we take a look at the forcing for adding a symmetric system of submodels by finite conditions. This forcing is either the first step or is subsumed in the first step in all our iterations. We show that this forcing adds  $\aleph_1$ -many reals but preserves CH. In Section 3 we prove Theorems 1.13 and 1.14. Finally, in Section 4 we deal with the Abraham–Cummings question. Most of our notation will be standard (see e.g. [12] or [15]) but we will also be introducing additional pieces of notation as we need them.

**1.1. Weak forms of Club Guessing.** A *ladder system* (also known as a *C-sequence*) is a sequence  $\langle C_\delta \mid \delta \in \text{Lim}(\omega_1) \rangle$ , where each  $C_\delta$  is a club of  $\delta$  of order type  $\omega$ . *Club Guessing* (CG) is the well-known weakening of  $\diamond$  saying that there is a *C-sequence*  $\langle C_\delta \mid \delta \in \text{Lim}(\omega_1) \rangle$  which guesses clubs  $C$  of  $\omega_1$ , in the sense that for every such  $C$  there is some  $\delta$  such that a tail of  $C_\delta$  is contained in  $C$ . In this subsection we focus our attention on certain weakenings of CG. The web of implications between these principles will be immediate. We will then point out how to prove several non-implications between these principles, with a focus on what can be obtained in the presence of forcing axioms for large families of dense sets. Finally we present our main separation theorems, to be proved in Section 3.<sup>1</sup>

---

<sup>1</sup>For further information on these (and other related) club-guessing principles, see for example [11], [10], and [17].

*Kunen's Axiom* (KA), also known as *Interval Hitting Principle* (see for example [10]), is the following statement first considered by Kunen: There is a  $C$ -sequence  $\langle C_\delta \mid \delta \in \text{Lim}(\omega_1) \rangle$  with the property that for every club  $C \subseteq \omega_1$  there is some  $\delta$  such that  $[C_\delta(n), C_\delta(n+1)) \cap C \neq \emptyset$  for co-finitely many  $n < \omega$  (where, here and throughout the paper,  $X(\xi)$  denotes the  $\xi$ -th member of  $X$  if  $X$  is a set of ordinals).

$\mathfrak{U}$  (mho), first defined by Todorćević ([22]), says that there is a sequence of continuous colourings  $g_\delta : \delta \rightarrow \omega$ , for  $\delta \in \text{Lim}(\omega_1)$ ,<sup>2</sup> such that for every club  $C \subseteq \omega_1$  there is some  $\delta$  with  $g_\delta^{-1}(\{n\}) \cap C \neq \emptyset$  for all  $n < \omega$ .

It is clear that CG implies KA and that KA implies  $\mathfrak{U}$ .

Weak Club Guessing (WCG), first defined by Shelah ([19]), says that there is a  $C$ -sequence  $\langle C_\delta \mid \delta \in \text{Lim}(\omega_1) \rangle$  such that every club of  $\omega_1$  has infinite intersection with some  $C_\delta$ . Very Weak Club Guessing (VWCG), also first considered by Shelah, says that there is a set  $\mathcal{X}$  of size  $\aleph_1$  consisting of subsets of  $\omega_1$  of order type  $\omega$  such that every club of  $\omega_1$  has infinite intersection with some member of  $\mathcal{X}$ .

One can weaken VWCG even further: Given a cardinal  $\lambda \geq \aleph_1$ ,  $\text{VWCG}_\lambda$  says that there is a set  $\mathcal{X}$  of size at most  $\lambda$  consisting of subsets of  $\omega_1$  of order type  $\omega$  and such that every club of  $\omega_1$  has infinite intersection with some member of  $\mathcal{X}$ .  $\neg \text{VWCG}_\lambda$  is called  $(*)_\lambda^\omega$  in [7] (Definition 1.10).

Obviously WCG implies VWCG and  $\text{VWCG}_\lambda$  implies  $\text{VWCG}_\mu$  whenever  $\aleph_1 \leq \lambda < \mu$ .

By a result of Shelah,  $\neg \text{WCG}$  is compatible with CH ([20]; see also [19]). On the other hand, CH obviously implies VWCG. In fact we have the following.<sup>3</sup>

**Fact 1.1.** (*Hrušák*)  $\mathfrak{b} \leq \lambda$  implies  $\text{VWCG}_\lambda$ .

*Proof.* Let  $\lambda' \leq \lambda$  and let  $(r_\alpha \mid \alpha < \lambda')$  be such that  $\{r_\alpha\}_{\alpha < \lambda'}$  is an  $<^*$ -unbounded subset of  ${}^\omega\omega$  (where  $f <^* g$  means that  $f(n) < g(n)$  for a tail of  $n$ ). Let  $\langle C_\delta \mid \delta \in \text{Lim}(\omega_1) \rangle$  be a ladder system and let  $h_\delta : \omega \rightarrow \delta$  be a bijection for each  $\delta \in \text{Lim}(\omega_1)$ . For all  $\alpha < \lambda'$  and  $\delta \in \text{Lim}(\omega_1)$  let

$$A_\delta^\alpha = \{C_\delta(n) \mid n < \omega\} \cup \bigcup \{(h_\delta^{-1}[0, r_\alpha(n))) \setminus C_\delta(n) \mid n < \omega\}$$

It is easy to check that for all  $\alpha, \delta$ ,  $A_\delta^\alpha$  has order type  $\omega$  and  $\sup(A_\delta^\alpha) = \delta$ . Given a club  $C \subseteq \omega_1$ , let  $\delta \in C$  be a limit point of  $C$  and let

<sup>2</sup>Where  $\delta$  and  $\omega$  are both endowed with the order topology.

<sup>3</sup>Thanks to M. Hrušák for pointing this out in the case  $\lambda = \aleph_1$  (the proof for general  $\lambda$  is the same).

$g_{C,\delta} : \omega \rightarrow \omega$  be the function given by

$$g_{C,\delta}(n) = \min\{m \mid h_\delta(m) \in C \setminus C_\delta(n)\}$$

Now let  $\alpha < \aleph'$  be such that  $r_\alpha(n) > g_{C,\delta}(n)$  for infinitely many  $n$ . It then follows that  $|A_\delta^\alpha \cap C| = \aleph_0$ .  $\square$

Let us consider the following notion of rank (see e.g. [5] and [6]).

**Definition 1.2.** Given a set  $X$  and an ordinal  $\delta$ , we define the *Cantor-Bendixson rank of  $\delta$  with respect to  $X$* ,  $\text{rank}(X, \delta)$ , by specifying that

- $\text{rank}(X, \delta) \geq 1$  if and only if  $\delta$  is a limit point of ordinals in  $X$ .
- If  $\mu > 1$ ,  $\text{rank}(X, \delta) \geq \mu$  if and only if for every  $\eta < \mu$ ,  $\delta$  is a limit of ordinals  $\epsilon$  with  $\text{rank}(X, \epsilon) \geq \eta$ .

Let us call an ordinal  $\delta$  *perfect* if  $\text{rank}(\delta, \delta) = \delta$ .

Given any uncountable cardinal  $\kappa$ , the set of perfect ordinals in  $\kappa$  form a club of order type  $\kappa$ . Hence, if  $\kappa \geq \omega_1$  is a regular cardinal and  $C \subseteq \kappa$  is a club, the set of perfect ordinals  $\delta$  in  $\kappa$  which are fixed points of the enumerating function of  $C$  forms a club of  $\kappa$ .

Given an ordinal  $\tau$ , we will say that a set  $X$  of ordinals is  $\tau$ -*thin* in case  $\text{rank}(X, \delta) \leq \tau$  for all ordinals  $\delta$ . One can strengthen  $\neg \text{VWCG}_\lambda$  even further in the following way.

**Definition 1.3.** ([7]) Given ordinals  $\tau$  and  $\lambda$ ,  $\tau < \omega_1$ ,  $(\cdot)_\lambda^\tau$  is the following statement: For every sequence  $(X_i)_{i < \lambda}$ , if  $X_i$  is a  $\tau$ -thin subset of  $\omega_1$  for all  $i < \lambda$ , then there is a club  $C \subseteq \omega_1$  such that  $|C \cap X_i| < \aleph_0$  for all  $i$ .

One can define the ‘strong’ form of the (weak) guessing principles we have been looking at by requiring that the relevant guessing occurs on a club of  $\delta$ ’s. For example, *Strong Club Guessing (Strong CG)* is the statement that there is a ladder system  $\langle C_\delta \mid \delta \in \text{Lim}(\omega_1) \rangle$  such that for every club  $C \subseteq \omega_1$ ,  $|C_\delta \setminus C| < \aleph_0$  for club-many  $\delta \in \text{Lim}(\omega_1)$ . Similarly we can define strong KA, strong  $\mathcal{U}$ , strong Weak Club Guessing, and so on. In particular, a strong  $\mathcal{U}$ -sequence is a sequence of continuous colourings  $g_\delta : \delta \rightarrow \omega$ , for  $\delta \in \text{Lim}(\omega_1)$ , such that for every club  $C \subseteq \omega_1$  there are club-many  $\delta < \omega_1$  with  $g_\delta^{-1}(\{n\}) \cap C \neq \emptyset$  for all  $n < \omega$ , and a strong WCG-sequence is a ladder system  $(C_\delta : \delta \in \text{Lim}(\omega_1))$  such that for every club  $C \subseteq \omega_1$  there are club-many  $\delta$  such that  $C_\delta \cap C$  is infinite. Note the following facts, which will be used in Section 3. In the statement of these facts, and throughout the paper, we adopt the convention of denoting by  $\delta_N$  the ordinal  $N \cap \omega_1$  if  $N$  is a set such that  $N \cap \omega_1$  is an ordinal.

**Fact 1.4.** *If  $N$  is a countable elementary substructure of  $H(\omega_2)$  containing a strong  $\mathcal{U}$ -sequence  $(g_\delta : \delta \in \text{Lim}(\omega_1))$ , then for every club  $C \subseteq \omega_1$  in  $N$  and every  $n \in \omega$  there is some  $\varepsilon \in C \cap \delta_N$  such that  $g_{\delta_N}(\varepsilon) = n$ .*

**Fact 1.5.** *If  $N$  is a countable elementary substructure of  $H(\omega_2)$  containing a strong WCG-sequence  $(C_\delta : \delta \in \text{Lim}(\omega_1))$ , then for every club  $C \subseteq \omega_1$  in  $N$ ,  $C_{\delta_N} \cap C$  is infinite.*

Of course Strong  $P$  implies  $P$  for all these guessing principles  $P$ , but the reverse implications do not hold. Also, Strong  $P_1$  implies Strong  $P_0$  whenever  $P_1$  implies  $P_0$ .<sup>4</sup> In particular, strong CG implies both strong  $\mathcal{U}$  and strong WCG. These strong guessing principles are consistent:

**Fact 1.6.** *(Folklore) CH + Strong CG can always be forced.*

*Proof.* Assume CH and let  $\mathcal{P}$  be the forcing for adding a  $C$ -sequence  $\vec{C}$  by initial segments. In  $V^{\mathcal{P}}$ , let  $\dot{Q}$  be a countable support iteration shooting all relevant clubs so as to make  $\vec{C}$  strongly club-guessing. It is easy to check that  $\mathcal{P} * \dot{Q}$  is a proper poset not adding reals and that after forcing with  $\dot{Q}$  over  $V^{\mathcal{P}}$ , the generic  $\vec{C}$  becomes strongly club-guessing.  $\square$

We fix some notation regarding forcing axioms: Given a partial order  $\mathcal{P}$  and a cardinal  $\lambda$ ,  $\text{FA}(\mathcal{P})_\lambda$  means: For every collection  $\{D_i \mid i < \lambda\}$  of dense subsets of  $\mathcal{P}$  there is a filter  $G \subseteq \mathcal{P}$  such that  $G \cap D_i \neq \emptyset$  for all  $i < \lambda$ . Given a class  $\Gamma$  of partial orders and a cardinal  $\lambda$ ,  $\text{FA}(\Gamma)_\lambda$  means  $\text{FA}(\mathcal{P})_\lambda$  for every  $\mathcal{P} \in \Gamma$ .

PFA (i.e.,  $\text{FA}(\{\mathcal{P} \mid \mathcal{P} \text{ proper}\})_{\aleph_1}$ ), and in fact BPFA, implies both  $\neg \text{VWCG}$  and  $\neg \mathcal{U}$  (by a standard argument using a natural poset for adding, by finite approximations, a club destroying the relevant guessing sequence).<sup>5</sup> On the other hand, every club of  $\omega_1$  in every c.c.c. extension contains a club in  $\mathbf{V}$ . In particular, all these guessing principles  $P$  are preserved by c.c.c. forcing, and so they are consistent with  $2^{\aleph_0}$  large. In particular, no forcing axiom  $\text{MA}_\lambda$  implies the negation of even Strong CG. Of course  $\text{MA}_{\omega_1}$  implies neither VWCG nor  $\mathcal{U}$  since  $\text{MA}_{\omega_1}$  follows from BPFA.

H. Sakai shows in [18] that the finite-support product  $\text{Add}(\omega, \theta)$  of  $\theta$  copies of Cohen forcing always preserves  $\neg \text{CG}$ . A refined version of

<sup>4</sup>One should be careful: Even if  $\diamond$  implies CG,  $\diamond^+$  (which is a ‘weakly strong’ form of  $\diamond$ ) does not imply Strong CG. This is a result of Ishii–P. Larson ([11]).

<sup>5</sup>We will deal with these posets in Section 3. There is also a (more) natural proper poset for adding, by initial segment, a club killing a potential  $\mathcal{U}$ -sequence, but in order to kill a VWCG-sequence one seems to need finite approximations.

this argument shows that  $\text{Add}(\omega, \theta)$  preserves also  $\neg\text{KA}$  (cf. Lemma 1.12). On the other hand, it is not hard to see that Cohen forcing adds both a WCG–sequence and a  $\mathcal{U}$ –sequence (cf. [13]). A consequence of these two facts together is that one can always force  $2^{\aleph_0}$  large +  $\text{FA}(\text{Add}(\omega, \lambda))_\mu$  for all  $\lambda, \mu < 2^{\aleph_0} + \neg\text{CG} + \text{WCG} + \mathfrak{b} = \omega_1 + \text{KA}$ . For this, start with a model of  $\text{KA} + \neg\text{CG}$  (cf. [10]) and add many Cohen reals to it. And of course one can also force  $2^{\aleph_0}$  large +  $\text{FA}(\text{Add}(\omega, \lambda))_\mu$  for all  $\lambda, \mu < 2^{\aleph_0} + \neg\text{KA} + \text{WCG} + \mathcal{U} + \mathfrak{b} = \omega_1$ : Start with a model of  $\neg\text{KA}$  and add many Cohen reals.

A forcing notion  $\mathcal{P}$  is  ${}^\omega\omega$ –bounding if any  $f : \omega \rightarrow \omega$  in any extension by  $\mathcal{P}$  is dominated by some  $g : \omega \rightarrow \omega$  in the ground model. It is not difficult to see that any c.c.c.  ${}^\omega\omega$ –bounding forcing preserves  $\neg\text{WCG}$  and  $\neg\text{VWCG}_\lambda$  for every  $\lambda$ :

**Lemma 1.7.** (*Hrušák*) *Given a limit ordinal  $\alpha < \omega_1$  and  $X \subseteq \alpha$ , a set of ordinals of order type  $\omega$  cofinal in  $\alpha$  in an extension by an  ${}^\omega\omega$ –bounding forcing notion,  $X$  is covered by a set  $Y$  in  $\mathbf{V}$  of order type  $\omega$  and with  $\text{sup}(Y) = \alpha$ . Hence, If  $\langle A_i \mid i < \lambda \rangle$  is a sequence of subsets of  $\omega_1$  of order type  $\omega$  in an extension by a c.c.c.  ${}^\omega\omega$ –bounding forcing notion, then there is a sequence  $\langle B_i \mid i < \lambda \rangle \in \mathbf{V}$  of subsets of  $\omega_1$  of order type  $\omega$  such that  $\text{sup}(B_i) = \text{sup}(A_i)$  and  $|A_i \setminus B_i| < \aleph_0$  for every  $i$ .*

*Proof.* For the first assertion, let  $(\alpha_n)_{n < \alpha} \in \mathbf{V}$  be a strictly increasing sequence converging to  $\alpha$  and let  $(h_n)_{n < \omega} \in \mathbf{V}$  be such that  $h_n : \omega \rightarrow [\alpha_n, \alpha_{n+1})$  is a surjection for every  $n$ . In the extension, let  $f : \omega \rightarrow \omega$  be given by  $f(n) = \min\{m \mid X \cap [\alpha_n, \alpha_{n+1}) \subseteq h_n \text{``}m\}$ , and let  $g : \omega \rightarrow \omega$  be a function in  $\mathbf{V}$  dominating  $f$ . It suffices to define  $Y$  such that  $Y \cap [\alpha_n, \alpha_{n+1}) = \{h_n(0), \dots, h_n(g(n))\}$ . Now we can prove the second assertion by a standard diagonalization argument using the c.c.c. □

It follows of course that we can always force  $2^{\aleph_0}$  large +  $\text{FA}(\mathcal{B}_\lambda)_\mu$  for all  $\lambda, \mu < 2^{\aleph_0} + \neg\text{VWCG}$ , where  $\mathcal{B}_\lambda$  denotes the measure algebra with  $\lambda$ –many generators: Start with a model of  $\neg\text{VWCG}$  and add lots of random reals to it.

At this point one would naturally ask whether  $\text{MA}_\lambda$ , for any  $\lambda > \aleph_1$ , implies any nontrivial weak club–guessing principle.<sup>6</sup> This question was answered negatively in [7]:

---

<sup>6</sup>Note that one cannot hope to use a finite support iteration of c.c.c. forcings to answer this question negatively since every such iteration will force WCG if it has length at least  $\omega$  (see [7]).

**Definition 1.8.** ([7]) A poset  $\mathcal{P}$  has the  $\aleph_{1.5}$ -c.c. if for every regular cardinal  $\lambda \geq |\text{TC}(\mathcal{P})|^+$  there is a club  $D \subseteq [H(\lambda)]^{\aleph_0}$  such that for every finite  $\{N_i : i < n\} \subseteq D$  and every  $p \in \mathcal{P}$ , if  $p \in N_j$  for some  $j$  such that  $\delta_{N_j} = \min\{\delta_{N_i} : i < n\}$ , then there is some condition extending  $p$  and  $(N_i, \mathcal{P})$ -generic for all  $i$ .

Also, given any cardinal  $\lambda$ ,  $\text{MA}_\lambda^{1.5}$  is  $\text{FA}(\{\mathcal{P} \mid \mathcal{P} \text{ has the } \aleph_{1.5}\text{-c.c.}\})_\lambda$ , and  $\text{MA}^{1.5}$  is  $\text{MA}_\lambda^{1.5}$  for all  $\lambda < 2^{\aleph_0}$ .

Every c.c.c. poset has the  $\aleph_{1.5}$ -c.c. and every  $\aleph_{1.5}$ -c.c. poset has the  $\aleph_2$ -c.c. and is proper. Also, for every cardinal  $\lambda$ ,  $\text{MA}_\lambda^{1.5}$  implies  $(\cdot)_\lambda^\tau$  for all  $\tau < \omega_1$ . These facts are proved in [7].

**Theorem 1.9.** ([7]) (CH) Let  $\kappa \geq \omega_2$  be a regular cardinal such that  $\mu^{\aleph_0} < \kappa$  for all  $\mu < \kappa$  and  $\diamond(\{\alpha < \kappa : \text{cf}(\alpha) \geq \omega_1\})$  holds. Then there is a proper forcing notion  $\mathcal{P}$  of size  $\kappa$  with the  $\aleph_2$ -c.c. such that the following statements hold in the generic extension by  $\mathcal{P}$ .

- (1)  $2^{\aleph_0} = \kappa$
- (2)  $\text{MA}^{1.5}$

One prominent  $\aleph_{1.5}$ -c.c. forcing notion is Baumgartner's forcing for adding a club of  $\omega_1$  with finite conditions ([8]). We will denote this poset by  $\mathbb{B}$ .<sup>7</sup> Conditions in  $\mathbb{B}$  are finite functions  $p \subseteq \omega_1 \times \omega_1$  that can be extended to a normal function  $F : \omega_1 \rightarrow \omega_1$ ,<sup>8</sup> and given two conditions  $p_0, p_1$ ,  $p_1$  is stronger than  $p_0$  iff  $p_0 \subseteq p_1$ .  $\mathbb{B}$  is  $\aleph_{1.5}$ -c.c., and in fact is a finitely proper poset of size  $\aleph_1$  (see [6] for the definition of finitely proper). On the other hand, it is nowhere c.c.c. (i.e., it is not c.c.c. below any condition) and, as Zapletal proved in [25], under PFA it is a minimal nowhere c.c.c. poset, in the sense that every nowhere c.c.c. poset adds a generic for  $\mathbb{B}$ . Also, if  $P = \{p_\alpha \mid \alpha < \omega_1\}$  is a proper nowhere c.c.c. forcing notion adding a club  $C \subseteq \omega_1$  such that for all  $\alpha \in C$ ,  $\dot{G} \cap \{p_\beta \mid \beta < \alpha\}$  is generic for  $\{p_\beta \mid \beta < \alpha\}$  (where  $\dot{G}$  denotes the generic filter), then  $\text{RO}(P) = \text{RO}(\mathbb{B})$  ([24]). Let  $\text{FA}(\mathbb{B})$  denote  $\text{FA}(\mathbb{B})_\lambda$  for all  $\lambda < 2^{\aleph_1}$ .

Given a partial order  $\mathcal{P}$ , let  $\mathfrak{m}(\mathcal{P})$  be the minimal size of a family  $\mathcal{D}$  of dense subsets of  $\mathcal{P}$  such that there is no filter  $G \subseteq \mathcal{P}$  intersecting all members of  $\mathcal{D}$ . There is a natural topology on the club filter  $\mathcal{C}_{\omega_1}$  on  $\omega_1$  such that  $\text{cov}(\mathcal{C}_{\omega_1}) = \mathfrak{m}(\mathbb{B})$  if  $\mathfrak{m}(\mathbb{B}) > \aleph_1$  (where  $\mathcal{C}_{\omega_1}$  is endowed with this topology and where for a topological space  $X$ ,  $\text{cov}(X)$ , the

<sup>7</sup>The observations on  $\mathbb{B}$  that follow are borrowed from [4].

<sup>8</sup>A function  $F : \omega_1 \rightarrow \omega_1$  is normal if it is strictly increasing and continuous. There is of course a natural correspondence between clubs of  $\omega_1$  and normal functions  $F : \omega_1 \rightarrow \omega_1$  which sends a club to its enumerating function and a normal function to its range.

covering number of  $X$ , is the minimal size of a collection of nowhere dense subsets of  $X$  whose union is  $X$ ). This is of course the topology whose basis is given by the conditions in  $\mathbb{B}$ ; in other words, a basis for this topology is given by  $\{B_p \mid p \in \mathbb{B}\}$ , where  $B_p$  is the set of all clubs  $C \subseteq \omega_1$  such that  $p \subseteq F$ , where  $F$  is the enumerating function of  $C$ . The proof of this is a standard translation exercise between topological notions and order-theoretical notions, and is identical to the proof that  $\mathfrak{m}(\text{Cohen})$  is the covering number of the meagre ideal for the Baire space. The assumption  $\mathfrak{m}(\mathbb{B}) > \aleph_1$  is needed only for the proof of  $\mathfrak{m}(\mathbb{B}) \leq \mathbf{cov}(\mathcal{C}_{\omega_1})$ : Given  $\kappa < \mathfrak{m}(\mathbb{B})$  and a collection  $\{X_i \mid i < \kappa\}$  of closed nowhere dense subsets of  $\mathcal{C}_{\omega_1}$ ,  $D_i = \{p \in \mathbb{B} \mid B_p \cap X_i = \emptyset\}$  is a dense open subset of  $\mathbb{B}$ . Let  $\{E_\nu \mid \nu < \omega_1\}$  be a set of dense subsets of  $\mathbb{B}$  such that  $\bigcup \text{range}(G)$  is a club of  $\omega_1$  for every filter  $G \subseteq \mathbb{B}$  meeting all  $E_\nu$ . Since  $\mathfrak{m}(\mathbb{B}) > \aleph_1$ , we can find a filter  $G \subseteq \mathbb{B}$  meeting all  $E_\nu$  and all  $D_i$ . It follows then that  $C = \bigcup \text{range}(G)$  is a club of  $\omega_1$  such that  $C \not\subseteq X_i$  of all  $i$ .

Of course there is nothing special about Cohen forcing or Baumgartner's forcing in any of these translations; in fact, a similar characterisation can be always obtained for  $\mathfrak{m}(\mathcal{P})$  for any poset  $\mathcal{P}$ . What is nice about Cohen forcing and Baumgartner's forcing (in the case  $\mathfrak{m}(\mathbb{B}) > \aleph_1$ ) is the appealing appearance of the topological side of the translation.

The assumption  $\mathfrak{m}(\mathbb{B}) > \aleph_1$  in the above characterisation is necessary:

**Fact 1.10.** *KA implies  $\mathbf{cov}(\mathcal{C}_{\omega_1}) = \aleph_0$ .*

*Proof.* Let  $\vec{C} = \langle C_\delta \mid \delta \in \text{Lim}(\omega_1) \rangle$  be a KA-sequence. For every  $n < \omega$ , let  $D_n$  be the set of  $\mathbb{B}$ -conditions  $p$  such that for every  $\delta \in \text{dom}(p)$ , if  $\delta$  is perfect and  $p(\delta) = \delta$ , then there is some  $m \geq n$  and some  $\xi \in \text{dom}(p) \cap \delta$  such that  $p(\xi) < C_\delta(m)$  and  $p(\xi + 1) > C_\delta(m + 1)$ .  $D_n$  is clearly a dense subset of  $\mathbb{B}$ , and therefore the set  $X_n$  of  $C \in \mathcal{C}_{\omega_1}$  such that  $p \not\subseteq \vec{C}$  for any  $p \in D_n$ , where  $\vec{C}$  is the enumerating function of  $C$ , is nowhere dense in  $\mathcal{C}_{\omega_1}$ . Now, if  $C \subseteq \omega_1$  is a club, then there is some  $n < \omega$  such that  $C \in X_k$  for every  $k \geq n$  since  $\vec{C}$  is a KA-sequence.  $\square$

Thus,  $\mathbf{cov}(\mathcal{C}_{\omega_1}) = \aleph_0$  can be regarded as a (very) weak club-guessing principle that might be worth investigating.

As we have seen, all forcing axioms of the form  $\text{MA}_\lambda$  are compatible with even the strongest club-guessing principle we are considering in this paper (namely Strong CG). This is not true for  $\text{FA}(\mathbb{B})_\lambda$ : Let us say that a set  $\mathcal{C}$  of subsets of  $\omega_1$  of order type  $\omega$  is a KA set if for every club  $D \subseteq \omega_1$  there is some  $C \in \mathcal{C}$  such that  $D \cap [C(n), C(n + 1)) \neq \emptyset$



for a tail of  $n < \omega$ . Given a cardinal  $\lambda$ , let  $\text{KA}_\lambda$  be the assertion that there is a KA set of size at most  $\lambda$ . It is not difficult to see that  $\mathbb{B}$  destroys every KA-sequence from the ground model. In particular we obtain the following.

**Proposition 1.11.** *For every cardinal  $\lambda$ ,  $\text{FA}(\mathbb{B})_\lambda$  implies  $\neg \text{KA}_\lambda$ .*

Given a cardinal  $\lambda$ , let  $\text{CG}_\lambda$  denote the weakening of CG saying that there is a set  $\mathcal{C}$  of size at most  $\lambda$  consisting of subsets of  $\omega_1$  of order type  $\omega$  and such that for every club  $D \subseteq \omega_1$  there is some  $C \in \mathcal{C}$  with  $C \subseteq D$ . Of course  $\text{CG}_\lambda$  implies  $\text{KA}_\lambda$ .

By essentially the same argument as in [18] one can show that  $\text{Add}(\omega, \theta)$  preserves  $\neg \text{CG}_\lambda$ . Also, by refining the argument from [18], one can establish the following preservation result ([4]), the proof of which we include here for completeness:

**Lemma 1.12.**  *$\text{Add}(\omega, \theta)$  preserves  $\neg \text{KA}_\lambda$ .*

*Proof.* Let  $X$  be any set and let  $P(X)$  be the set of finite functions  $p \subseteq (X \times \omega) \times 2$  ordered by reverse inclusion. It suffices to show that forcing with  $P(X)$  preserves  $\neg \text{KA}_\lambda$ . Let  $\langle \dot{A}_i \mid i < \lambda \rangle$  be a sequence of  $P(X)$ -names for subsets of  $\omega_1$  of order type  $\omega$ . The first observation is that for every  $i$  there is a countable  $Y_i \subseteq X$  such that  $\dot{A}_i$  is in fact a  $P(Y_i)$ -name and such that for every  $\alpha < \omega_1$  there is some  $p \in P(X)$  such that  $p \Vdash_{P(X)} \check{\alpha} \in \dot{A}_i$  if and only if there is some  $p \in P(Y)$  such that  $p \Vdash_{P(Y)} \check{\alpha} \in \dot{A}_i$ .

For every  $i$  let  $(p_n^i)_{n < \omega}$  be an enumeration of  $P(Y_i)$ . Also, for every  $n < \omega$ , if there is some  $\sigma \in \omega_1$  such that  $p_n^i \Vdash_{P(Y_i)} \text{sup}(\dot{A}_i) = \sigma$ , then let  $\mathcal{X}_n^i$  be a set of pairwise compatible  $P(Y_i)$ -conditions extending  $p_n^i$  and such that  $\{\xi < \sigma \mid p' \Vdash_{P(Y_i)} \xi \in \dot{A}_i \text{ for some } p' \in \mathcal{X}_n^i\}$  is cofinal in  $\sigma$ .<sup>9</sup>

Given any club  $C \subseteq \omega_1$  there are  $i < \lambda$ ,  $\sigma < \omega_1$ ,  $\gamma < \sigma$  and  $n^C < \omega$  such that  $p_{n^C}^i$  forces in  $P(Y_i)$  that  $\text{sup}(\dot{A}_i) = \sigma$  and that  $[\alpha, \alpha'] \setminus C \neq \emptyset$  for even two consecutive points  $\alpha, \alpha'$  of  $\dot{A}_i$  above  $\gamma$ . Let  $n^* < \omega$  be such that the set  $\mathcal{C}$  of clubs  $C$  such that  $n^C = n^*$  is  $\subseteq$ -dense in the set of all clubs of  $\omega_1$ . For every  $i < \lambda$ , if there is a  $\sigma$  such that  $p_{n^*}^i \Vdash_{P(Y_i)} \text{sup}(\dot{A}_i) = \sigma$ , then let  $B_i = \{\xi < \sigma \mid p \Vdash_{P(Y_i)} \xi \in \dot{A}_i \text{ for some } p \in \mathcal{X}_{n^*}^i\}$ .

By  $\subseteq$ -density of  $\mathcal{C}$  it suffices to show that if  $C \in \mathcal{C}$ , then there is some  $i < \lambda$  such that  $p_{n^*}^i \Vdash \text{sup}(\dot{A}_i) = \sigma$  for some  $\sigma$ , and such that  $[\beta, \beta'] \cap C \neq \emptyset$  for every two consecutive points  $\beta < \beta'$  in  $B_i$  above

<sup>9</sup>The introduction of the  $\mathcal{X}_n^i$ 's is the new ingredient with respect to the proof in [18].

some  $\gamma < \sigma$ . But this is true by the previous paragraph and the definition of  $B_i$  since all conditions in  $\mathcal{X}_{n^*}^i$  extend  $p_{n^*}^i$  and are pairwise compatible.  $\square$

It follows that if, say, GCH holds and  $\omega_2 \leq \kappa \leq \lambda$  are successor cardinals, then there is a proper  $\aleph_2$ -c.c. poset producing a model of  $2^{\aleph_0} = \lambda + \text{FA}(\text{Add}(\omega, \mu))_{\lambda'}$  for all  $\lambda', \mu < \lambda + \neg \text{KA}_{\kappa'}$  for any  $\kappa' < \kappa + \text{WCG} + \mathfrak{U} + \mathfrak{b} = \omega_1$ .<sup>10</sup> For this, first force  $2^{\aleph_0} = \kappa + \text{MA}^{1.5}$  by a proper  $\aleph_2$ -c.c. poset as in Theorem 1.9 and then force with  $\text{Add}(\omega, \lambda)$ . The first forcing kills  $\text{KA}_{\kappa'}$  for all  $\kappa' < \kappa$ , and the second forcing preserves  $\neg \text{KA}_{\kappa'}$  for all  $\kappa' < \kappa$  and forces  $2^{\aleph_0} = \lambda + \text{WCG} + \mathfrak{b} = \omega_1$ .

In this paper we focus on the construction of models separating club-guessing principles in which  $2^{\aleph_0}$  is arbitrarily large and in which some relatively large fragment of  $\text{MA}^{1.5}$  (say, comprising  $\text{MA} + \text{FA}(\mathbb{B})$ ) holds. Our main theorems are the following.

**Theorem 1.13.** (CH) *Suppose there is a strong  $\mathfrak{U}$ -sequence  $\vec{\mathcal{G}}$ . Let  $\kappa$  be a regular cardinal such that  $2^{<\kappa} = \kappa$ . Then there exists a proper poset  $\mathcal{P}$  with the  $\aleph_2$ -c.c. such that the following statements hold in  $V^{\mathcal{P}}$ .*

- (1)  $\vec{\mathcal{G}}$  is a  $\mathfrak{U}$ -sequence.
- (2)  $(\cdot)_{\lambda}^{\tau}$  for all  $\tau < \omega_1$  and  $\lambda < 2^{\aleph_0}$ .
- (3) MA
- (4)  $\text{FA}(\mathbb{B})$
- (5)  $2^{\aleph_0} = \kappa$

**Theorem 1.14.** (CH) *Suppose there is a strong WCG-sequence  $\vec{C}$ . Let  $\kappa$  be a regular cardinal such that  $2^{<\kappa} = \kappa$ . Then there exists a proper poset  $\mathcal{P}$  with the  $\aleph_2$ -c.c. such that the following statements hold in  $V^{\mathcal{P}}$ .*

- (1)  $\vec{C}$  is a WCG-sequence.
- (2)  $\neg \mathfrak{U}$
- (3) MA
- (4)  $\text{FA}(\mathbb{B})$
- (5)  $2^{\aleph_0} = \kappa$

By Fact 1.6 and the paragraph before Fact 1.6, the hypotheses of Theorems 1.13 and 1.14 can always be forced.

Both theorems above can be proved starting from the weaker hypothesis that  $\vec{\mathcal{G}}$  (resp.,  $\vec{C}$ ) is defined only on a stationary subset of  $\omega_1$  (for Theorem 1.13 and Theorem 1.14, respectively). Here, the relevant object  $\vec{\mathcal{A}}$  being strongly guessing in the appropriate sense means

<sup>10</sup>This result will be partially improved in Theorem 1.14.

of course that for every club  $C$  the guessing occurs for all ordinals in  $D \cap \text{dom}(\vec{\mathcal{A}})$  for some club  $D \subseteq \omega_1$ .

Finally, we should point out that other separations of consequences of  $\text{MA}^{1.5}$ , in the presence of  $2^{\aleph_0}$  large, have been obtained also by other people using iterations with symmetric systems of structures as side conditions and considering instances from a restricted class of  $\aleph_{1.5}$ -c.c. forcings only (i.e., as in the proof of Theorem 1.13). Specifically, Yorioka ([23]) builds a model of  $\neg\mathfrak{U} + \neg\text{WCG} + \mathfrak{p} = \text{add}(\mathcal{N}) = 2^{\aleph_0} + 2^{\aleph_0}$  large in which there is a destructible  $(\omega_1, \omega_1^*)$ -gap in  $({}^\omega\omega, <^*)$ .

## 2. FORCING WITH SYMMETRIC SYSTEMS

For this section, let  $\kappa$  be a cardinal such that  $\text{cf}(\kappa) \geq \omega_2$ .

**Definition 2.1.** ([6]; cf. [21], [14], [1]) Let  $P \subseteq H(\kappa)$ , and let  $\mathcal{N}$  be a collection of countable subsets of  $H(\kappa)$ . We will say that  $\mathcal{N}$  is a *P-symmetric system* if

- (A) For every  $N \in \mathcal{N}$ ,  $(N, \in, P)$  is an elementary substructure of  $(H(\kappa), \in, P)$ .
- (B) Given  $N, N' \in \mathcal{N}$ , if  $\delta_N = \delta_{N'}$ , then there is a (unique) isomorphism

$$\Psi_{N, N'} : (N, \in, P) \longrightarrow (N', \in, P)$$

Furthermore,  $\Psi_{N, N'}$  is the identity on  $N \cap N'$ .

- (C) For all  $M, N$  in  $\mathcal{N}$ , if  $\delta_M < \delta_N$ , then there is some  $N' \in \mathcal{N}$  such that  $\delta_{N'} = \delta_N$  and  $M \in N'$ .
- (D) For all  $M, N$  and  $N'$  in  $\mathcal{N}$ , if  $M \in N$  and  $\delta_{N'} = \delta_N$ , then  $\Psi_{N, N'}(M) \in \mathcal{N}$ .

In (A) in the above definition, and elsewhere, we will tend to refer to structures  $(N, \in, P \cap N)$  by the simpler expression  $(N, \in, P)$ . Also, given any two structures  $N, N'$ , if there is a (unique) isomorphism between  $(N, \in)$  and  $(N', \in)$ , then we denote this isomorphism by  $\Psi_{N, N'}$ .

Given  $P \subseteq H(\kappa)$ , there is a natural forcing notion  $\mathcal{S}_P$  for adding, by initial approximations, a symmetric  $P$ -system  $\mathcal{N}$  such that  $\bigcup \mathcal{N} = H(\kappa)$ :  $\mathcal{S}_P$  is just the set of finite  $P$ -symmetric systems, ordered by reverse inclusion.

The following lemma is proved in [6] (Lemma 2.3).

**Lemma 2.2.** *Let  $P \subseteq H(\kappa)$ , let  $\mathcal{N}$  be a  $P$ -symmetric system, and let  $N \in \mathcal{N}$ . Then the following holds.*

- (i)  $\mathcal{N} \cap N$  is a  $P$ -symmetric system.
- (ii) If  $\mathcal{W} \subseteq N$  is a  $P$ -symmetric system and  $\mathcal{N} \cap N \subseteq \mathcal{W}$ , then

$$\mathcal{V} := \mathcal{N} \cup \{\Psi_{N, N'}(W) : W \in \mathcal{W}, N' \in \mathcal{N}, \delta_{N'} = \delta_N\}$$

is a  $P$ -symmetric system.

**Corollary 2.3.**  $\mathcal{S}_P$  is proper. In fact, suppose that  $\theta$  is a regular cardinal and  $N^*$  is a countable elementary substructure of  $H(\theta)$  such that  $P \in N^*$ . Then, letting  $N = N^* \cap H(\kappa)$ , the following conditions hold for every  $\mathcal{N} \in \mathcal{S}_P$ .

- (1) If  $\mathcal{N} \in N$ , then  $\mathcal{N} \cup \{N\} \leq_{\mathcal{S}_P} \mathcal{N}$ .
- (2) If  $N \in \mathcal{N}$ , then  $\mathcal{N}$  is  $(N^*, \mathcal{S}_P)$ -generic.

*Proof.* (1) is obvious. For (2), let  $E$  be a dense subset of  $\mathcal{S}_P$  in  $N^*$ . It suffices to show that there is some condition in  $E \cap N^*$  compatible with  $\mathcal{N}$ . Notice that  $\mathcal{N} \cap N^* \in \mathcal{S}_P$  by Lemma 2.2 (i). Hence, we may find a condition  $\mathcal{N}^\circ \in E \cap N^*$  extending  $\mathcal{N} \cap N^*$ . Now let

$$\mathcal{N}^* = \mathcal{N} \cup \{\Psi_{N, \bar{N}}(M) : M \in \mathcal{N}^\circ, \bar{N} \in \mathcal{N}, \delta_{\bar{N}} = \delta_N\}$$

By Lemma 2.2 (ii) we have that  $\mathcal{N}^*$  is a condition in  $\mathcal{S}_P$  extending both  $\mathcal{N}$  and  $\mathcal{N}^\circ$ .  $\square$

The following is essentially Lemma 2.4 in [6].

**Lemma 2.4.** Let  $P \subseteq H(\kappa)$  and let  $\mathcal{N}_0 = \{N_i^0 : i < \mu\}$  and  $\mathcal{N}_1 = \{N_i^1 : i < \mu\}$  be  $P$ -symmetric systems. Suppose that  $(\bigcup \mathcal{N}_0) \cap (\bigcup \mathcal{N}_1) = X$  and that there is an isomorphism  $\Psi$  between the structures

$$\langle \bigcup_{i < \mu} N_i^0, \in, P, X, N_i^0 \rangle_{i < \mu}$$

and

$$\langle \bigcup_{i < \mu} N_i^1, \in, P, X, N_i^1 \rangle_{i < \mu}$$

which is the identity on  $X$ . Then  $\mathcal{N}_0 \cup \mathcal{N}_1$  is a  $P$ -symmetric system.

Recall that, given a cardinal  $\lambda$ , a poset  $\mathbb{P}$  is  $\lambda$ -Knaster if for every  $X \in [\mathbb{P}]^\lambda$  there is some  $Y \in [X]^\lambda$  consisting of pairwise compatible conditions. The following lemma is easy to prove using Lemma 2.4 together with standard  $\Delta$ -system arguments (cf. [6], Lemma 3.9).

**Lemma 2.5.** (CH) Let  $P \subseteq H(\kappa)$ . If there is a bijection  $\varphi : H(\kappa) \rightarrow \kappa$  definable in  $(H(\kappa), \in, P)$ , then  $\mathcal{S}_P$  is  $\aleph_2$ -Knaster.

It is not difficult to see that forcing with  $\mathcal{S}_P$  adds  $\aleph_1$ -many new reals.

**Fact 2.6.** For every  $P \subseteq H(\kappa)$ ,  $\mathcal{S}_P$  adds  $\aleph_1$ -many Cohen reals.

*Proof.* Let  $S \subseteq \omega_1$  be stationary and co-stationary. Say that a countable  $N$  such that  $(N, \in, P) \preceq (H(\kappa), \in, P)$  is *minimal* if there is some  $x \in N$  with the property that either there is no countable  $M \in N$  such

that  $x \in M$ ,  $\delta_M \in S$  and  $(M, \in, P) \preceq (N, \in, P)$  or there is no countable  $M \in N$  such that  $x \in M$ ,  $\delta_M \notin S$  and  $(M, \in, P) \preceq (N, \in, P)$ . Let  $G$  be  $\mathcal{S}_P$ -generic and let  $\mathcal{N}_G = \bigcup G$ . Given any  $\bar{\delta} < \omega_1$  and  $\bar{N} \in \mathcal{N}_G$  such that  $\delta_{\bar{N}} = \bar{\delta}$ , let  $f_{\bar{\delta}} : \omega \rightarrow 2$  be the function sending  $n$  to 0 if and only if there are minimal  $N_0 \in \dots \in N_n$  in  $\mathcal{N}_G$  such that  $\bar{N} \in N_0$ , such that, letting  $N_{-1} = \bar{N}$ , for all  $i \leq n$ ,  $0 \leq i$ , it holds that there is no minimal  $M \in \mathcal{N}_G$  such that  $N_{i-1} \in M \in N_i$ , and such that  $\delta_{N_n} \in S$ . It is easy to see that  $f_{\bar{\delta}}$  is a well-defined function defined on all of  $\omega$  and that it is (the numerating function of) a Cohen real over  $\mathbf{V}$ . It is also not difficult to see that  $f_{\bar{\delta}_0} \neq f_{\bar{\delta}_1}$  if  $\bar{\delta}_0 = \delta_{N_0} < \bar{\delta}_1 = \delta_{N_1}$  and  $N_0, N_1 \in \mathcal{N}_G$  are both non-minimal. But then we are done since by density there are arbitrarily high  $\bar{\delta} < \omega_1$  such that  $\bar{\delta} = \delta_N$  for some non-minimal  $N \in \mathcal{N}_G$ .  $\square$

What is perhaps more surprising is that if CH holds, then  $\mathcal{S}_P$  does not add more than  $\aleph_1$ -many reals.

**Proposition 2.7.** (CH) *Let  $P \subseteq H(\kappa)$ . If there is a bijection  $\varphi : H(\kappa) \rightarrow \kappa$  definable in  $(H(\kappa), \in, P)$ , then  $\mathcal{S}_P$  preserves CH.*

*Proof.* Suppose  $\dot{r}_\alpha$  (for  $\alpha < \omega_2$ ) are  $\mathcal{S}_P$ -names for members of  ${}^\omega 2$  and  $\mathcal{N} \in \mathcal{S}_P$  forces  $\dot{r}_\alpha \neq \dot{r}_{\alpha'}$ , for all  $\alpha < \alpha' < \omega_2$ . By the  $\aleph_2$ -c.c. of  $\mathcal{S}_P$  we may assume that each  $\dot{r}_\alpha$  is in  $H(\kappa)$ . Let  $\theta$  be a regular cardinal such that  $\mathcal{S}_P \in H(\theta)$ . For each  $\alpha$  let  $N_\alpha$  be such that  $\{\mathcal{N}, \dot{r}_\alpha\} \in N_\alpha$  and  $N_\alpha$  is a countable elementary substructure of  $(H(\kappa), \in, P, \mathcal{S}_P)$ . We may also assume that for each  $\alpha$  there is a countable  $N_\alpha^* \prec H(\theta)$  such that  $N_\alpha = H(\kappa) \cap N_\alpha^*$ . By CH we may find distinct  $\alpha, \alpha'$  such that  $(N_\alpha, \in, P, \mathcal{S}_P, \mathcal{N}, \dot{r}_\alpha)$  and  $(N_{\alpha'}, \in, P, \mathcal{S}_P, \mathcal{N}, \dot{r}_{\alpha'})$  are isomorphic, and we may also assume that the unique isomorphism  $\Psi$  between these structures fixes  $N_\alpha \cap N_{\alpha'}$ . By Lemma 2.4 we know that  $\mathcal{N} \cup \{N_\alpha, N_{\alpha'}\}$  is an  $\mathcal{S}_P$ -condition, and by Corollary 2.3,  $\mathcal{N} \cup \{N_\alpha, N_{\alpha'}\}$  is  $(N_\alpha^*, \mathcal{S}_P)$ -generic and  $(N_{\alpha'}^*, \mathcal{S}_P)$ -generic. Note that for every  $n < \omega$  and every condition  $\mathcal{N}' \mathcal{S}_P$ -extending  $\mathcal{N} \cup \{N_\alpha, N_{\alpha'}\}$  there are conditions  $\mathcal{N}''$ ,  $\mathcal{M}$  and  $\epsilon \in 2$  such that  $\mathcal{M} \in N_\alpha$ ,  $\mathcal{M} \Vdash_{\mathcal{S}_P} \dot{r}_\alpha(n) = \epsilon$ , and  $\mathcal{N}''$  is a common  $\mathcal{S}_P$ -extension of  $\mathcal{M}$  and  $\mathcal{N}'$  (this is true since  $\mathcal{N} \cup \{N_\alpha, N_{\alpha'}\}$  is  $(\mathcal{S}_P, N_\alpha^*)$ -generic). By clause (D) in Definition 2.1,  $\mathcal{N}''$  also  $\mathcal{S}_P$ -extends  $\Psi(\mathcal{M}) \in N_{\alpha'}$ . By correctness of  $\Psi$  with respect to the predicate  $\mathcal{S}_P$ ,  $\Psi(\mathcal{M}) \Vdash_{\mathcal{S}_P} \Psi(\dot{r}_\alpha)(n) = \dot{r}_{\alpha'}(n) = \epsilon$ . Since  $\mathcal{N}'$  and  $n$  were arbitrary this shows that  $\mathcal{N} \cup \{N_\alpha, N_{\alpha'}\}$  forces that  $\dot{r}_\alpha = \dot{r}_{\alpha'}$ , which is a contradiction.  $\square$

### 3. PROVING THEOREMS 1.13 AND 1.14

The following notation will be used throughout the paper: If  $q$  is an ordered pair, we denote the first and the second components of  $q$  by

$F_q$  and  $\Delta_q$ , respectively. Also, if  $q$  is an ordered pair such that  $F_q$  is a function and  $\Delta_q$  is a binary relation with  $\text{range}(\Delta_q) \subseteq \text{Ord}$  and  $\xi$  is an ordinal, *the restriction of  $q$  to  $\xi$* , denoted by  $q|_\xi$ , is defined as the pair

$$q|_\xi := (F_q \upharpoonright \xi, \{(N, \min\{\beta, \xi\}) : (N, \beta) \in \Delta_q\})$$

Similarly, if  $q$  is an ordered pair such that  $F_q$  is a function and  $\Delta_q$  is a set of triples  $(M, \vec{W}, \beta)$  with  $\beta \in \text{Ord}$  and  $\xi$  is an ordinal, *the restriction of  $q$  to  $\xi$* , again denoted by  $q|_\xi$ , is defined as the pair

$$q|_\xi := (F_q \upharpoonright \xi, \{(N, \vec{W}, \min\{\beta, \xi\}) : (N, \vec{W}, \beta) \in \Delta_q\})$$

We will be using instances of the following forcing for adding, by initial approximations, a club of  $\omega_1$  avoiding a given collection of subsets of  $\omega_1$  in the sense of  $(\cdot)_\lambda^\tau$  (cf. [7]).

**Definition 3.1.** Let  $\tau < \omega_1$  and let  $\mathcal{A} = (A_i : i < \lambda) \subseteq \mathcal{P}(\omega_1)$  be such that each  $A_i$  is  $\tau$ -thin. Let  $\mathbb{B}_{\mathcal{A}}^\tau$  consist of all pairs  $(f, b)$  such that

- (a)  $f \subseteq \omega_1 \times \omega_1$  is a finite strictly increasing function such that  $\text{rank}(f(\nu), f(\nu)) \geq \max\{\tau, \nu\}$  for every  $\nu \in \text{dom}(f)$ ,
- (b)  $b$  is finite function with  $\text{dom}(b) \subseteq \lambda$  and  $b(i) \in [\text{range}(f)]^{<\omega}$  for every  $i \in \text{dom}(b)$ , and
- (c) for every  $i \in \text{dom}(b)$ ,  $\text{range}(f) \cap A_i = b(i)$ .

Given  $\mathbb{B}_{\vec{C}}$ -conditions  $(f_0, b_0)$  and  $(f_1, b_1)$ ,  $(f_1, b_1)$  extends  $(f_0, b_0)$  if  $f_0 \subseteq f_1$  and  $b_0 \subseteq b_1$ .

**Remark 3.2.** If  $N$  is a countable elementary substructure of  $H(\theta)$ , for a large enough  $\theta$ , and  $(f, b) \in N \cap \mathbb{B}_{\vec{C}}$ , then  $(f \cup \{(\delta_N, \delta_N)\}, b)$  is  $(N, \mathbb{B}_{\vec{C}})$ -generic.

Let  $\Phi : \kappa \rightarrow H(\kappa)$  be such that  $\Phi^{-1}(x)$  is unbounded in  $\kappa$  for every  $x \in H(\kappa)$ .<sup>11</sup> Let  $\langle \theta_\alpha : \alpha \leq \kappa \rangle$  be the strictly increasing sequence of regular cardinals defined as  $\theta_0 = |2^\kappa|^+$  and  $\theta_\alpha = |2^{\text{sup}\{\theta_\beta : \beta < \alpha\}}|^+$  if  $\alpha > 0$ . For each  $\alpha \leq \kappa$  let  $\mathcal{M}_\alpha^*$  be the collection of all countable elementary substructures of  $H(\theta_\alpha)$  containing  $\Phi$  and  $\langle \theta_\beta : \beta < \alpha \rangle$ . Let also  $\mathcal{M}_\alpha = \{N^* \cap H(\kappa) : N^* \in \mathcal{M}_\alpha^*\}$  and note that if  $\alpha < \beta$ , then  $\mathcal{M}_\alpha^* \in N^*$  for all  $N^* \in \mathcal{M}_\beta^*$  such that  $\alpha \in N^*$ .

The forcing  $\mathcal{P}$  witnessing Theorem 1.13 will be  $(\mathcal{P}_\kappa, \leq_\kappa)$ , where the sequence  $\langle (\mathcal{P}_\alpha, \leq_\alpha) : \alpha \leq \kappa \rangle$  is defined as follows. To start with,  $\mathcal{P}_0$  consists of all pairs of the form

- (a)  $(\emptyset, \{(N_i, 0) : i < m\})$ , where  $\{N_i : i < m\}$  is a finite  $\Phi$ -symmetric system.

<sup>11</sup>This function  $\Phi$  exists by  $2^{<\kappa} = \kappa$ .

Given  $\mathcal{P}_0$ -conditions  $q$  and  $p$ ,  $q \leq_0 p$  iff  $\Delta_p \subseteq \Delta_q$ .

Let  $\alpha = \sigma + 1$ . Suppose that  $\Phi(\sigma)$  is a  $\mathcal{P}_\sigma$ -name for either a c.c.c. forcing or that there is some  $\tau < \omega_1$  and some  $\mathcal{P}_\sigma$ -name  $\dot{A}$  for a subset of  $\mathcal{P}(\omega_1)$  all of whose members are  $\tau$ -thin and such that  $\Phi(\sigma)$  is a  $\mathcal{P}_\sigma$ -name for  $\mathbb{B}_{\dot{A}}^\tau$ . Then we let  $\dot{Q}_\sigma = \Phi(\sigma)$ . If  $\alpha = \sigma + 1 > 0$  and  $\Phi(\sigma)$  is not as above, then we let  $\dot{Q}_\sigma$  be a  $\mathcal{P}_\sigma$ -name for (say) trivial forcing on  $\{0\}$ .

If  $\alpha \leq \kappa$  (regardless of whether  $\alpha$  is a successor or a nonzero limit ordinal), the definition of  $\mathcal{P}_\alpha$  is as follows. Conditions in  $\mathcal{P}_\alpha$  are pairs of the form

$$q = (F_q, \Delta_q)$$

with the following properties.

- (b0)  $F_q$  is a finite function with  $\text{dom}(F_q) \subseteq \alpha$ .
- (b1)  $\Delta_q$  is of the form  $\{(N_i, \beta_i) : i < m\}$  where, for all  $i < m$ ,  $\beta_i \leq \alpha \cap \text{sup}(N_i \cap \kappa)$ .
- (b2) For all  $\xi < \alpha$ ,  $q|_\xi \in \mathcal{P}_\xi$ .
- (b3) If  $\xi \in \text{dom}(F_q)$ , then  $q|_\xi$  forces (in  $\mathcal{P}_\xi$ ) that  $F_q(\xi) \in \dot{Q}_\xi$ .
- (b4) If  $\xi \in \text{dom}(F_q)$ ,  $\dot{Q}_\xi$  is a  $\mathcal{P}_\xi$ -name for a poset of the form  $\mathbb{B}_{\dot{A}}^\tau$  for some  $\tau < \omega_1$  and  $\mathcal{A} \subseteq \mathcal{P}(\omega_1)$ ,  $(N, \beta) \in \Delta_q$ ,  $\beta \geq \xi + 1$ , and  $N \in \mathcal{M}_{\xi+1}$ , then  $\delta_N$  is a fixed point of the first component of  $F_q(\xi)$ .

Given conditions

$$q^\epsilon = (F_\epsilon, \{(N_i^\epsilon, \beta_i^\epsilon) : i < m_\epsilon\})$$

(for  $\epsilon \in \{0, 1\}$ ) in  $\mathcal{P}_\alpha$ , we will say that  $q^1 \leq_\alpha q^0$  if and only if the following holds.

- (c1) For all  $\xi < \alpha$ ,  $q^1|_\xi \leq_\xi q^0|_\xi$ .
- (c2)  $\text{dom}(F_0) \subseteq \text{dom}(F_1)$  and, for all  $\xi \in \text{dom}(F_0)$ ,

$$q^1|_\xi \Vdash_\xi F_1(\xi) \leq_{\dot{Q}_\xi} F_0(\xi)$$

- (c3) For all  $i < m_0$  there is some  $\tilde{\beta}_i \geq \beta_i^0$  such that  $(N_i^0, \tilde{\beta}_i) \in \Delta_{q^1}$ .

Note that if  $\alpha < \beta \leq \kappa$ , then  $\mathcal{P}_\alpha \subseteq \mathcal{P}_\beta$ ; in fact, every  $\mathcal{P}_\beta$ -condition  $q = (F, \{(N_j, \beta_j) : j < m\})$  such that  $\text{dom}(q) \subseteq \alpha$  and  $\beta_j \leq \alpha$  for all  $j$  is also a  $\mathcal{P}_\alpha$ -condition and is in fact its restriction to  $\alpha$ .

Also note that if  $\alpha \leq \kappa$  and  $q \in \mathcal{P}_\alpha$ , then  $\text{dom}(\Delta_q) \subseteq \text{dom}(\Delta_{q|_0})$  and  $q|_0 \in \mathcal{P}_0$ .

Finally note that if  $\alpha$  is a nonzero limit ordinal, then a pair  $q = (F_q, \Delta_q)$  is a  $\mathcal{P}_\alpha$ -condition if and only if it satisfies (b0)–(b2).

For every  $\mathcal{P}_\kappa$ -condition  $q$  and every  $\xi \in \text{dom}(F_q)$ , if  $\dot{Q}_\xi$  is a  $\mathcal{P}_\xi$ -name for a forcing of the form  $\mathbb{B}_{\mathcal{A}}^\tau$ , for some  $\tau < \omega_1$  and  $\mathcal{A} \subseteq \mathcal{P}(\omega_1)$ , we denote by  $f_{q,\xi}$  and  $b_{q,\xi}$  the first and second components of  $F_q(\xi)$ , respectively.

The following lemmas are proved essentially in [6].

**Lemma 3.3.** *Let  $\alpha < \beta \leq \kappa$ , let  $r \in \mathcal{P}_\beta$  and let  $q \in \mathcal{P}_\alpha$  be such that  $q \leq_\alpha r|_\alpha$ . Then  $(F_q \cup (F_r \upharpoonright [\alpha, \beta]), \Delta_q \cup \Delta_r)$  is a condition in  $\mathcal{P}_\beta$  extending both of  $q$  and  $r$ . In particular,  $\mathcal{P}_\alpha$  is a complete suborder of  $\mathcal{P}_\beta$ .*

**Lemma 3.4.** (CH) *For every ordinal  $\alpha \leq \kappa$ ,  $\mathcal{P}_\alpha$  has the  $\aleph_2$ -c.c.*

**Lemma 3.5.**  *$\mathcal{P}_\kappa$  forces  $2^{\aleph_0} = \kappa$ .*

**Definition 3.6.** Given  $\alpha \leq \kappa$ , a condition  $q \in \mathcal{P}_\alpha$ , and a countable elementary substructure  $N \prec H(\kappa)$ , we will say that  $q$  is  $(N, \mathcal{P}_\alpha)$ -pre-generic in case

- $\alpha < \kappa$  and  $(N, \alpha) \in \Delta_q$ , or else
- $\alpha = \kappa$  and the pair  $(N, \text{sup}(N \cap \kappa)) \in \Delta_q$ .

**Lemma 3.7.** *Suppose  $\alpha \leq \kappa$  and  $N^* \in \mathcal{M}_\alpha^*$ . Let  $N = N^* \cap H(\kappa)$ . Then the following conditions hold.*

- (1) $_\alpha$  *For every  $q \in N$  there is some  $q' \leq_\alpha q$  such that  $q'$  is  $(N, \mathcal{P}_\alpha)$ -pre-generic.*
- (2) $_\alpha$  *If  $\mathcal{P}_\alpha \in N^*$  and  $q \in \mathcal{P}_\alpha$  is  $(N, \mathcal{P}_\alpha)$ -pre-generic, then  $q$  is  $(N^*, \mathcal{P}_\alpha)$ -generic.*

**Corollary 3.8.** *For every  $\alpha \leq \kappa$ ,  $\mathcal{P}_\alpha$  is proper.*

It is easy to see that forcing with any instance of  $\mathbb{B}_{\mathcal{A}}^\tau$  as in Definition 3.1 adds a generic for  $\mathbb{B}$ . By standard book-keeping arguments together with Lemma 3.4 it follows then that  $\mathcal{P}$  forces  $\text{MA} + \text{FA}(\mathbb{B}) + (\cdot)_\lambda^\tau$  for all  $\tau < \omega_1$  and  $\lambda < \kappa$ .

In order to complete the proof of Theorem 1.13 it therefore suffices to prove the following lemma.

**Lemma 3.9.** *Let  $\alpha < \kappa$  and suppose  $\mathcal{G} = \langle g_\delta : \delta \in \text{Lim}(\omega_1) \rangle$  is a strong  $\mathcal{U}$ -sequence. Let  $n < \omega$  and suppose that  $M^*$  and  $M$  satisfy the following:*

- (1)  $M^* \in \mathcal{M}_{\alpha+2}^*$  and  $M = M^* \cap H(\kappa)$ .
- (2)  $\dot{C} \in M$  is a  $\mathcal{P}_\alpha$ -name for a club of  $\omega_1$ .
- (3)  $(M, \alpha) \in \Delta_q$  for some  $q \in \mathcal{G}$ .

*Then  $q$  forces that there is some  $\varepsilon \in \dot{C} \cap \delta_M$  such that  $g_{\delta_M}(\varepsilon) = n$ .*



*Proof.* The proof is by induction on  $\alpha$ . The case  $\alpha = 0$  follow immediately from Fact 1.4.

For the case when  $\alpha = \sigma + 1$ , let  $q' \in \mathcal{P}_\alpha$  be an extension of  $q$ . Without loss of generality we may assume that  $\sigma \in F_{q'}$  and that there is some  $\tau < \omega_1$  and some  $\mathcal{P}_\sigma$ -name  $\dot{A}$  for a subset of  $\mathcal{P}(\omega_1)$  all of whose members are  $\tau$ -thin and such that  $\dot{Q}_\sigma$  is a  $\mathcal{P}_\sigma$ -name for  $\mathbb{B}_{\dot{A}}^\tau$  (the proof in the other cases is easier). So,  $\delta_M$  is a fixed point of  $f_{q',\sigma}$ . Let  $G$  be a  $\mathcal{P}_\sigma$ -generic with  $q'|_\sigma \in G$  and let  $\dot{A}_G = (A_i : i < \lambda)$ . By Lemma 3.7,  $G$  is also generic over  $M^*$  and we may therefore assume  $\tau \in M$ . We may assume that  $f_{q',\sigma} \upharpoonright \delta_M \neq \emptyset$ . Let  $\delta = \max(\text{dom}(f_{q',\sigma} \upharpoonright \delta_M))$ . Now we consider the following function  $F$  with domain the open interval  $(f_{q',\sigma}(\delta), \omega_1)$ . Suppose  $\varrho$  is in this interval. Assume that there exists a condition  $t$  in  $\mathcal{P}_\alpha$  with the following properties.

- (a)  $t|_\sigma \in G$
- (b)  $(f_{q',\sigma} \upharpoonright \delta_M) \cup \{\langle \delta + 1, \varrho \rangle\} \subseteq f_{t,\sigma}$
- (c)  $f_{t,\sigma} \upharpoonright (\delta + 1) = f_{q',\sigma} \upharpoonright \delta_M$
- (d) There is some  $\nu \in \text{dom}(f_{t,\sigma})$ ,  $\nu > \delta + 1$ , such that  $f_{t,\sigma}(\nu) = \delta_N$  for some  $N \in \mathcal{M}_{\alpha+1}$  such that  $\dot{C} \in N$  and  $(N, \alpha) \in \Delta_t$  (so, by Lemma 3.7,  $t$  forces  $f_{t,\sigma}(\nu) \in \dot{C}$ ).

In this case we fix such a  $t$  and set  $F(\varrho) = f_{t,\sigma}(\nu)$  for a  $\nu$  as in (d). Otherwise, we set  $F(\varrho) = 0$ .

Now, let  $E(F)$  be the club of all  $\varepsilon$  such that  $\text{rank}(\varepsilon, \varepsilon) > \tau$  and  $F \restriction \varepsilon \subseteq \varepsilon$ . Note that  $F$  and  $E(F)$  are definable from parameters in  $M^*[G]$ , and therefore they are in  $M^*[G]$  as well. Moreover, by the  $\aleph_2$ -c.c., we can assume that  $E(F)$  is in  $M[G]$ . Using the induction hypothesis, we know that there is some  $\varepsilon \in \delta_M$  which is a limit point of  $E(F)$  and such that  $g_{\delta_M}(\varepsilon) = n$ . Let  $\varrho < \eta < \varepsilon$  be such that  $\text{rank}(\varrho, \varrho) \geq \delta + 1$ ,  $\eta \in E(F)$ ,  $g_{\delta_M} \restriction [\varrho, \eta) = \{n\}$ , and  $[\varrho, \eta) \cap A_i = \emptyset$  for every  $i \in \text{dom}(b_{q',\sigma})$ . These ordinals can be found since  $g_{\delta_M}$  is continuous and since  $\varepsilon$  is a limit of ordinals in  $E(F)$  and  $\text{rank}(\varepsilon, \varepsilon) > \tau$ , together with the fact that  $\text{rank}(A_i, \varepsilon) \leq \tau$  for every  $i \in \text{dom}(b_{q',\sigma})$ . Let  $t \in \mathcal{P}_\alpha \in M^*[G]$  witness (a)–(d) for  $\varrho$ . The existence of such a  $t$  follows from the fact that  $q''$  witnesses (a)–(d) in  $\mathbf{V}[G]$ , where  $\text{dom}(F_{q''}) = \text{dom}(F_{q'})$  and  $\Delta_{q''} = \Delta_{q'}$ ,  $F_{q''}(\xi) = F_{q'}(\xi)$  for all  $\xi \in \text{dom}(F_{q'})$ ,  $\xi \neq \sigma$ , and where  $F_{q''}(\sigma) = (f_{q',\sigma} \cup \{\langle \delta + 1, \varrho \rangle\}, b_{q',\sigma})$ . It follows from (d) for  $t$  that there is some  $\nu \in \text{dom}(f_{t,\sigma})$  such that  $\nu > \delta + 1$  and such that  $t$  forces  $f_{t,\sigma}(\nu) \in \dot{C}$ . Since we may assume that  $f_{t,\sigma}(\nu) \in [\varrho, \eta)$  by the definition of  $E(F)$  and the fact that  $\eta \in E(F)$ , we have that  $g_{\delta_M}(f_{t,\sigma}(\nu)) = n$ . By extending  $q'|_\sigma$  if necessary we may assume that  $q'|_\sigma$  decides the above facts holding in  $\mathbf{V}[G]$ . Finally, by the choice of  $\varrho$  such that  $[\varrho, \eta) \cap A_i = \emptyset$  for all  $i \in \text{dom}(b_{q',\sigma})$ , we may amalgamate  $q'$  and  $t$  into

a  $\mathcal{P}_\alpha$ -condition  $q^\dagger$ . Since  $g_{\delta_M}(f_{t,\sigma}(\nu)) = n$ ,  $q^\dagger$  forces  $f_{t,\sigma}(\nu) \in \dot{C}$ , and  $q'$  was an arbitrary extension of  $q$ , we are done in this case.

Now assume that  $\alpha$  is a limit ordinal. Let  $q' \in \mathcal{P}_\alpha$  be an extension of  $q$  and let  $\sigma \in M$  be an ordinal with  $\sigma > \max(\text{dom}(F_{q'}) \cap M)$ . Let  $G \subseteq \mathcal{P}_\sigma$  be a generic filter with  $q'|_\sigma \in G$ . By Lemma 3.7,  $G$  is also generic over  $M^*$ .

Note that the set  $D$  of all  $\delta < \omega_1$  for which there is some  $N \in \mathcal{M}_{\alpha+1}$  such that

- (e)  $\delta = \delta_N$ ,
- (f)  $\dot{C} \in N$ , and
- (g)  $(N, \alpha) \in \Delta_t$  for some  $t \in \mathcal{P}_\alpha$  such that  $t|_\sigma \in G$  and  $\text{dom}(F_t) \subseteq \sigma$

is definable in  $M^*[G]$ .

Note also that  $D$  is cofinal in  $\omega_1$  since  $\delta_M \in D$  as witnessed by  $q|_\sigma$ . So, if  $C$  is the set of the accumulation points of  $D$ , then  $C$  is a club definable in  $M^*[G]$ . Moreover, by the  $\aleph_2$ -c.c., we may assume  $C \in M[G]$ . Using the induction hypothesis, we know that there is some  $\varepsilon \in C \cap \delta_M$  such that  $g_{\delta_M}(\varepsilon) = n$ . Take  $\eta < \varepsilon$  such that  $g_\delta "[\eta, \varepsilon) = \{n\}$ . We may now find  $\delta \in (\eta, \varepsilon) \cap D$ . Let now  $N \in \mathcal{M}_{\alpha+1}$  and  $t$  satisfy (e)–(g) for  $\delta$ , and note that  $t$  forces  $\delta = \delta_N \in \dot{C}$  by Lemma 3.7. As in the previous case, by extending  $q'|_\sigma$  if necessary we may assume that  $q'|_\sigma$  forces the above facts holding in  $\mathbf{V}[G]$ . Since  $t$  forces  $\delta = \delta_N \in \dot{C}$  by Lemma 3.7, it suffices to find a common extension  $q^\dagger$  of  $q'$  and  $t$ . But  $q'$  and  $t$  can indeed be amalgamated into an extension  $q^\dagger$  by (g) for  $t$ . Since  $q'$  was an arbitrary extension of  $q$ , we are done in this case too.<sup>12</sup>  $\square$

We will next prove Theorem 1.14. We start by defining a natural forcing for destroying a potential  $\mathcal{U}$ -sequence (cf. [6] and [7]).

**Definition 3.10.** Let  $\mathcal{G} = \langle g_\delta : \delta \in \text{Lim}(\omega_1) \rangle$  be such that each  $g_\delta$  is a continuous function from  $\delta$  into  $\omega$  with respect to the order topology. Let  $\mathbb{B}_\mathcal{G}$  be the forcing notion consisting of all pairs  $(f, \langle b_\xi : \xi \in D \rangle)$  satisfying the following conditions.

- (1)  $f \subseteq \omega_1 \times \omega_1$  is a finite strictly increasing function such that  $\text{rank}(f(\nu), f(\nu)) \geq \nu$  for every  $\nu \in \text{dom}(f)$ .
- (2)  $b$  is a function with  $\text{dom}(b) \subseteq \text{dom}(f)$  and such that for each  $\xi$  in  $\text{dom}(b)$ ,
  - (2.1)  $b(\xi) \in \omega$ ,
  - (2.2)  $g_{f(\xi)}$  “  $\text{range}(f) \subseteq \omega \setminus \{b(\xi)\}$ , and

<sup>12</sup>The proof in the successor case could be simplified a bit by looking, in the definition of  $F$ , at conditions  $t$  such that  $\sigma \notin \text{dom}(F_t)$  (similarly as what we do in the limit case).

$$(2.3) \text{ rank}(\{\gamma < f(\xi) : g_{f(\xi)}(\gamma) \neq b(\xi)\}, f(\xi)) = \text{rank}(f(\xi), f(\xi)).$$

Given  $\mathbb{B}_{\mathcal{G}}$ -conditions  $c^\epsilon = (f^\epsilon, b^\epsilon)$  for  $\epsilon \in \{0, 1\}$ ,  $c^1$  extends  $c^0$  if and only if  $f^0 \subseteq f^1$ , and  $b^0 \subseteq b^1$ .

The forcing  $\mathcal{P}$  witnessing Theorem 1.14 is  $(\mathcal{P}_\kappa, \leq_\kappa)$ , where the sequence  $\langle (\mathcal{P}_\alpha, \leq_\alpha) : \alpha \leq \kappa \rangle$  will be defined soon by recursion. Let  $\vec{C} = \langle C_\delta : \delta \in \text{Lim}(\omega_1) \rangle$  be a strong WCG-sequence. For every  $\delta \in \text{Lim}(\omega_1)$ , let  $(C_\delta(m))_{m < \omega}$  be the strictly increasing enumeration of  $C_\delta$  and define  $C_m^\delta := C_\delta \setminus C_\delta(m)$ .

**Definition 3.11.** Let  $N$  and  $\mathcal{U}$  be two sets. We will say that  $\mathcal{U}$  is  $N$ -unbounded in case for every  $x \in N$  there is some  $M \in \mathcal{U} \cap N$  such that  $x \in M$ .

Given a set  $N$ , let us say that  $N$  is *sufficiently closed* if for every finite set  $x$ ,  $x \subseteq N$  iff  $x \in N$ .

**Remark 3.12.** If  $\mathcal{U}$  is  $N$ -unbounded,  $\mathcal{U}' \subseteq \mathcal{U}$  and every member of  $\{N\} \cup \mathcal{U}'$  is sufficiently closed, then at least one of  $\mathcal{U}'$ ,  $\mathcal{U} \setminus \mathcal{U}'$  is  $N$ -unbounded.

Given an  $\omega$ -sequence  $\vec{\mathcal{U}}$ , we will use  $\mathcal{U}_n$  to denote the  $n$ th member of  $\vec{\mathcal{U}}$ ; that is,  $\vec{\mathcal{U}} = (\mathcal{U}_n)_{n < \omega}$ . Given two  $\omega$ -sequences  $\vec{\mathcal{U}}, \vec{\mathcal{V}}$  of sets, by  $\vec{\mathcal{V}} \subseteq \vec{\mathcal{U}}$  we will mean that  $\mathcal{V}_n \subseteq \mathcal{U}_n$  for all  $n$ . Finally, an  $\omega$ -sequence  $\vec{\mathcal{U}}$  is said to be  $\subseteq$ -decreasing if  $\mathcal{U}_m \subseteq \mathcal{U}_n$  whenever  $n < m < \omega$ .

**Lemma 3.13.** Let  $\mathcal{G} = \langle g_\delta : \delta \in \omega_1 \rangle$  be such that each  $g_\delta$  is a continuous function from  $\delta$  into  $\omega$  and let us denote by  $\mathcal{M}$  the set of all sufficiently closed countable subsets  $N$  of  $H(\kappa)$  such that  $N \models \text{ZFC}^*$ ,  $\delta_N$  exists and  $\omega_1^N = \omega_1$ . Suppose  $p < \omega$  and  $\{N_i : i < p\}$ ,  $\{\vec{\mathcal{U}}_i : i < p\}$  are such that for all  $i < p$ ,  $n < \omega$  and  $m < \omega$ , the following hold.

- (o)  $N_i \in \mathcal{M}$ .
- (o)  $\delta_{N_{i'}} = \delta_{N_i}$  for all  $i'$ .
- (o)  $\vec{\mathcal{U}}_i$  is a  $\subseteq$ -decreasing  $\omega$ -sequence.
- (o)  $\mathcal{U}_{i,n} \subseteq N_i \cap \mathcal{M}$ .
- (o)  $\{M \in \mathcal{U}_{i,n} : \delta_M \in C_m^{\delta_{N_i}}\}$  is  $N_i$ -unbounded.

Then, for each  $s$  in  $[\omega]^{p+1}$  and each  $i < p$  there is some  $s_i \in [s]^p$  such that

$$\{M \in \mathcal{U}_{i,n} : \delta_M \in C_m^{\delta_{N_i}} \text{ and } g_{\delta_{N_i}}(\delta_M) \notin s_i\}$$

is  $N_i$ -unbounded for all  $n, m < \omega$ . In particular, for each  $s$  in  $[\omega]^{p+1}$  there is at least one colour  $j \in s$  such that

$$\{M \in \mathcal{U}_{i,n} : \delta_M \in C_m^{\delta_{N_i}} \text{ and } g_{\delta_{N_i}}(\delta_M) \neq j\}$$

is  $N_i$ -unbounded for all  $i < p$ ,  $n < \omega$  and  $m < \omega$ .

*Proof.* It suffices to note that for each pair  $\{n_0, n_1\} \in [s]^2$ , the above remark (together with the fact that  $\vec{\mathcal{U}}_i$  is a  $\subseteq$ -decreasing  $\omega$ -sequence) ensures the existence of  $j \in 2$  such that

$$\{M \in \mathcal{U}_{i,n} : \delta_M \in C_m^{\delta_{N_i}} \text{ and } g_{\delta_{N_i}}(\delta_M) \neq n_j\}$$

is  $N_i$ -unbounded for co-finally many  $n$ ,  $m < \omega$ , and therefore for all  $n, m < \omega$ .  $\square$

**Notation 3.14.** If  $\mathcal{W}$  is a set of ordered pairs, then the projection  $\{a : \langle a, b \rangle \in \mathcal{W} \text{ for some } b\}$  will be denoted by  $\pi_0(\mathcal{W})$ . Similarly,  $\pi_1(\mathcal{W})$  denotes the projection  $\{b : \langle a, b \rangle \in \mathcal{W} \text{ for some } a\}$

Let us fix  $\Phi$  and  $\theta_\alpha$ ,  $\mathcal{M}_\alpha^*$  and  $\mathcal{M}_\alpha$  (for  $\alpha \leq \kappa$ ) exactly as in the proof of Theorem 1.13. We proceed to the definition of  $\langle \mathcal{P}_\alpha : \alpha \leq \kappa \rangle$  now.  $\mathcal{P}_0$  consist of all pairs of the form  $(\emptyset, \{(N_i, \vec{\mathcal{W}}_i, 0) : i < m\})$  with the following properties.

- (a1)  $\{N_i : i < m\}$  is a finite  $\Phi$ -symmetric system.
- (a2) For every  $i < m$ ,  $\vec{\mathcal{W}}_i = (\mathcal{W}_{i,n})_{n < \omega}$  is a  $\subseteq$ -decreasing  $\omega$ -sequence such that each  $\mathcal{W}_{i,n}$  is a set of ordered pairs in  $N_i$ .
- (a3) For every  $i < m$  and  $n < \omega$ , every member of  $\pi_0(\mathcal{W}_{i,n})$  is a countable  $M \preceq (H(\kappa), \in, \Phi)$ .

Given  $\mathcal{P}_0$ -conditions  $q$  and  $p$ ,  $q \leq_0 p$  iff  $\Delta_p \subseteq \Delta_q$ .

Let  $\alpha = \sigma + 1$ . Suppose  $\Phi(\sigma)$  is a  $\mathcal{P}_\sigma$ -name for a c.c.c. poset or there is a  $\mathcal{P}_\sigma$ -name  $\dot{\mathcal{G}}$  for a sequence of continuous functions  $g_\delta : \delta \rightarrow \omega$  ( $\delta \in \text{Lim}(\omega_1)$ ) such that  $\Phi(\sigma)$  is a  $\mathcal{P}_\sigma$ -name for  $\mathbb{B}_{\dot{\mathcal{G}}}$ . Then we let  $\dot{\mathcal{Q}}_\sigma = \Phi(\sigma)$ . In the other case we let  $\Phi(\sigma)$  be a  $\mathcal{P}_\sigma$ -name for, say, trivial forcing on  $\{0\}$ .

If  $\alpha = \sigma + 1$  and  $\vec{\mathcal{W}} \in V$  is an  $\omega$ -sequence whose members are sets of ordered pairs, then  $\vec{\mathcal{W}}^{\dot{\mathcal{G}}_\sigma}$  denotes the canonical  $\mathcal{P}_\sigma$ -name for the  $\omega$ -sequence whose  $n$ -th member is defined as

$$\mathcal{W}_n^{\dot{\mathcal{G}}_\sigma} := \{(M, \vec{\mathcal{Z}}) \in \mathcal{W}_n : (M, \vec{\mathcal{Z}}, \sigma + 1) \in \Delta_u \text{ for some } u \in \dot{\mathcal{G}}_\sigma\}$$

Note that if  $\vec{\mathcal{W}}$  is  $\subseteq$ -decreasing, then  $\mathcal{P}_\sigma$  forces that so is  $\vec{\mathcal{W}}^{\dot{\mathcal{G}}_\sigma}$ .

If  $\alpha \leq \kappa$  (regardless of whether  $\alpha$  is a successor or a nonzero limit ordinal), the definition of  $\mathcal{P}_\alpha$  is as follows. Conditions in  $\mathcal{P}_\alpha$  are pairs of the form

$$q = (F_q, \Delta_q)$$

with the following properties.

- (b0)  $F_q$  is a finite function with  $\text{dom}(F_q) \subseteq \alpha$ .
- (b1)  $\Delta_q$  is of the form  $\{(N_i, \vec{\mathcal{W}}_i, \beta_i) : i < m\}$  where, for all  $i < m$ ,  $\beta_i \leq \alpha \cap \text{sup}(N_i \cap \kappa)$ .

- (b2) For all  $\xi < \alpha$ , the restriction of  $q$  to  $\xi$  is a condition in  $\mathcal{P}_\xi$ .
- (b3) If  $\xi \in \text{dom}(F_q)$ , then  $q|_\xi$  forces (in  $\mathcal{P}_\xi$ ) that  $F_q(\xi) \in \dot{Q}_\xi$ .
- (b4) Suppose  $\xi \in \text{dom}(F_q)$  and there is some  $\mathcal{P}_\xi$ -name  $\dot{G}$  for a sequence of continuous functions  $g_\delta : \delta \rightarrow \omega$  (for  $\delta \in \text{Lim}(\omega_1^{\mathbf{V}})$ ) such that  $\Phi(\xi)$  is a  $\mathcal{P}_\xi$ -name for  $\mathbb{B}_{\dot{G}}$ . Suppose  $(N, \vec{W}, \beta) \in \Delta_q$ ,  $\beta \geq \xi + 1$ ,  $N \in \mathcal{M}_{\xi+1}$ , and

$$\{M \in \pi_0(\mathcal{W}_n) : \delta_M \in C_m^{\delta_N}\}$$

is  $N$ -unbounded for all  $n, m \in \omega$ . Then  $\delta_N$  is a fixed point of  $f_{q,\xi}$ ,  $b_{q,\xi}(\delta_N)$  is defined, and  $q|_\xi$  forces that

$$\{M \in \pi_0(\mathcal{W}_n^{\dot{G}_\xi}) : \dot{g}_{\xi,\delta_N}(\delta_M) \neq b_{q,\xi}(\delta_N) \text{ and } \delta_M \in C_m^{\delta_N}\}$$

is  $N$ -unbounded for all  $n, m \in \omega$ .<sup>13</sup>

Given conditions

$$q^\epsilon = (F_\epsilon, \{(N_i^\epsilon, \vec{W}_i^\epsilon, \beta_i^\epsilon) : i < m_\epsilon\})$$

(for  $\epsilon \in \{0, 1\}$ ) in  $\mathcal{P}_\alpha$ , we will say that  $q^1 \leq_\alpha q^0$  if and only if the following holds.

- (c1) For all  $\xi < \alpha$ ,  $q^1|_\xi \leq_\xi q^0|_\xi$ .
- (c2)  $\text{dom}(F_0) \subseteq \text{dom}(F_1)$  and, for all  $\xi \in \text{dom}(F_0)$ ,

$$q^1|_\xi \Vdash_\xi F_1(\xi) \leq_{\dot{Q}_\xi} F_0(\xi)$$

- (c3) For all  $i < m_0$  there is some  $\tilde{\beta}_i \geq \beta_i^0$  such that  $(N_i^0, \vec{W}_i^0, \tilde{\beta}_i) \in \Delta_{q^1}$ .

As in the proof of Theorem 1.13,<sup>14</sup> we can show that the corresponding versions of Lemmas 3.3, 3.4 and 3.5 are true for our present iteration. The proof of the corresponding form of Lemma 3.7 will need to be a bit more elaborate.

**Definition 3.15.** Given an ordinal  $\alpha \leq \kappa$  and a pair  $(N, \vec{W})$  such that  $N = N^* \cap H(\kappa)$  for some  $N^* \in \mathcal{M}_\alpha^*$  and such that  $\vec{W}$  is a  $\subseteq$ -decreasing  $\omega$ -sequence of sets consisting of ordered pairs in  $N$ , we will say that  $\vec{W}$  is  $N$ - $\alpha$ -large whenever the following conditions hold.

- (1) For all  $n < \omega$ , every member of  $\pi_0(\mathcal{W}_n)$  is a countable elementary substructure of  $(H(\kappa), \in, \Phi)$ . If  $\alpha = 0$ , we also require that  $\pi_1(\mathcal{W}_n) = \{\emptyset\}$
- (2) For all  $n, m < \omega$ ,  $\{M \in \pi_0(\mathcal{W}_n) : \delta_M \in C_m^{\delta_N}\}$  is  $N$ -unbounded.

<sup>13</sup>Here, and throughout the proof of Theorem 1.14, we are using the obvious notational conventions when writing things like  $f_{q,\xi}$  and  $b_{q,\xi}$  (as in the proof of Theorem 1.13).

<sup>14</sup>I.e., as in [6].

- (3) For every nonzero ordinal  $\beta \in N \cap \alpha$  there is some  $n < \omega$  with the property that for every  $(M, \vec{W}') \in \mathcal{W}_n$ ,
- (o)  $M = M^* \cap H(\kappa)$  for some  $M^* \in \mathcal{M}_\beta^*$  with  $\mathcal{P}_\beta \in M^*$  and
  - (o)  $\vec{W}'$  is  $M$ - $\beta$ -large.

**Lemma 3.16.** *Let  $\alpha \leq \kappa$  and let  $N^* \in \mathcal{M}_\alpha^*$  be such that  $\vec{C} \in N^*$  and  $(\mathcal{P}_\beta)_{\beta < \alpha} \in N^*$ . Let  $N = N^* \cap H(\kappa)$ . Then there is a  $\subseteq$ -decreasing  $\omega$ -sequence  $\vec{W}$  such that  $\vec{W}$  is  $N$ - $\alpha$ -large.*

*Proof.* The proof is by induction on  $\alpha$ . We first show that for every  $m \in \omega$ , every  $x$  in  $N$  and every finite set  $s \subseteq N \cap \alpha$  there is some  $(M, \vec{W}') \in N$  such that  $M \preceq (H(\kappa), \in, \Phi)$ ,  $|M| = \aleph_0$ ,  $M \in \mathcal{M}_{\max(s)}$  if  $s \neq \emptyset$ ,  $x \in M$ , and such that  $\delta_M \in C_m^{\delta N}$ . If  $s$  is empty, we simply define  $\vec{W}'$  as the empty set. Otherwise, we also require that  $\vec{W}'$  is  $M$ - $\max(s)$ -large and  $M = M^* \cap H(\kappa)$  for some  $M^* \in \mathcal{M}_{\max(s)}^*$  with  $\{s, \mathcal{P}_{\max(s)}\} \subseteq M^*$ .

If  $s \neq \emptyset$ , let  $C \in N^*$  be a club of  $\omega_1$  such that for every  $\delta \in C$  there is some  $M^* \in \mathcal{M}_{\max(s)}^*$  such that  $\{x, s, \mathcal{P}_{\max(s)}\} \subseteq M^*$  and  $\delta_{M^*} = \delta$ . (If  $s = \emptyset$ , we take  $C \in N^*$  to be a club of  $\omega_1$  such that for every  $\delta \in C$  there is some countable  $M \preceq (H(\kappa), \in, \Phi)$  such that  $x \in M$  and  $\delta_M = \delta$ .) Since  $\vec{C} \in N^*$ , by Fact 1.5, we may fix  $\delta \in C \cap C_m^{\delta N}$ . By correctness of  $N^*$  there is then some  $M^* \in \mathcal{M}_{\max(s)}^* \cap N^*$  such that  $\{x, s, \mathcal{P}_{\max(s)}\} \subseteq M^*$  and such that  $\delta_{M^*} = \delta$ , if  $s \neq \emptyset$ , and a countable  $M \preceq (H(\kappa), \in, \Phi)$ ,  $M \in N^*$ , such that  $x \in M$  and  $\delta_M = \delta$  if  $s = \emptyset$ . Let  $M = M^* \cap H(\kappa)$  (if  $s \neq \emptyset$ ). By induction hypothesis there is some  $\omega$ -sequence  $\vec{W}'$  such that  $\vec{W}'$  is  $M$ - $\max(s)$ -large if  $s \neq \emptyset$ , and by correctness of  $N^*$  there is some such  $\vec{W}'$  in  $N$ . Now  $(M, \vec{W}')$  is as desired.

For every  $m < \omega$ ,  $x \in N$ , and every finite subset of  $N \cap \alpha$ , choose a pair  $(M, \vec{W}')$  as above and let  $\mathcal{W}_0$  be the set of all these pairs. If  $\alpha = 0$ , we let  $\mathcal{W}_{n+1} = \mathcal{W}_0$  for every  $n$ . If  $\alpha > 0$ , we fix a surjection  $f : \omega \rightarrow N \cap \alpha$ . In this case,  $\mathcal{W}_{n+1}$  is defined as the set of those pairs  $(M, \vec{W}')$  in  $\mathcal{W}_0$  such that, letting  $\mu = \max(\text{range}(f \upharpoonright n+1))$ ,  $\vec{W}'$  is  $M$ - $\mu$ -large and  $M = M^* \cap H(\kappa)$  for some  $M^* \in \mathcal{M}_\mu^*$  with  $\{\text{range}(f \upharpoonright n+1), \mathcal{P}_\mu\} \subseteq M^*$ . Since  $\vec{W} = (\mathcal{W}_n)_{n < \omega}$  is  $\subseteq$ -decreasing, it is clear that this  $\omega$ -sequence is  $N$ - $\alpha$ -large.  $\square$

We consider the following natural notion of pre-properness (cf. Definition 3.6).

**Definition 3.17.** Given  $\alpha \leq \kappa$ , a condition  $q \in \mathcal{P}_\alpha$ , and a pair  $(N, \vec{W})$ , we say that  $q$  is  $\mathcal{P}_\alpha$ -pre-generic for  $(N, \vec{W})$  in case

- $\alpha < \kappa$  and  $(N, \vec{\mathcal{W}}, \alpha) \in \Delta_q$ , or else
- $\alpha = \kappa$  and  $(N, \vec{\mathcal{W}}, \sup(N \cap \kappa)) \in \Delta_q$ .

Our properness lemma now is the following.

**Lemma 3.18.** *Suppose  $\alpha \leq \kappa$  and  $N^* \in \mathcal{M}_\alpha^*$ . Let  $N = N^* \cap H(\kappa)$  and assume that  $\vec{\mathcal{W}}$  is a  $\subseteq$ -decreasing  $\omega$ -sequence consisting of ordered pairs in  $N$  and such that for all  $n < \omega$ , every member of  $\pi_0(\mathcal{W}_n)$  is a countable elementary substructure of  $(H(\kappa), \in, \Phi)$ . Then the following conditions hold.*

- (1) $_\alpha$  For every  $q \in N$  there is  $q' \leq_\alpha q$  such that  $q'$  is  $\mathcal{P}_\alpha$ -pre-generic for  $(N, \vec{\mathcal{W}})$ .
- (2) $_\alpha$  If
  - (o)  $\mathcal{P}_\alpha \in N^*$  and  $\vec{\mathcal{W}}$  is  $N$ - $\alpha$ -large and
  - (o)  $q \in \mathcal{P}_\alpha$  is  $\mathcal{P}_\alpha$ -pre-generic for  $(N, \vec{\mathcal{W}})$ ,
 then  $q$  is  $(N^*, \mathcal{P}_\alpha)$ -generic.
- (3) $_\alpha$  For all  $q \in \mathcal{P}_\alpha$ ,  $m^*$ ,  $n^* < \omega$ , and  $x \in N$ , if
  - (o)  $q$  is  $\mathcal{P}_\alpha$ -pre-generic for  $(N, \vec{\mathcal{W}})$  and
  - (o)  $\{M \in \pi_0(\mathcal{W}_n) : \delta_M \in C_m^{\delta_N}\}$  is  $N$ -unbounded for all  $m$ ,  $n < \omega$ ,
 then there is some condition  $q' \in \mathcal{P}_\alpha$  extending  $q$  and some  $(M, \vec{\mathcal{W}}') \in \mathcal{W}_{n^*}$  such that
  - (a)  $x \in M$ ,
  - (b)  $\delta_M \in C_{m^*}^{\delta_N}$ , and
  - (c)  $q'$  is  $\mathcal{P}_\alpha$ -pre-generic for  $(M, \vec{\mathcal{W}}')$ .

*Proof.* The proof will be by induction on  $\alpha$ . The case  $\alpha = 0$  follows from Corollary 2.3 (the proof of (3) $_0$  is like the proof of that corollary, using the fact that  $\{M \in \pi_0(\mathcal{W}_n) : \delta_M \in C_m^{\delta_N}\}$  is  $N$ -unbounded for all  $m < \omega$  and  $n < \omega$ ).

Now we prove the conclusion for the case  $\alpha = \sigma + 1$ . We may assume that there is some  $\mathcal{P}_\sigma$ -name  $\dot{\mathcal{G}}$  for a sequence of continuous functions  $g_\delta : \delta \rightarrow \omega$  (for  $\delta \in \text{Lim}(\omega_1^{\mathbf{V}})$ ) such that  $\Phi(\sigma)$  is a  $\mathcal{P}_\sigma$ -name for  $\mathbb{B}_{\dot{\mathcal{G}}}$  (otherwise the proof is easier). Let also  $\dot{g}_\delta$  be a name for  $g_\delta$  for every  $\delta$  as above. We start with the proof of (1) $_\alpha$ . By (1) $_\sigma$  we may assume, by extending  $q|_\sigma$  if necessary, that  $q|_\sigma$  is  $\mathcal{P}_\sigma$ -pre-generic for  $(N, \vec{\mathcal{W}})$ . If the hypothesis of (b4) fails for  $q$  and  $\sigma$ , then (b4) holds vacuously for this pair. So, assume that  $\sigma \in \text{dom}(F_q)$  and  $\{M \in \pi_0(\mathcal{W}_n) : \delta_M \in C_m^{\delta_N}\}$  is  $N$ -unbounded for all  $n, m \in \omega$ . By (3) $_\sigma$ ,  $q|_\sigma$  forces that  $\{M \in \pi_0(\mathcal{W}_n^{\dot{\mathcal{G}}_\sigma}) : \delta_M \in C_m^{\delta_N}\}$  is  $N$ -unbounded for all  $n, m \in \omega$ . By Lemma 3.13, we may assume, by extending  $q|_\sigma$  if necessary, that there is a colour  $j \in \omega$  such that  $q|_\sigma$  forces that  $j$  is not

in the range of  $\dot{g}_{\sigma, \delta_N} \upharpoonright \text{range}(f_{q, \sigma})$  and

$$\{M \in \pi_0(\mathcal{W}_n^{\dot{G}_\sigma}) : \dot{g}_{\sigma, \delta_N}(\delta_M) \neq j \text{ and } \delta_M \in C_m^{\delta_N}\}$$

is  $N$ -unbounded for all  $n, m \in \omega$ . In this case, it suffices to define  $q'$  in such a way that  $\Delta_{q'} = \Delta_q \cup \{(N, \vec{\mathcal{W}}, \alpha)\}$ ,  $F_{q'} \upharpoonright \sigma = F_q \upharpoonright \sigma$ ,  $f_{q', \sigma} = f_{q, \sigma} \cup \{(\delta_N, \delta_N)\}$  and  $b_{q', \sigma} = b_{q, \sigma} \cup \{(\delta_N, j)\}$ .

Given  $q \in \mathcal{P}_\kappa$ , we will say that a countable structure  $M$  is *relevant for  $q$  at stage  $\xi$*  if there are some  $\vec{\mathcal{W}}$  and  $\beta$  such that  $(M, \vec{\mathcal{W}}, \beta) \in \Delta_q$ ,  $\beta \geq \xi + 1$ ,  $N \in \mathcal{M}_{\xi+1}$  and  $\{M \in \pi_0(\mathcal{W}_n) : \delta_M \in C_m^{\delta_N}\}$  is  $N$ -unbounded for all  $n, m \in \omega$ . The following claim can be proved by essentially repeating the same argument as above (based on  $(3)_\sigma$  together with Lemma 3.13) finitely many times.

**Claim 3.19.** *The set of  $q \in \mathcal{P}_\alpha$  such that  $\sigma \in \text{dom}(F_q)$  is a dense subset of  $\mathcal{P}_\alpha$ . Moreover, if  $q \in \mathcal{P}_\alpha$ ,  $\sigma \notin \text{dom}(F_q)$ ,  $q|_\alpha$  forces that  $(f, b)$  is in  $\dot{\mathcal{Q}}_\sigma$ , and  $\max(\text{range}(f)) < \min\{\delta_M : M \text{ is relevant at stage } \sigma\}$ , then there is a condition  $r$  extending  $q$  such that  $\sigma \in \text{dom}(F_r)$ ,  $f \subseteq f_{r, \sigma}$  and  $b \subseteq b_{r, \sigma}$ .*

Let us proceed to the proof of  $(3)_\alpha$  now. By the above claim, we may assume that  $\sigma \in \text{dom}(F_q)$ . In particular,  $\delta_N$  is a fixed point of  $f_{q, \sigma}$ . Since  $\dot{\mathcal{G}}$  is forced to be a sequence of continuous functions, we may assume, by extending  $q|_\sigma$  if necessary, that there is an ordinal  $\eta < \delta_N$  such that  $q|_\sigma$  forces that for every  $\mu \in [\eta, \delta_N]$  and every  $\nu \in \text{dom}(b_{q, \sigma}) \setminus (\delta_N + 1)$ ,  $\dot{g}_\nu(\mu) \neq b_{q, \sigma}(\nu)$ . By condition (b4) in the definition of  $\mathcal{P}_\alpha$  and extending  $q|_\sigma$  if necessary, we can find some  $(M, \vec{\mathcal{W}}') \in \mathcal{W}_{n^*}$  such that  $\{\eta, x, f_{q, \sigma} \upharpoonright \delta_N\} \subseteq M$ ,  $\delta_M \in C_m^{\delta_N}$ ,  $(M, \vec{\mathcal{W}}', \sigma) \in \Delta_{q|_\sigma}$  and such that  $q|_\sigma$  forces that  $\dot{g}_{\delta_N}(\delta_M) \neq b_{q, \sigma}(\delta_N)$ . So,  $q|_\sigma$  forces that for all  $\nu \in \text{dom}(b_{q, \sigma}) \setminus \delta_M$ ,  $\dot{g}_\nu(\delta_M) \neq b_{q, \sigma}(\nu)$ . If  $M \notin \mathcal{M}_\alpha$  or  $\{M' \in \pi_0(\mathcal{W}'_n) : \delta_{M'} \in C_m^{\delta_M}\}$  fails to be  $M$ -unbounded for any  $m < \omega$  and  $n < \omega$ , then the pair

$$q' = (F_q, \Delta_q \cup \{(M, \vec{\mathcal{W}}', \alpha)\})$$

is a condition in  $\mathcal{P}_\alpha$  extending  $q$ , so we are done in this case. Otherwise, we can find  $q'$  using  $(2)_\sigma$  together with Lemma 3.13 (in the extension via  $\mathcal{P}_\sigma$ ) as in the relevant case of the proof of  $(1)_\alpha$ .

Now let us prove  $(2)_\alpha$ . Let  $A$  be a maximal antichain of  $\mathcal{P}_\alpha$  in  $N^*$ . By extending  $q$  if necessary we may assume that  $q$  extends a condition in  $A$ . We want to show that such a condition is in  $N$  and for this it will suffice to find a member of  $A \cap N$  compatible with  $q$ . Let  $G_\sigma$  be a  $\mathcal{P}_\sigma$ -generic filter over  $V$  with  $q|_\sigma \in G_\sigma$ . By  $(2)_\sigma$  we have that  $G_\sigma$  is also generic over  $N^*$ .



By Claim 3.19 we may assume that  $\sigma \in \text{dom}(F_q)$ . In particular,  $\delta_N$  is a fixed point of  $f_{q,\sigma}$ . As in the proof of  $(3)_\alpha$ , we may assume, by extending  $q|_\sigma$  if necessary, that there is an ordinal  $\eta < \delta_N$  such that  $q|_\sigma$  forces that for every  $\mu \in [\eta, \delta_N]$  and every  $\nu \in \text{dom}(b_{q,\sigma}) \setminus (\delta_N + 1)$ ,  $\dot{g}_\nu(\mu) \neq b_{q,\sigma}(\nu)$ . Let  $\rho = \max(\text{range}(f_{q,\sigma} \upharpoonright \delta_N))$  and define  $A^+$  as the (partially defined) function sending each  $p$  in  $A$  to the  $\Phi$ -first  $\mathcal{P}_\alpha$ -condition  $p^+$  such that  $p^+|_\sigma \in G_\sigma$ ,  $f_{p,\sigma} \upharpoonright \delta_N \subseteq f_{p^+,\sigma}$ ,  $\text{range}(f_{p,\sigma}) \cap \rho = \text{range}(f_{p^+,\sigma}) \cap \rho$  and  $b_{p,\sigma} \upharpoonright \delta_N \subseteq b_{p^+,\sigma}$ . By the  $\aleph_2$ -c.c.,  $A^+$  is not only in  $N^*[G_\sigma]$ , but in  $N[G_\sigma]$  (i.e., we can find in  $N$  a  $\mathcal{P}_\alpha$ -name for  $A^+$ ). Since  $\vec{W}$  is  $N$ - $\alpha$ -large, by condition (b4) we can find  $n \in \omega$  and  $(M, \vec{W}') \in \mathcal{W}_n$  such that

- (i)  $\vec{W}'$  is  $M$ - $\sigma$ -large and  $M = M^* \cap H(\kappa)$  for some  $M^* \in \mathcal{M}_\sigma^*$  such that  $\mathcal{P}_\sigma \in M^*$ ,
- (ii)  $(M, \vec{W}', \sigma) \in \Delta_u$  for some  $u \in G_\sigma$ ,
- (iii)  $\dot{g}_{\delta_N}(\delta_M) \neq b_{q,\sigma}(\delta_N)$ , and
- (iv)  $\{\eta, \rho, A^+\} \subseteq M[G_\sigma]$ .

Since  $q|_\sigma$  and  $u$  are both in  $G_\sigma$ , we may assume that  $q|_\sigma$  extends  $u$ . We may also assume that there is an ordinal  $\eta' \in (\eta, \delta_M)$  such that  $q|_\sigma$  forces that for every ordinal  $\mu \in [\eta', \delta_M]$ ,  $\dot{g}_{\delta_N}(\mu) \neq b_{q,\sigma}(\delta_N)$ . Now, in  $M[G_\sigma]$ , there is  $r \in \text{range}(A^+)$  such that  $\min(\text{range}(f_{r,\sigma}) \setminus \rho) > \eta'$ . Note that  $\max(\text{range}(f_{r,\sigma})) < \delta_M$  since  $\delta_M = \delta_{M[G_\sigma]}$  (this equality follows from the conjunction of (i), (ii) and  $(2)_\sigma$ ). Since  $r|_\sigma$  and  $q|_\sigma$  are both in  $G_\sigma$ , we may assume, by extending  $q|_\sigma$  if necessary, that  $q|_\sigma$  decides  $r$  and extends its corresponding restriction. Finally note that the inequalities  $\eta < \eta' < \min(\text{range}(f_{r,\sigma}) \setminus \rho) \leq \max(\text{range}(f_{r,\sigma})) < \delta_M$  ensure that  $q|_\sigma$  forces that  $F_q(\sigma)$  and  $F_r(\sigma)$  are compatible in  $\dot{Q}_\sigma$ . So,  $q$  and  $r$  are compatible.

It remains to prove the lemma for the case when  $\alpha$  is a nonzero limit ordinal. The proof of  $(1)_\alpha$  (resp.,  $(3)_\alpha$ ) is trivial using  $(1)_\beta$  (resp.,  $(3)_\beta$ ) for all  $\beta < \alpha$ , together with the fact that  $\text{dom}(F_r)$  is bounded in  $\alpha$  for any  $\mathcal{P}_\alpha$ -condition  $r$ . For  $(2)_\alpha$ , let  $E \subseteq \mathcal{P}_\alpha$  be dense and open,  $E \in N^*$ , and let  $q$  satisfy the hypothesis of  $(2)_\alpha$ . We want to find a condition in  $E \cap N^*$  compatible with  $q$ . We may assume that  $q \in E$ .

Suppose first that  $\text{cf}(\alpha) = \omega$ . In this case we may take  $\sigma \in N^* \cap \alpha$  above  $\text{dom}(F_q)$ . Let  $G$  be  $\mathcal{P}_\sigma$ -generic with  $q|_\sigma \in G$ . In  $N^*[G]$  it is true that there is a condition  $q^\circ \in \mathcal{P}_\alpha$  such that

- (a)  $q^\circ \in E$  and  $q^\circ|_\sigma \in G$ , and
- (b)  $\text{dom}(F_{q^\circ}) \subseteq \sigma$ .

(the existence of such a  $q^\circ$  is witnessed in  $\mathbf{V}[G]$  by  $q$ .)

Since  $q|_\sigma$  is  $(N^*, \mathcal{P}_\sigma)$ -generic by induction hypothesis,  $q^\circ \in N^*$ . By extending  $q$  below  $\sigma$  if necessary, we may assume that  $q|_\sigma$  decides  $q^\circ$  and extends  $q^\circ|_\sigma$ . But now, if  $q = (p, \Delta_q)$ , the natural amalgamation  $(p, \Delta_q \cup \Delta_{q^\circ})$  of  $q$  and  $q^\circ$  is a  $\mathcal{P}_\alpha$ -condition extending them.

Finally, suppose  $\text{cf}(\alpha) \geq \omega_1$ . We may assume that  $\text{dom}(F_q)$  is not bounded by  $\text{sup}(N \cap \alpha)$  as otherwise we can argue as in the  $\text{cf}(\alpha) = \omega$  case. The crucial observation now is that if  $N' \in \text{dom}(\Delta_q)$  and  $\delta_{N'} < \delta_N$ , then  $\text{sup}(N' \cap N \cap \alpha) \leq \text{sup}(\Psi_{\bar{N}, N}(N') \cap \alpha) \in N \cap \alpha$  whenever  $\bar{N} \in \text{dom}(\Delta_q)$  is such that  $\delta_{\bar{N}} = \delta_N$  and  $N' \in \bar{N}$ . To see this, recall that  $\Psi_{\bar{N}, N}$  fixes  $\bar{N} \cap N \cap \kappa$ . Also,  $\text{sup}(\Psi_{\bar{N}, N}(N') \cap \alpha) \in N \cap \alpha$  since  $\Psi_{\bar{N}, N}(N')$  is countable in  $N$  and  $\alpha$  has uncountable cofinality. The symmetry of the systems  $\text{dom}(\Delta_q)$  is needed precisely to derive the conclusion that  $\text{sup}(N' \cap N \cap \alpha) < \text{sup}(N \cap \alpha)$  for every  $N' \in \text{dom}(\Delta_q)$  with  $\delta_{N'} < \delta_N$ .

Hence we may fix  $\sigma \in N \cap \alpha$  such that:

- (i)  $\text{sup}(N' \cap N \cap \alpha) < \sigma$  for all  $N' \in \text{dom}(\Delta_q)$  with  $\delta_{N'} < \delta_N$ , and
- (ii) if  $\eta \in \text{dom}(F_q)$  and  $\eta < \text{sup}(\alpha \cap N)$ , then  $\eta < \sigma$ .

As in the case  $\text{cf}(\alpha) = \omega$ , if  $G_\sigma$  is  $\mathcal{P}_\sigma$ -generic with  $q|_\sigma \in G_\sigma$ , then in  $N^*[G_\sigma]$  we can find a condition  $q^\circ \in \mathcal{P}_\alpha$  such that  $q^\circ \in E$  and  $q^\circ|_\sigma \in G_\sigma$ , and such a  $q^\circ$  will necessarily be in  $N^*$  by  $(2)_\sigma$ . By extending  $q$  below  $\sigma$  we may assume that  $q|_\sigma$  decides  $q^\circ$  and extends  $q^\circ|_\sigma$ . The proof of  $(2)_\alpha$  in this case will be finished if we can show that there is a condition  $q^\dagger$  extending  $q$  and  $q^\circ$ . The condition  $q^\dagger$  can be built by recursion on  $\text{dom}(F_{q^\circ}) \setminus \sigma$  (note that  $\text{sup}(N \cap \alpha) \leq \min(\text{dom}(F_q) \setminus \sigma)$  by the choice of  $\sigma$ , and therefore  $\min(\text{dom}(F_q) \setminus \sigma) > \max(\text{dom}(F_{q^\circ}))$ ). The details of this construction, which for the sake of completeness we will sketch here, appear in [6].

Suppose  $|\text{dom}(F_{q^\circ}) \setminus \sigma| = n$  and let  $\mu = \max(\text{dom}(F_{q^\circ}))$ . Applying  $n$  times Claim 3.19, we get a condition  $q^\Delta \in \mathcal{P}_{\mu+1}$  extending both  $q|_{\mu+1}$  and  $q^\circ|_{\mu+1}$  (note that if  $\xi \in \text{dom}(F_{q^\circ}) \setminus \sigma$  and  $M$  is relevant for  $q$  at stage  $\xi$ , then the choice of  $\sigma$  ensures that  $\delta_M \geq \delta_N$ ). Finally note that if  $\xi \in (\mu, \alpha) \cap \text{dom}(F_q)$  and  $M$  is relevant either for  $q$ , or for  $q^\circ$  or for  $q^\Delta$  at stage  $\xi$ , then  $M$  must be relevant for  $q$  (the choice of  $\sigma$  shows in fact that  $M$  cannot be relevant for  $q^\circ$ . On the other hand, every  $M$  associated to  $q^\Delta$  is relevant at most at stage  $\mu + 1$  since  $q^\Delta \in \mathcal{P}_{\mu+1}$ ). Therefore,

$$q^\dagger = (F_{q^\Delta} \cup (F_q \upharpoonright [\mu + 1, \alpha)), \Delta_{q^\Delta} \cup \Delta_{q^\circ} \cup \Delta_q)$$

is a condition in  $\mathcal{P}_\alpha$  extending both  $q$  and  $q^\circ$ . □

The following corollary follows immediately from Lemmas 3.16 and 3.18.

**Corollary 3.20.**  $\mathcal{P}_\kappa$  is proper.

**Corollary 3.21.**  $\mathcal{P}_\kappa$  forces that  $\vec{C}$  is a WCG-sequence.

*Proof.* Let  $\dot{C}$  be a  $\mathcal{P}_\kappa$ -name for a club of  $\omega_1$  and let  $q$  be a  $\mathcal{P}_\kappa$ -condition. By the  $\aleph_2$ -c.c. of  $\mathcal{P}_\kappa$  there is some  $\alpha < \kappa$  such that  $\dot{C}$  is a  $\mathcal{P}_\alpha$ -name and such that  $q$  is a condition in  $\mathcal{P}_\alpha$ . We may also assume that  $\dot{C}$  is in  $H(\kappa)$ . Let us take a countable  $N^* \preceq H(\theta)$  (for a large enough  $\theta$ ) containing everything relevant (this includes  $\dot{C}$ ). Let  $N = N^* \cap H(\kappa)$ .

By Lemma 3.16 there is a  $\subseteq$ -decreasing  $\omega$ -sequence  $\vec{W}$  such that  $\vec{W}$  is  $N$ - $(\alpha + 1)$ -large. By Lemma 3.18 (1) $_\alpha$  there is a condition  $q'$  in  $\mathcal{P}_\alpha$  extending  $q$  and  $\mathcal{P}_\alpha$ -pre-generic for  $(N, \vec{W})$ . It suffices to show that  $q'$  forces  $\dot{C} \cap C_m^{\delta_N} \neq \emptyset$  for every  $m \in \omega$ .

Notice that the hypothesis of (3) $_\alpha$  in Lemma 3.18 is realized by  $q'$ . Hence, for every  $n, m < \omega$ , and every  $q''$  extending  $q'$  there is some  $\mathcal{P}_\alpha$ -condition  $q^*$  extending  $q''$  and there is some  $(M, \vec{W}')$  in  $\mathcal{W}_n$  such that  $\dot{C} \in M$ ,  $\delta_M \in C_m^{\delta_N}$ , and such that  $q^*$  is  $\mathcal{P}_\alpha$ -pre-generic for  $(M, \vec{W}')$ . But by (3) in the definition of  $N$ - $(\alpha + 1)$ -large we may take  $n$  large enough so that  $\vec{W}'$  is necessarily  $M$ - $\alpha$ -large. Then, by Lemma 3.18 (2) $_\alpha$  we have that  $q^*$  is  $(M^*, \mathcal{P}_\alpha)$ -generic for any  $M^* \in \mathcal{M}_\alpha$  containing  $\mathcal{P}_\alpha$  such that  $M^* \cap H(\kappa) = M$ . But then  $q^*$  forces  $\delta_M \in \dot{C}$ .  $\square$

Finally, by arguing as in the proof of Theorem 1.13 (i.e., by the usual book-keeping arguments using the  $\aleph_2$ -c.c. of  $\mathcal{P}_\kappa$ ) we can prove that  $\mathcal{P}_\kappa$  forces  $\neg\mathfrak{U}$ , MA and FA( $\mathbb{B}$ ). This completes the proof of Theorem 1.14.

#### 4. ON POLYCHROMATIC RAMSEY THEORY

Polychromatic Ramsey theory<sup>15</sup> (see [9], [2] and [1]) is the study of colourings  $f$  of sets of the form  $[X]^n$ , focusing on the existence or non-existence of large sets  $Y \subseteq X$  such that  $c \upharpoonright [Y]^n$  is a one-to-one function (such a set  $Y$  is usually called a *rainbow for  $c$* ). Here we consider the following negative polychromatic partition relation (see [1]).

**Definition 4.1.**  $\omega_2 \not\rightarrow_{\text{poly}} (\omega_1)_{2\text{-bd}}^2$  means that there is a function  $f$  with the following properties.

- (a)  $\text{dom}(f) = [\omega_2]^2$
- (b)  $f$  is 2-bounded (which means that  $|f^{-1}(x)| \leq 2$  for every  $x$ ).

<sup>15</sup>Also known as anti-Ramsey theory

- (c)  $f$  does not have rainbows of order-type  $\omega_1$  (i.e., there is no  $Y \subseteq \omega_2$  of order type  $\omega_1$  such that the restriction of  $f$  to  $[Y]^2$  is one to one).

In [1], Abraham and Cummings show the consistency of the relation  $\omega_2 \dashrightarrow_{\text{poly}} (\omega_1)_{2\text{-bd}}^2$ . In their model,  $2^{\aleph_0} = \aleph_2$ . One of the questions asked in [1] is whether  $\omega_2 \dashrightarrow_{\text{poly}} (\omega_1)_{2\text{-bd}}^2$  is consistent together with  $2^{\aleph_0} > \aleph_2$ . Here we give an affirmative answer to this question. In fact, we show that if CH holds and  $\lambda$  is any cardinal, then there is a proper poset with the  $\aleph_2$ -c.c. forcing  $\omega_2 \dashrightarrow_{\text{poly}} (\omega_1)_{2\text{-bd}}^2$  together with  $\text{FA}(\mathbb{B})_\lambda$ . Our result is the following.

**Theorem 4.2.** (CH) *If  $\kappa$  is a regular cardinal such that  $2^{<\kappa} = \kappa$ , then there exists a proper forcing notion  $\mathcal{P}$  with the  $\aleph_2$ -chain condition such that  $\text{FA}(\mathbb{B})$ ,  $2^\lambda = \kappa$  for every infinite  $\lambda < \kappa$ , and  $\omega_2 \dashrightarrow_{\text{poly}} (\omega_1)_{2\text{-bd}}^2$  hold in the generic extension by  $\mathcal{P}$ .*

Starting from CH and using symmetric systems,<sup>16</sup> Abraham and Cummings generically add a function  $c$  such that

- (1)  $c : [\omega_2]^2 \longrightarrow \omega_1$ ,
- (2) There are no  $\alpha_0 < \alpha_1 < \alpha_2 < \beta < \omega_2$  such that  $c(\alpha_0, \beta) = c(\alpha_1, \beta) = c(\alpha_2, \beta)$ , and
- (3) For every  $X \subseteq \omega_2$  of order type  $\omega_1$  there are  $\alpha_0 < \alpha_1 < \beta$  in  $X$  such that  $c(\alpha_0, \beta) = c(\alpha_1, \beta)$ .

It is straightforward to check that  $f(\alpha, \beta) = (c(\alpha, \beta), \beta)$  witnesses  $\omega_2 \dashrightarrow_{\text{poly}} (\omega_1)_{2\text{-bd}}^2$ . Let us assume the hypotheses of Theorem 4.2 and let  $\Phi : \kappa \longrightarrow H(\kappa)$  be a bijection. Let also  $(e_\alpha)_{\alpha < \omega_2}$  be  $\Phi$ -first such that each  $e_\alpha$  is a bijection between  $\alpha$  and  $|\alpha|$ . The following is (essentially) the forcing in [1] for adding  $c$ , which we will call  $(\mathcal{P}_0, \leq_0)$ .

**Definition 4.3.** Conditions in  $\mathcal{P}_0$  are triples  $p = (\emptyset, c_p, \Delta_p)$  with the following properties.

- (1)  $c_p$  is a finite partial function from  $[\omega_2]^2$  to  $\omega_1$ .
- (2) There are no  $\alpha_0 < \alpha_1 < \alpha_2 < \beta$  in the domain of  $c_p$  such that  $c_p(\alpha_0, \beta) = c_p(\alpha_1, \beta) = c_p(\alpha_2, \beta)$ .
- (3) For every  $(\alpha, \beta) \in \text{dom}(c_p)$ ,  $c_p(\alpha, \beta) \geq e_\beta(\alpha)$ .
- (4)  $\Delta_p$  is a finite set of pairs of the form  $(N, 0)$  such that  $\text{dom}(\Delta_p)$  is a  $\Phi$ -symmetric system.

Given  $p$  and  $q$  conditions in  $\mathcal{P}_0$ ,  $q \leq_0 p$  whenever the following holds:

- (i)  $c_p \subseteq c_q$ ,

<sup>16</sup>In our terminology.

- (ii)  $\Delta_p \subseteq \Delta_q$ , and
- (iii) for every  $(\alpha, \beta) \in \text{dom}(c_q) \setminus \text{dom}(c_p)$  and every pair  $(N, 0) \in \Delta_p$ , if  $\{\alpha, \beta\} \in N$ , then  $c_q(\alpha, \beta) \in N$ .

In [1], what we have called here  $\Delta_q$  is an actual  $\Phi$ -symmetric system (i.e., there is no mention of 0's). Here we add the 'marker' 0 to the  $N$ 's occurring in  $\Delta_q$  – as well as the first vacuous component  $\emptyset$ , which of course does not show up in [1] either – since we are going to incorporate this forcing into an iteration with symmetric systems of structures and markers as in the other constructions in this paper. The restriction to 0 of any condition in  $\mathcal{P}_\alpha$ , for any  $\alpha > 0$ , will be a condition in  $\mathcal{P}_0$  by definition, and this is the need for the  $\emptyset$ 's and 0's in Definition 4.3. Another difference between the original definition and the poset in Definition 4.3 is that the  $N$ 's occurring in conditions in [1] are countable elementary substructures of  $H(\omega_2)$  and not of  $H(\kappa)$ . These differences have no effect whatsoever on the proofs of the following lemmas from [1].

**Lemma 4.4.** *Let  $p = (\emptyset, c_p, \Delta_p) \in \mathcal{P}_0$  and let  $\alpha_0 \leq \alpha_1 < \beta < \omega_2$  be such that*

- (1) none of  $\{\alpha_0, \beta\}$ ,  $\{\alpha_1, \beta\}$  is in  $\text{dom}(c_p)$ , and
- (2) for every  $N \in \text{dom}(\Delta_p)$ ,  $\{\alpha_0, \beta\} \in N$  if and only if  $\{\alpha_1, \beta\} \in N$ .

*Then there is  $q = (\emptyset, c_q, \Delta_q) \leq_0 p$  such that  $\{\alpha_0, \beta\}$ ,  $\{\alpha_1, \beta\} \in \text{dom}(c_q)$  and  $c_q(\alpha_0, \beta) = c_q(\alpha_1, \beta)$ .*

**Lemma 4.5.**  *$\mathcal{P}_0$  has the  $\aleph_2$ -c.c.*

**Lemma 4.6.** *Let  $\theta > |\mathcal{P}_0|$  be a regular cardinal,  $p = (\emptyset, c_p, \Delta_p) \in \mathcal{P}_0$ , and  $N^*$  a countable elementary substructure of  $H(\theta)$  such that  $p$  and  $\Phi$  are in  $N^*$ . Let  $N = N^* \cap H(\kappa)$  and let  $p^+ = (\emptyset, c_p, \Delta_p \cup \{(N, 0)\})$ . Then the following holds.*

- (1)  $p^+ \leq_0 p$ ,
- (2) If  $q = (\emptyset, c_q, \Delta_q) \leq_0 p^+$  and  $r = (\emptyset, c_q \upharpoonright N, \Delta_q \cap N)$ , then the following are true:
  - (2.1)  $r \in N \cap \mathcal{P}_0$
  - (2.2) If  $s \in N \cap \mathcal{P}_0$  and  $s \leq_0 r$ , then  $q$  and  $s$  are  $\leq_0$ -compatible.
- (3)  $p^+$  is an  $(N^*, \mathcal{P}_0)$ -generic condition.

Lemma 4.6 implies of course that  $\mathcal{P}_0$  is strongly proper.

The forcing  $\mathcal{P}$  witnessing Theorem 4.2 is  $(\mathcal{P}_\kappa, \leq_\kappa)$ , where  $\mathcal{P}_0$  is the poset in Definition 4.3 and  $\langle (\mathcal{P}_\alpha, \leq_\alpha) : 1 \leq \alpha \leq \kappa \rangle$  is defined as follows.

Let  $\alpha$ ,  $1 \leq \alpha \leq \kappa$ , and suppose  $\mathcal{P}_\xi$  has been defined for all  $\xi < \alpha$ . Conditions in  $\mathcal{P}_\alpha$  are triples of the form  $q = (F_q, c_q, \Delta_q)$  with the following properties.

- (b0)  $F_q$  is a finite function with  $\text{dom}(F_q) \subseteq \alpha$ .
- (b1)  $\Delta_q$  is a finite set of pairs of the form  $(N, \gamma)$ , with  $N$  a countable elementary substructure of  $(H(\kappa), \in, \Phi)$  and  $\gamma \leq \alpha \cap \text{sup}(N \cap \kappa)$ .
- (b2) For all  $\xi < \alpha$ , the restriction of  $q$  to  $\xi$ , denoted<sup>17</sup>  $q|_\xi$ , is in  $\mathcal{P}_\xi$ . This restriction is defined as the triple

$$q|_\xi := (F_q \upharpoonright \xi, c_q, \{(N, \min\{\gamma, \xi\}) : (N, \gamma) \in \Delta_q\})$$

- (b3) If  $\xi \in \text{dom}(F_q)$ , then  $F_q(\xi) \in \mathbb{B}$ .
- (b4) If  $\xi \in \text{dom}(F_q)$ ,  $(N, \beta) \in \Delta_q$ ,  $\beta \geq \xi + 1$ , and  $N \in \mathcal{M}_{\xi+1}$ , then  $\delta_N$  is a fixed point of  $F_q(\xi)$ .

Given conditions  $q^\epsilon = (F_\epsilon, c_\epsilon, \Delta_\epsilon)$  (for  $\epsilon \in \{0, 1\}$ ) in  $\mathcal{P}_\alpha$ ,  $q^1 \leq_\alpha q^0$  if and only if

- (c1)  $q^1|_0 \leq_0 q^0|_0$ ,
- (c2)  $\text{dom}(F_0) \subseteq \text{dom}(F_1)$  and, for all  $\xi \in \text{dom}(F_0)$ ,  $F_1(\xi) \supseteq F_0(\xi)$ , and
- (c3)  $\Delta_0 \subseteq \Delta_1$

The proofs of the corresponding versions of Lemmas 3.3, 3.4 3.5 and 3.7 are identical to the proofs in [6]. One can also show that  $\mathcal{P}_\kappa$  forces  $\text{FA}(\mathbb{B})$  by a standard argument using Lemmas 3.3 and 3.4 (cf. [6]).

It remains to prove that  $\mathcal{P}_\kappa$  forces  $\omega_2 \dashv\vdash_{\text{poly}} (\omega_1)_{2\text{-bd}}^2$ . This is an immediately consequence of Lemma 4.7. This lemma is a variant of a corresponding lemma in [1].

**Lemma 4.7.** *Let  $p \in \mathcal{P}_\kappa$  and let  $\dot{X}$  be a  $\mathcal{P}_\kappa$ -name. Assume that  $p$  forces that  $\dot{X}$  is a subset of  $\omega_2$  of order type  $\omega_1$ . Then there are  $t \leq_\kappa p$  and ordinals  $\alpha_0 < \alpha_1 < \beta$  in  $\text{dom}(c_t)$  such that*

- (1)  $t$  forces that  $\alpha_0, \alpha_1$  and  $\beta$  are in  $\dot{X}$ , and
- (2)  $c_t(\alpha_0, \beta) = c_t(\alpha_1, \beta)$ .

*Proof.* By extending  $p$  if necessary, we may start assuming that there is some  $\gamma$  such that  $p$  forces that  $\text{sup}(\dot{X}) = \gamma$ . Let  $\theta$  be a sufficiently large regular cardinal and let  $N^* \preceq H(\theta)$  be countable and such that  $p$ ,  $\mathcal{P}_\kappa$ ,  $\Phi$  and  $\dot{X}$  are in  $N^*$ . Let  $N = N^* \cap H(\kappa)$  and let  $p_0^+ \in \mathcal{P}_\kappa$  extend  $p$  and such that  $(N, \text{sup}(N \cap \kappa)) \in \Delta_{p_0^+}$ . By the relevant form of Lemma 3.7,  $p_0^+$  is  $(N^*, \mathcal{P}_\alpha)$ -generic. Since  $\text{cf}(\gamma) = \omega_1$ ,  $N \cap \gamma$  is bounded in  $\gamma$ . By extending  $p_0^+$  if necessary we may assume that there is an ordinal  $\beta$  such that  $\text{sup}(N \cap \gamma) < \beta < \gamma$  and  $p_0^+$  forces  $\beta \in \dot{X}$ . Let  $m < \omega$  be

<sup>17</sup>As usual.

bigger than the collection of all subsets of  $\Delta_{p_0^+}$ , let  $F_0$  be the function with domain  $\text{dom}(F_{p_0^+}) \cap N$  and such that  $F_0(\xi) = F_{p_0^+}(\xi) \upharpoonright \delta_N$  for all  $\xi \in \text{dom}(F_0)$ , and let  $p_0 = (F_0, c_{p_0^+} \cap N, \Delta_{p_0^+} \cap N)$ .

Now we can build increasing sequences of conditions  $(p_i^+)_{1 \leq i \leq m}$  and  $(p_i)_{1 \leq i \leq m}$ , together with a sequence  $(\alpha_i)_{1 \leq i \leq m}$  of distinct ordinals such that for all positive  $i \leq m$ ,

- (a)  $p_i \in N$  extends  $p_{i-1}$  and forces  $\alpha_i \in \dot{X}$ ,
- (b)  $p_i^+$  is a condition extending  $p_{i-1}^+$ ,
- (c)  $\Delta_{p_i^+} \setminus N = \Delta_{p_{i-1}^+} \setminus N$ , and
- (d)  $\text{dom}(F_{p_i}) = \text{dom}(F_{p_i^+}) \cap N$  and  $F_{p_i}(\xi) = F_{p_i^+}(\xi) \upharpoonright \delta_N$  for all  $\xi \in \text{dom}(F_{p_i})$ ,  $c_{p_i} = c_{p_i^+} \cap N$ , and  $\Delta_{p_i} = \Delta_{p_i^+} \cap N$ .

Given a positive  $i \leq m$ , if  $p_i^+$  and  $p_i$  – namely, its restriction to  $N$  – have been defined, we can find  $p_i \in N$  by correctness of  $N$  since  $p_{i-1}^+$  forces  $\beta \in \dot{X}$  and  $\beta \notin N$ . But then we can easily find  $p_i^+$ . The point is that the forcings we are plugging in at nonzero stages  $\xi$  of the iteration are  $\mathbb{B}$ , so that, for every such  $\xi$  in  $\text{dom}(F_{p_i})$ , it is indeed easy to find an extension of  $F_{p_i}(\xi)$  which is generic for the relevant structures  $N'$  which are not in  $N$  simply by adding  $\delta_{N'}$  as fixed point. The point is of course that we do not need to go to any extension of  $p_i \upharpoonright \xi$  to do that.

In the end we can find two  $i, i' \leq m$  and an extension  $q$  of  $p_m^+$  such that  $c_q(\{\alpha_i, \beta\}) = c_q(\{\alpha_{i'}, \beta\})$  as in the proof (cf. [1]) of Lemma 4.4. More specifically, we may find distinct  $i, i'$  such that for all  $M \in \text{dom}(\Delta_{p_0^+})$ ,  $\{\alpha_i, \beta\} \in M$  if and only if  $\{\alpha_{i'}, \beta\} \in M$ , and then we may fix some  $\nu > e_\beta(\max\{\alpha_i, \alpha_{i'}\})$  such that  $\nu \in M$  for all  $M \in \text{dom}(\Delta_{p_0^+})$  such that  $\{\alpha_i, \beta\} \in M$  (equivalently, such that  $\{\alpha_{i'}, \beta\} \in M$ ). Then we may let  $c_q(\{\alpha_i, \beta\}) = c_q(\{\alpha_{i'}, \beta\}) = \nu$ . This finishes the proof.  $\square$

It is worth pointing out that if we were to force the forcing axiom for Cohen forcing rather than  $\text{FA}(\mathbb{B})$  (together with  $2^{\aleph_0} = \kappa$ ), the construction could be made a bit simpler, in that we could dispense with the markers  $\gamma$  – in  $(N, \gamma) \in \Delta_q$ , for a condition  $q$  – in the definition of our iteration. Such a construction would of course suffice also to answer the question in [1].

## REFERENCES

- [1] U. Abraham and J. Cummings, *More results in polychromatic Ramsey theory*, Central European Journal of Math., vol. 10, 3 (2012), 1004–1016.
- [2] U. Abraham, J. Cummings and C. Smyth, *Some results in polychromatic Ramsey theory*, J. Symbolic Logic, vol. 72, 3 (2007), 865–896.
- [3] D. Asperó, *The consistency of a club-guessing failure at the successor of a regular cardinal*, submitted.

- [4] D. Asperó, *Adding many Baumgartner clubs*, submitted.
- [5] D. Asperó and S. Friedman, *Large cardinals and locally defined well-orders of the universe*, *Ann. Pure and Applied Logic*, vol. 157, 1 (2009), 1–15.
- [6] D. Asperó and M.A. Mota, *Forcing consequences of PFA together with the continuum large*, *Trans. of the American Math. Soc.*, to appear (2013).
- [7] D. Asperó and M.A. Mota, *A generalization of Martin's Axiom*, submitted.
- [8] J.E. Baumgartner, *Applications of the Proper Forcing Axiom*, in *Handbook of Set Theoretic Topology*, K. Kunen and J.E. Vaughan (Eds.), North Holland Publishing Co., Amsterdam (1984).
- [9] F. Galvin, *Advanced problems: 6034*, *American Math. Monthly*, 1975, 82 (5), 529.
- [10] T. Ishiu, *The comparison of various club guessing principles*, submitted.
- [11] T. Ishiu and P. Larson, *Some results about (+) proved by iterated forcing*, *J. Symbolic Logic* vol. 77, no. 2 (2012), 515–531.
- [12] T. Jech, *Set Theory: The Third Millenium Edition, Revised and Expanded*, Springer, Berlin (2002).
- [13] I. Juhasz, *A weakening of  $\clubsuit$ , with applications to topology*, *Commentationes Mathematicae Universitatis Carolinae*, vol. 29, 4 (1988), 767–773.
- [14] P. Koszmider, *On strong chains of uncountable functions*, *Israel Journal of Math.*, vol. 118 (2000), 289–315.
- [15] K. Kunen, *Set Theory, An introduction to independence proofs*, North-Holland Publishing Company, Amsterdam (1980).
- [16] J.T. Moore, *Set Mapping Reflection*, *Journal of Math. Logic*, vol. 5, 1 (2005), 87–98.
- [17] P. Nyikos, *Antidiamond and anti-PFA axioms and topological applications [incomplete preliminary version]*, unpublished note.
- [18] H. Sakai, *Preservation of  $\neg$ CG by finite support product of Cohen forcing*, unpublished note.
- [19] S. Shelah, *Proper and improper forcing*, Springer, Berlin (1998).
- [20] S. Shelah, *NNR revisited*, [Sh:656] (2000), arXiv:math.LO/0003115.
- [21] S. Todorćević, *Directed sets and cofinal types*, *Trans. American Math. Society*, vol. 290, 2 (1985), 711–723.
- [22] S. Todorćević, *Partition problems in topology*, *Contemporary Math.*, vol. 84, Providence, RI (1989).
- [23] T. Yorioka, *A comment on Asperó–Mota iteration*, preprint (2013).
- [24] J. Zapletal, *Characterization of the club forcing*, *Papers on general topology and applications (Gorham, ME, 1995)*, *Ann. of the New York Academy of Sciences*, vol. 806, 1996, pp. 476484.
- [25] J. Zapletal, *A dichotomy for forcing notions*, *Math. Research Letters*, vol. 5 (1998), 213–226.

DAVID ASPERÓ, SCHOOL OF MATHEMATICS, UNIVERSITY OF EAST ANGLIA,  
NORWICH NR4 7TJ, UK

*E-mail address:* d.aspero@uea.ac.uk

MIGUEL ANGEL MOTA, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF  
TORONTO, TORONTO, ONTARIO, CANADA M5S 2E4

*E-mail address:* motagaytan@gmail.com