



Significant text differences from Bakker et al. (2014) are shown in italic.

A multi-decade record of high-quality fCO₂ data in version 3 of the Surface Ocean CO₂ Atlas (SOCAT)

Dorothee C. E. Bakker¹, Benjamin Pfeil^{2,3}, Camilla S. Landa^{2,3}, Nicolas MetzI⁴, Kevin M. O'Brien^{5,6}, Are
5 Olsen^{2,3}, Karl Smith^{5,6}, Cathy Cosca⁵, Sumiko Harasawa⁷, Stephen D. Jones⁸, Shin-ichiro Nakaoka⁷,
Yukihiro Nojiri⁷, Ute Schuster⁸, Tobias Steinhoff⁹, Colm Sweeney^{10,11}, Taro Takahashi¹², Bronte
Tilbrook^{13,14}, Chisato Wada⁷, Rik Wanninkhof¹⁵, Simone R. Alin⁵, Carlos F. Balestrini¹⁶, Leticia
Barbero^{15,17}, Nicholas R. Bates^{18,19}, Alejandro A. Bianchi¹⁶, Frédéric Bonou²⁰, Jacqueline Boutin⁴, Yann
Bozec²¹, Eugene F. Burger⁵, Wei-Jun Cai²², Robert D. Castle¹⁵, Liqi Chen^{23,24}, Melissa Chierici^{25,26}, Kim
10 Currie²⁷, Wiley Evans^{5,28,29}, Charles Featherstone¹⁵, Richard A. Feely⁵, Agneta Fransson³⁰, Catherine
Goyet^{31,32}, Naomi Greenwood³³, Luke Gregor³⁴, Steven Hankin^{5,6}, Nick J. Hardman-Mountford³⁵, Jérôme
Harlay³⁶, Judith Hauck³⁷, Mario Hoppema³⁷, Matthew P. Humphreys¹⁹, Christopher W. Hunt³⁸, Betty
Huss¹⁵, J. Severino P. Ibánhez^{39,20}, Truls Johannessen^{2,3,40}, Ralph Keeling⁴¹, Vassilis Kitidis⁴², Arne
Körtzinger⁹, Alex Kozyr⁴³, Evangelia Krasakopoulou⁴⁴, Akira Kuwata⁴⁵, Peter Landschützer⁴⁶, Siv K.
15 Lauvset^{40,2}, Nathalie Lefèvre⁴, Claire Lo Monaco⁴, Ansley Manke⁵, Jeremy T. Mathis⁵, Liliane Merlivat⁴,
Frank J. Millero⁴⁷, Pedro M. S. Monteiro³⁴, David R. Munro⁴⁸, Akihiko Murata⁴⁹, Timothy
Newberger^{10,11}, Abdirahman M. Omar^{40,2}, Tsuneo Ono⁵⁰, Kristina Paterson¹³, David Pearce³³, Denis
Pierrot^{15,17}, Lisa L. Robbins⁵¹, Shu Saito⁵², Joe Salisbury³⁸, Reiner Schlitzer³⁷, Bernd Schneider⁵³, Roland
Schweitzer⁵⁴, Rainer Sieger³⁷, Ingunn Skjelvan^{40,2}, Kevin F. Sullivan^{15,17}, Stewart C. Sutherland¹²,
20 Adrienne J. Sutton^{5,6}, Kazuaki Tadokoro⁴⁵, Maciej Telszewski⁵⁵, Matthias Tuma⁵⁶, Steven M. A. C. van
Heuven⁵⁷, Doug Vandemark³⁸, Brian Ward⁵⁸, Andrew J. Watson⁸, Suqing Xu²³

¹Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, United Kingdom

²Geophysical Institute, University of Bergen, 5020 Bergen, Norway

25 ³Bjerknes Centre for Climate Research, 5007 Bergen, Norway

⁴Sorbonne Universités (UPMC, Univ Paris 06), CNRS, IRD, MNHN, LOCEAN Laboratory, 4, Place Jussieu, F-75005 Paris, France

⁵Pacific Marine Environmental Laboratory, National Oceanic and Atmospheric Administration, Seattle, WA 98115, USA

⁶Joint Institute for the Study of the Atmosphere and Ocean, University of Washington, Seattle, WA 98105, USA

30 ⁷National Institute for Environmental Studies, Tsukuba, Ibaraki, 305-8506, Japan

⁸College of Life and Environmental Sciences, University of Exeter, Exeter EX4 4QE, United Kingdom

⁹GEOMAR, Helmholtz Centre for Ocean Research, 24105 Kiel, Germany

¹⁰Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO 80309, USA

¹¹Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO 80305-3337, USA

35 ¹²Lamont Doherty Earth Observatory, Columbia University, Palisades, NY 10964, USA

¹³CSIRO Oceans and Atmosphere, Hobart, Tasmania 7001, Australia

¹⁴Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania 7001, Australia

¹⁵Atlantic Oceanographic and Meteorological Laboratory, National Oceanic and Atmospheric Administration, Miami, FL 33149, USA

40 ¹⁶Departemento Oceanografía, Servicio de Hidrografía Naval, C1270ABV Buenos Aires, Argentina



- ¹⁷Cooperative Institute for Marine and Atmospheric Studies, Rosenstiel School for Marine and Atmospheric Science, University of Miami, Miami, FL 33149-1098, USA
- ¹⁸Bermuda Institute of Ocean Sciences, Ferry Reach, St. Georges, GE01, Bermuda
- ¹⁹Ocean and Earth Science, University of Southampton, Southampton SO14 3ZH, United Kingdom
- 5 ²⁰Centro de Estudos e Ensaio em Risco e Modelagem Ambiental, Universidade Federal de Pernambuco, 50740-550 Recife, Brazil
- ²¹Sorbonne Universités, UPMC Univ Paris 06, CNRS, Adaptation et Diversité en Milieu Marin (UMR7144), Station Biologique de Roscoff, 29680 Roscoff, France
- ²²School of Marine Science and Policy, University of Delaware, Newark, DE 19716, USA
- 10 ²³Key Laboratory of Global Change and Marine-Atmospheric Chemistry, Third Institute of Oceanography, State Oceanic Administration, Xiamen, 361005, P. R. China
- ²⁴Chinese Arctic and Antarctic Administration, Beijing, 100860, P. R. China
- ²⁵Institute of Marine Research, 9294 Tromsø, Norway
- ²⁶Department of Marine Sciences, University of Gothenburg, 40530 Gothenburg, Sweden
- 15 ²⁷National Institute of Water and Atmospheric Research, Dunedin 9054, New Zealand
- ²⁸Ocean Acidification Research Center, University of Alaska Fairbanks, Fairbanks, AK 99775-7220, USA
- ²⁹Hakai Institute, British Columbia V0P 1H0, Canada
- ³⁰Norwegian Polar Institute, Fram Centre, 9296 Tromsø, Norway
- ³¹IMAGES_ESPACE-DEV, Université de Perpignan, 66860 Perpignan, France
- 20 ³²UMR ESPACE-DEV, Maison de la teledétection, 34000 Montpellier, France
- ³³Centre for Environment, Fisheries and Aquaculture Science, Lowestoft NR33 0HT, United Kingdom
- ³⁴Ocean Systems and Climate, CSIR-CHPC, Cape Town, 7700, South Africa
- ³⁵CSIRO Oceans and Atmosphere, Floreat WA 6014, Australia
- ³⁶University of Hawaii at Manoa, Department of Oceanography, Honolulu, HI 96822, USA
- 25 ³⁷Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, 27515 Bremerhaven, Germany
- ³⁸Ocean Process Analysis Laboratory, University of New Hampshire, Durham, NH 03824, USA
- ³⁹IRD - Institut de Recherche pour le Développement, Lago Sul, 71640-230 Brasilia, Brazil
- ⁴⁰Uni Research Climate, Bjerknes Centre for Climate Research, 5007 Bergen, Norway
- ⁴¹University of California, San Diego, CA 92093, USA
- 30 ⁴²Plymouth Marine Laboratory, Plymouth PL1 3DH, United Kingdom
- ⁴³Carbon Dioxide Information Analysis Center, Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831-6290, USA
- ⁴⁴University of the Aegean, Department of Marine Sciences, 81100, Mytilene, Lesvos Island, Greece
- ⁴⁵Tohoku National Fisheries Research Institute, Fisheries Research Agency, Shiogama, Miyagi, 985-0001, Japan
- 35 ⁴⁶Environmental Physics, Institute of Biogeochemistry and Pollutant Dynamics, ETH Zürich, 8092 Zürich, Switzerland
- ⁴⁷Department of Ocean Sciences, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, FL 33149-1031, USA
- ⁴⁸Department of Atmospheric and Oceanic Sciences and Institute of Arctic and Alpine Research, University of Colorado, Boulder, CO 80309-0450, USA
- 40 ⁴⁹Japan Agency for Marine-Earth Science and Technology, Yokosuka, Kanagawa, 237-0061, Japan
- ⁵⁰National Research Institute for Fisheries Science, Fisheries Research Agency, Yokohama 236-8648, Japan
- ⁵¹US Geological Survey, Saint Petersburg, FL 33701, USA
- ⁵²Marine Division, Global Environment and Marine Department, Japan Meteorological Agency, Tokyo 100-8122, Japan
- ⁵³Leibniz Institute for Baltic Sea Research, D-18119 Rostock (Warnemünde), Germany
- 45 ⁵⁴WeatherTop Consulting LLC, College Station, TX 77845, USA
- ⁵⁵International Ocean Carbon Coordination Project, Institute of Oceanology of the Polish Academy of Sciences, 81-712 Sopot, Poland
- ⁵⁶WCRP Joint Planning Staff, World Meteorological Organization, CH-1211 Geneva 2, Switzerland
- ⁵⁷Royal Netherlands Institute for Sea Research, 1797 SZ 't Horntje, Texel, The Netherlands
- 50 ⁵⁸AirSea Laboratory, School of Physics and Ryan Institute, National University of Ireland, Galway, Ireland



Correspondence to: Dorothee C. E. Bakker (d.bakker@uea.ac.uk)

Abstract. The Surface Ocean CO₂ Atlas (SOCAT) is a synthesis of quality-controlled fCO₂ (fugacity of carbon dioxide) values for the global surface oceans and coastal seas with regular updates. *Version 3 of SOCAT has 14.5 million fCO₂ values from 3646 data sets covering the years 1957 to 2014. This latest version has an additional 4.4 million fCO₂ values relative to version 2 and extends the record from 2011 to 2014. Version 3 also significantly increases the data availability for 2005 to 2013. SOCAT has an average of approximately 1.2 million surface water fCO₂ values per year for the years 2006 to 2012. Quality and documentation of the data has improved. A new feature is the data set quality control (QC) flag of E for data from alternative sensors and platforms. The accuracy of surface water fCO₂ has been defined for all data set QC flags. Automated range checking has been carried out for all data sets during their upload into SOCAT. The upgrade of the interactive Data Set Viewer (previously known as the Cruise Data Viewer) allows better interrogation of the SOCAT data collection and rapid creation of high-quality figures for scientific presentations. Automated data upload has been launched for version 4 and will enable more frequent SOCAT releases in the future. High-profile scientific applications of SOCAT include quantification of the ocean sink for atmospheric carbon dioxide and its long-term variation, detection of ocean acidification, as well as evaluation of coupled-climate and ocean-only biogeochemical models. Users of SOCAT data products are urged to acknowledge the contribution of data providers, as stated in the SOCAT Fair Data Use Statement. This ESSD (Earth System Science Data) ‘Living Data’ publication documents the methods and data sets used for the assembly of this new version of the SOCAT data collection and compares these with those used for earlier versions of the data collection (Pfeil et al., 2013; Sabine et al., 2013; Bakker et al., 2014).*

Data coverage

20 Repository-References:

Individual data set files and synthesis product: doi:10.1594/PANGAEA.849770

Gridded products: doi:10.3334/CDIAC/OTG.SOCAT_V3_GRID

Available at: <http://www.socat.info/>

Coverage: 79°S to 90°N; 180°W to 180°E

25 Location Name: Global Oceans and Coastal Seas

Date/Time Start: 21 October 1957

Date/Time End: 4 October 2014



1 Introduction

The oceans represent a vast reservoir for carbon, mainly in the form of dissolved inorganic carbon (DIC), made up of the species bicarbonate, carbonate and dissolved carbon dioxide (CO_2). This carbon reservoir is in contact with the much smaller reservoir of CO_2 in the atmosphere via air-sea gas exchange.

5 Emissions of CO_2 by human activity, such as fossil fuel burning, cement manufacturing and changes in land use, are rapidly increasing the atmospheric concentration of the long-lived greenhouse gas. The oceans are taking up about 26% of these global CO_2 emissions with ocean uptake estimated at $2.6 \pm 0.5 \text{ Pg C yr}^{-1}$ for the time period 2005 to 2014 (Le Quéré et al., 2015b). This ocean carbon sink slows down the rate of climate change caused by human activity. Ocean carbon uptake changes ocean carbonate chemistry, notably by reducing ocean pH and carbonate ion concentration, a process known as
10 ocean acidification and sometimes referred to as ‘the other CO_2 problem’ (Turley, 2005; Henderson, 2006; Doney et al., 2009a). These changes in ocean chemistry are expected to affect key physiological processes of marine organisms, such as calcification, growth, development and survival (Kroeker et al., 2013). Ocean acidification is likely to have far reaching impacts on marine organisms and marine biodiversity with the effects expected to first being felt in the polar oceans (Orr et al., 2005).

15 The annual change in marine carbonate chemistry resulting from net ocean carbon uptake is small in comparison to its natural variation. A mean annual increase of $1.5 \mu\text{atm}$ has been estimated in surface ocean $f\text{CO}_2$ (fugacity of CO_2) for the period from 1970 to 2007 (Takahashi et al., 2009), which is superimposed on large seasonal variation, here defined as the difference between winter and summer values, of, for example, $120 \mu\text{atm}$ in the seasonally ice covered Southern Ocean and $160 \mu\text{atm}$ in Georgia Basin (Jones et al., 2015a). The annual increase also occurs against a background of large spatial
20 variation, of e.g. $140 \mu\text{atm}$ in different regions of the Southern Ocean in spring (Bakker et al., 2008; Jones et al., 2015a). Similarly, seasonal variation of 0.04 in pH in the subtropical North Atlantic Ocean (González-Dávila et al., 2007) is 20 times the mean annual decrease in surface ocean pH at a rate of -0.002 year^{-1} (Feely et al., 2009; Lauvset et al., 2015).

Seasonal and spatial variation in surface water $f\text{CO}_2$ and pH tend to be larger in coastal waters than in the open ocean, as illustrated by an $f\text{CO}_2$ decrease of $250 \mu\text{atm}$ from winter to summer at a coastal site near Antarctica (Legge et al., 2015)
25 and spatial variation of up to $200 \mu\text{atm}$ within the North Sea (Thomas et al., 2004; Omar et al., 2010). Arctic coastal and shelf seas equally have large spatial ($>500 \mu\text{atm}$ within the region in summer), seasonal ($300 \mu\text{atm}$) and year-to-year variation ($100 \mu\text{atm}$) in surface water $f\text{CO}_2$ (Fransson et al., 2006, 2009). Surface water $f\text{CO}_2$ may range from less than 200 to $800 \mu\text{atm}$ (or even $1200 \mu\text{atm}$) over short time (days) and space scales (less than 10 nm) in the upwelling system of the US West Coast (Hales et al., 2005, 2012; Harris et al., 2013, supplemental figure.)

30 The annual changes in surface ocean $f\text{CO}_2$ and pH exhibit spatial and temporal variation. Basin-specific rates in the $f\text{CO}_2$ increase vary from 1.2 to $2.1 \mu\text{atm year}^{-1}$ for the years 1970 to 2007 (Takahashi et al., 2009) with higher rates of 2.3 to $3.3 \mu\text{atm year}^{-1}$ at different mooring sites in the equatorial Pacific Ocean for the more recent period of 1997 to 2011 (Sutton et al., 2014). The annual pH decreases at rates of $-0.0013 \text{ year}^{-1}$ in the South Pacific Ocean (for 1998 to 2012) to $-0.0026 \text{ year}^{-1}$



in the Irminger Sea (for 1982 to 2006) (Bates et al., 2014), while annual pH changes vary from -0.0018 to -0.0026 year⁻¹ for moorings in the Equatorial Pacific Ocean for 1997 to 2011 (Sutton et al., 2014). Here it is worth noting that such rates of change vary with the start date and period used for the calculation as a result of interannual to decadal variability (McKinley et al., 2011).

5 Modelling has long been a primary tool for quantification of the ocean carbon sink (e.g. Le Quéré et al., 2014) and ocean acidification (Orr et al., 2005). The availability of large surface ocean CO₂ data synthesis products, such as the Lamont Doherty Earth Observatory (LDEO) surface ocean pCO₂ (partial pressure of CO₂) data-base (Takahashi et al., 2009, 2015) and the Surface Ocean CO₂ Atlas (SOCAT) (Pfeil et al., 2013; Sabine et al., 2013; Bakker et al., 2014; this study), now enables data-based estimates of the ocean carbon sink, as well as direct model-to-data comparison for surface ocean fCO₂ and ocean
10 carbon sink estimates (Le Quéré et al., 2014, 2015a, 2015b; Séférian et al., 2014; Turi et al., 2014). A challenge for data-based estimates of the ocean carbon sink is the gap-filling required for times and locations without surface ocean fCO₂ data. Different techniques and assumptions are applied for doing this, however, the resulting estimates of the ocean carbon sink differ considerably between the methods, especially in data sparse regions, such as the South Pacific Ocean (Rödenbeck et al., 2015). Recent data-based studies highlight large year-to-year, decadal and longer-term variation in surface ocean fCO₂
15 with consequent variation in the global ocean CO₂ sink (Fay and McKinley, 2013; Fay et al., 2014; Landschützer et al., 2014, 2015; Rödenbeck et al., 2014, 2015). Several model-to-data comparison studies suggest that models underestimate the spatial and temporal variation in surface ocean fCO₂ and the ocean carbon sink (Séférian et al., 2014; Turi et al., 2014; Rödenbeck et al., 2015). Such results could only be achieved because of the huge progress that has been made in data collection efforts like SOCAT.

20 The Global Carbon Budget provides an annual estimate of the carbon sinks and sources for the atmosphere (Le Quéré et al., 2014, 2015a, 2015b). The land carbon sink is determined as a residual of the other terms in the budget, namely the atmospheric and ocean components. Thus, quantification of the ocean carbon sink is critical to resolving the Global Carbon Budget.

25 The above highlights the need for long-term sustained, accurate observations over the entire surface ocean and synthesis of the marine carbonate chemistry measurements for quantification of trends in the ocean carbon sink and ocean acidification. This has been eloquently expressed for in situ observations of the climate system by Carl Wunsch and colleagues (Wunsch et al., 2013):

30 ‘No substitute exists for adequate observations. [...] Models will evolve and improve, but, without data, will be untestable, and observations not taken today are lost forever. [...] Today’s climate models will likely prove of little interest in 100 years. But adequately sampled, carefully calibrated, quality controlled, and archived data for key elements of the climate system will be useful indefinitely.’

In 2007, the international marine carbon community decided to create a quality controlled, publicly available synthesis product of surface ocean CO₂ for the global oceans and coastal seas (IOCCP, 2007; Doney et al., 2009b). The Surface Ocean CO₂ Atlas provides regular updates of:



1. A synthesis product of surface ocean $f\text{CO}_2$ measurements,
2. A gridded product of surface ocean $f\text{CO}_2$ values (without interpolation to grid cells with no measurements).

Both data products cover the global oceans and coastal seas. The Surface Ocean CO_2 Atlas (<http://www.socat.info/>) provides a key synthesis data set of surface ocean $f\text{CO}_2$ for global and regional scientific studies of the ocean carbon sink and ocean acidification.

The SOCAT data collection only contains original surface water CO_2 data, as reported by the data originator, as input values. Thus, the SOCAT data collection does not contain CO_2 values processed by secondary data sources. The SOCAT data products only contain surface water $f\text{CO}_2$ values from $x\text{CO}_2$ (mole fraction) measurements (Pfeil et al., 2013). SOCAT does not include surface water $f\text{CO}_2$ calculated from the other seawater carbonate system parameters, such as pH, dissolved inorganic carbon or total alkalinity. Almost all $f\text{CO}_2$ values were determined on ships by determination of the CO_2 concentration in the headspace of an equilibrator with a continuous seawater flow (Pfeil et al., 2013; Bakker et al., 2014). Shipboard systems for equilibrators normally use gas chromatography or infrared detection to determine the CO_2 concentration in headspace air (Pierrot et al., 2009). SOCAT versions 2 and 3 also have data sets from fixed moorings and drifting buoys with measurements made by an equilibrator system with infrared detection and a membrane spectrophotometer. The SOCAT data collection includes a small number of historical, discrete surface water $f\text{CO}_2$ measurements.

Version 1 of SOCAT was made available in 2011 (Pfeil et al., 2013; Sabine et al., 2013), followed by the release of version 2 in 2013 (Bakker et al., 2014). *SOCAT version 3 was made public during the SOCAT and SOCOM (Surface Ocean $p\text{CO}_2$ Mapping Intercomparison) Event on 7 September 2015 (SOCAT and SOCOM, 2015). The event was part of the Surface Ocean Lower Atmosphere Study (SOLAS) Open Science Conference in Kiel, Germany.*

This contribution documents SOCAT version 3, while highlighting the key differences with respect to version 2 (Sect. 2). The SOCAT Fair Data Use Statement is presented in Section 3. This is followed by a description of data upload, quality control (Sect. 4) and the data products available for version 3 (Sect. 5). We also look forward towards ongoing developments affecting future SOCAT versions, notably automated data upload, inclusion of additional parameters and annual releases (Sect. 6). The article ends with an assessment of the impact and scientific applications of SOCAT to date (Sect. 7) and concluding remarks (Sect. 8). This publication will be updated regularly using the format of the ESSD (Earth System Science Data) ‘Living Data’ to document the SOCAT versions and significant changes in the data collection, data upload, quality control and data products. This is the first version of the SOCAT ‘Living Data’ and is closely associated with earlier ESSD publications describing SOCAT versions 1 (Pfeil et al., 2013; Sabine et al., 2013) and 2 (Bakker et al., 2014).

2 Characteristics of SOCAT version 3 and key differences with version 2

Version 3 of the Surface Ocean CO_2 Atlas includes 14.5 million surface water $f\text{CO}_2$ values over the time period 1957 to 2014 for the oceans and coastal seas around the world (Figs. 1 and 2; Table 1). The $f\text{CO}_2$ values are from 3646 data sets, collected on ships (3504 cruises), moorings (123) and drifters (19). The 3646 data sets include 3640 data sets with a WOCE flag of 2



(good), available in all data products, as well as 6 data sets with a WOCE of 3 (questionable), only available in some data products, if selected. Version 3 is an update of version 2 with an additional 4.4 million $f\text{CO}_2$ values from 986 data sets. Version 3 takes the start of the data record backwards from 1968 to 1957 by adding 4 historic cruises. It also extends the data collection forward by adding 1.8 million $f\text{CO}_2$ values for 2012 and 2013, as well as a small number of values from 2014 (Fig. 2). Version 3 also increases the number of $f\text{CO}_2$ values for many years between 1989 and 2011. It adds 50% more $f\text{CO}_2$ values for 2008 to 2010, while doubling the available data for 2011. The year 2006 has the largest number of $f\text{CO}_2$ values, closely followed by 2009 and 2011.

New in version 3 is an accuracy criterion for all surface ocean $f\text{CO}_2$ values, described by data set quality control (QC) flags of A to E, for accuracies of 2 (A, B), 5 (C, D) and 10 μatm (E) (Table 2) (Sect 4.4) (Wanninkhof et al., 2013b; Olsen et al., 2015). Flag A now also requires a high-quality crossover with another data set. The introduction of a lower accuracy, data set quality control flag of E (accuracy of $f\text{CO}_2$ values better than 10 μatm) enables the inclusion of well-calibrated $f\text{CO}_2$ measurements made by alternative sensors and on alternative platforms (Wanninkhof et al., 2013b; Olsen et al., 2015). Version 3 has significantly more data sets from fixed moorings (123 data sets) and drifting buoys (19) than version 2 (28 and 3 data sets, respectively). These measurements were made by an equilibrator system with infrared detection (e.g. Johengen, 2010; Sutton et al., 2014) and a membrane spectrophotometer (e.g. Boutin and Merlivat, 2009; Merlivat et al., 2015).

The quality and documentation of the data has improved relative to version 2, with 49% of the data sets receiving a quality control flag of A or B, 23% a flag of C and 27% a flag of D, in comparison to 41%, 22% and 37%, respectively, in version 2. The reduction in the relative amount of data sets with a flag of D, which implies incomplete metadata, highlights an improvement in the documentation accompanying data sets. A total of 41 data sets (1%) received a flag of E, most of these are sensor data, but they also include a small number of valuable historic data sets with an accuracy deemed better than 10 μatm .

Version 3 represents a major step towards the automation of the SOCAT data and metadata upload and quality control in future versions. A new interface, the SOCAT Upload Dashboard, hosts data and metadata upload, (re-)calculation of $f\text{CO}_2$, automated data checks, data visualisation and submission to the quality control system into a single application (Table 1). A prototype of the SOCAT Upload Dashboard was used for data upload for version 3 (Sect. 4.1) and (re-)calculation of $f\text{CO}_2$ (Sect. 4.2). All data sets were run across a newly developed, automated data checker for identification of values that were out of range (Sect. 4.3). As a result, issues identified during data upload were already corrected prior to entry on the quality control system. The search capabilities and graphical interface of the quality control system and the associated Data Set Viewer (previously known as the Cruise Data Viewer) were upgraded (Sects. 4.4 and 5.4). Version 4 will see enhanced implementation of SOCAT automation by enabling data providers to upload their data using the SOCAT Upload Dashboard and submission onto the SOCAT QC Editor (Sect. 6.1).

The publicly accessible, user-friendly and interactive Data Set Viewer now allows selection of $f\text{CO}_2$ values by data set, year, month, region, data provider, vessel or platform name, country of the vessel's or platform's flag, data set quality control flag, WOCE flag and SOCAT version, as well as to set limits on data ranges. The graphical tools of the Data Set Viewer



(access via <http://www.socat.info/>) for SOCAT version 3 have been extended (Figs. 1, 3 and 4). Users can now set fixed colour scales and create high-quality, publishable images.

A small error was detected in the gridded data products of SOCAT versions 1 and 2 (Sect. 5.5). In short, the data set-weighted $f\text{CO}_2$ values (formerly known as cruise-weighted $f\text{CO}_2$ values) in these products were found to have missing values for a small number of grid cells, as a result of an inconsistency between the algorithms used for computing the weighted and unweighted gridded products. This was both in time and in position. This error was corrected in the gridded data products for version 3. Note that this error remains present in the gridded products for versions 1 and 2.

In summary, SOCAT version 3 is a significant update of version 2. It provides a 58-year record (1957-2014) of 14.5 million surface ocean $f\text{CO}_2$ values for the global oceans and coastal seas. It has higher quality data with better documentation than version 2. Addition of a flag of E has enabled inclusion of well-calibrated $f\text{CO}_2$ values from alternative sensors and platforms. All surface ocean $f\text{CO}_2$ values now have an accuracy estimate, embedded in the data set QC flag. Automated quality control checks during version 3 data upload have identified outliers. The graphical interface of the Data Set Viewer has been vastly improved. These characteristics of version 3 are described in more detail in Sections 4 to 6.

3 Fair Data Use Statement for SOCAT version 3

The Surface Ocean CO_2 Atlas provides access to a vast amount of surface ocean CO_2 data from the global oceans and coastal seas, painstakingly collected by marine carbon scientists around the world over 58 years. These data sets represent an important scientific output by these scientists. Individual researchers and the marine carbon community make these data public in SOCAT, such that they are available for scientific research and for informing policy (Sect. 7 and 8). Nonetheless, it is important that the data providers receive credit for the data that they collected. This will provide data providers with vital evidence of how their data are being used, enabling successful funding applications for future data collection.

Furthermore, the assembly, quality control and archiving of SOCAT data products involve many data managers and scientists (Tables 3 and 4). Planning meetings and community events have proved effective in informing SOCAT contributors and users, in discussing SOCAT progress and in setting SOCAT strategy (Table 5).

The SOCAT Fair Data Use Statement therefore contains an urgent request to generously acknowledge the contribution by SOCAT data contributors and investigators. Ideally users will invite large data providers to contribute to regional studies and, if they do, to co-author relevant papers. Citation of relevant scientific articles by data providers is good scientific practice. The following Fair Data Use Statement applies to SOCAT data products (SOCAT, 2016): ‘The synthesis and gridded SOCAT products are a result of scientific effort by data providers, data managers and quality controllers. It is important that users of the SOCAT products fairly acknowledge this effort. This will help generate funding for continuation of observational products and promote further sharing of data.’

We expect that users of SOCAT data products:



1. **Generously acknowledge the contribution of SOCAT data providers and investigators** in the form of invitation to co-authorship, reference to relevant scientific articles by data providers or by naming the data providers in the acknowledgements. Specifically, in regional studies invite large data providers, who frequently possess valuable expert knowledge on data and region, to collaborate at an early stage, which may lead to an invitation of co-authorship. We
5 recognize that co-authorship is only justified in case of a significant scientific contribution to a publication and that provision of data on its own does not warrant co-authorship.

2. Cite SOCAT and its data products as:

Version 3: *This study*;

Version 2: *Bakker et al. (2014)*;

10 Version 1 (synthesis data products): *Pfeil et al. (2013)*;

Version 1 (gridded data product): *Sabine et al. (2013) and Pfeil et al. (2013)*.

3. Include in the acknowledgements: ‘The Surface Ocean CO₂ Atlas (SOCAT) is an international effort, endorsed by the International Ocean Carbon Coordination Project (IOCCP), the Surface Ocean Lower Atmosphere Study (SOLAS), and the Integrated Marine Biogeochemistry and Ecosystem Research program (IMBER), to deliver a uniformly quality-
15 controlled surface ocean CO₂ database. The many researchers and funding agencies responsible for the collection of data and quality control are thanked for their contributions to SOCAT.’

4. Report problems to submit@socat.info.

5. Inform submit@socat.info of publications in which SOCAT is used.’

20 The Fair Data Use Statement (SOCAT, 2016) replaces the earlier ‘SOCAT Data Policy’ (SOCAT, 2013a; Bakker et al., 2014). The text has been phrased more strongly and examples of the application of the Fair Data Use Statement have been added. The Fair Data Use Statement is available in full on the SOCAT web pages (e.g. http://www.socat.info/SOCAT_fair_data_use_statement.htm). The revision follows concerns raised by SOCAT data providers and discussions among SOCAT scientists at two recent community events (SOCAT, 2014a; SOCAT and SOCOM, 2015).

4 Data assembly and quality control in version 3

25 4.1 Data retrieval and data upload on the SOCAT Upload Dashboard

In version 3, new and updated data sets were obtained from the Carbon Dioxide Information Analysis Centre (CDIAC), PANGAEA® and public websites. In addition, many data sets were directly submitted to SOCAT. As well as 887 new data sets, version 3 also contains 1258 updated version of data sets previously submitted to versions 1 and 2, with revised metadata or data. Some of these were updates of data sets previously suspended from SOCAT (e.g. Table 10 in Bakker et al., 2014).

30 As in previous versions, all new and updated data sets were put in a uniform format (Pfeil et al., 2013). Similar to version 2, an expocode was assigned to all data sets, including moorings and drifters (Bakker et al., 2014). In general, an expocode consists of 12 characters, describing the country, the vessel or platform and the data set start day (Swift, 2008). The expocode



320620090306, for example, indicates a data set collected on the US (32) ship R. V. Nathaniel B. Palmer (06) with the first day of the cruise on 6 March 2009. There are a few exceptions to this. If two American mooring data sets (they always start with 3164) have the same start date, they will end with '-1' and '-2', corresponding to an expocode of 14 characters.

In version 3, the SOCAT data managers used the new SOCAT Upload Dashboard for upload of data and metadata (Table 1). All data sets previously included in versions 1 and 2 were also uploaded, automatically screened for obvious outliers and added to version 3 via the SOCAT Upload Dashboard (Table 1). This new capability is an important step in the ongoing SOCAT automation effort and integrates data and metadata upload, (re-)calculation of $f\text{CO}_2$, automated data checks, data visualisation and data submission into a single application which is tightly coupled to the SOCAT QC editor. Once fully operational in version 4, the Upload Dashboard will allow data providers to upload, verify and submit their data for SOCAT quality control.

Not all data sets had time stamps which included seconds. In such cases, multiple occurrences of a time stamp were often present. Artificial seconds were added to data sets with 50 or more duplicate time stamps. For these data sets, evenly distributed artificial seconds were added for each equal time stamp. However, because of time constraints it was not possible to add artificial seconds to all data sets. Therefore, if there were less than 50 duplicate times in a data set, a WOCE flag of 4 was generated for the corresponding $f\text{CO}_2$ value during the automated data checks (Sect. 4.3).

4.2 (Re-)calculation of $f\text{CO}_2$

Data providers reported CO_2 values as $x\text{CO}_2$, $p\text{CO}_2$ and/or $f\text{CO}_2$, at the equilibration temperature (T_{equ}) and/or the sea surface temperature (SST or intake temperature). Surface water $f\text{CO}_2$ values at sea surface temperature were recalculated from the reported CO_2 values using a strict calculation protocol with the following procedure (quoting Pfeil et al., 2013):

1. When possible, (re-)calculate $f\text{CO}_2$;
2. The preferred starting point for the calculations is $x\text{CO}_2$, next $p\text{CO}_2$, and finally $f\text{CO}_2$;
3. Minimize the use of external data required to complete the calculations.'

In total, 14 algorithms were used for calculating these 'recommended' $f\text{CO}_2$ ($f\text{CO}_{2\text{rec}}$) values (Table 6). The term 'recommended' $f\text{CO}_2$ values is used here to distinguish the $f\text{CO}_2$ values (re-)calculated by the SOCAT algorithms from the $x\text{CO}_2$, $p\text{CO}_2$ and/or $f\text{CO}_2$ values reported by the data providers. The algorithm used for a given data set is listed in the data products (Sect. 5). Equations recommended by Dickson et al. (2007) were applied for the conversion of the dry CO_2 mole fraction to $p\text{CO}_2$, for the calculation of the water vapour pressure and for the correction of $p\text{CO}_2$ to $f\text{CO}_2$ (Pfeil et al., 2013). The temperature correction suggested by Takahashi et al. (1993) was used to correct for temperature change between the seawater intake and the equilibrator. Atmospheric pressure from reanalysis and climatological values of salinity were used in the calculation, if *in situ* values had not been reported (Table 6). The 2014 version of the atmospheric pressure data product was used (NCEP, 2014), which is an update of the 2012 data product used in the previous SOCAT version (NCEP, 2012).

An important change relative to earlier versions is that the (re-)calculation in version 3 took place using Ferret scripts on the new SOCAT Upload Dashboard after data upload (Sect. 4.1), rather than in Matlab before the bulk upload of the data



(Table 1). The Matlab code used for the (re-)calculation in versions 1 and 2 was transferred to Ferret scripts on the Upload Dashboard for version 3. The new Ferret scripts were checked by comparing $f\text{CO}_2\text{rec}$ values in version 2 calculated using Matlab and new values calculated using Ferret. Almost all new values were within $0.01 \mu\text{atm}$ of the value calculated in Matlab, if not identical to it. Significant changes (smaller than $5 \mu\text{atm}$) for less than 200 data points were attributed to changes in atmospheric pressure from reanalysis (Table 1).

4.3 Automated data checks

A newly developed, automated data checker performed checks on parameters directly influencing the position, time or calculation of $f\text{CO}_2\text{rec}$ values (Tables 1 and 7). A WOCE flag of 4 (meaning a bad data point) was assigned to all $f\text{CO}_2\text{rec}$ values with an incorrect position or time stamp or otherwise identified as inaccurate. These automated checks were carried out on all data in version 3 after (re-)calculation of $f\text{CO}_2\text{rec}$ and before submission to the quality control system.

Unintentionally WOCE flags of 4 were also assigned for values which were out of range in parameters which do not directly affect $f\text{CO}_2\text{rec}$ values, such as wind speed and ship direction (Table 7). This resulted in a WOCE flag of 4 being given to some good quality $f\text{CO}_2\text{rec}$ values in newly added and updated data sets in version 3. The criteria for the automated checks will be reconsidered for version 4.

Automated data checks were also performed for data sets previously included in versions 1 and 2 (and not updated in version 3). For these data sets all WOCE flags of 4 assigned by the automated data checker, other than for duplicate time stamps, were removed, to preserve the data sets as reported for version 2.

4.4 Secondary quality control

Secondary quality control is a key part of the creation of a high-quality data synthesis product. During secondary quality control, scientists, also known as quality controllers, assess the quality of each new and updated data set by following a checklist of specific criteria, while also examining the documentation of the data, known as metadata, for completeness. The quality controllers assign a data set quality control flag to each data set, based on their findings (Table 2).

The SOCAT quality control system has been upgraded (Table 1), as part of the ongoing SOCAT automation. In particular, the ease of use, search options and visualisation tools have been improved. Other modifications are that the quality control criteria used for setting the data set quality control flag now must be specified (by a tick box system) and that a comment needs to be entered when assigning a WOCE flag (Table 1). Text relating to the tick boxes and the comments accompanying WOCE flags are incorporated in the quality control comments.

The definitions of the data set quality control flags in version 3 have been revised relative to versions 1 and 2 (Tables 1 and 2) (Wanninkhof et al., 2013b; Olsen et al., 2015). These revised QC criteria were applied to all new and updated data sets in version 3, but not retrospectively to data sets included in earlier versions, unless data providers had updated these. Version 3 has data set quality control flags of A to E and WOCE flags of 2, 3 and 4 for individual $f\text{CO}_2\text{rec}$ values. For a data set to obtain a data set quality control flag, it needs to meet all the criteria of that specific data set flag (Table 2).



All data set flags now have an accuracy requirement for the fCO_{2rec} values. Previously, flags of C and D did not have an accuracy requirement (Pfeil et al., 2013; Bakker et al., 2014). In version 3, requirements are an accuracy of better than $2 \mu atm$ for flags of A and B, of better than $5 \mu atm$ for flags of C and D and of better than $10 \mu atm$ for a flag of E (Table 2). The accuracy requirement takes precedent over the criteria that follow (Wanninkhof et al., 2013b; Olsen et al., 2015), implying that, if the accuracy requirement is not met, a data set is given a data set flag with a lower accuracy requirement, appropriate to the accuracy of the data set.

Seven approved methods or SOP (standard operating procedure) criteria need to be met for a data set quality control flag of A and B (after Pfeil et al., 2013):

1. The data are based on xCO_2 analysis, not fCO_2 calculated from the other carbon parameters pH, total alkalinity and dissolved inorganic carbon;
2. Continuous CO_2 measurements have been made, not discrete CO_2 measurements;
3. The CO_2 detection is based on an equilibrator system and is performed by infrared analysis or gas chromatography;
4. The calibration has included at least two non-zero gas standards, traceable to World Meteorological Organisation (WMO) standards;
5. The equilibrator temperature has been measured to within $0.05^\circ C$ accuracy;
6. The intake seawater temperature has been measured to within $0.05^\circ C$ accuracy;
7. The equilibrator pressure has been measured to within 2.0 hPa accuracy.

The requirement regarding the accuracy of the equilibrator pressure has been relaxed to an accuracy of 2.0 hPa in version 3, replacing the earlier requirement of 0.5 hPa, as an accuracy of 2.0 hPa in pressure is sufficient for achieving an accuracy of $2.0 \mu atm$ in fCO_2 (Wanninkhof et al., 2013b; Olsen et al., 2015). The six other criteria are the same in all SOCAT versions.

In version 3, a high-quality cross-over has become a pre-requisite for a data set flag of A, replacing the earlier requirement of 'an acceptable comparison (or cross-over) with other data' (Wanninkhof et al., 2013b; Olsen et al., 2015). As in previous versions, a cross-over is defined by an equivalent distance of less than 80 km between two data sets (Pfeil et al., 2013). This criterion combines distance and time as $([\Delta x^2 + (\Delta t * 30)^2]^{0.5}) \leq 80$ with distance x in kilometres and time t in days. One day (or 24 hours) of separation in time is equivalent (heuristically) to 30 km of separation in space. According to this definition, the maximum time separation (at a spatial distance of 0 km) is about 64 hours for a cross-over to occur. The new definition of a high-quality cross-over between two data sets requires that differences in sea surface temperature and fCO_{2rec} between the data sets do not exceed $0.3^\circ C$ and $5 \mu atm$, respectively. 'Inconclusive' cross-overs, where differences in temperature or fCO_{2rec} exceed these values, do not qualify for a data set flag of A in version 3.

It is worth noting that meaningful high-quality cross-overs are rarely found in coastal waters, near sea ice and in regions of freshwater influence (ROFIs), as a result of high spatial variation in sea surface temperature and fCO_{2rec} , not for lack of measurement quality. Even if a small number of sea surface temperature and fCO_{2rec} values are within $0.3^\circ C$ and $5 \mu atm$, this tends to be a coincidence rather than a meaningful correspondence between data sets. This can be illustrated for the US



research ships Nathaniel B. Palmer and the Lawrence M. Gould, which have frequent high-quality cross-overs in the open Southern Ocean, but few high-quality cross-overs near Palmer station, where they both make port calls.

In version 3, a data set with a flag of C 'did or did not follow approved methods or SOP criteria' (Wanninkhof et al., 2013b; Olsen et al., 2015). This is an amendment from the earlier requirement that the data set 'did not follow approved methods or SOP criteria' (Pfeil et al., 2013; Bakker et al., 2014). The new flag of E enables inclusion of $f\text{CO}_2$ values from well-calibrated alternative sensors and platforms (Wanninkhof et al., 2013b; Olsen et al., 2015). A flag of E requires complete metadata and a demonstrable accuracy for $f\text{CO}_{2\text{rec}}$ of better than $10 \mu\text{atm}$ by in situ calibration of the sensor. The WOCE flags for individual $f\text{CO}_{2\text{rec}}$ values are defined as 2 (good), 3 (questionable) and 4 (bad) in versions 1, 2 and 3 (Pfeil et al., 2013). New is the requirement to add a comment when assigning WOCE flags of 3 and 4 (Table 1).

As in version 2, five additional guidelines were considered, but not systematically applied, for open ocean $f\text{CO}_{2\text{rec}}$ values, away from sea ice. The guidelines were used for assigning data set quality control flags and WOCE flags (after Pfeil et al., 2013 and Bakker et al., 2014):

1. Warming between the seawater intake and the equilibrator should be less than 3°C ;
2. Warming rate should be less than 1°C h^{-1} , unless a sharp temperature front is apparent;
3. Warming outliers should be less than 0.3°C , compared to background data;
4. Cooling between the seawater intake and the equilibrator is unlikely in high-latitude oceans for an indoor measurement system;
5. Zero or constant temperature difference between the equilibrator and seawater intake usually indicates the absence of SST values.

As for SOCAT version 2, quality controllers were organised in eight regions, each with a group lead (Table 4). The eight regions included the coastal and marginal seas, the Arctic Ocean, the North and Tropical Atlantic, the North and Tropical Pacific, the Indian Ocean and the Southern Ocean. The quality controllers gave data sets a quality control flag for each region they crossed. As a final step, the data set quality control flags for the different regions had to be reconciled.

5 Data products in version 3

5.1 Overview of data products

In essence, the data products and data platforms are the same as for earlier SOCAT versions with some modifications (Table 8). Improvements include a major upgrade of the search and visualisation capabilities of the Data Set Viewer (previously known as the Cruise Data Viewer) and uniform contents for the files downloadable from the Data Set Viewer (Tables 1 and 9). Access to the data products is via the SOCAT website (<http://www.socat.info/>) and the web addresses for the individual data platforms (Table 8).

Quality controlled recommended surface ocean $f\text{CO}_2$ measurements in a uniform format are available in individual data set files, in regional and global synthesis files and in gridded form (Table 8). These three data products can be accessed via the



user-friendly, interactive online Data Set Viewer and Gridded Data Viewer, by downloading data files or in Ocean Data View (Schlitzer, 2015). Similar to earlier versions, data sets with a quality control flag of A to D and recommended $f\text{CO}_2$ values with a WOCE flag of 2 (good) are included in the synthesis files and gridded products. *Data sets with a flag of E are available in a separate synthesis file. Data set flags of A to E and a WOCE flag of 2 for $f\text{CO}_2$ values is the default setting for the Data Set Viewer.* Quality control comments can be accessed via the Data Set Viewer (Table 8). While the SOCAT data products include seawater temperature and salinity, as these are required for (re-)calculation of $f\text{CO}_2$, these two parameters have not been quality controlled to the high standards required by the physical oceanographic community (SOCAT, 2014a).

As in earlier versions, each individual data set has a digital object identifier (doi), which provides a direct link to the metadata, including the name and affiliation of the data provider. This doi for the data set is available for each recommended surface ocean $f\text{CO}_2$ value in the synthesis files. This enables users to easily identify the data provider and to gain access to the original data set and to detailed information on the data set, including any relevant peer-reviewed journal articles that we are aware of. *The Data Set Viewer now enables to search the data collection by data provider. Data providers are also prominently displayed in the Table of Datasets (access via the Data Set Viewer) (Table 8).* A more detailed description of the data products follows.

5.2. Individual data set files

Individual data set files are available for all data sets with flags of A, B, C, D and E. Each individual data set has a doi. The files contain all original CO_2 measurements and recommended $f\text{CO}_2$ values with a WOCE flag of 2, 3 and 4 (Table 8), as set by the data originator, by the automated range checker or during the secondary quality control. The files also contain other parameters, such as atmospheric pressure from re-analysis, climatological salinity and the atmospheric CO_2 mole fraction. Metadata reported by the data provider accompany the files and links to the original data sets are provided. The files are available in text format at PANGAEA® (<https://doi.org/10.1594/PANGAEA.849770>).

5.3. Global synthesis product

The global and regional synthesis files contain recommended $f\text{CO}_2$ values with a WOCE flag of 2 (good) for data sets with flags of A, B, C and D (Table 8). A separate synthesis file is available for data sets with a flag of E. Each line of the global and regional synthesis files contains the doi for the corresponding individual data set, as archived at PANGAEA®, thus enabling retrieval of metadata, name of the data provider and the original CO_2 values reported by the data provider (Table 9) (Sect. 5.2). Global and regional files are available as compressed zip text files via CDIAC (<http://cdiac.ornl.gov/ftp/oceans/SOCATv3/>). Matlab code is available for reading these text files. Regional files for the SOCAT regions (Table 4) only contain data for a specific region with no overlap, so that many data sets on moving ships are split between several regional files. The global synthesis product for data sets with flags of A to D is also available in Ocean Data View format (https://odv.awi.de/en/data/ocean/socat_fCO2_data) (Schlitzer, 2015).



5.4 Subsetting the global synthesis product

The interactive Data Set Viewer (http://ferret.pmel.noaa.gov/SOCAT_Data_Viewer/) has powerful search capabilities and an attractive graphical interface, following the upgrade for version 3 (Tables 1 and 8). The SOCAT Data Viewer now hosts the Data Set Viewer and the Gridded Data Viewer on a single software platform. The move of the Data Set Viewer onto this platform in version 3 streamlines access to the SOCAT synthesis and gridded products via a Live Access Server (LAS). The move and upgrade of the Data Set Viewer accompany that of the closely associated SOCAT quality control system (Sects. 2 and 4.4).

The Data Set Viewer enables subsetting of the global SOCAT data collection. *The default setting is for data sets with flags of A to E and ‘valid’ fCO₂ values with a WOCE flag of 2 for years 1957 to 2014, corresponding to 3640 data sets for version 3. Recommended fCO₂ values with flags of 3 and 4 can also be selected. In the Data Set Viewer, the user can select data sets by, for example, year, month, region, platform/vessel, ‘valid’ values, data provider, data set flag, WOCE flag and SOCAT version. It is also possible to define limits for the values shown. Maps of surface ocean fCO₂ demonstrate the data distribution, as well as temporal and spatial variation in surface ocean fCO₂ for the selected data sets (Figs. 1, 3 and 4). High-quality figures can be rapidly created for scientific presentations to fellow scientists, funding agencies and policy makers. Property-property plots, available via the Correlation Viewer, can be used for further investigation of the data sets.*

The data shown on the Data Set Viewer have been subsampled for system efficiency, such that only part of the data are shown. *Visual display of these data sets on maps in the Data Set Viewer is subject to further improvement, as the interpolation of sparse data ignores topographic features. As a result cruise tracks occasionally appear to cross land. This issue does not affect the data sets themselves. The Table of Datasets (previously known as the Table of Cruises) can be accessed from the Data Set Viewer. It provides access to the original CO₂ measurements, fCO₂ values with a WOCE flag of 2, 3 and 4, metadata, comments entered during quality control and thumbnail plots (Table 8) (Sect. 4.4). Thumbnail plots consist of a series of property-property plots for key parameters in an individual data set and are useful for obtaining a quick overview of a data set. Both the Data Set Viewer and the Table of Datasets allow download of data sets in NetCDF and text format (Tables 8 and 9). All downloadable files now contain the same parameters (Table 9).*

The performance speed of the Data Set Viewer may be slower if the full SOCAT data collection is accessed. Subsetting the data collection by decade or region considerably improves the system speed of the Data Set Viewer. *Updates of web browsers occasionally result in less than perfect web access to the Data Set Viewer. In such cases, another web browser may provide better access. The web manager (socat.support@noaa.gov) may have useful advice.*

5.5 Gridded products

The protocol for the creation of gridded fCO₂ products in version 3 follows that for versions 1 and 2, as described by Sabine et al. (2013). The gridded products have a 1° latitude by 1° longitude resolution with a higher resolution of 0.25° latitude by 0.25° longitude for coastal seas. Recommended surface ocean fCO₂ values from 1970 to 2014 with a WOCE flag of 2 from



data sets with flags of A to D have been used for the gridded products. The gridded products have no interpolation, i.e. there is no gap-filling and grid cells without $f\text{CO}_2$ values are empty. No correction is made for the long-term increase in surface ocean $f\text{CO}_2$.

Gridded $f\text{CO}_2$ values are reported as unweighted means and as data set-weighted means (Sabine et al., 2013). In an unweighted mean, all $f\text{CO}_2$ values in a grid cell have equal weight for calculating the mean. In a data set-weighted mean, averages of the $f\text{CO}_2$ values are calculated per data set for each grid cell, before calculating averages of these data set means. *In version 3, a small error was corrected in the procedure for creating the gridded data products. This resulted in a small reduction in the number of grid cells with data in the data set-weighted product for versions 1 and 2. This problem was corrected in gridded files in version 3 with the revised gridded data set made public on 2 November 2015.*

Gridded products are available per decade, per year and monthly per year (Table 10). *A monthly climatological $f\text{CO}_2$ product has not been made available for version 3, out of concern, that such a product without a correction for the long-term increase in $f\text{CO}_2$ could be misinterpreted.* Gridded $f\text{CO}_2$ values may have temporal bias, for example, if only summer time $f\text{CO}_2$ values are available for a grid cell in the annual gridded product. Several auxiliary variables are reported per grid cell, for example, the number of data sets and observations and the standard deviation in the unweighted and weighted $f\text{CO}_2$ mean values (Table 10).

Gridded products are available in NetCDF format at CDIAC (http://cdiac.ornl.gov/ftp/oceans/SOCATv3/SOCATv3_Gridded_Dat/) (Table 8). Matlab code is available for reading the files. The Gridded Data Viewer (<http://www.socat.info/>, Select Gridded Data Viewer) provides easy access to the gridded data products, as well as comparison to gridded products from earlier versions. Figures 5 and 6 have been made with the gridded data product.

6 Future developments

6.1 Direct data upload and annual SOCAT releases

The SOCAT automation system was formally launched on 7 September 2015 (SOCAT and SOCOM, 2015). Data providers can now directly upload, check and submit their data on the SOCAT quality control system for future SOCAT versions. The SOCAT automation was first discussed at the 2011 Data2Flux Workshop in Paris (SOCAT, 2011). The automation system was designed at the 2012 Automation Planning Meeting (SOCAT, 2012a) and approved shortly afterwards by global and regional group leads (SOCAT, 2012b) (Table 5). The automation system has been implemented in the background, with all the work for the bi-annual SOCAT releases of versions 2 and 3 taking place in the foreground (Bakker et al., 2014, this study). This considerable achievement has been made possible by the hard work and planning of the NOAA-PMEL and University of Washington Live Access Server team and other members of the SOCAT automation team (Table 3).

The new automation system allows data providers to upload their data, to check their data with the automated data checker and to visualize their data. Finally, if the data provider deems the data of good quality, he or she can submit them to



the SOCAT quality control system. The automation system will enable annual SOCAT releases. The time-table for future SOCAT versions envisages that data upload will end in late January of each year and quality control in late March for a release in summer later that year. With the new system it is now possible to upload and submit data to SOCAT, while quality control of previously submitted data sets is in progress. Thus, both data upload and quality control can now be carried out in parallel. Data upload and quality control for the next SOCAT version will start as soon as they have finished for the preceding version. Thus, the automation system will enable rolling, continuous data upload and quality control, as well as annual SOCAT releases. The system for automated data upload is under continuous improvement. Metadata templates and upload will be integrated into the SOCAT data upload system. Other planned improvements include searchable information for funding agency and entry of preliminary data set flags by the data provider. A number of additional features are being considered for future SOCAT versions, some of which may be implemented as early as version 4. These are discussed below.

6.2 Atmospheric CO₂ values

Data providers can now submit measurements of atmospheric CO₂ mole fraction, made in parallel to surface water fCO₂. A separate WOCE flag will be created for measurements of the atmospheric CO₂ mole fraction in future SOCAT versions. Once quality control has been carried out on the atmospheric CO₂ measurements, such values will be included in the SOCAT data products.

In future, atmospheric fCO₂ will be calculated from atmospheric xCO₂ values, both from the measurements and from GLOBALVIEW-CO₂ (2014) values. New graphics will enable comparison of surface ocean fCO₂ values to atmospheric fCO₂ values. The graphs will become an important quality control tool. Future data products will contain atmospheric fCO₂ values calculated from atmospheric measurements and from GLOBALVIEW-CO₂, in addition to the atmospheric mole fractions from GLOBALVIEW-CO₂ already part of the SOCAT data products (Table 9).

6.3 Additional surface water parameters

In 2014, SOCAT scientists decided to allow inclusion of additional surface water parameters accompanying surface water fCO₂ values in SOCAT data output files (SOCAT, 2014a). Such additional parameters might include dissolved inorganic carbon, total alkalinity, pH, nutrients, methane (CH₄) and nitrous oxide (N₂O) concentrations. SOCAT scientists will not carry out quality control on these additional parameters, but would welcome collaboration with other communities taking responsibility for this. These additional parameters will be made public in parallel to the official SOCAT releases. The extra parameters will be posted in separate data files, to emphasize that they have not been quality controlled. A SOCAT and MEMENTO (Marine Methane and Nitrous Oxide; Bange et al., 2009) working group is considering the way forward for surface water CH₄ and N₂O measurements (SOCAT and SOCOM, 2015).



7 Impact and scientific highlights of SOCAT

7.1 A multi-decade record of surface ocean fCO₂ values

SOCAT provides a record of the history of surface ocean CO₂ research (Fig. 3). Initial, exploratory surface water CO₂ measurements in the late 1950s, 1960s and 1970s, were followed by more frequent CO₂ data collection on research ships in the 1980s and large (inter-)national research programs, such as the World Ocean Circulation Experiment (WOCE), the Joint Global Ocean Flux Study (JGOFS) and the Tropical Atmosphere Ocean (TAO) network in the 1990s. The operation of CO₂ instruments on Carbon Voluntary Observing Ships (Carbon VOS), also referred to as Ships Of Opportunity Programme (SOOP), strongly increased the number of available fCO₂ values from the 1990s onwards. Data availability in the SOCAT collection has increased fourfold from 0.2 to 0.4 million fCO₂ values per year for the years 1995 to 2000 to 1.0 to 1.2 million values per year for 2005 to 2012. Nevertheless, large gaps are notable in the data collection since the year 2000, e.g. in the Indian Ocean, the South Pacific Ocean, the Mediterranean Sea, the East China Sea, the Malay Archipelago and the Sea of Okhotsk. Elsewhere, in the Arctic Ocean, measurements are being reported for the first time.

The seasonal distribution of surface ocean fCO₂ values in the relatively data-rich decade from 2000 to 2009 is shown in Figure 4. This figure highlights the lack of winter data in the high-latitude oceans, as well as the opposing seasonal cycle of surface ocean fCO₂ in the subtropical and temperate oceans (Takahashi et al., 2002). The distribution of surface ocean fCO₂ values per decade clearly shows the long-term increase in surface ocean fCO₂ (Fig. 5), while suggesting that surface ocean fCO₂ has increased slower than the atmospheric CO₂ concentration since the 1990s. Figure 6 visualizes the data availability as the number of months in each 1° latitude by 1° longitude grid cell with fCO₂ values since 1970, both as unique months and as total months.

7.2 Impact of SOCAT

SOCAT and its data products are cited or named in influential international reports, in more than 100 peer-reviewed scientific publications, PhD and Master theses, book chapters and numerous other publications, as listed on the SOCAT website (<http://www.socat.info/publications.html>). Figure 7 shows the rapid increase in such publications, since the initiation of SOCAT in 2007 (IOCCP, 2007) and the first SOCAT release in 2011 (Pfeil et al., 2013; Sabine et al., 2013). The SOCAT data collection forms the basis of several data products (<http://www.socat.info/products.html>) and diverse scientific applications. These include a dozen mapping products of surface ocean pCO₂ and air-sea CO₂ fluxes for the global oceans (see overview in Rödenbeck et al., 2015). The SOCAT gridded product and one data product based on SOCAT (Landschützer et al., 2015) are integrated with the ESMValTool (Eyring et al., 2015) for routine evaluation of Earth System Models. For the same purpose, the SOCAT gridded product is currently being integrated into the Obs4MIPs (Observations for Model Intercomparison Projects) data repository (Ferraro et al., 2015). Citation of SOCAT in high-impact reports, scientific applications of SOCAT and scientific findings based on SOCAT are discussed below.



The importance of the SOCAT synthesis is highlighted by its citation in three categories of high-impact reports, notably reports on ocean observing systems, assessments of climate change and global carbon budgeting, including carbon observing strategies, and ocean acidification studies.

- Reports on ocean observing systems include publications from OceanObs'09 (Borges et al., 2010; Monteiro et al., 2010), the Framework for Ocean Observing (FOO, 2012), the Tropical Pacific Observing System 2020 (Mathis et al., 2014) and the 2nd International Indian Ocean Expedition (Hood et al., 2015).
- Assessments of climate change and global carbon budgeting citing SOCAT are the 2013 IPCC (Intergovernmental Panel on Climate Change) report (Ciais et al., 2013) and the State of the Climate in 2014 (Blunden and Arndt, 2015). Four reports describing a global carbon or climate observing system highlight SOCAT, notably the GEO (Group on Earth Observations) Carbon Strategy (Ciais et al., 2010), the Carbon Strategy for Carbon Observations from Space (CEOS, 2014), the GEO Carbon Flagship Work Programme (GEO, 2015) and Status of the Global Observing System for Climate (GCOS, 2015).
- A number of ocean acidification studies cite SOCAT, notably reports by the International Council for the Exploration of the Sea (ICES, 2013), the Joint OSPAR / ICES Ocean Acidification Study Group (ICES, 2014), the Global Ocean Acidification Observing Network (Newton et al., 2014) and the Secretariat of the Convention on Biodiversity (2014).

7.3 Scientific applications of SOCAT

SOCAT is used for a variety of scientific applications (Fig. 7b), some of which imply a wider relevance for SOCAT data products than envisaged during the creation of SOCAT (IOCCP, 2007). Scientific applications of SOCAT include:

- Figures of surface ocean CO₂ observations,
- Use of SOCAT tools and protocols,
- Use of surface ocean fCO₂ in diverse environmental studies,
- Model-data comparison, model evaluation and data assimilation,
- Detection of ocean acidification trends,
- Regional process studies of surface ocean fCO₂,
- Quantification of coastal ocean carbon sinks and sources,
- Quantification of the ocean carbon sink and its variation,
- Quantification of the land carbon sink.

These applications are roughly listed in order of the increasing importance of the SOCAT synthesis for the studies. The use of the SOCAT data collection in peer-reviewed, scientific publications is evolving. Initial publications made reference to the ongoing synthesis activity. Actual use of the SOCAT data collection started as soon as version 1 was released in 2011 (Pfeil et al., 2013; Sabine et al., 2013). Studies that heavily rely on SOCAT data products, such as modelling, ocean acidification



trend analysis and carbon budgeting, represent one third to half of the scientific publications citing or naming SOCAT from 2013 onwards.

5 *Examples of scientific applications of SOCAT are given below. There is no strict separation between the different types of applications identified here, with several studies belonging to more than one type of application. Many of the studies use surface ocean $p\text{CO}_2$ values, derived from the $f\text{CO}_2$ values reported in SOCAT data products.*

Figures of surface ocean CO_2 observations: Newly created figures based on the SOCAT data collection and existing figures from SOCAT publications have been used in scientific publications. Such figures generally highlight the availability or lack of surface ocean CO_2 data in specific regions or seasons or over time (Chierici et al., 2012; Regnier et al., 2013; Wanninkhof et al., 2013a; Ciais et al., 2014; Majkut et al., 2014a; Brévière et al., 2015; Hofmann et al., 2015).

10 **Use of SOCAT tools and protocols:** A variety of tools and protocols has been developed in SOCAT. One of these is the definition of a continental margin mask which defines coastal waters as waters within 400 km from land (Pfeil et al., 2013). Evans and Mathis (2013) and Evans et al. (2015) use this continental margin mask. Other studies have adopted SOCAT protocols for calculation of $f\text{CO}_2$ (Ulfsbo et al., 2014) and quality control (Sutton et al., 2014).

15 **Use of surface ocean $f\text{CO}_2$ in diverse environmental studies:** Regional $f\text{CO}_2$ values from SOCAT are used in diverse environmental studies with topics ranging from ocean acidification, to genomics, gas transfer velocity and evaluation of independent measurements (Blomquist et al., 2014; Larsen et al., 2014; Holding et al., 2015; Marrec et al., 2015; Bonou et al., 2016; Reum et al., 2016). Reum et al. (2016) assess the co-variance between $p\text{CO}_2$, pH and other environmental parameters with the aim to improve the design of future ocean acidification incubation experiments. Larsen et al. (2014) establish a significant correlation between gene expression for the relative turnover (synthesis or consumption) of CO_2 and surface ocean $f\text{CO}_2$. SOCAT $f\text{CO}_2$ values are also used for evaluation of surface ocean $f\text{CO}_2$ estimates from eddy correlation (Blomquist et al., 2014) or from other carbonate parameters (Bonou et al., 2016) and for evaluation of regression parameterisations (Marrec et al., 2015; Xu et al., 2016).

25 **Model-to-data comparison, model evaluation and data assimilation:** SOCAT data products are used for model-to-data comparison, model evaluation and data assimilation in coupled and ocean-only biogeochemical models. Model-to-data comparisons of surface water $f\text{CO}_2$ have been carried out for seasonal (Tjiputra et al., 2012; Arruda et al., 2015) to multi-year time scales (Tjiputra et al., 2014; McKinley et al., 2016). In several studies, model data are subsampled to surface ocean $p\text{CO}_2$ observations from SOCAT (Séférian et al., 2014; Tjiputra et al., 2014; Turi et al., 2014). Cooley et al. (2015) evaluate surface ocean $p\text{CO}_2$ values from an integrated assessment model with $p\text{CO}_2$ observations from SOCAT and other sources. SOCAT data products are supporting model evaluation in context of the Coupled Model Intercomparison Project (CMIP) and beyond (Eyring et al., 2015). The SOCAT data collection is used for assimilation of surface ocean $p\text{CO}_2$ values in global ocean biogeochemical models (While et al. 2012; Simon and Bertino, 2013, as cited in Gehlen et al., 2015). Ocean biogeochemical models have many applications, such as quantification and attribution of trends in the ocean carbon sink (Le Quéré et al., 2014, 2015a, 2015b; Séférian et al., 2014) and forecasting population dynamics of sea scallops, basis of important commercial fisheries (Cooley et al., 2015).



Detection of ocean acidification trends: A number of studies estimate trends in surface ocean pH or the carbonate concentration by combining SOCAT $f\text{CO}_2$ values with another carbonate parameter (Lauvset and Gruber, 2014; Freeman and Lovenduski, 2015; Lauvset et al., 2015).

5 **Regional process studies of surface ocean $f\text{CO}_2$:** Several authors investigate regional processes driving temporal or spatial variation in surface ocean $f\text{CO}_2$ and CO_2 air-sea fluxes. Examples are for the Subantarctic Indian Ocean (Lourantou and Metzl, 2011) and the Eastern Equatorial Pacific Ocean (Walker Brown et al., 2015).

Quantification of coastal ocean carbon sinks and sources: SOCAT data products are used for quantification of CO_2 sources and sinks in coastal seas. Such studies are regional or global in extent (Chen et al., 2013; Signorini et al., 2013; Laruelle et al., 2014, 2015).

10 **Quantification of the ocean carbon sink and its variation:** An important application of the SOCAT data collection is quantification of the ocean carbon sink on seasonal to multi-year time scales with a mapping or gap-filling method. Such studies may be regional or global in extent. Studies tend to be either for the coastal seas (Signorini et al., 2013) or for the open ocean (Rödenbeck et al., 2015). The studies interpolate sparse $p\text{CO}_2$ data from a SOCAT or LDEO synthesis product in time and space by a gap-filling method. Approaches include statistical interpolation (Rödenbeck et al., 2013; Goddijn-Murphy et al., 2015; Jones et al., 2015b), multiple linear regression (Schuster et al., 2013; Signorini et al., 2013; Iida et al., 2015), neural network approaches (Landschützer et al., 2013, 2014; Nakaoka et al., 2013; Sasse et al., 2013; Zeng et al., 2014) and model-based regression and tuning (Valsala and Maksyutov, 2010; Majkut et al., 2014b). Mapping methods may be specific to individual regions ('biomes') (Signorini et al., 2013; Landschützer et al., 2014) or may apply to the full (global) domain (e.g. Rödenbeck et al., 2013; Jones et al., 2015b). Most of these approaches use additional parameters with good data coverage
15 during the gap-filling process, for example satellite-derived sea surface temperature and chlorophyll a, as well as sea surface salinity and mixed layer depth from reanalysis. Many mapping methods use a time-dependent variable, such as time itself or the steadily increasing atmospheric CO_2 mole fraction, in order to be able to reproduce a long-term increase in surface ocean $p\text{CO}_2$.

20 **The Surface Ocean $p\text{CO}_2$ Mapping Intercomparison** (<http://www.bgc-jena.mpg.de/SOCOM/>) compares the surface ocean $p\text{CO}_2$ distribution and air-sea CO_2 fluxes in 14 data-based mapping products, ten of them using SOCAT (Rödenbeck et al., 2015). The methods vary in their characteristics, making them suitable for different space and time scales. The SOCOM initiative aims to quantify uncertainties and to identify common features in the gap-filling methods. The first SOCOM results highlight considerable differences between mapping products, especially in data-sparse regions (Rödenbeck et al., 2015).

30 **The high-profile Global Carbon Budget** uses ocean biogeochemical models for estimating trends in the global ocean carbon sink (Le Quéré et al., 2014, 2015a, 2015b). Recent budgets also consider observation-based estimates of the ocean carbon sink using the LDEO and SOCAT synthesis products (Park et al., 2010; Landschützer et al., 2014, 2015; Rödenbeck et al., 2014). The 2015 Global Carbon Budget assesses the uncertainty in the ocean carbon sink by comparing model results to observation-based estimates (Le Quéré et al., 2015b).



Quantification of the land carbon sink: *Quantification of the ocean carbon sink is critical to resolving the Global Carbon Budget and underpins the estimate of the land carbon sink (Le Quéré et al., 2014, 2015a, 2015b). In addition, quantification of ocean-atmosphere CO₂ fluxes in space and time provides priors for atmospheric inversion, thus improving estimates of the land carbon sink (Rödenbeck et al., 2014; Van der Laan et al., 2014; Jones et al., 2015b).*

5 7.4 Scientific findings obtained using the SOCAT data collection

This section provides an overview of scientific findings obtained using the SOCAT data collection.

Model-to-data comparison: *Schuster et al. (2013) carry out a comparison of CO₂ air-sea fluxes for the Atlantic Ocean from data-based methods, ocean biogeochemical models, ocean inversion, and atmospheric inversions. The seasonal cycle and year-to-year variation in the fluxes differ between the various methods for most Atlantic regions.*

10 *Two studies subsample model pCO₂ data to surface ocean pCO₂ observations derived from SOCAT. The authors conclude that ocean biogeochemical models on average underestimate the spatial and temporal variation in regional and global surface ocean pCO₂ by 10 to 40 % (Séférian et al., 2014; Turi et al., 2014). This corroborates the SOCOM finding that ocean biogeochemical models underestimate the year-to-year and decadal variation in the global air-sea CO₂ flux (Rödenbeck et al., 2015). However, at least one model-to-data comparison study concludes that the Community Earth System Model captures*
15 *the annual to 30-year variability in the ocean carbon cycle at regional to global scales (McKinley et al., 2016). Landschützer et al. (2015) demonstrate how ocean carbon observations are delivering new insights into large and globally significant decadal changes in the ocean carbon sink. The variability in regions like the Southern Ocean is not apparent in modelled estimates of ocean carbon uptake or from atmospheric inverse calculations, which can show considerable differences for regions (e.g. Lenton et al., 2013).*

20 **Detection of ocean acidification trends:** *SOCAT-based research indicates a decrease in global surface ocean pH at a rate of $-0.0018 \pm 0.0004 \text{ yr}^{-1}$ for 1991 to 2011 with significant decreases in 70% of all ocean regions (Lauvset et al., 2015).*

Data-based carbon budgeting: *Using SOCAT and other data sources, Regnier et al. (2013) estimate that anthropogenic activities may have increased open ocean outgassing of land-derived carbon by 0.1 Pg C yr⁻¹. The global CO₂ sink in continental shelf seas has been estimated as 0.4 Pg C yr⁻¹ (Chen et al., 2013) and $0.19 \pm 0.05 \text{ Pg C yr}^{-1}$ (Laruelle et al., 2014).*

25 *Several mapping studies highlight large year-to-year variation in air-sea CO₂ fluxes in the tropical Pacific Ocean (Landschützer et al., 2014; Rödenbeck et al., 2014, 2015). This variation is closely related to the El Niño-Southern Oscillation (ENSO) (Feely et al., 1999, 2002; Inoue et al., 2001; Rödenbeck et al., 2014). The variation in the equatorial Pacific Ocean roughly corresponds to 40% of the interannual variation in the global ocean carbon sink (Rödenbeck et al., 2014), which has been estimated as 0.31 Pg C yr⁻¹ (Rödenbeck et al., 2015).*

30 *The SOCOM comparison of mapping methods identifies an increase in global ocean carbon sink by 1 Pg C decade⁻¹ since 2000 (Rödenbeck et al., 2015). About half of this increase in the global ocean carbon sink originates south of 35°S in the Southern Ocean (Landschützer et al., 2014, 2015).*



8 Conclusions

SOCAT version 3 represents an important release of the SOCAT data collection, by creating a 58-year data record and by adding many additional data sets for recent years. This article provides an ESSD ‘Living Data’ update of SOCAT version 3. The launch of the SOCAT automation system will enable annual SOCAT releases from version 4 onwards.

5 *The rapid growth of scientific publications using SOCAT (Fig. 7) demonstrates the importance of this synthesis activity by the international marine carbon community. The SOCAT data collection is being used in high-impact, scientific applications such as evaluation of ocean biogeochemical models, carbon budgeting and trend analysis of the ocean carbon sink and ocean acidification. SOCAT-based studies have informed the Paris climate negotiations, as the 2015 Global Carbon Budget was released at the 21st Conference of the Parties of the United Nations Framework Convention on Climate Change (Le Quéré et al., 2015b).*

10 However, despite much progress in data synthesis, major uncertainties remain in observation-based studies of the ocean carbon sink and ocean acidification due to 1) inadequate spatial and seasonal data coverage, 2) short data records, 3) uncertainty in the correction for ‘natural’, pre-industrial oceanic outgassing of land-derived CO₂ (Jacobsen et al., 2007) and any anthropogenic perturbation of this outgassing (Regnier et al., 2013). Data coverage is particularly poor in the Indian Ocean, the southern hemisphere oceans and coastal seas and in the high-latitude oceans, notably in ice-covered regions and in winter (Figs. 3, 4 and 6).

15 The above reinforces the need for the continuing collection and synthesis of accurate, well-calibrated and well-documented observations. Adequate resources need to continue to be made available for data collection, quality control and data synthesis. Systems should be automated whenever possible. The SOCAT data synthesis highlights the success of a bottom-up approach with buy-in from the international marine carbon community and endorsement by IOCCP, SOLAS and IMBER.

20 **Acknowledgements.** *Research vessel Tiglax in Columbia Bay, Alaska, is shown on the website for SOCAT version 3. The Columbia Glacier can be seen at the head of the bay, as well as calved ice from the glacier. The photo was taken by Wiley Evans. Pete Brown (National Oceanography Centre Southampton, UK) designed the SOCAT logo. IOCCP (via the US National Science Foundation grant (OCE- 124 3377) to the Scientific Committee on Oceanic Research), IOC-UNESCO (International Oceanographic Commission of the United Nations Educational, Scientific and Cultural Organization), SOLAS and IMBER provided travel and meeting support. Funding was received from the University of East Anglia (UK), the Bjerknes Centre for Climate Research (Norway), the Geophysical Institute at the University of Bergen (Norway) and the University of Washington (US). The US National Oceanic and Atmospheric Administration (NOAA) made important financial contributions via the Climate Observation Division of the Climate Program Office, the NOAA Ocean Acidification Program, the NOAA Pacific Marine Environmental Laboratory (PMEL), the NOAA Atlantic Oceanographic and Meteorological Laboratory (AOML) and the NOAA Earth System Research Laboratory. Funding was also received from Oak Ridge National Laboratory (US), PANGAEA® Data Publisher for Earth and Environmental Science (Germany), the Alfred Wegener Institute Helmholtz*



Centre for Polar and Marine Research (Germany), the Antarctic Climate and Ecosystems Cooperative Research Centre (Australia), the National Institute for Environmental Studies (Japan) and Uni Research (Norway). Research projects making SOCAT possible included the European Union projects CarboChange (FP7 264879), GEOCARBON (FP7 283080) and AtlantOS (633211), the UK Ocean Acidification Research Programme (NE/H017046/1; funded by the Natural Environment Research Council (NERC) and the Departments for Energy and Climate Change and for Environment, Food and Rural Affairs (Defra)) and the UK Shelf Sea Biogeochemistry Blue Carbon project (NE/K00168X/1; funded by NERC and Defra). Numerous government and funding agencies financially supported SOCAT notably the Australian International Marine Observing System, the European Space Agency, the German Federal Ministry of Education and Research (BMBF projects 01LK1224J, 01LK1101C, 01LK1101E, ICOS-D), the Japanese Ministry of the Environment, the Royal Society of New Zealand via the New Zealand - Germany Science and Technology Programme, the Norwegian Research Council (SNACS, 229752), the Swedish Research Council (project 2004-4034) and the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (Formas, project 2004-797). This is PMEL contribution number 4441.

References

- Antonov, J. I., Locarnini, R. A., Boyer, T. P., Mishonov, A. V., and Garcia, H. E.: World Ocean Atlas 2005, in: Volume 2: Salinity, edited by Levitus, S., NOAA Atlas NESDIS 62, US Government Printing Office, Washington, DC, 182 pp., 2006.
- Arruda, R., Calil, P. H. R., Bianchi, A. A., Doney, S. C., Gruber, N., Lima, I., and Turi, G.: Air-sea CO₂ fluxes and the controls on ocean surface pCO₂ variability in coastal and open-ocean southwestern Atlantic Ocean: A modeling study, *Biogeosciences*, 12, 5793-5809, doi:10.5194/bg-12-5793-2015, 2015.
- Bakker, D. C. E., Hoppema, M., Schröder, M., Geibert, W., and De Baar, H. J. W.: A rapid transition from ice covered CO₂-rich waters to a biologically mediated CO₂ sink in the eastern Weddell Gyre. *Biogeosciences*, 5, 1373-1386, doi:10.5194/bg-5-1373-2008, 2008.
- Bakker, D. C. E., Pfeil, B., Smith, K., Hankin, S., Olsen, A., Alin, S. R., Cosca, C., Harasawa, S., Kozyr, A., Nojiri, Y., O'Brien, K. M., Schuster, U., Telszewski, M., Tilbrook, B., Wada, C., Akl, J., Barbero, L., Bates, N. R., Boutin, J., Bozec, Y., Cai, W.-J., Castle, R. D., Chavez, F. P., Chen, L., Chierici, M., Currie, K., De Baar, H. J. W., Evans, W., Feely, R. A., Fransson, A., Gao, Z., Hales, B., Hardman-Mountford, N. J., Hoppema, M., Huang, W.-J., Hunt, C. W., Huss, B., Ichikawa, T., Johannessen, T., Jones, E. M., Jones, S., Jutterstrøm, S., Kitidis, V., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Manke, A. B., Mathis, J. T., Merlivat, L., Metzl, N., Murata, A., Newberger, T., Omar, A. M., Ono, T., Park, G.-H., Paterson, K., Pierrot, D., Ríos, A. F., Sabine, C. L., Saito, S., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Sieger, R., Skjelvan, I., Steinhoff, T., Sullivan, K. F., Sun, H., Sutton, A. J., Suzuki, T., Sweeney, C., Takahashi, T., Tjiputra, J., Tsurushima, N., Van Heuven, S. M. A. C., Vandemark, D., Vlahos, P., Wallace, D. W. R., Wanninkhof, R., and Watson, A. J.: An update to the Surface Ocean CO₂ Atlas (SOCAT version 2). *Earth Syst. Sci. Data*, 6, 69-90, doi:10.5194/essd-6-69-2014, 2014.



- Bange, H. W., Bell, T. G., Cornejo, M., Freing, A., Uher, G., Upstill-Goddard, R. C., Zhang, G.: MEMENTO: A proposal to develop a database of marine nitrous oxide and methane measurements. *Environ Chemistry* 6, 195–197, 2009.
- Bates, N.R., Astor, Y. M., Church, M. J., Currie, K., Dore, J. E., González-Dávila, M., Lorenzoni, L., Muller-Karger, F., Olafsson, J., and Santana-Casiano, J. M.: A time-series view of changing ocean chemistry due to ocean uptake of anthropogenic
- 5 CO₂ and ocean acidification. *Oceanography*, 27 (1), 126–141, doi:10.5670/oceanog.2014.16, 2014.
- Blomquist, B. W., Huebert, B. J., Fairall, C. W., Bariteau, L., Edson, J. B., Hare, J. E., and McGillis, W. R.: Advances in air–sea CO₂ flux measurement by eddy correlation. *Boundary-Layer Meteorol.*, 152, 245–276, doi:10.1007/s10546-014-9926-2, 2014.
- Blunden, J., and Arndt, D. S. (eds.): State of the Climate in 2014. *B. Am. Meteorol. Soc.*, 96 (7), S1–S267, available at:
- 10 <http://journals.ametsoc.org/doi/pdf/10.1175/2015BAMSSStateoftheClimate.1> (last access: 8 December 2015), 2015.
- Bonou, F. K., Noriega, C., Lefèvre, N., and Araujo, M.: Distribution of CO₂ parameters in the Western Tropical Atlantic Ocean, *Dynamics of Atmospheres and Oceans*, 73, 47–60, doi:10.1016/j.dynatmoce.2015.12.001, 2016.
- Borges, A. V., Alin, S. R., Chavez, F. P., Vlahos, P., Johnson, K. S., Holt, J. T., Balch, W. M., Bates, N., Brainard, R., Cai, W.-J., Chen, C. T. A., Currie, K., Dai, M., Degrandpré, M., Delille, B., Dickson, A., Evans, W., Feely, R. A., Friederich, G.
- 15 E., Gong, G.-C., Hales, B., Hardman-Mountford, N., Hendee, J., Hernandez-Ayon, J. M., Hood, M., Huertas, E., Hydes, D., Ianson, D., Krasakopoulou, E., Litt, E., Luchetta, A., Mathis, J., McGillis, W. R., Murata, A., Newton, J., Ólafsson, J., Omar, A., Perez, F. F., Sabine, C., Salisbury, J. E., Salm, R., Sarma, V. V. S. S., Schneider, B., Sigler, M., Thomas, H., Turk, D., Vandemark, D., Wanninkhof, R., and Ward, B.: A global sea surface carbon observing system: Inorganic and organic carbon dynamics in coastal oceans, in: *Proceedings of OceanObs'09: Sustained Ocean Observations and Information for Society*,
- 20 edited by: Hall, J., Harrison, D. E., and Stammer, D., Vol. 2, Venice, Italy, 21–25 September 2009, ESA Publication WPP-306. doi:10.5270/OceanObs09.cwp.07, 2010.
- Boutin, J., and Merlivat, L.: New in situ estimates of carbon biological production rates in the Southern Ocean from CARIOCA drifter measurements, *Geophys. Res. Lett.*, 36, L13608, doi:10.1029/2009GL038307, 2009.
- Brévière, E. H. G., Bakker, D. C. E., Bange, H. W., Bates, T. S., Bell, T. G., Boyd, P. W., Duce, R. A., Garçon, V., Johnson, M. T., Law, C. S., Marandino, C. A., Olsen, A., Quack, B., Quinn, P. K., Sabine, C. L., and Saltzman, E.: Surface ocean -
- 25 lower atmosphere study: Scientific synthesis and contribution to Earth System science, *Anthropocene*, 12, 54–68, doi.org/10.1016/j.ancene.2015.11.001, 2015.
- CEOS: Carbon strategy for carbon observations from space, The Committee on Earth Observation Satellites (CEOS) response to the Group on Earth Observations (GEO) Carbon Strategy, available at:
- 30 http://ceos.org/document_management/Publications/WGClimate_CEOS-Strategy-for-Carbon-Observations-from-Space_Apr2014.pdf, (last access: 8 December 2015), 2014.
- Chen, C.-T. A., Huang, T.-H., Chen, Y.-C., Bai, Y., He, X., and Kang, Y.: Air–sea exchanges of CO₂ in the world's coastal seas, *Biogeosciences*, 10, 6509–6544, doi:10.5194/bg-10-6509-2013, 2013.



- Chierici, M., Signorini, S.R., Mattsdotter-Björk, M., Fransson, A., and Olsen, A.: Surface water fCO₂ algorithms for the high-latitude Pacific sector of the Southern Ocean, *Remote Sens. Environm.*, 119, 184-196, doi:10.1016/j.rse.2011.12.020, 2012.
- Ciais, P., Dolman, A. J., Dargaville, R., Barrie, L., Bombelli, A., Butler, J., Canadell, P., and Moriyama, T.: GEO Carbon Strategy, GEO (Group on Earth Observations) Secretariat Geneva / Food and Agriculture Organization of the United Nations, Rome, 48 pp, available at: http://www.globalcarbonproject.org/global/pdf/GEO_CARBNSTRATEGY_20101020.pdf (last access: 8 December 2015), 2010.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S. and Thornton, P.: Carbon and other biogeochemical cycles, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V. and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, available at <http://www.ipcc.ch/report/ar5/wg1/> (last access: 3 January 2015), 2013.
- Ciais, P., Dolman, A. J., Bombelli, A., Duren, R., Peregon, A., Rayner, P. J., Miller, C., Gobron, N., Kinderman, G., Marland, G., Gruber, N., Chevallier, F., Andres, R. J., Balsamo G., Bopp, L., Bréon, F.-M., Broquet, G., Dargaville, R., Battin, T. J., Borges, A., Bovensmann, H., Buchwitz, M., Butler, J., Canadell, J. G., Cook, R. B., DeFries, R., Engelen, R., Gurney, K. R., Heinze, C., Heimann, M., Held, A., Henry, M., Law, B., Luysaert, S., Miller, J., Moriyama, T., Moulin, C., Myneni, R. B., Nussli, C., Obersteiner, M., Ojima, D., Pan, Y., Paris, J.-D., Piao, S. L., Poulter, B., Plummer, S., Quegan, S., Raymond, P., Reichstein, M., Rivier, L., Sabine, C., Schimel, D., Tarasova, O., Valentini, R., Wang, R., Van der Werf, G., Wickland, D., Williams, M., and Zehner, C.: Current systematic carbon-cycle observations and the need for implementing a policy-relevant carbon observing system, *Biogeosciences*, 11, 3547-3602, doi:10.5194/bg-11-35477-2014, 2014.
- Cooley, S. R., Rheuban, J. E., Hart, D. R., Luu, V., Glover, D. M., Hare, J. A., Doney, S. C.: An Integrated assessment model for helping the United States sea scallop (*Placopecten magellanicus*) fishery plan ahead for ocean acidification and warming, *PLoS ONE* 10(5), e0124145, doi:10.1371/journal.pone.0124145, 2015.
- Dickson, A. G., Sabine, C. L., and Christian, J. R. (Eds.): Guide to best practices for ocean CO₂ measurements, PICES Special Publication 3, IOCCP Report 8, 191 pp., 2007.
- Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean Acidification: The other CO₂ problem. *Ann. Rev. Mar. Sci.*, 1, 169-192, doi:10.1146/annurev.marine.010908.163834, 2009a.
- Doney, S. C., Tilbrook, B., Roy, S., Metzl, N., Le Quéré, C., Hood, M., Feely, R. A., and Bakker, D. C. E.: Surface Ocean CO₂ Variability and Vulnerability. *Deep-Sea Res. II*, 56, 504-511, doi:10.1016/j.dsr2.2008.12.016, 2009b.
- ETOPO2: 2-minute Gridded Global Relief Data (ETOPO2v2), National Geophysical Data Center, National Oceanic and Atmospheric Administration, US Dept. of Commerce, available at: <http://www.ngdc.noaa.gov/mgg/global/global.html> (last access: 1 September 2015), 2006.
- Evans, W., and Mathis, J. T.: The Gulf of Alaska coastal ocean as an atmospheric CO₂ sink, *Cont. Shelf Res.*, 65, 52-63, doi:10.1016/j.csr.2013.06.013, 2013.



- Evans, W., Mathis, J. T., Cross, J. N., Bates, N. R., Frey, K. E., Else, B. G. T., Papkyriakou, T. N., DeGrandpre, M. D., Islam, F., Cai, W.-J., Chen, B., Yamamoto-Kawai, M., Carmack, E., Williams, W. J., and Takahashi, T.: Sea-air CO₂ exchange in the western Arctic coastal ocean, *Global Biogeochem. Cy.*, 29, 1–20, doi:10.1002/2015GB005153, 2015.
- Eyring, V., Righi, M., Evaldsson, M., Lauer, A., Wenzel, S., Jones, C., Anav, A., Andrews, O., Cionni, I., Davin, E. L., Deser, C., Ehbrecht, C., Friedlingstein, P., Gleckler, P., Gottschaldt, K.-D., Hagemann, S., Jukes, M., Kindermann, S., Krasting, J., Kunert, D., Levine, R., Loew, A., Mäkelä, J., Martin, G., Mason, E., Phillips, A., Read, S., Rio, C., Roehrig, R., Senftleben, D., Sterl, A., Van Ulft, L. H., Walton, J., Wang, S., Williams, K. D.: ESMValTool (v1.0) – A community diagnostic and performance metrics tool for routine evaluation of Earth System Models in CMIP. *Geosci. Model Development Disc.*, 8, 7541–7661, doi:10.5194/gmdd-8-7541-2015, 2015.
- 5 C., Ehbrecht, C., Friedlingstein, P., Gleckler, P., Gottschaldt, K.-D., Hagemann, S., Jukes, M., Kindermann, S., Krasting, J., Kunert, D., Levine, R., Loew, A., Mäkelä, J., Martin, G., Mason, E., Phillips, A., Read, S., Rio, C., Roehrig, R., Senftleben, D., Sterl, A., Van Ulft, L. H., Walton, J., Wang, S., Williams, K. D.: ESMValTool (v1.0) – A community diagnostic and performance metrics tool for routine evaluation of Earth System Models in CMIP. *Geosci. Model Development Disc.*, 8, 7541–7661, doi:10.5194/gmdd-8-7541-2015, 2015.
- 10 Fay, A. R., and McKinley, G. A.: Global trends in surface ocean pCO₂ from in situ data. *Global Biogeochem. Cy.*, 27, 541–557, doi:10.1002/gbc.20051, 2013.
- Fay, A. R., McKinley, G. A., and Lovenduski, N. S.: Southern Ocean carbon trends: Sensitivity to methods. *Geophys. Res. Lett.*, 41, 6833–6840, doi:10.1002/2014GL061324, 2014.
- Feely, R., Boutin, J., Cosca, C., Dandonneau, Y., Etcheto, J., Inoue, H., Ishii, M., Le Quéré, C. L., Mackey, D., McPhaden, M., Metzl, N., Poisson, A., and Wanninkhof, R.: Seasonal and interannual variability of CO₂ in the Equatorial Pacific, *Deep-Sea Res. Pt. II*, 49, 2443–2469, doi:10.1016/S0967-0645(02)00044-9, 2002.
- 15 Feely, R. A., Doney, S. C., and Cooley, S. R.: Ocean acidification: Present conditions and future changes in a High-CO₂ world. *Oceanography*, 22 (4), 36–47, doi:10.5670/oceanog.2009.95, 2009.
- Feely, R. A., Wanninkhof, R., Takahashi, T., and Tans, P.: Influence of El Niño on the equatorial Pacific contribution to atmospheric CO₂ accumulation, *Nature*, 398, 597–601, 1999.
- 20 Ferraro, R., Waliser, D. E., Gleckler, P., Taylor, K. E., and Eyring, V.: Evolving Obs4MIPs to support Phase 6 of the Coupled Model Intercomparison Project (CMIP6). *Bull. Amer. Meteor. Soc.*, 96, ES131–ES133, doi:10.1175/BAMS-D-14-00216.1, 2015.
- Fransson A, Chierici, M., and Nojiri, Y.: Increased net CO₂ outgassing in the upwelling region of the southern Bering Sea in a period of variable marine climate between 1995 and 2001, *J. Geophys. Res.*, 111, C08008, doi:10.1029/2004JC002759, 2006
- 25 Fransson, A., Chierici, M., and Nojiri, Y.: New insights into the spatial variability of the surface water CO₂ in varying sea ice conditions in the Arctic Ocean, *Cont. Shelf Res.*, 29, 1317–1328, doi:10.1016/j.csr.2009.03.008, 2009.
- Freeman, N. M., and Lovenduski, N. S.: Decreased calcification in the Southern Ocean over the satellite record, *Geophys. Res. Lett.*, 42, 1834–1840, doi:10.1002/2014GL062769, Supplement, 2015.
- 30 FOO: A Framework for Ocean Observing, Task Team for an Integrated Framework for Sustained Ocean Observing, UNESCO 2012, IOC/INF-1284 rev., doi:10.5270/OceanObs09-FOO, available at: http://www.jodc.go.jp/info/ioc_doc/INF/211260e.pdf (last access: 28 November 2015), 2012.
- GCOS: Status of the Global Observing System for Climate. GCOS-195. 358 pp, available at: http://www.wmo.int/pages/prog/gcos/Publications/GCOS-195_en.pdf (last access: 09 February 2016), 2015.



- Gehlen, M., Barciela, R., Bertino, L., Brasseur, P., Butenschön, M., Chai, F., Crise, A., Drillet, Y., Ford, D., Lavoie, D., Lehodey, P., Perruche, C., Samuelsen, A., and Simon, E.: Building the capacity for forecasting marine biogeochemistry and ecosystems: Recent advances and future developments, *J. Operational Oceanography*, 8(1), 168-187, doi:10.1080/1755876X.2015.1022350, 2015.
- 5 GEO: 2016 Work Programme. Document 15, Submitted to the GEO-XII Plenary for decision, Group on Earth Observations, available at: https://www.earthobservations.org/documents/geo_xii/GEO-XII_15_2016%20Work%20Programme.pdf (last access: 8 December 2015), 2015.
- GLOBALVIEW-CO₂: Cooperative Atmospheric Data Integration Project – Carbon Dioxide, 2012 version, Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, Colorado, USA, available at: 10 <http://www.esrl.noaa.gov/gmd/ccgg/globalview/> (last access: 26 June 2013), 2012.
- GLOBALVIEW-CO₂: Cooperative Atmospheric Data Integration Project – Carbon Dioxide, 2014 version, Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, Colorado, USA, available at: <http://www.esrl.noaa.gov/gmd/ccgg/mbl/data.php> (last access: 29 October 2015), 2014.
- 15 Goddijn-Murphy, L. M., Woolf, D. K., Land, P. E., Shutler, J. D., and Donlon, C.: The OceanFlux Greenhouse Gases methodology for deriving a sea surface climatology of CO₂ fugacity in support of air–sea gas flux studies, *Ocean Sci.*, 11, 519-541, doi:10.5194/os-11-519-2015, 2015.
- González-Dávila, M., Santana-Casiano, J. M., and González-Dávila, E. F.: Interannual variability of the upper ocean carbon cycle in the northeast Atlantic Ocean. *Geophys. Res. Lett.*, 34, L07608, doi:10.1029/2006GL028145, 2007.
- Hales, B., Strutton, P. G., Saraceno, M., Letelier, R., Takahashi, T., Feely, R., Sabine, C., and Chavez, F.: Satellite-based 20 prediction of pCO₂ in coastal waters of the eastern North Pacific, *Prog. Oceanography*, 103, 1-15, doi:10.1016/j.pocean.2012.03.001, 2012.
- Hales, B., Takahashi, T., and Bandstra, L.: Atmospheric CO₂ uptake by a coastal upwelling system. *Global Biogeochem. Cy.*, 19(1), GB1009, doi:10.1029/2004GB002295, 2005.
- Harris, K. E., DeGrandpre, M. D., and Hales, B.: Aragonite saturation state dynamics in a coastal upwelling zone, *Geophys. 25 Res. Lett.* 40(11), 2720-2725, doi:10.1002/grl.50460, 2013.
- Henderson, C.: Ocean acidification: The other CO₂ problem, *New Scientist*, 2 August 2006, available at: <http://environment.newscientist.com/article/mg19125631.200> (last access: 19 January 2016), 2006.
- Hofmann, E., Bundy, A., Drinkwater, K., Piola, A. R., Avril, B., Robinson, C., Murphy, E., Maddison, L., Svendsen, E., Hall, J., and Xu, Y.: IMBER research for marine sustainability: Synthesis and the way forward, *Anthropocene*, 12, 42-53, 30 doi:10.1016/j.ancene.2015.12.002, 2015.
- Holding J. M., Duarte, C. M., Sanz-Martin, M., Mesa, E., Arrieta, J. M., Chierici, M., Hendriks, I. E., Garcia-Corral, L. S., Regaudie-de-Gioux, A., Delgado, A., Reigstad, M., Wassmann, P., and Agusti, S.: Temperature dependence of CO₂-enhanced primary production in the European Arctic Ocean, *Nature Climate Change*, 5, 1079-1082, doi:10.1038/nclimate2768, 2015.



- Hood, R. R., Bange, H. W., Beal, L., Beckley, L. E., Burkill, P., Cowie, G. L., D'Adamo, N., Ganssen, G., Hendon, H., Hermes, J., Honda, M., McPhaden, M., Roberts, M., Singh, S., Urban, E., Yu, W.: Science Plan of the Second International Indian Ocean Expedition (IIOE-2): A Basin-Wide Research Program. Scientific Committee on Oceanic Research, Newark, Delaware, USA, available at: <http://www.iioe-2.incois.gov.in/IIOE-2/pdfviewer.jsp?docname=IIOE-2-SciencePlan-2015-2020.pdf> (last access: 25 February 2016), 2015.
- ICES (International Council for the Exploration of the Sea): Chemical aspects of ocean acidification monitoring in the ICES marine area, ICES Cooperative Research Report, 319, 1-86, 2013.
- ICES (International Council for the Exploration of the Sea): Final Report to OSPAR of the Joint OSPAR / ICES Ocean Acidification Study Group (SGOA), ICES CM 2014 / ACOM: 67, 141 pp, available at: http://ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2014/SGOA/sgoa_finalOSPAR_2015.pdf (last access: 18 January 2016), 2014.
- Iida, Y., Kojima, A., Takatani, Y., Nakano, T., Sugimoto, H., Midorikawa, T., and Ishii, M.: Trends in pCO₂ and sea-air CO₂ flux over the global open oceans for the last two decades. *J. Oceanography*, 71, 637-661, doi:10.1007/s10872-015-0306-4, 2015.
- Inoue, H., Ishii, M., Matsueda, H., Saito, S., Aoyama, M., Tokieda, T., Midorikawa, T., Nemoto, K., Kawano, T., Asanuma, I., Ando, K., Yano, T., and Murata, A.: Distributions and variations in the partial pressure of CO₂ in surface waters (pCO_{2w}) of the central and western equatorial Pacific during the 1997/1998 El Niño event, *Mar. Chem.*, 76, 59-75, 2001.
- IOCCP: Surface Ocean CO₂ Variability and Vulnerabilities Workshop. UNESCO, Paris, France, 11-14 April 2007, IOCCP Report 7, available at: <http://www.ioccp.org/> (last access: 1 May 2013), 2007.
- Jacobson, A. R., Mikaloff Fletcher, S. E., Gruber, N., Sarmiento, J., and Gloor, M.: A joint atmosphere-ocean inversion for surface fluxes of carbon dioxide: 2. Regional results, *Global Biogeochem. Cy.*, 21, GB1020, doi:10.1029/2006GB002703, 2007.
- Johengen, T.: Performance demonstration statement PMEL MAPCO₂/Battelle Seaology pCO₂ monitoring system, Alliance for Coastal Technologies report, ACT DS10-02, available at: http://www.act-us.info/Download/Evaluations/pCO2/PMEL_MAPCO2_Battelle_Seaology/ (last access: 10 January 2014), 2010.
- Jones, E. M., Bakker, D. C. E., Venables, H. J., and Hardman-Mountford, N.: Seasonal cycle of CO₂ from the sea ice edge to island blooms in the Scotia Sea, Southern Ocean, *Mar. Chem.*, 177, 490-500, doi:10.1016/j.marchem.2015.06.031, 2015a.
- Jones, S. D., Le Quéré, C., Rödenbeck, C., Manning, A. C., Olsen, A.: A statistical gap-filling method to interpolate global monthly surface ocean carbon dioxide data. *J. Adv. Modeling Earth Syst.*, 7, 1554-1575, doi:10.1002/2014MS000416, 2015b.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *Bull. Am. Meteorological Soc.*, 77, 437-470, 1996.



- Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M., and Gattuso, J.-P.: Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Global Change Biology*, 19, 1884-1896, doi:10.1111/gcb.12179, 2013.
- Landschützer, P., Gruber, N., Bakker, D. C. E., Haumann, F. A., Rödenbeck, C., Van Heuven, S., Hoppema, M., Metzl, N., Sweeney, C., Takahashi, T., Tilbrook, B., and Wanninkhof, R.: The reinvigoration of the Southern Ocean carbon sink, *Science*, 5 349 (6253), 1221-1224, doi:10.1126/science.aab2620, 2015.
- Landschützer, P., Gruber, N., Bakker, D. C. E., and Schuster, U.: Recent variability of the global ocean carbon sink, *Global Biogeochem. Cy.*, 28, 927-949, doi:10.1002/2014GB004853, 2014.
- Landschützer, P., Gruber, N., Bakker, D.C.E., Schuster, U., Nakaoka, S. Payne, M.R., Sasse, T., and Zeng, J.: A neural network-based estimate of the seasonal to inter-annual variability of the Atlantic Ocean carbon sink, *Biogeosciences*, 10, 7793-7815, doi:10.5194/bg-10-7793-2013, 2013.
- Larsen, P. E., Scott, N., Post, A. F., Field, D., Knight, R., Hamada, Y., and Gilbert, J. A.: Satellite remote sensing data can be used to model marine microbial metabolite turnover, *The ISME Journal*, 9 (1), 166-179, doi:10.1038/ismej.2014.107, 2014.
- Laruelle, G. G., Lauerwald, R., Pfeil, B., and Regnier, P.: Regionalized global budget of the CO₂ exchange at the air-water interface in continental shelf seas. *Global Biogeochem. Cy.*, 28, 1199-1214, doi:10.1002/2014GB004832, 2014.
- Laruelle, G. G., Lauerwald, R., Rotschi, J., Raymond, P. A., Hartmann, J., and Regnier, P.: Seasonal response of air-water CO₂ exchange along the land-ocean aquatic continuum of the North East American coast, *Biogeosciences*, 12, 1447-1458, doi:10.5194/bg-12-1447-2015, 2015.
- Lauvset, S. K., and Gruber, N.: Long-term trends in surface ocean pH in the North Atlantic. *Mar. Chem.*, 162, 71-76, doi:10.1016/j.marchem.2014.03.009, 2014.
- Lauvset, S. K., Gruber, N., Landschützer, P., Olsen, A., and Tjiputra, J.: Trends and drivers in global surface ocean pH over the past 3 decades, *Biogeosciences*, 12, 1285-1298, doi:10.5194/bg-12-1285-2015, 2015.
- Legge, O. J., Bakker, D. C. E., Johnson, M. T., Meredith, M. P., Venables, H. J., Brown, P. J. and Lee, G. A.: The seasonal cycle of ocean-atmosphere CO₂ flux in Ryder Bay, West Antarctic Peninsula. *Geophys. Res. Lett.*, 42 (8), 2934-2942, doi:10.1002/2015GL063796, 2015.
- Le Quéré, C., Peters, G. P., Andres, R. J., Andrew, R. M., Boden, T., Ciais, P., Friedlingstein, P., Houghton, R. A., Marland, G., Moriarty, R., Sitch, S., Tans, P., Arneth, A., Arvanitis, A., Bakker, D. C. E., Bopp, L., Canadell, J. G., Chini, L. P., Doney, S. C., Harper, A., Harris, I., House, J. I., Jain, A. K., Jones, S. D., Kato, E., Keeling, R. F., Klein Goldewijk, K., Körtzinger, A., Koven, C., Lefèvre, N., Maignan, F., Omar, A., Ono, T., Park, G.-H., Pfeil, B., Poulter, B., Raupach, M. R., Regnier, P., Rödenbeck, C., Saito, S., Schwinger, J., Segschneider, J., Stocker, B. D., Takahashi, T., Tilbrook, B., Van Heuven, S., Viovy, N., Wanninkhof, R., Wiltshire, A., and Zaehle, S.: Global Carbon Budget 2013, *Earth Syst. Sci. Data*, 6, 235-263, doi:10.5194/essd-6-235-2014, 2014.
- Le Quéré, C., Moriarty, R., Andrew, R. M., Peters, G. P., Ciais, P., Friedlingstein, P., Jones, S. D., Sitch, S., Tans, P., Arneth, A., Boden, T. A., Bopp, L., Bozec, Y., Canadell, J. G., Chini, L. P., Chevallier, F., Cosca, C. E., Harris, I.,



- Hoppema, M., Houghton, R. A., House, J. I., Jain, A. K., Johannessen, T., Kato, E., Keeling, R. F., Kitidis, V., Klein Goldewijk, K., Koven, C., Landa, C. S., Landschützer, P., Lenton, A., Lima, I. D., Marland, G., Mathis, J. T., Metzl, N., Nojiri, Y., Olsen, A., Ono, T., Peng, S., Peters, W., Pfeil, B., Poulter, B., Raupach, M. R., Regnier, P., Rödenbeck, C., Saito, S., Salisbury, J. E., Schuster, U., Schwinger, J., Séférian, R., Segsneider, J., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Werf, G. R., Viovy, N., Wang, Y.-P., Wanninkhof, R., Wiltshire, A., and Zeng, N.: Global Carbon Budget 2014. *Earth Syst. Sci. Data*, 7, 47-85, doi:10.5194/essd-7-47-2015, 2015a.
- Le Quéré, C., Moriarty, R., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Friedlingstein, P., Peters, G. P., Andres, R. J., Boden, T. A., Houghton, R. A., House, J. I., Keeling, R. F., Tans, P., Arneeth, A., Bakker, D. C. E., Barbero, L., Bopp, L., Chang, J., Chevallier, F., Chini, L. P., Ciais, P., Fader, M., Feely, R., Gkritzalis, T., Harris, I., Hauck, J., Ilyina, T., Jain, A. K., Kato, E., Kitidis, V., Klein Goldewijk, K., Koven, C., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lima, I. D., Metzl, N., Millero, F., Munro, D. R., Murata, A., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y., O'Brien, K., Olsen, A., Ono, T., Pérez, F. F., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Rödenbeck, C., Saito, S., Schuster, U., Schwinger, J., Séférian, R., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., Van der Laan-Luijckx, I. T., Van der Werf, G. R., Van Heuven, S., Vandemark, D., Viovy, N., Wiltshire, A., Zaehle, S., and Zeng, N.: Global Carbon Budget 2015, *Earth Syst. Sci. Data* 7, 349-396, doi:10.5194/essd-7-349-2015, 2015b.
- Lenton, A., Tilbrook, B., Law, R. M., Bakker, D. C. E., Doney, S. C., Gruber, N., Ishii, M., Hoppema, M., Lovenduski, N. S., Matear, R. J., McNeil, B. I., Metzl, N., Mikaloff-Fletcher, S., Monteiro, P. M. S., Rodenbeck, C., Sweeney, C., and Takahashi, T.: Sea-air CO₂ fluxes in the Southern Ocean for the period 1990–2009, *Biogeosciences*, 10, 4037-4054, doi:10.5194/bg-10-4037-2013, 2013.
- Lourantou, A., and Metzl, N.: Decadal evolution of carbon sink within a strong bloom area in the subantarctic zone, *Geophys. Res. Lett.*, 38, L23608, doi:10.1029/2011GL049614, 2011.
- Majkut, J. D., Carter, B. R., Frölicher, T. L., Dufour, C. O., Rodgers, K. B., Sarmiento, J. L.: An observing system simulation for Southern Ocean carbon dioxide uptake, *Phil. Trans. Roy. Soc. A*, 372, 20130046, doi:10.1098/rsta.2013.0046, 2014a.
- Majkut, J. D., Sarmiento, J. L., and Rodgers, K. B.: A growing oceanic carbon uptake: Results from an inversion study of surface pCO₂ data, *Global Biogeochem. Cy.*, 28, 335–351, doi:10.1002/2013GB004585, 2014b.
- Marrec, P., Cariou, T., Macé, E., Morin, P., Salt, L. A., Vernet, M., Taylor, B., Paxman, K., and Bozec, Y.: Dynamics of air-sea CO₂ fluxes in the northwestern European shelf based on voluntary observing ship and satellite observations, *Biogeosciences*, 12, 5371-5391, doi:10.5194/bg-12-5371-2015, 2015.
- Mathis, J. T., Feely, R. A., Sutton, A., Carlson, C., Chai, F., Chavez, F., Church, M., Cosca, C., Ishii, M., Mordy, C., Murata, A., Resing, J., Stratton, P., Takahashi, T., and Wanninkhof, R.: White paper 6: Tropical Pacific biogeochemistry: Status, implementation and gaps. Tropical Pacific Observing System 2020, available at: http://tpos2020.org/wp-content/uploads/WP06_Tropical-Pacific_biogeochemistry.pdf (last access: 3 January 2015).
- McKinley, G. A., Fay, A. R., Takahashi, T., and Metzl, N.: Convergence of atmospheric and North Atlantic carbon dioxide trends on multidecadal timescales, *Nat. Geosci.*, 4, 606–610, doi:10.1038/NGEO1193, 2011.



- McKinley, G. A., Pilcher, D. J., Fay, A. R., Lindsay, K., Long, M. C., and Lovenduski, N. S.: Timescales for detection of trends in the ocean carbon sink, *Nature*, 530, 469-472. doi:10.1038/nature16958, 2016.
- Merlivat, L., Boutin, J., and Antoine, D.: Roles of biological and physical processes in driving seasonal air-sea CO₂ flux in the Southern Ocean: New insights from CARIOCA pCO₂, *J. Mar. Syst.*, 147, 9-20, doi:10.1016/j.jmarsys.2014.04.015, 2015.
- 5 Monteiro, P. M. S., Schuster, U., Hood, M., Lenton, A., Metzl, N., Olsen, A., Rogers, K., Sabine, C. L., Takahashi, T., Tilbrook, B., Yoder, J., Wanninkhof, R., and Watson, A. J.: A global sea surface carbon observing system: Assessment of changing sea surface CO₂ and air-sea CO₂ fluxes, in: *Proceedings of OceanObs'09: Sustained Ocean Observations and Information for Society*, edited by: Hall, J., Harrison, D. E., and Stammer, D., Vol 2, Venice, Italy, 21-25 September 2009, ESA Publication WPP-306. doi:10.5270/OceanObs09.cwp.64, 2010.
- 10 Nakaoka, S., Telszewski, M., Nojiri, Y., Yasunaka, S., Miyazaki, C., Mukai, H., and Usui, N.: Estimating temporal and spatial variation of ocean surface pCO₂ in the North Pacific using a self organizing map neural network technique, *Biogeosciences*, 10, 6093–6106, doi:10.5194/bg-10-6093-2013, 2013.
- NCEP: 40-Year Reanalysis Project on a 6-hourly, global, 2.5° latitude by 2.5° longitude grid (Kalnay et al., 1996) by the National Centers for Environmental Prediction and the National Center for Atmospheric Research, available at
- 15 <http://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis.surface.html> (last access: 1 May 2013), 2012.
- NCEP: 40-Year Reanalysis Project on a 6-hourly, global, 2.5° latitude by 2.5° longitude grid (Kalnay et al., 1996) by the National Centers for Environmental Prediction and the National Center for Atmospheric Research, available at
- <http://www.esrl.noaa.gov/psd/data/gridded/data.ncep.reanalysis.surface.html> (last access: 29 October 2015), 2014.
- Newton, J.A., Feely, R. A., Jewett, E. B., Williamson, P., and Mathis, J.: Global Ocean Acidification Observing Network:
- 20 Requirements and Governance Plan, 60 pp, available at: http://goa-on.org/docs/GOA-ON_plan_print.pdf (last access: 23 December 2015), 2014.
- Olsen, A., Bakker, D. C. E., and Pfeil, B.: Second version of the Surface Ocean CO₂ Atlas (SOCAT) to be released this summer for climate research, *SOLAS News*, 15, 21, 2013.
- Olsen, A., Metzl, N., Bakker, D. C. E., and O'Brien, K.: SOCAT quality control cookbook - For SOCAT Version 3, available
- 25 at: http://www.socat.info/upload/2015_SOCAT_QC_Cookbook_v3.pdf (last access: 11 December 2015), 2015.
- Omar, A. M., Olsen, A., Johannessen, T., Hoppema, M., Thomas, H., and Borges, A. V.: Spatiotemporal variations of fCO₂ in the North Sea, *Ocean Science*, 6, 77–89, doi:10.5194/os-6-77-2010, 2010.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G.-K., Rodgers, K.
- 30 B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M.-F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681-686, 2005.



- Park, G.-H., Wanninkhof, R., Doney, S. C., Takahashi, T., Lee, K., Feely, R. A., Sabine, C. L., Triñanes, J., and Lima, I. D.: Variability of global net sea-air CO₂ fluxes over the last three decades using empirical relationships, *Tellus B*, 62, 352–368, 2010.
- Pfeil, B., Olsen, A., Bakker, D. C. E., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metzl, N., Sabine, C. L., Akl, J., Alin, S. R., Bates, N., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F. P., Chen, A., Cosca, C., Fassbender, A. J., Feely, R. A., González-Dávila, M., Goyet, C., Hales, B., Hardman-Mountford, N., Heinze, C., Hood, M., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Jones, S. D., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Santana-Casiano, J. M., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Tjiputra, J., Vandemark, D., Veness, T., Wanninkhof, R., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: A uniform, quality controlled Surface Ocean CO₂ Atlas (SOCAT), *Earth Syst. Sci. Data*, 5, 125-143, doi:10.5194/essd-5-125-2013, 2013.
- Pierrot, D., Neill, C., Sullivan, K., Castle, R., Wanninkhof, R., Lüger, H., Johannessen, T., Olsen, A., Feely, R. A., and Cosca, C. E.: Recommendations for autonomous underway pCO₂ measuring systems and data reduction routines, *Deep-Sea Res. Pt. II*, 56 (8-10), 512-522, doi:10.1016/j.dsr2.2008.12.005, 2009.
- Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F. T., Gruber, N., Janssens, I. A., Laruelle, G. G., Lauerwald, R., Luysaert, S., Andersson, A. J., Arndt, S., Arnosti, C., Borges, A. V., Dale, A. W., Gallego-Sala, A., Goddérís, Y., Goossens, N., Hartmann, J., Heinze, C., Ilyina, T., Joos, F., LaRowe, D. E., Leifeld, J., Meysman, F. J. R., Munhoven, G., Raymond, P. A., Spahni, R., Suntharalingam, P., and Thullner, M.: Anthropogenic perturbation of the carbon fluxes from land to ocean. *Nat. Geosci.*, 6, 597-607, doi:10.1038/NGEO1830, 2013.
- Reum, J. C. P., Alin, S. R., Harvey, C. J., Bednaršek, N., Evans, W., Feely, R. A., Hales, B., Lucey, N., Mathis, J. T., McElhany, P., Newton, J., and Sabine, C. L.: Interpretation and design of ocean acidification experiments in upwelling systems in the context of carbonate chemistry co-variation with temperature and oxygen, *ICES J. Mar. Sci.*, 73 (3), 582-595, doi:10.1093/icesjms/fsu231, 2016.
- Rödenbeck, C., Keeling, R. F., Bakker, D. C. E., Metzl, N., Olsen, A., Sabine, C. L., and Heimann, M.: Global surface-ocean pCO₂ and sea-air CO₂ flux variability from an observation-driven ocean mixed-layer scheme. *Ocean Sci.*, 9, 193-216, doi:10.5194/os-9-193-2013, 2013.
- Rödenbeck, C., Bakker, D.C.E., Metzl, N., Olsen, A., Sabine, C.L., Cassar, N., Reum, F., Keeling, R.F., and Heimann, M.: Interannual sea-air CO₂ flux variability from an observation-driven ocean mixed-layer scheme. *Biogeosciences*, 11, 4599-4613, doi:10.5194/bg-11-4599-2014, 2014.
- Rödenbeck, C., Bakker, D. C. E., Gruber, N., Iida, Y., Jacobson, A.R., Jones, S., Landschützer, P., Metzl, N., Nakaoka, S., Olsen, A., Park, G.-H., Peylin, P., Rodgers, K. B., Sasse, T. P., Schuster, U., Shutler, J. D., Valsala, V., Wanninkhof, R., and



- Zeng, J.: Data-based estimates of the ocean carbon sink variability – First results of the Surface Ocean pCO₂ Mapping intercomparison (SOCOM), *Biogeosciences*, 12, 7251-7278, doi:10.5194/bg-12-7251-2015, 2015.
- Sabine, C. L., Hankin, S., Koyuk, H., Bakker, D. C. E., Pfeil, B., Olsen, A., Metzl, N., Kozyr, A., Fassbender, A., Manke, A., Malczyk, J., Akl, J., Alin, S. R., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F. P., Chen, A.,
5 Cosca, C., Feely, R. A., González-Dávila, M., Goyet, C., Hardman-Mountford, N., Heinze, C., Hoppema, M., Hunt, C. W.,
Hydes, D., Ishii, M., Johannessen, T., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A.,
Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S.,
Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F.,
Salisbury, J., Santana-Casiano, J. M., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I.,
10 Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Vandemark, D., Veness, T.,
Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: Surface Ocean CO₂ Atlas (SOCAT) gridded data products,
Earth Syst. Sci. Data, 5, 145-153, doi:10.5194/essd-5-145-2013, 2013.
- Sasse, T. P., McNeil, B. I., and Abramowitz, G.: A new constraint on global air-sea CO₂ fluxes using bottle carbon data,
Geophys. Res. Lett., 40, 1594–1599, doi:10.1002/grl.50342, 2013.
- 15 Schlitzer, R.: Data analysis and visualization with Ocean Data View, *CMOS Bulletin SCMO* 43 (1), 9-13, Canadian
Meteorological and Oceanographic Society / Société canadienne de météorologie et d’océanographie, available at
http://cmos.ca/site/cmos_bulletin (last access: 6 March 2016), 2015.
- Schuster, U., McKinley, G. A., Bates, N., Chevallier, F., Doney, S. C., Fay, A. R., González-Dávila, M., Gruber, N., Jones, S.,
Krijnen, J., Landschützer, P., Lefèvre, N., Manizza, M., Mathis, J., Metzl, N., Olsen, A., Rios, A. F., Rödenbeck, C., Santana-
20 Casiano, J. M., Takahashi, T., Wanninkhof, R., and Watson, A. J.: Atlantic and Arctic sea-air CO₂ fluxes, 1990–2009,
Biogeosciences, 10, 607-627, doi:10.5194/bg-10-607-2013, 2013.
- Secretariat of the Convention on Biological Diversity: An updated synthesis of the impacts of ocean acidification on marine
biodiversity. Hennige, S., Roberts, J. M., and Williamson, P. (eds.), Montreal, Canada, CBD Technical Series, 75, 99 pp,
available at: <https://www.cbd.int/doc/publications/cbd-ts-75-en.pdf> (last access: 23 December 2015), 2014.
- 25 Séférian, R., Ribes, A., and Bopp, L.: Detecting the anthropogenic influences on recent changes in ocean carbon uptake,
Geophys. Res. Lett., 41, 1-10, doi:10.1002/2014GL061223, 2014.
- Signorini, S. R., Mannino, A., Najjar Jr., R. G., Friedrichs, M. A. M., Cai, W.-J., Salisbury, J., Wang, Z. A., Thomas, H., and
Shadwick, E.: Surface ocean pCO₂ seasonality and sea-air CO₂ flux estimates for the North American east coast. *J. Geophys.*
Res.-Oceans, 118, 5439–5460, doi:10.1002/jgrc.20369, 2013.
- 30 Simon, E., and Bertino, L.: Optimized parameters of mathematical formulation with respect to observations. First attempt to
optimize biogeochemical-governing parameters through EnKF, SKD Biofeedback project deliverable D2.3, Bjerknes Center,
Bergen, Norway, 2013.



- SOCAT: Future for the Surface Ocean CO₂ Atlas: Data quality, management and products, Data2Flux Workshop, Unesco, Paris, 12-13 September 2011, SOCAT Report, available at: <http://www.socat.info/meetings.html> (last access: 24 June 2013), 2011.
- SOCAT: Surface Ocean CO₂ Atlas (SOCAT) automation planning meeting, NOAA-PMEL, Seattle, USA, 10-11 May 2012, SOCAT Report, available at: <http://www.socat.info/meetings.html> (last access: 1 May 2013), 2012a.
- 5 SOCAT: Surface Ocean CO₂ Atlas (SOCAT) progress meeting, Epochal Tsukuba, Tsukuba, Japan, 3-5 July 2012, SOCAT Report, available at: <http://www.socat.info/meetings.html> (last access: 1 May 2013), 2012b.
- SOCAT: Data policy for the SOCAT public release, available at: http://www.socat.info/SOCAT_data_policy_public_release_v2.htm (last access: 1 May 2013), 2013a.
- 10 SOCAT: Surface Ocean CO₂ Atlas side event: Release of version 2 and science highlights, SOCAT side event during the 9th International Carbon Dioxide Conference, Beijing International Convention Center, Beijing, China, 4 June 2013, SOCAT Report, available at: http://www.socat.info/upload/2013_SOCAT_Release_version_2_and_Science_Highlights_04June2013.pdf (last access: 29 November 2013), 2013b.
- 15 SOCAT: The Surface Ocean CO₂ Atlas (SOCAT) Community Event. Workshop 10 of the IMBER Open Science Conference, Bergen, Norway, on 23 June 2014, available at: http://www.socat.info/upload/2014_SOCAT_Community_Event_Report_15082014.pdf (last access: 11 December 2014), 2014a.
- SOCAT: Surface Ocean CO₂ Atlas (SOCAT) automation workshop, Seattle, WA, USA, 21-23 October 2014, available at: http://www.socat.info/upload/SOCAT_2014_automation_meeting_report_final.pdf (last access: 11 December 2014), 2014b.
- 20 SOCAT: Fair Data Use Statement for SOCAT, available at: http://www.socat.info/SOCAT_fair_data_use_statement.htm (last access: 1 March 2016), 2016.
- SOCAT and SOCOM: SOCAT (Surface Ocean CO₂ Atlas) and SOCOM (Surface Ocean pCO₂ Mapping Intercomparison) Event, SOLAS (Surface Ocean Lower Atmosphere Study) Open Science Conference, University of Kiel, Kiel, Germany, 7 September 2015, available at http://www.socat.info/upload/2015_SOCAT_and_SOCOM_Event_Report.pdf (last access: 10 December 2015), 2015.
- 25 Sutton, A. J., Feely, R. A., Sabine, C. L., McPhaden, M. J., Takahashi, T., Chavez, F. P., Friederich, G. E., and Mathis, J. T.: Natural variability and anthropogenic change in equatorial Pacific surface ocean pCO₂ and pH, *Global Biogeochem. Cy.*, 28, 131-145, doi:10.1002/2013GB004679, 2014.
- 30 Sutton, A. J., Sabine, C. L., Maenner-Jones, S., Lawrence-Slavas, N., Meinig, C., Feely, R. A., Mathis, J. T., Musielewicz, S., Bott, R., McLain, P. D., Fought, H. J., and Kozyr, A.: A high-frequency atmospheric and seawater pCO₂ data set from 14 open-ocean sites using a moored autonomous system, *Earth Syst. Sci. Data*, 6, 353–366, doi:10.5194/essd-6-353-2014, 2014.



- Swift, J.: A guide to submitting CTD/hydrographic/tracer data and associated documentation to the CLIVAR and Carbon Hydrographic Data Office, version of 22 April 2008, Scripps Institution of Oceanography, University of California, San Diego, 37 pp., 2008.
- Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., and Sutherland, S. C.: Seasonal variation of CO₂ and nutrients in the high-latitude surface oceans: a comparative study, *Global Biogeochem. Cy.*, 7 (4), 843-878, 1993.
- Takahashi, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., Bates, N., Wanninkhof, R. H., Feely, R. A., Sabine, C. L., Olafsson, J., and Nojiri, Y.: Global sea-air CO₂ flux based on climatological surface ocean pCO₂, and seasonal biological and temperature effects, *Deep-Sea Res. Pt. II*, 49, 1601-1622, 2002
- Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C. L., Watson, A. J., Bakker, D. C. E., Schuster, U., Metzl, N., Inoue, H. Y., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen T., Olsen, A., Bellerby, R. G. J., Wong, C. S., Delille B., Bates, N. R., and de Baar, H. J. W.: Climatological mean and decadal change in surface ocean pCO₂, and net sea-air CO₂ flux over the global oceans, *Deep-Sea Res. Pt. II.*, 56, 544-577, doi:10.1016/j.dsr2.2008.12.009, 2009.
- 15 Takahashi, T., Sutherland, S. C., and Kozyr, A. (2015) Global ocean surface water partial pressure of CO₂ database: Measurements performed during 1957-2014 (Version 2014). ORNL/CDIAC-161, NDP-088(V2014). Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee, doi:10.3334/CDIAC/OTG.NDP088(V2014), available at: http://cdiac.ornl.gov/oceans/LDEO_Underway_Database/ (last access: 4 September 2015), 2014.
- 20 Tans, P. and Keeling, R.: Annual mean atmospheric CO₂ values for Mauna Loa from Pieter Tans, NOAA/ESRL (<http://www.esrl.noaa.gov/gmd/ccgg/trends/>) and Ralph Keeling, Scripps Institution of Oceanography (scrippsco2.ucsd.edu/), available at: <http://www.esrl.noaa.gov/gmd/ccgg/trends/data.html> (last access: 14 January 2016), 2016.
- Thomas, H., Bozec, Y., Elkalay, K., and de Baar, H. J. W.: Enhanced open ocean storage of CO₂ from shelf sea pumping, *Science*, 304 (5673), 1005-1008, doi:10.1126/science.1095491, 2004.
- 25 Tjiputra, J. F., Olsen, A., Assmann, K., Pfeil, B., and Heinze, C.: A model study of the seasonal and long-term North Atlantic surface pCO₂ variability, *Biogeosciences*, 9, 907-923, doi:10.5194/bg-9-907-2012, 2012.
- Tjiputra, J. F., Olsen, A., Bopp, L., Lenton, A., Pfeil, B., Roy, T., Segschneider, J., and Totterdell, I.: Long-term surface pCO₂ trends from observations and models, *Tellus B* 66, 23083, doi:10.3402/tellusb.v66.23083, 2014.
- Turi, G., Lachkar, Z., and Gruber, N.: Spatiotemporal variability and drivers of pCO₂ and air-sea CO₂ fluxes in the California Current System: an eddy-resolving modelling study, *Biogeosciences*, 11, 671-690, doi:10.5194/bg-11-671-2014, 2014.
- 30 Turley, C.: The other CO₂ problem: Open Democracy, available at: http://www.acamedia.info/sciences/sciliterature/globalw/reference/carol_turley (last access: 21 January 2016), 2005



- Ulfso, A., Cassar, N., Korhonen, M., Van Heuven, S., Hoppema, M., Kattner, G., and Anderson, L. G.: Late summer net community production in the central Arctic Ocean using multiple approaches, *Global Biogeochem. Cy.*, 28 (10), 1129-1148, doi:10.1002/2014GB004833, 2014.
- Valsala, K. V., and Maksyutov, S.: Simulation and assimilation of global ocean pCO₂ and air-sea CO₂ fluxes using ship observations of surface ocean pCO₂ in a simplified biogeochemical offline model, *Tellus*, 62B, 821–840, 2010.
- 5 Van der Laan, S., Van der Laan-Luijkx, I. T., Rödenbeck, C., Varlagin, A., Shironya, I., Neubert, R. E. M., Ramonet, M., and Meijer, H. A. J.: Atmospheric CO₂, delta(O₂/N₂), APO and oxidative ratios from aircraft flask samples over Fyodorovskoye, Western Russia. *Atmospheric Environment*, 97, 174-181, doi:10.1016/j.atmosenv.2014.08.022, 2014.
- Walker Brown, C., Boutin, J., and Merlivat, L.: New insights into fCO₂ variability in the tropical eastern Pacific Ocean using SMOS SSS, *Biogeosciences*, 12, 7315-7329, doi:10.5194/bg-12-7315-2015, 2015.
- 10 Wang, X., Murtugudde, R., Hackert, E., Wang, J., and Beauchamp, J.: Seasonal to decadal variations of sea surface pCO₂ and sea-air CO₂ flux in the equatorial oceans over 1984–2013: A basin-scale comparison of the Pacific and Atlantic Oceans, *Global Biogeochem. Cy.*, 29(5), 597-609, doi:10.1002/2014GB005031, 2015.
- Wanninkhof, R., Park, G.-H., Takahashi, T., Sweeney, C., Feely, R., Nojiri, Y., Gruber, N., Doney, S. C., McKinley, G. A., Lenton, A., Le Quéré, C., Heinze, C., Schwinger, J., Graven, H., and Khatiwala, S.: Global ocean carbon uptake: magnitude, variability and trends, *Biogeosciences*, 10, 1983-2000, doi:10.5194/bg-10-1983-2013, 2013a.
- 15 Wanninkhof, R., Bakker, D. C. E., Bates, N., Olsen, A., Steinhoff, T., and Sutton, A. J. : Incorporation of alternative sensors in the SOCAT database and adjustments to dataset Quality Control flags, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, available at: <http://cdiac.ornl.gov/oceans/Recommendationnewsensors.pdf> (last access : 23 December 2015), doi:10.3334/CDIAC/OTG.SOCAT_ADQCF, 2013b.
- 20 While, J., Totterdell, I., and Martin, M.: Assimilation of pCO₂ data into a global coupled physical-biogeochemical ocean model, *J. Geophys. Res.*, 117, C03037, doi:10.1029/2010JC006815, 2012.
- Wunsch, C., Schmitt, R. W. Baker, D. J.: Climate change as an intergenerational problem. *Proc. Nat. Acad. Sci. USA*, 110 (12), 4435-4436, doi:10.1073/pnas.1302536110, 2013.
- 25 Xu, S., Chen, L., Chen, H., Li, J., Lin, W., and Qi, D.: Sea-air CO₂ fluxes in the Southern Ocean for the late spring and early summer in 2009, *Rem. Sens. Env.*, 175, 158-166, doi:10.1016/j.rse.2015.12.049, 2016.
- Zeng, J., Nojiri, Y., Landschützer, P., Telszewski, M., and Nakaoka, S.: A global surface ocean fCO₂ climatology based on a feed forward neural network, *J. Atmos. Ocean Technol.*, 31, 1838–1849, 2014.



Table 1: Key differences between SOCAT versions 2 and 3. See text and Table 2 for further detail and an explanation of cross-overs and Standard Operating Procedures (SOP).

	Version 2	Version 3
Description	Bakker et al., 2014	<i>This study.</i>
Fair Data Use Statement	Data Policy on web pages.	<i>Renamed to Fair Data Use Statement. Phrased more strongly and given more prominence on the SOCAT web sites.</i>
Data coverage	1968 to 2011, 10.1 million surface water fCO_2 values, 2660 data sets with a WOCE flag of 2.	<i>1957 to 2014, 14.5 million surface water fCO_2 values from 3646 data sets (3640 with a WOCE flag of 2 and 6 with a flag of 3).</i>
Time stamp	The time stamp includes seconds for all data sets. When equal time stamps occurred, evenly distributed artificial seconds were added to time stamps.	<i>Artificial seconds were added for concurrent entries. A WOCE flag of 4 was given to duplicate times in data sets with less than 50 equal time stamps (Table 7).</i>
Upload Dashboard	Not available.	<i>Single platform for data upload, fCO_2rec calculation and automated data checks.</i>
Data upload	Bulk data upload on quality control system.	<i>All data sets in versions 1, 2 and 3 were uploaded on the Upload Dashboard.</i>
Calculation of fCO_2rec	In Matlab, prior to bulk data upload.	<i>On the Upload Dashboard with Ferret scripts for all data in versions 1, 2 and 3.</i>
Automated data checks	Not available.	<i>Automated checks after calculation of fCO_2rec for all new and updated data sets. WOCE flags of 4 were assigned in specific cases (Table 7).</i>
Quality Control Editor	As in version 1.	<i>After automated checks. Upgraded search options and graphical interface. Data set QC flag needs to match QC criteria (tick boxes).</i>
Data set QC flags in data products	Flags of A-D.	<i>Flags of A-E. Revised data set QC criteria (Table 2) applied to all new and updated data sets.</i>
Flag A	Needs a cross-over (an acceptable comparison with other data).	<i>Needs a high-quality cross-over.</i>
Flags A, B	Accuracy equilibrator pressure ≤ 0.5 hPa. Six other SOP criteria apply.	<i>Accuracy equilibrator pressure ≤ 2 hPa. Six other SOP criteria apply.</i>
Flag C	Did not follow approved methods or SOP criteria	<i>Did or did not follow approved methods or SOP criteria.</i>
Flags C, D	Accuracy fCO_2rec not specified.	<i>Accuracy $fCO_2rec \leq 5 \mu atm$.</i>
Flag E	Not available.	<i>Accuracy $fCO_2rec \leq 10 \mu atm$, mainly for alternative sensors and platforms with in situ calibration and full documentation.</i>
WOCE flags for fCO_2rec	Flag of 2 (good) as a default. Manual entry of flags of 3 (questionable) and 4 (bad).	<i>Flag of 2 as a default. Flags of 4 given during automated data checks (Table 7). Quality control comment added during manual entry of flags of 3 and 4.</i>
Parameter	NCEP (2012) atmospheric pressure, atmospheric CO_2 mole fraction from GLOBALVIEW-CO2 (2012).	<i>NCEP (2014) atmospheric pressure, atmospheric CO_2 mole fraction from GLOBALVIEW-CO2 (2014).</i>
Synthesis products	Data sets with flags of A-D and fCO_2rec with a WOCE flag of 2 in synthesis and gridded files and as default elsewhere.	<i>Data sets with flags of A-E made public (Table 8). Data sets with flags of A-D and fCO_2rec with a WOCE flag of 2 in synthesis and gridded files. Data sets with a flag of E and fCO_2rec with a flag of 2 in a separate synthesis file. Contents of files downloadable from the Data Set Viewer have been streamlined (Table 9).</i>
Gridded products	Missing grid cells in cruise weighted gridded products (versions 1 and 2). A gridded product of means per climatological month is available.	<i>Correction of data set-weighted gridded products (version 3). No gridded product of means per climatological month.</i>
Data Set and Gridded Data Viewers	On different software platforms.	<i>On a single software platform with a powerful graphical interface, following the move of the Data Set Viewer.</i>
Terminology	Terms in version 1: Cruises, ships, Cruise Data Viewer, Table of Cruises, cruise weighted means.	<i>New terms to accommodate sensors and platforms: Data sets, platforms, Data Set Viewer, Table of Datasets, data set-weighted means.</i>



Table 2: Data set quality control (QC) flags in version 3 (Wanninkhof et al., 2013b; Olsen et al., 2015). All criteria need to be met for assigning a flag of A to E. Data sets with flags of A to E have been made public. Data sets with a flag of A to D are included in the global synthesis and gridded products (Table 8). Changes relative to versions 1 and 2 are in bold.

Flag (ID)	Criteria for version 3
A (11)	(1) Accuracy of calculated fCO_2rec (at SST) is better than $2 \mu atm$. (2) A high-quality cross-over ^{1,2} with another data set is available. (3) Followed approved methods/SOP ³ criteria. (4) Metadata documentation complete. (5) Data set QC was deemed acceptable.
B (12)	(1) Accuracy of calculated fCO_2rec (at SST) is better than $2 \mu atm$. (2) Followed approved methods/SOP criteria. (3) Metadata documentation complete. (4) Data set QC was deemed acceptable.
C (13)	(1) Accuracy of calculated fCO_2rec (at SST) is better than $5 \mu atm$. (2) Did or did not follow approved methods/SOP criteria. (3) Metadata documentation complete. (4) Data set QC was deemed acceptable.
D (14)	(1) Accuracy of calculated fCO_2rec (at SST) is better than $5 \mu atm$. (2) Did or did not follow approved methods/SOP criteria. (3) Metadata documentation incomplete. (4) Data set QC was deemed acceptable.
E (17)	Primarily for alternative sensors (1) Accuracy of calculated fCO_2rec (at SST) is better than $10 \mu atm$. (2) Did not follow approved methods/SOP criteria. (3) Metadata documentation complete. (4) Data set QC was deemed acceptable.
S (15) (Suspend)	(1) More information is needed for data set before flag can be assigned. (2) Data set QC revealed non-acceptable data. (3) Data are being updated (part or the entire data set).
X (15) (Exclude)	The data set duplicates another data set in SOCAT.
N (No flag)	No data set flag has yet been given to this data set.
U (Update)	The data set has been updated. No data set flag has yet been given to the revised data.

5 ¹A cross-over between two data sets is defined as an equivalent distance of less than 80 (Pfeil et al., 2013). This criterion combines distance and time as $([\Delta x^2 + (\Delta t * 30)^2]^{0.5}) \leq 80$ with distance x in kilometres and time t in hours. One day of separation in time is equivalent (heuristically) to 30 km of separation in space.

²A high-quality cross-over is defined as a cross-over between two data sets with a maximum cross-over equivalent distance of 80 km, a maximum difference in sea surface temperature of $0.3^\circ C$ and a maximum fCO_2rec difference of $5 \mu atm$. Inconclusive cross-overs with the temperature or fCO_2rec difference between the data sets exceeding $0.3^\circ C$ or $5 \mu atm$, respectively, do not receive a flag of A. High-quality cross-overs are rare in coastal waters, near sea ice and in regions of freshwater influence, as a result of high spatial variation, not for lack of measurement quality (Sect 4.4).

³Seven approved methods or SOP (Standard Operating Procedures) criteria need to be fulfilled for a data set quality control flag of A and B (Sect. 4.4) (after Pfeil et al., 2013). In version 3, the accuracy requirement for equilibrator pressure has been relaxed to $2.0 hPa$ from $0.5 hPa$ in earlier SOCAT versions. The six other criteria are the same in SOCAT versions 1, 2 and 3.



Table 3: Activities and participants in SOCAT version 3 and the automation (after Bakker et al., 2014). Regional group leads are in Table 4.

Activity	Participants
Global group	Bakker (chair), Currie, Kozyr, Metzl, O'Brien, Olsen, Pfeil, Pierrot, Telszewski
Data retrieval, upload, fCO ₂ calculation	<i>Landa</i> , Pfeil, Olsen, Smith
Live Access Server for data upload, quality control and Data Viewers	Smith, O'Brien, Manke, Hankin, <i>Schweitzer</i>
<i>Inclusion of sensors</i>	<i>Wanninkhof, Steinhoff, Bakker, Bates, Boutin, Olsen, Sutton</i>
<i>Automation (version 3)</i>	<i>O'Brien, Smith, S. D. Jones, Landa, Manke, Olsen, Pfeil, Schweitzer, Bakker</i>
<i>Automation (metadata, version 4)</i>	<i>As automation for version 3, plus: Shrestha, Ranjeet</i>
Quality control	Alin, Bakker, Barbero, <i>Bonou</i> , Castle, Cosca, <i>Currie</i> , Evans, <i>Featherstone</i> , <i>Greenwood</i> , Harasawa, <i>Hauck</i> , <i>Humphreys</i> , Hunt, <i>Ibáñez</i> , Lefèvre, Metzl, Nakaoka, Paterson, Schuster, Skjelvan, Steinhoff, Sullivan, Sutton, Tilbrook, Wada
Data products, archiving	Pfeil, Smith, Kozyr, Manke, O'Brien, Schlitzer, Sieger
Matlab code for reading products	Pierrot, Landschützer
Website	Pfeil, Bakker, <i>Landa</i> , Metzl
Meetings	Bakker, Cosca, O'Brien, Steinhoff, Telszewski


Table 4: Regions with their leads in version 3 (after Bakker et al., 2014). The regions are the same as in version 2.

Region	Definition	Lead(s)
Coastal and marginal seas	<400 km from land; 70°N to 30°S for 100°W to 43°E; 66°N to 30°S elsewhere	Alin
Arctic Ocean	North of 70°N for 100°W to 43°E; north of 66°N elsewhere, incl. coastal waters	Mathis
North Atlantic	70°N to 30°N	Schuster
North Pacific	66°N to 30°N	Nojiri
Tropical Atlantic	30°N to 30°S	Lefèvre
Tropical Pacific	30°N to 30°S	Cosca
Indian Ocean	North of 30°S	Sarma
Southern Ocean	South of 30°S, incl. coastal waters	Tilbrook, Metzl



Table 5: Meetings for SOCAT version 3 and the ongoing SOCAT automation. The meeting reports are available on the SOCAT website (<http://www.socat.info/meetings.html>).

Timing	Meeting	Location	Reference
05/2012	Automation planning meeting	NOAA-PMEL, Seattle, USA	SOCAT, 2012a
07/2012	Progress meeting	Epochal Centre, Tsukuba, Japan	SOCAT, 2012b
06/2013	SOCAT side event, Release of version 2	9 th International Carbon Dioxide Conference, Beijing, China	SOCAT, 2013b
06/2014	<i>Community Event</i>	<i>IMBER Open Science Conference, Bergen, Norway</i>	<i>SOCAT, 2014a</i>
10/2014	<i>Automation meeting</i>	<i>NOAA-PMEL, Seattle, USA</i>	<i>SOCAT, 2014b</i>
09/2015	<i>SOCAT and SOCOM Event, Release of version 3, Launch of automation system, SOCOM science.</i>	<i>SOLAS Open Science Conference, Kiel, Germany</i>	<i>SOCAT and SOCOM, 2015</i>



5

Table 6: Algorithms and surface water CO₂ parameters used in the calculation of recommended fCO₂ (fCO₂rec) at sea surface temperature in version 3 (after Pfeil et al., 2013). Algorithm 1 was the preferred method, followed by algorithm 2 and so forth. The algorithm used for each data set is stated in the output files (Table 9). In case of incomplete reporting, NCEP (National Centres for Environmental Prediction) atmospheric pressure (Kalnay et al., 1996; NCEP, 2014) and WOA (World Ocean Atlas) salinity (Antonov et al., 2006) were applied.

Algo rith m	CO ₂ parameter	Unit	Data set percentage (%)	Extra variable
1	xCO ₂ water_equi_dry	μmol mol ⁻¹	59.1	-
2	xCO ₂ water_SST_dry	μmol mol ⁻¹	12.5	-
3	pCO ₂ water_equi_wet	μatm	7.2	-
4	pCO ₂ water_SST_wet	μatm	3.0	-
5	fCO ₂ water_equi	μatm	0.4	-
6	fCO ₂ water_SST_wet	μatm	12.2	-
7	pCO ₂ water_equi_wet ¹	μatm	0.4	NCEP pressure
8	pCO ₂ water_SST_wet ¹	μatm	6.1	NCEP pressure
9	xCO ₂ water_equi_dry ²	μmol mol ⁻¹	2.9	WOA salinity
10	xCO ₂ water_SST_dry ²	μmol mol ⁻¹	3.1	WOA salinity
11	xCO ₂ water_equi_dry ¹	μmol mol ⁻¹	0.3	NCEP pressure
12	xCO ₂ water_SST_dry ¹	μmol mol ⁻¹	0.5	NCEP pressure
13	xCO ₂ water_equi_dry ^{1,2}	μmol mol ⁻¹	0.05	NCEP pressure, WOA salinity
14	xCO ₂ water_SST_dry ^{1,2}	μmol mol ⁻¹	0.2	NCEP pressure, WOA salinity

¹ Atmospheric pressure was not reported in the original data file.

² Salinity was not reported in the original data file.



Table 7: Criteria for the automated data checks and the action taken in version 3. In case of duplicate time stamps, artificial seconds were generated. If there were less than 50 duplicate times in the data set, a WOCE flag of 4 was given. For other parameters, a flag of 4 was automatically assigned to the $f\text{CO}_2\text{rec}$ value, if their values were outside a specified range. Criteria not directly affecting $f\text{CO}_2\text{rec}$ values will be revised for version 4 (Sect. 4.3).

Parameter	Unit	Criteria	Action
<i>Time</i>	-	<i>Duplicate times</i>	<i>Artificial seconds added; flag of 4, if <50 duplicate times in data set.</i>
<i>Sampling depth, water</i>	<i>m</i>	<i><-20 or > 20</i>	<i>flag of 4</i>
<i>Salinity</i>	-	<i><0 or >50</i>	<i>flag of 4</i>
<i>Sea surface temperature</i>	<i>°C</i>	<i><-8 or >50</i>	<i>flag of 4</i>
<i>Equilibrator temperature</i>	<i>°C</i>	<i><-10 or >45</i>	<i>flag of 4</i>
<i>Atmospheric pressure</i>	<i>mbar</i>	<i><800 or >1200</i>	<i>flag of 4</i>
<i>Equilibrator pressure</i>	<i>mbar</i>	<i><800 or >1200</i>	<i>flag of 4</i>
<i>$x\text{CO}_2$, $p\text{CO}_2$, $f\text{CO}_2$ water</i>	<i>$\mu\text{mol mol}^{-1}$ or μatm</i>	<i><0 or >10,000</i>	<i>flag of 4</i>
<i>$x\text{CO}_2$, $p\text{CO}_2$, $f\text{CO}_2$ air</i>	<i>$\mu\text{mol mol}^{-1}$ or μatm</i>	<i><0 or >10,000</i>	<i>flag of 4</i>
<i>$\Delta x\text{CO}_2$, $\Delta p\text{CO}_2$, $\Delta f\text{CO}_2$</i>	<i>$\mu\text{mol mol}^{-1}$ or μatm</i>	<i><-10,000 or >10,000</i>	<i>flag of 4</i>
<i>$x\text{H}_2\text{O}$ equilibration</i>	<i>mmol mol^{-1}</i>	<i><0 or >200</i>	<i>flag of 4</i>
<i>WOCE flag, from PI</i>	-	<i><1 or >9</i>	<i>flag of 4</i>
<i>Air temperature</i>	<i>°C</i>	<i><-35 or >60</i>	<i>flag of 4</i>
<i>Relative humidity</i>	<i>%</i>	<i><0 or >100</i>	<i>flag of 4</i>
<i>Specific humidity</i>	-	<i><0 or >40</i>	<i>flag of 4</i>
<i>Wind direction</i>	<i>°</i>	<i><0 or >360</i>	<i>flag of 4</i>
<i>Wind speed</i>	<i>m s^{-1}</i>	<i><0 or >50</i>	<i>flag of 4</i>
<i>Ship direction</i>	<i>°</i>	<i><0 or >360</i>	<i>flag of 4</i>
<i>Ship speed, from PI</i>	<i>km h^{-1}</i>	<i><0 or >100</i>	<i>flag of 4</i>
<i>Ship speed, calculated</i>	<i>km h^{-1}</i>	<i>>720</i>	<i>flag of 4 for following point</i>



Table 8: Key characteristics of the SOCAT data products in version 3 (Sect. 5) (after Bakker et al., 2014). Data products differ in whether they include data sets with flags of A to D or A to E and fCO₂rec values with a WOCE flag of 2 only or 2 to 4. Two data products provide access to metadata. Quality control comments are available via the Table of Datasets in the Data Set Viewer. The SOCAT website (<http://www.socat.info>) and the web links in the footnotes provide access to the data products.

5

Data product	Characteristics	Data set QC flag	WOCE flag	Meta-data	QC entries	Format and access
Individual data set files	All original CO ₂ values and (re-)calculated fCO ₂ values for data sets with flags of A-E.	A-E	2-4	Yes	No	Text files ¹
Synthesis data set (i)	Global and regional synthesis files.	A-D only	2 only	No	No	Text files ²
<i>Synthesis data set (ii)</i>	<i>Global synthesis file.</i>	<i>E only</i>	<i>2 only</i>	<i>No</i>	<i>No</i>	<i>Text file²</i>
Synthesis data set (iii)	Global synthesis file.	A-D only	2 only	Optional	No	ODV ³
Subset of synthesis data set (j)	Interactive. Data sets with flags of A-E and fCO ₂ values with flags of 2-4 can be selected.	A-E <i>default</i>	2 default; 2-4 if selected	No	No	Text and NetCDF files ⁴
Subset of synthesis data set (jj)	All original CO ₂ values and calculated fCO ₂ values for data sets with flags of A-E.	A-E <i>default</i>	2 default; 2-4 if selected	Yes	Yes	Text and NetCDF files ⁵
Gridded data set	Gridded unweighted and data set-weighted means of fCO ₂ values on a 1° x 1° grid without any interpolation. Means are per decade, per year and per month. A monthly 0.25° x 0.25° gridded data set exists for coastal regions.	A-D only	2 only	No	No	NetCDF files ^{6,7} , ODV ³

¹PANGAEA®: <https://doi.org/10.1594/PANGAEA.849770>.

²CDIAC: <http://cdiac.ornl.gov/ftp/oceans/SOCATv3/>.

³Ocean Data View: https://odv.awi.de/en/data/ocean/socat_fCO2_data/.

10 ⁴Data Set Viewer: http://ferret.pmel.noaa.gov/SOCAT_Data_Viewer/, Select Data Set, Select Cruise data, Select SOCAT v3 data collection.

⁵Table of Datasets via the Data Set Viewer⁴: Select Table of Datasets.

⁶CDIAC: http://cdiac.ornl.gov/ftp/oceans/SOCATv3/SOCATv3_Gridded_Dat/.

⁷Gridded Data Viewer: http://ferret.pmel.noaa.gov/SOCAT_Data_Viewer/, Select Data Set, Select Current Version Gridded
 15 (v3).



Table 9: Content of the individual data set files (IF) and the synthesis files in SOCAT version 3 (after Bakker et al., 2014). The global synthesis product is available as zip text files (ZIP) at CDIAC and in Ocean Data View (ODV) format (Table 8). Subsets of the global synthesis data set can be created via the Data Set Viewer (DSV), both in the main menu and via the Table of Datasets. The first column lists column headers for the parameters in the files.

Column header	IF	ZIP	DSV	ODV	Unit	Description
Expocode/ Cruise	-	√	√	√	-	12-character expocode
Version	-	√	-	√	-	Most recent SOCAT version in which data set was added (N) or updated (U).
SOCAT_DOI	-	√	√	√	-	Digital object identifier for the individual data set and metadata
QC_Flag	-	√	√	√	-	Data set quality control flags A, B, C, D and E
Date/Time / Datetime	√	-	√	√	-	yyyy-mm-dd / hh:mm:ss (ISO8859 and other formats)
yr / Year	-	√	√	-	Year	Year (UTC)*
mon	-	√	√	-	Month	Month (UTC)*
day	-	√	√	-	Day	Day (UTC)*
hh / Hour	-	√	√	-	Hour	Hour (UTC)*
mm / Minute	-	√	√	-	Minute	Minute (UTC)*
ss / Second	-	√	√	-	Seconds	Seconds (may include decimals)*
Day of Year	-	-	-	√	Day of Year	Day of Year (UTC) with 1 January 00:00 as 1.0.
Longitude	√	√	√	√	°E, °W	Longitude (0 to 360 / -180 to 180)*
Latitude	√	√	√	√	°N, °S	Latitude (-90 to 90)*
Depth water / Depth	√	√	√	√	m	Water sampling depth* ¹
Sal / Salinity	√	√	√	√	-	Salinity on Practical Salinity Scale*
Temp / SST	√	√	√	√	°C	Sea surface temperature*
Tequ / Temperature_Equi	√	√	√	√	°C	Equilibrator chamber temperature*
PPPP / Pressure_Atm	√	√	√	√	hPa	Atmospheric pressure*
Pequ / Pressure_Equi	√	√	√	√	hPa	Equilibrator chamber pressure*
Sal interp / WOA_SSS	√	√	√	√	-	Salinity from WOA ²
PPPP interp / NCEP_SLP	√	√	√	√	hPa	NCEP Atmospheric pressure ³
Bathy_depth / ETOPO2_depth	√	√	√	√	m	ETOPO2 Bathymetry ⁴
Distance / dist-to-land	√	√	√	√	km	Distance to major land mass
xCO2air_interp / GVCO2	√	√	√	√	μmol/mol	Atmospheric xCO ₂ from GLOBALVIEW-CO2 (2014) ⁵
xCO2water_equ_dry	√	-	-	-	μmol/mol	xCO ₂ (water) at equilibrator temperature (dry air)*
fCO2water_SST_wet	√	-	-	-	μatm	fCO ₂ (water) at sea surface temperature (air at 100% humidity)*
pCO2water_SST_wet	√	-	-	-	μatm	pCO ₂ (water) at sea surface temperature (air at 100% humidity)*
xCO2water_SST_dry	√	-	-	-	μmol/mol	xCO ₂ (water) at sea surface temperature (dry air)*
fCO2water_equ_wet	√	-	-	-	μatm	fCO ₂ (water) at equilibrator temperature (air at 100% humidity)*
pCO2water_equ_wet	√	-	-	-	μatm	pCO ₂ (water) at equilibrator temperature (air at 100% humidity)*
fCO2water_SST_wet / fCO2rec	√	√	√	√	μatm	Recommended fCO ₂ calculated following the SOCAT protocol
Algorithm / fCO2_source	√	√	√	√	-	Algorithm for calculating fCO ₂ rec (0: not generated; algorithm 1-14, Table 6)
Flag / WOCE_CO2_Water	√	√	√	-	-	WOCE flag for fCO ₂ rec (2: good, 3: questionable, 4: bad) ⁶
fCO2 in wet air	-	-	-	√	μatm	fCO ₂ (air) calculated for sea surface temperature and 100% humidity from GVCO2
Ocean – Air fCO2 Difference	-	-	-	√	μatm	fCO ₂ difference between water and air
Vessel	-	-	-	√	-	Name of vessel or platform



√ Available.

(√) Available upon selection of parameter.

* If reported by the data originator.

¹If the intake depth has not been reported by the data originator, an intake depth of 5 m has been assumed.

5 ²Sea surface salinity on the Practical Salinity Scale interpolated from the World Ocean Atlas (WOA) 2005 (Antonov et al., 2006), available at <http://www.nodc.noaa.gov/OC5/WOA05/woa05nc.html>, using the data set s0112an1.nc from the "monthly" link at <http://data.nodc.noaa.gov/opendap/woa/WOA05nc/> (last access: 1 September 2015). This data set is identical to that SOCAT version 2.

10 ³Atmospheric pressure interpolated from the NCEP/NCAR (National Centers for Environmental Prediction/ National Center for Atmospheric Research) 40-Year Reanalysis Project on a 6-hourly, global, 2.5° latitude by 2.5° longitude grid (Kalnay et al., 1996; NCEP, 2014). This is an update relative to the 2012 data set (NCEP, 2012) used in SOCAT version 2.

⁴Bathymetry interpolated from ETOPO2 (2006) 2-minute Gridded Global Relief Data. This data set is identical to that in SOCAT version 2.

15 ⁵GLOBALVIEW-CO2 (2014), downloading the "surface" reference type gives the sine function of latitude versus time for the Reference Marine Boundary Layer. This is an update relative to the 2012 version used in SOCAT version 2.

⁶Individual data set files contain all fCO₂rec data. Synthesis files at CDIAC and via ODV contain data sets with a flag of A-D and fCO₂rec values with a WOCE flag of 2 (Table 6).




Table 10: Gridded products and parameters reported for each grid cell in SOCAT version 3 (after Sabine et al., 2013). Version 3 does not have a monthly climatology.


Parameter	Unit	Decadal Mean	Annual Mean	Monthly Mean	Monthly 1/4° x 1/4° Coastal
Number of data sets	-	√	√	√	√
Number of observations	-	√	√	√	√
fCO ₂ unweighted mean	µatm	√	√	√	√
fCO ₂ data set-weighted mean	µatm	√	√	√	√
fCO ₂ max	µatm	√	√	√	√
fCO ₂ min	µatm	√	√	√	√
fCO ₂ stdev unweighted	µatm	-	-	√	√
fCO ₂ stdev weighted	µatm	-	-	√	√
Latitudinal average offset from cell centre	°N	-	-	√	√
Longitudinal average offset from cell centre	°E	-	-	√	√




Figures




5 *Figure 1: Global distribution of a) all and b) newly added surface water $f\text{CO}_2$ values (μatm) and c) the timing of the newly added data sets in SOCAT version 3 with data set flags of A to E. Version 3 has data sets from 1957 to 2014.*




5 *Figure 2: a) Number of surface water fCO_2 values per year and b) the base 10 logarithm of this number per year for 1957 to 2014 in SOCAT versions 1, 2 and 3 (after Bakker et al., 2014).*




5 *Figure 3: Decadal distribution of surface water $f\text{CO}_2$ (μatm) for the global oceans and coastal seas in SOCAT version 3: a) 1957 through 1969, b) 1970s, c) 1980s, d) 1990s, e) 2000s, f) 2010 through 2014. Similar figures are available for versions 1 and 2 (Pfeil et al., 2013; Brévière et al., 2015).*



5 *Figure 4: Seasonal distribution of surface water $f\text{CO}_2$ (μatm) for the months a) January through March, b) April through June, c) July through September and d) October through December in the years 2000 through 2009 in SOCAT version 3 for data sets with flags of A to E (after Bakker et al., 2014).*




5 *Figure 5: Bar plots of the number of decadal mean $f\text{CO}_2$ values per $4 \mu\text{atm}$ range for data set-weighted gridded $f\text{CO}_2$ values in version 3. Red bars indicate the mean atmospheric value ($\mu\text{mol mol}^{-1}$) at Mauna Loa, Hawaii, for each decade (Tans and Keeling, 2016). Note the changing scale on the y-axis. Similar figures have been made for versions 1 and 2 (Olsen et al., 2013; Sabine et al., 2013).*



5

Figure 6: a) Number of unique months and b) total number of months with $f\text{CO}_2$ values per $1^\circ \times 1^\circ$ grid cell for 1970 through 2014 in SOCAT version 3. Similar figures are available for versions 1 and 2 (Sabine et al., 2013; Bakker et al., 2014). The higher resolution of $0.25^\circ \times 0.25^\circ$, available for coastal seas (Sect 5.5), is not shown.



5

Figure 7: a) Number of publications citing or naming SOCAT per year by type of publication and b) scientific applications of SOCAT in peer-reviewed, scientific articles. The number of publications in 2016 only includes publications before 22 April 2016. Types of publications are: peer-reviewed, scientific articles, PhD and MSc theses, high-impact reports, book chapters and all other publications. Scientific applications in peer-reviewed, scientific articles are grouped as reference (only) to the SOCAT data synthesis, use of figures or tools based on SOCAT, use of surface ocean $f\text{CO}_2$ values for various environmental studies, modelling, trend analysis in ocean acidification studies, $f\text{CO}_2$ process studies and carbon budgeting of coastal seas, open ocean and land systems. These scientific applications are discussed in Sections 7.2 and 7.3. A list of publications citing or naming SOCAT is available on the SOCAT website (www.socat.info/publications.html).

10